

RISK FACTOR LBW

by M Zen Rahfiludin

Submission date: 03-May-2019 02:58AM (UTC+0700)

Submission ID: 1123760075

File name: al_Kesmas_FKM_UI_Nop_2018_dengan_pak_Yudhi_1719-8194-1-PB_1.pdf (349.97K)

Word count: 4408

Character count: 22949

Risk Factors Associated with Low Birth Weight

Faktor-faktor Risiko Bayi dengan Berat Badan Lahir Rendah

Mohammad Zen Rahfiludin*, Yudhy Dharmawan**

*Public Health Nutrition Department, Faculty of Public Health Diponegoro University, Semarang, Indonesia,
**Biostatistics & Population Studies Department, Faculty of Public Health, Diponegoro University, Semarang, Indonesia

Abstract

Neonatal deaths are associated with preterm birth complications. The aim of this study was to analyze risk factors associated with LBW. This was a cross-sectional study held in Bulu Primary Health Care, Temanggung, Central Java, Indonesia. The sample size required for this study was 69 based on the Slovin formula. Data were collected using questionnaires and semi-quantitative Food Frequency Questionnaire forms. Data on infant birth weight was taken from midwives' delivery cohort records. Mid upper arm circumference (MUAC), hemoglobin level, blood pressure, maternal age, parity, nutritional intake, and serum transferrin receptor data were taken from the infant's mother using a MUAC tape, automatic blood pressure monitor and blood laboratory analysis by Prodia. Data analysis procedures were carried out with quantitative methods. Descriptive statistics were analyzed as means and standard deviations. Inferential statistics used the chi-square test for bivariate analysis and binary logistic regression for multivariate analysis. The results of this study showed that mean infant birth weight was 2917.68 ± 374.673 kg. Inferential analysis showed that MUAC and pregnancy at a risky age were significant risk factors associated with LBW, while serum transferrin receptor levels, anemia, parity, energy and protein consumption levels, and systolic and diastolic blood pressure were nonsignificant risk factors. The probability of LBW in pregnant women with LILA under 23.5 cm and pregnancy at a risky age was 68.9%.

Keywords: Low birth weight, risk factors, Central Java

Abstrak

Kematian neonatal terkait dengan komplikasi kelahiran prematur. Penelitian ini bertujuan menganalisis faktor risiko bayi dengan berat badan lahir rendah (BBLR). Penelitian ini merupakan penelitian potong lintang dan dilakukan di Puskesmas Bulu, Temanggung, Jawa Tengah, Indonesia. Sampel penelitian berjumlah 69 sampel berdasarkan rumus perhitungan sampel Slovin. Data dikumpulkan dengan menggunakan kuesioner dan formulir semi kuantitatif *Food Frequency Questionnaire*. Data berat lahir bayi diambil dari rekaman kohort persalinan bidan. Lingkar lengan atas (LILA), kadar hemoglobin, tekanan darah, usia ibu, paritas, asupan gizi dan data reseptor transferrin serum diambil dari ibu bayi menggunakan pita LILA, Monitor Tekanan Darah Otomatis dan analisis laboratorium darah oleh Prodia. Prosedur analisis data dilakukan dengan metode kuantitatif. Statistik deskriptif dianalisis dengan mean dan standar deviasi. Statistik inferensial menggunakan uji kai kuadrat untuk analisis bivariat dan regresi logistik biner untuk analisis multivariat. Hasil penelitian menunjukkan bahwa berat lahir bayi rata-rata $2917,68 \pm 374,673$ kg. Analisis inferensial menunjukkan LILA dan usia kehamilan berisiko secara signifikan sebagai faktor risiko yang menyebabkan BBLR, sedangkan reseptor serum transferin, anemia, paritas, tingkat konsumsi energi dan protein, tekanan darah sistolik dan diastolik bukanlah faktor risiko yang signifikan. Kemungkinan ibu hamil BBLR dengan LILA di bawah 23,5 cm dan kehamilan pada usia berisiko sebesar 68,9%.

Kata kunci: Berat bayi lahir rendah, faktor risiko, Jawa Tengah

How to Cite: Rahfiludin MZ, Dharmawan Y. Risk factors associated with low birth weight. *Kesmas: National Public Health Journal*. 2018; 13(2): 75-80 (doi:10.21109/kesmas.v13i2.1719)

Correspondence: Mohammad Zen Rahfiludin, Faculty of Public Health Diponegoro University, Prof. Sudharto Street, Semarang, Indonesia, Phone: +6224-7460044, E-mail: yudrahfiludin@yahoo.com

Received: September 11th 2017

Revised: November 29th 2017

Accepted: January 04th 2018

Introduction

Child mortality is a core indicator of child health and well-being. The proposed Sustainable Development Goals target for child mortality represents a renewed commitment to the world's children under 5 years of all age, with all countries aiming to reduce neonatal mortality to 12 deaths per 1000 live births or lower and under-5 mortality to 25 deaths per 1000 live births or lower. The worldwide neonatal mortality rate in 2015 was 19 deaths per 1000 live births.¹ This number shows that the worldwide neonatal mortality rate is still high. Among the top 10 countries that contribute 67% of neonatal deaths in the world, Indonesia contributes 2%.²

Neonatal deaths are caused by intrapartum complications and severe infections, and the leading cause of death in all regions of the world is preterm birth complications.³ Preterm birth is also related to low birth weight (LBW). LBW is defined as a weight at birth of less than 2500 gram. Not only is LBW a major predictor of prenatal mortality and morbidity, but recent studies have found that low birth weight also increases the risk of non-communicable diseases, such as diabetes and cardiovascular disease, later in life.⁴ There are multiple causes of LBW, including low income, maternal age under 20 years or over 35 years, heavy physical work, low maternal education, maternal complications (placenta previa, pregnancy-induced hypertension, premature rupture of membranes), anemia, malaria, inadequate antenatal care, and maternal nutritional factors.⁵⁻⁶

National Basic Health Research in 2015 showed that LBW occurs in Indonesia at a rate of 10.2%. In Central Java, the rate of LBW was 9.7%.⁷ In Temanggung, specifically in the working area of Bulu Primary Health Care, the incidence of LBW in 2013, 2014, and 2015 was 6.94%, 7.09%, and 7.88%, respectively.⁸

The aim of this study was to analyze risk factors of LBW, including mid upper arm circumference (MUAC), hemoglobin levels, blood pressure, maternal age, parity, nutritional intake, and serum transferrin receptor levels. There have been many studies on risk factors associated with LBW, but only a few studies have examined the relation between the serum transferrin receptor and LBW. The transferrin receptor is the best indicator of iron deficiency in pregnant woman. The serum level of this receptor can be measured easily by conventional techniques and presents a large distinction between iron deficiency anemia and chronic anemia disease.⁹

Method

This was a cross-sectional study held in Bulu Primary Health Care, Temanggung, Central Java, Indonesia. The study population consisted of 114 pregnant women in the second and the third trimesters. The required sample size for this study was 69 samples based on the Slovin

formula.¹⁰ A random sampling technique was applied. MUAC, maternal age, serum transferrin receptor, hemoglobin, parity, and systolic and diastolic blood pressure data were collected using a questionnaire, while energy and protein consumption data were collected using semi-quantitative food frequency questionnaire forms. MUAC was measured using a MUAC Tape accurate to 1 mm. Blood samples were taken for measurement of hemoglobin and serum transferrin receptor levels according to standard protocols in the Prodia Laboratory. Hemoglobin was measured using cyanmethemoglobin, and serum transferrin receptor was measured using the Quantikine IV D, human sTfR Immunoassay, R&D systems, Minneapolis, MN, USA.¹¹⁻¹² Anemia is defined as a hemoglobin level of less than 11 gram/dl or a transferrin receptor level of more than 21 nmol/l. Data on the infant's birth weight was taken from the midwives in Bulu Primary Health Care, by copying the infant's weight recorded by the midwives in a baby cohort. The data were collected a week after delivery. Data analysis procedures were carried out with quantitative methods. Descriptive statistics were analyzed by mean and standard deviation. Categorical data were analyzed by cross tabulation. Inferential statistics used the chi-square test for bivariate analysis and binary logistic regression for multivariate analysis. A significant correlation between independent and dependent variables was shown at p value = 0.05. Ethical clearance was obtained from the Commission of Ethics of Medical and Public Health Research, Faculty of Public Health, Diponegoro University (approval no. 252/EC/FKM/2016).

Results

The mean (SD) birth weight was 2917.68 (374.673) kg, and the mean age of the pregnant women was 26.71 (5.806) years; furthermore, the mean and standard deviation of each variable are presented in Table 1. The results of this study, shown in Table 2, indicated that MUAC and pregnancy at a risky age were significant risk factors causing LBW, while serum transferrin receptor, anemia, parity, energy and protein consumption levels, and systolic and diastolic blood pressure were nonsignificant risk factors.

Multivariate analysis with multiple logistic regression showed a significance of 0.905 for the Hosmer and Lemeshow Test, indicating an acceptable goodness of fit to the model tested. A Nagelkerke R Square of 0.528 showed that risky age and MUAC variable models are risk factors for LBW at 32.8%. The values of B and the exponential B, along with their significance, can be seen in Table 3.

The odds ratio for maternal age was 3.7 and that of MUAC was 15.38 for the incidence of LBW. Based on the table above, Exp(B) can be used to construct a logistic

Table 1. Infant Birth Weight According to Selected Maternal Factors

Variable	Category	Birth Weight		Mean	SD
		Low	Normal		
Anemia	Anemia (<11 gr/dl)	4 (11.4%)	51 (88.6%)	10.90	1.02
	Normal (11 gr/dl)	5 (8.8%)	51 (91.2%)		
Mid upper arm circumference	Chronic Energy Deficiency (< 25.5 cm)	4 (50%)	4 (50%)	26.84	2.98
	Normal (> 25.5cm)	5 (4.9%)	58 (54.8%)		
Serum transferrin receptor	Iron Deficiency (> 21 nmol/l)	6 (12.5%)	42 (87.5%)	24.80	7.29
	Normal (<21 nmol/l)	1 (4.8%)	20 (95.2%)		
Maternal age at pregnancy	Risky Age (<19 years or >35 years)	5 (30%)	7 (70%)	26.71	5.80
	Normal Age (19-35 years)	4 (6.8%)	55 (89.9%)		
Parity	Multipara (> 2 live births)	5 (9.6%)	47 (90.4%)	2.14	0.93
	Primipara (1 live birth)	2 (11.8%)	15 (88.2%)		
Level of energy consumption	Less (<80%)	1 (7.7%)	12 (92.3%)	96.09	12.66
	Normal (>80%)	6 (10.7%)	50 (89.3%)		
Level of protein consumption	Less (<80%)	2 (7.4%)	25 (92.6%)	90.54	26.77
	Normal (>80%)	5 (11.9%)	37 (88.1%)		
Systolic blood pressure	Prehypertension (>120 mmHg)	1 (3.6%)	27 (96.4%)	114.51	12.66
	Normal (<120 mmHg)	6 (14.6%)	55 (85.4%)		
Diastolic blood pressure	Prehypertension (>80 mmHg)	1 (6.7%)	14 (93.3%)	72.68	10.50
	Normal (<80 mmHg)	6 (11.1%)	48 (88.9%)		

Notes:
SD = Standard Deviation

Table 2. Prevalence Risk of Maternal Factors (n = 69)

Variable	PR	95% CI		p Value
		Lower	Upper	
Anemia	1.555	0.275	6.457	0.72
Mid upper arm circumference	19.553*	5.172	117.855	0.0001*
Serum transferrin receptor	2.857	0.522	25.350	0.527
Maternal age at pregnancy	5.895*	1.806	51.698	0.025*
Parity	0.798	0.140	4.545	0.799
Level of energy consumption	0.684	0.076	6.325	0.745
Systolic blood pressure	0.216	0.025	1.905	0.155
Diastolic blood pressure	0.571	0.065	5.155	0.614

Notes:
PR = Prevalence Risk, CI = Confidence Interval, Significant < 0.05

Table 3. Variables in the Equation

Variable	B	S.E.	Wald	df	p Value	Exp(B)
Maternal Age at Pregnancy	1.512	1.010	1.687	1	0.194	5.715
MUAC	2.752	.955	8.176	1	0.004	15.558
Constant	-5.212	.662	25.551	1	0.000	0.040

Equation 1. Probability of LBW

$$\text{Probability of LBW: } 1 + e^{(-3.2+1.3 \text{ Maternal Age Pregnancy}+2.7 \text{ MUAC} < 23.5)}$$

regression equation to determine the probability of occurrence of LBW if pregnancy occurred at a risky age and MUAC was <25.5 cm, as follows Equation 1.

Based on this model, if there was a mother with a risky age at pregnancy (<19 or >35 years old) and she had MUAC less than 25.5 cm, the probability of LBW was 68.2%.

Discussion

Anemia is the most common nutritional deficiency in the world. The World Health Organization has estimated that the prevalence of anemia is 51% in developing countries.¹⁵ In Southeast Asia, anemia during pregnancy has a 48.2% prevalence, equal to 18.1 million cases. In Indonesia, the prevalence of anemia during pregnancy is 50.9%.¹⁴ In this study, anemia in pregnant women was nonsignificant as a risk factor for having a LBW infant. Anemia in pregnant mothers associated with adverse outcomes may be related to other maternal complications, such as hemoglobinopathy, malnutrition, chronic infection, and inadequate access to prenatal care, and anemia itself may be a marker of these underlying conditions. A study by Xiong, *et al.*¹⁵ also showed that anemia was not statistically significantly associated with infant LBW.

MUAC is another anthropometric measure used to evaluate adult nutritional status that has been found to be particularly effective in determining malnutrition in adults in developing countries.¹⁶ This study used an MUAC measure threshold of 25.5 for pregnancy outcome, infant morbidity, and mortality.¹⁷ The results showed that a MUAC of less than 25.5 cm increased the risk of LBW by a factor of 19. Women with MUAC less than 25.5 cm had more LBW infants than those with MUAC of 25 cm and more. A study by Assefa,¹⁸ also showed a statistically significant association between LBW and MUAC less than 25 cm.

Serum transferrin receptor was a nonsignificant as a risk factor in LBW. Transferrin is the main iron transport protein found in the blood and plays a role in maintaining cellular iron homeostasis through regulation of cellular

iron intake. Serum transferrin receptor is largely derived from developing red blood cells. Assessment of serum transferrin receptor levels has been used to distinguish iron deficiency anemia from anemia of chronic disease because the receptors are generally unaffected by concurrent infection or inflammation.¹⁹ The results of the present study showed that the serum transferrin receptor was not significant as risk factor in LBW. Although many studies found an association between iron deficiency and pregnancy outcomes.²⁰⁻²² A previous study by Khambalia *et al*,²³ did not detect a significant association between iron deficiency and preterm birth. The inconsistencies may have been caused by study population differences and confounders.

Maternal age affects fertility. Fertility starts to decrease at the age of 20 years and decreases rapidly after the age of 35 years. Getting pregnant at a young age is also a risk factor, as the endometrium has not yet matured, whereas the endometrium is less fertile after the age of 35 years. This will increase the likelihood of having congenital syndrome and affecting maternal and child health during pregnancy.²⁴ A previous study indicated that mothers at the younger and older ends of the childbearing age range are at increased risk for LBW. This study showed that maternal age increase the risk of LBW by a factor of 5. A study by Fraser,²⁵ found that adolescent mothers aged 13 to 17 years had a significantly higher risk (p value < 0.001) than mothers aged 20 to 24 years of delivering an infant with LBW (relative risk, 1.7; 95% CI, 1.5 to 2.0). Kartasurya,²⁶ also found that pregnancy at a risky age (<20 or >35 years) was also a risk factor associated with LBW (<20 or >35 years; OR = 1.95 with CI = 1.16-3.36) in Batang District, Central Java Province.

Based on the result, parity was nonsignificant as a risk factor in LBW. Parity and maternal age have been shown to increase the risk of adverse neonatal outcomes, such as intrauterine growth restriction, prematurity, and mortality.^{22, 27, 28} Nulliparity may confer risk through complications during childbirth, such as obstructed labor, whereas high parity has been linked to an increased risk of hypertension, placenta previa, and uterine rupture.²⁹ A meta-analysis by Shah,³⁰ concludes that multiparity, although it is associated with reduced birth weight, is not associated with LBW or preterm birth, in which multiparity is often confounded by socioeconomic status. Higher complications in terms of birth outcome associated with multiparity are valid for communities with poor socioeconomic status, low levels of education, and inadequate access to health care.

In this study, energy consumption was found to be nonsignificant as a risk factor in LBW. It should be that deficient energy consumption or weight during pregnancy can cause impaired fetal growth and increase the risk of LBW in newborns.³¹ Furthermore, LBW infants have

serious health problems, such as cerebral palsy, mental retardation, and even cardiovascular disease, when they become adults.³² In study by Karima,³³ pre-pregnancy weight, weight gain during pregnancy, maternal age, and birth order were factors affecting birth weight significantly, with pre-pregnancy weight as the dominant factor (OR = 6.643, CI: 2.3-18.8). Therefore, it is important to pay more attention to undernourished women who are planning a pregnancy.³³ Pre-pregnancy weight and maternal weight gain in the first, second, and third trimesters have a moderate power relation and positive pattern.³⁴

Protein is important for fetal development because one of its roles is to form fetal cells and tissues. Protein consumption is increased during pregnancy to prevent protein deficiency and malnutrition.³⁵ In this study, the level of protein consumption was nonsignificant as risk factor in LBW. Even though a previous study by Ramakrishnan,³⁶ stated that poor nutrition is one of factors associated with LBW babies, this inconsistency may be due to the limitation of sample size in this study (n = 69) with only seven infants with LBW.

The maternal cardiovascular system undergoes progressive adaptations throughout pregnancy, including decreased vascular resistance, increased blood volume, and other metabolic changes. Systolic blood pressure (SBP) and diastolic blood pressure (DBP) decreased from the first to second trimester and then increased up to the postpartum period.³⁷ The causes of LBW are multifactorial, and one such factor is blood pressure. This study showed that SBP and DBP were nonsignificant as risk factors in LBW infants. A study by Walker *et al*,³⁸ indicated an association between blood pressure and infants with LBW. This inconsistency may be due to the inclusion of subjects with pre-hypertension, whose blood pressures were not very high (<120 mmHg for SBP and <80 mmHg for DBP).

LBW is one of the risk factors for perinatal death. LBW has a PAR value of 14.90. This means that if the focus of the intervention program is on decreasing the perinatal mortality rate by decreasing the prevalence of LBW infants, there will be a 15% reduction in the risk of perinatal death out of all live births in the population.³⁹ Interventions for pregnancy at a risky age and mothers with MUAC less than 23.5 cm are indispensable for reducing LBW occurrence, as confirmed in a study by Kartasurya.²⁶

Conclusion

It can be concluded that MUAC and age are risk factors associated with LBW in newborns, with a probability of 68.2%. It is suggested that public education on the importance of nutrition during pregnancy be increased to increase MUAC and avoid pregnancy under the age of

19 years or above the age of 35 years to reduce the incidence of infants with LBW.

10

Acknowledgment

The authors would like to thank the Directorate of Research and Community Services and the Ministry of Research, Technology and Higher Education of the Republic of Indonesia, as well as the Research Institute of Diponegoro University that facilitated and funded this study.

References

- You D, Hug L, Ejdemyr S. Level and trends in child mortality. United State of America: United Nations Children's Fund; 2015.
- Lawn J, Cousens S, Zupan J. 4 Million neonatal deaths: When? Where? Why? *Lancet Global Health*. 2005; 3(5): 891-900.
- Oza S, Lawn JE, Hogan DR, Cousens SN. Neonatal cause-of-death estimates for the early and late neonatal periods for 194 countries: 2000–2013. 2015; 93(1): 19-28.
- World Health Organization. WHA global nutrition targets 2025: low birth weight policy brief. Geneva: World Health Organization; 2012.
- Pallewatta N. Study on some risk factors affecting low birth weight born at base hospital, Gampaha. Sri Lanka: Post Graduate Institute of Medicine; 1998.
- Siza J. Risk factors associated with low birth weight of neonates among pregnant women attending a referral hospital in northern Tanzania. *Tanzan Jurnal Health Research*. 2008; 10: 1-8.
- Badan Penelitian dan Pengembangan Kementerian Kesehatan Republik Indonesia. Riset kesehatan dasar 2015. Jakarta: Badan Penelitian dan Pengembangan Kementerian Kesehatan Republik Indonesia; 2015.
- Dinas Kesehatan Kabupaten Temanggung. Rekapitulasi laporan jumlah kasus BBLR Kabupaten Temanggung tahun 2013-2015. Temanggung: Dinas Kesehatan Kabupaten Temanggung; 2015.
- López-Sierra M, Calderón S, Gómez J, Pilleux L. Prevalence of anaemia and evaluation of transferrin receptor (sTfR) in the diagnosis of iron deficiency in the hospitalized elderly patients: anaemia clinical studies in Chile. *Anemia*. 2012; 6 pages. <http://dx.doi.org/10.1155/2012/646201>.
- Sevilla C, Phunsalan A, Regala G, Uriarte PG. 1995. Pengantar metode penelitian. Jakarta: UI Press; 2013.
- Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, et al. The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: The JNC 7 report. *Jama*. 2003; 289(19): 2560-71.
- Benoist BD, McLean E, Egli I. Worldwide prevalence of anaemia report: 1995–2005. Geneva: World Health Organization; 2008.
- Kalaivani K. Prevalence & consequences of anaemia in pregnancy. 2009; 130(5): 627-33.
- Departemen Kesehatan Republik Indonesia. Profil kesehatan Indonesia 2005. Jakarta: Departemen Kesehatan Republik Indonesia; 2006.
- Xiong X, Buekens P, Alexander S, Demianczuk N, Wollast E. Anemia during pregnancy and birth outcome: a meta-analysis. *American Journal of Perinatology*. 2000; 17(3): 137-46.
- James W, Norgan N, Mascie-Taylor G, Bistrian B, Shetty P, Ferro-Luzzi A. The value of arm circumference measurements in assessing chronic energy deficiency in Third World adults. *European Journal of Clinical Nutrition*. 1994; 48(12): 883-94.
- Ariyani DE, Achadi EL, Irawati A. Validitas lingkar lengan atas mende-teksi risiko kekurangan energi kronis pada wanita Indonesia. *Kesmas: National Public Health Journal*. 2012; 7(2): 83-90.
- Assefa N, Berhane Y, Worku A. Wealth status, mid upper arm circumference (MUAC) and antenatal care (ANC) are determinants for low birth weight in Kersa, Ethiopia. *PLoS One*. 2012; 7(6): e39957. doi:10.1371/journal.pone.0039957.
- World Health Organization. Serum transferrin receptor levels for the assessment of iron status and iron deficiency in populations. Geneva: World Health Organization; 2013.
- Allen LH. Anemia and iron deficiency: effects on pregnancy outcome. *American Journal of Clinical Nutrition*. 2000; 71(5): 1280s-4s.
- Zhou L-M, Yang W-W, Hua J-Z, Deng C-Q, Tao X, Stoltzfus RJ. Relation of hemoglobin measured at different times in pregnancy to pre-term birth and low birth weight in Shanghai, China. *American Journal of Epidemiology*. 1998; 148(10): 998-1006.
- Scholl T, Hediger M, Huang J, Johnson F, Smith W, Ances I. Young maternal age and parity. Influences on pregnancy outcome. *Annual Epidemiology*. 1992; 2(5): 565-75.
- Khambalia AZ, Collins CE, Roberts CL, Morris JM1, Powell KL5, Tasevski V, et al. Iron deficiency in early pregnancy using serum ferritin and soluble transferrin receptor concentrations are associated with pregnancy and birth outcomes. *European Journal of Clinical Nutrition*. 2016; 70(3): 558-63.
- Engle WA, Kominarek MA. Late preterm infants, early term infants , and timing of elective deliveries. 2008; 35(2): 325-41. doi:10.1016/j.clp.2008.03.003.
- Fraser A, Brockert J, Ward R. Association of young maternal age with adverse reproductive outcomes. *National England Journal of Medicine*. 1995; 332(17): 1113-7.
- Kartasurya MI, Dharmawan Y, Widjanarko B, Handayani N. The prediction model for low birth weight in Batang district, central Java, Indonesia. *Advance Science Letter*. 2017; 23(4): 3442-4. doi:10.1166/asl.2017.9123.
- Kiely J, Paneth N, Susser M. An assessment of the effects of maternal age and parity in different components of perinatal mortality. *American Journal of Epidemiology*. 1986; 123(3): 444-54.
- Gibbs C, Wendt A, Peters S, Hogue C. The impact of early age at first childbirth on maternal and infant health. *Paediatric Perinatal Epidemiology*. 2012; 26(1): 259-84.
- Lee A, Mullany L, Tielsch J, et al. Community-based stillbirth rates and risk factors in rural Sarlahi, Nepal. *International Journal Gynecology Obstetric*. 2011; 113(5): 199-204.
- Shah PS. Parity and low birth weight and preterm birth: a systematic review and meta-analyses. *Acta Obstetric Gynecologic Scand*. 2010; 89(7): 862-75.
- Keefe CJ, Couch SC, Philipson E. *Handbook Of Nutrition and Pregnancy*. Maryland: Humana Press; 2008.
- Keram A, Abdulaziz A. Low birth weight prevalence, risk factors, outcomes in primary health care setting : A cross-sectional study. *Obstetric Gynecologic International Journal*. 2016; 5(5): 00176. doi:10.15406/ogij.2016.05.00176.

Kesmas: *National Public Health Journal*, 2018; 13 (2): 75-80

33. Karima K, Achadi EL. Status gizi ibu dan berat badan lahir bayi. *Kesmas: National Public Health Journal*. 2012; 7(3): 111-9.
34. Sari M, Sudiarti T. Model prediksi berat lahir bayi berdasarkan berat badan ibu hamil. *Kesmas: National Public Health Journal*. 2013; 7(8): 339-45.
35. Morgan JB. *Nutrition in early life*. England: Wiley; 2003.
36. Ramakrishnan U. Nutrition and low birth weight: from research to practice. *The American Journal of Clinical Nutrition*. 2004; 79(1): 17-21.
37. Rebelo F, Farias DR, Mendes RH, Schlüssel MM, Kac G. Blood Pressure Variation Throughout Pregnancy According to Early Gestational BMI : A Brazilian Cohort. *Arq Bras Cardiol*. 2012; 104(4): 284-91 doi:10.5935/abc.20150007.
38. Walker BR, McConnachie A, Noon JP, Webb DJ, Watt GCM. Contribution of parental blood pressures to association between low birth weight and adult high blood pressure: cross sectional study. *British Medical Journal*. 1998; 316(7134): 834-7.
39. Prameswari MF. Kematian perinatal di Indonesia dan faktor yang berhubungan, tahun 1997-2003. *Kesmas: National Public Health Journal*. 2007; 1(4): 154-60.

RISK FACTOR LBW

ORIGINALITY REPORT

16%

SIMILARITY INDEX

%

INTERNET SOURCES

16%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Kozuki, Naoko, Anne CC Lee, Mariangela F Silveira, Ayesha Sania, Joshua P Vogel, Linda Adair, Fernando Barros, Laura E Caulfield, Parul Christian, Wafaie Fawzi, Jean Humphrey, Lieven Huybregts, Aroonsri Mongkolchat, Robert Ntozini, David Osrin, Dominique Roberfroid, James Tielsch, Anjana Vaidya, Robert E Black, and Joanne Katz. "The associations of parity and maternal age with small-for-gestational-age, preterm, and neonatal and infant mortality: a meta-analysis", BMC Public Health, 2013.
Publication 1%
- 2** Assefa, Nega, Yemane Berhane, and Alemayehu Worku. "Wealth Status, Mid Upper Arm Circumference (MUAC) and Ante Natal Care (ANC) Are Determinants for Low Birth Weight in Kersa, Ethiopia", PLoS ONE, 2012.
Publication 1%
- 3** Praba Ginandjar, Lintang Dian Saraswati, Bagoes Widjanarko. "Profile of glycated- 1%

hemoglobin, antioxidant vitamin and cytokine levels in pulmonary tuberculosis patients: A cross sectional study at Pulmonary Diseases Center Semarang City, Indonesia", Biomedical Journal, 2016

Publication

4

Xiong, X.. "Anemia during pregnancy in a Chinese population", International Journal of Gynecology and Obstetrics, 200311

1%

Publication

5

Dennis, Jeff A., and Stefanie Mollborn. "Young maternal age and low birth weight risk: An exploration of racial/ethnic disparities in the birth outcomes of mothers in the United States", The Social Science Journal, 2013.

1%

Publication

6

Semira Ahmed, Kalkidan Hassen, Tolassa Wakayo. "A health facility based case-control study on determinants of low birth weight in Dassie town, Northeast Ethiopia: the role of nutritional factors", Nutrition Journal, 2018

1%

Publication

7

Gedion Asnake Azeze, Kelemu Abebe Gelaw, Natnael Atnafu Gebeyehu, Molalegn Mesele Gesese, Taklu Marama Mokonnon. "Exclusive Breastfeeding Practice and Associated Factors among Mothers in Boditi Town, Wolaita Zone,

1%

Southern Ethiopia, 2018: A Community-Based Cross-Sectional Study", International Journal of Pediatrics, 2019

Publication

8

Yutaka Kohgo, Yoshihiro Torimoto, Junji Kato. "Transferrin Receptor in Tissue and Serum: Updated Clinical Significance of Soluble Receptor", International Journal of Hematology, 2002

Publication

9

"Handbook of Nutrition and Pregnancy", Springer Nature, 2008

Publication

10

Imas Sukaesih Sitanggang, Mirza Rakhmadiani, Husnul Khotimah. "Association patterns of hotspot sequence with socio-economic aspects in peatland in Sumatra", 2016 International Conference on Computer, Control, Informatics and its Applications (IC3INA), 2016

Publication

11

Daniale Tekelia Ekubagewargies, Destaye Guadie Kassie, Wubet Worku Takele. "Maternal HIV infection and preeclampsia increased risk of low birth weight among newborns delivered at University of Gondar specialized referral hospital, Northwest

1%

1%

1%

1%

Ethiopia, 2017", Italian Journal of Pediatrics,
2019

Publication

12

Joses Muthuri Kirigia, Germano Mwigu Mwabu.
"The Monetary Value of Disability-Adjusted-
Life-Years Lost in the East African Community
in 2015", Modern Economy, 2018

Publication

1%

13

A Z Khambalia, C E Collins, C L Roberts, J M
Morris, K L Powell, V Tasevski, N Nassar. "Iron
deficiency in early pregnancy using serum
ferritin and soluble transferrin receptor
concentrations are associated with pregnancy
and birth outcomes", European Journal of
Clinical Nutrition, 2015

Publication

1%

14

"2009 International BiIron Society Meeting",
American Journal of Hematology, 08/2009

Publication

1%

15

Abdulaziz Aljohani. "Low Birth Weight
Prevalence, Risk Factors, Outcomes in Primary
Health Care Setting: A Cross-Sectional Study",
Obstetrics & Gynecology International Journal,
2016

Publication

<1%

16

Marthe-Lise Næss-Andresen, Åse Ruth
Eggemoen, Jens Petter Berg, Ragnhild Sørum

<1%

Falk, Anne Karen Jenum. "Serum ferritin, soluble transferrin receptor, and total body iron for the detection of iron deficiency in early pregnancy: a multiethnic population-based study with low use of iron supplements", *The American Journal of Clinical Nutrition*, 2019

Publication

17

Sultana, Tania, Md. Nazmul Karim, Tahmeed Ahmed, and Md. Iqbal Hossain. "Assessment of Under Nutrition of Bangladeshi Adults Using Anthropometry: Can Body Mass Index Be Replaced by Mid-Upper-Arm-Circumference?", *PLoS ONE*, 2015.

Publication

<1%

18

Bo-sin Tang, Lennon Hung-tat Choy. "Modelling planning control decisions: a logistic regression analysis on office development applications in urban Kowloon, Hong Kong", *Cities*, 2000

Publication

<1%

19

Jaqueline Lepsch, Dayana Rodrigues Farias, Ilana Eshriqui, Fernanda Rebelo et al. "Serum fatty acids are positively associated with changes in systemic blood pressure throughout pregnancy", *Pregnancy Hypertension*, 2018

Publication

<1%

20

Seyed Abdolvahab Taghavi, Najmeh Tehranian, Reza Jamhiri, Shahintaj Aramesh et

<1%

al. "Relation between total Iron intake and gestational diabetes: a cohort study", Hormozgan Medical Journal, 2018

Publication

21

Lei Jin, Lorraine F Yeung, Mary E Cogswell, Rongwei Ye, Robert J Berry, Jianmeng Liu, Dale J Hu, Li Zhu. "Prevalence of anaemia among pregnant women in south-east China, 1993–2005", Public Health Nutrition, 2010

Publication

22

Joy E. Lawn, Anne CC Lee, Mary Kinney, Lynn Sibley, Wally A. Carlo, Vinod K. Paul, Robert Pattinson, Gary L. Darmstadt. "Two million intrapartum-related stillbirths and neonatal deaths: Where, why, and what can be done?", International Journal of Gynecology & Obstetrics, 2009

Publication

23

Simegn Alemu, Abdulhalik Workicho, Mamo Nigatu, Tariku Bokila, Tsedeke Wolde. "Determinants of low birth weight in public health facilities, of Kambata Tembaro zone, South Ethiopia", PONS - medicinski casopis, 2018

Publication

24

A. Ren, J. Wang, R.W. Ye, S. Li, J.M. Liu, Z. Li. "Low first-trimester hemoglobin and low birth

<1%

<1%

<1%

<1%

weight, preterm birth and small for gestational age newborns", International Journal of Gynecology & Obstetrics, 2007

Publication

25

Ary I Savitri, Peter Zuithoff, Joyce L Browne, Dwirani Amelia, Mohammad Baharuddin, Diederick E Grobbee, Cuno S P M Uiterwaal. "Does pre-pregnancy BMI determine blood pressure during pregnancy? A prospective cohort study", BMJ Open, 2016

Publication

<1%

26

Getnet Asmare, Nigusie Berhan, Mengistu Berhanu, Animut Alebel. "Determinants of low birth weight among neonates born in Amhara Regional State Referral Hospitals of Ethiopia: unmatched case control study", BMC Research Notes, 2018

Publication

<1%

27

Gunnar Johanson. "Toxicity Review of Ethylene Glycol Monomethyl Ether and its Acetate Ester", Critical Reviews in Toxicology, 2008

Publication

<1%

28

Wondim Aboye, Tesfay Berhe, Taddis Birhane, Hadgu Gerensea. "Prevalence and associated factors of low birth weight in Axum town, Tigray, North Ethiopia", BMC Research Notes, 2018

Publication

<1%

29

Xu Xiong. "ANEMIA DURING PREGNANCY AND BIRTH OUTCOME: A META-ANALYSIS", American Journal of Perinatology, 2000

Publication

<1%

30

Kedir, Haji, Yemane Berhane, and Alemayehu Worku. "Magnitude and determinants of malnutrition among pregnant women in eastern Ethiopia: evidence from rural, community-based setting : Malnutrition during pregnancy in eastern Ethiopia", Maternal and Child Nutrition, 2014.

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

RISK FACTOR LBW

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
