



UNIVERSITI PUTRA MALAYSIA

***A COLORIMETRIC ASSAY TO DETECT MERCURY (II) IONS IN WATER
SOURCES USING CONJUGATED GOLD-NANOPARTICLES***

NURFATINI IDAYU BINTI BUSAIRI

FBSB 2018 42



**A COLORIMETRIC ASSAY TO DETECT MERCURY (II) IONS IN WATER
SOURCES USING CONJUGATED GOLD-NANOPARTICLES**

By

NURFATINI IDAYU BINTI BUSAIRI

**Thesis Submitted to School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

April 2018

All materials contained within the thesis including without limitation text, logos, icons, photographs and all other artworks are copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from copyright holder. Commercial use of materials may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

A COLORIMETRIC ASSAY TO DETECT MERCURY (II) IONS IN WATER SOURCES USING CONJUGATED GOLD-NANOPARTICLES

By

NURFATINI IDAYU BINTI BUSAIRI

April 2018

Chairman: Amir Syahir bin Amir Hamzah, PhD
Faculty: Biotechnology and Science Biomolecules

Rapid industrial progress has led to the elevation of heavy metal concentration in the environment, which raised tacit concerns. Among these metals, mercury (II) ion, Hg^{2+} is one of the most detrimental to the human health due to its neurotoxic and nephrotoxic properties. To date, methods to determine the presence of Hg^{2+} ions require expensive equipment and sophisticated operation. This is impractical for a constant monitoring of Hg^{2+} in water sources. Thus, it is necessary to develop a simple yet effective bio sensing method to monitor Hg^{2+} level in our environment. Recently, gold-nanoparticles (AuNPs) have attracted considerable amount of attention due to their nature of strong plasmonic resonance which can be exploited for a simple and rapid colorimetric assay. As the distance between nanoparticles decreases, the plasmon energy band is lowered and consequently it turns the colour of AuNPs from red to blue. The key of utilizing this phenomenon as a bio sensing mechanism is to control the AuNPs aggregation via surface chemistry approach. Here, a sensitive and practical colorimetric assay for *in-situ* detection of Hg^{2+} ions in water using cysteine functionalized gold nanoparticles (Cys-AuNPs) together with the addition of polyamidoamine (PAMAM), G2 dendrimer is reported. In this study, Cys-AuNPs and PAMAM dendrimer specifically capture Hg^{2+} ions and induce colour changes of the solution. The mechanism of Hg^{2+} interaction with the system was investigated using UV-vis spectrophotometer, dynamic light scattering (DLS) and transmission electron microscope (TEM). To increase the practicality value of the system, a custom-made chamber to be integrated with a mobile phone was fabricated. Colour changes can be monitored using a mobile app. The newly developed detection system can detect Hg^{2+} as low as 0.5 part per billion (ppb) in a laboratory prepared sample or in a real river water sample. The results were validated using direct thermal decomposition mercury analyser. This cost-effective colorimetric approach is practical to determine the presence of Hg^{2+} water sources.

Abstrak tesis yang dikemukakan kepada Senat of Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

KOLORIMETRIK ASSAY UNTUK MENGESAN MERKURI (II) ION DALAM SUMBER AIR MENGGUNAKAN KONJUGAT EMAS NANOPARTIKEL

Oleh

NURFATINI IDAYU BINTI BUSAIRI

April 2018

Pengerusi: Amir Syahir Bin Amir Hamzah, PhD
Fakulti: Bioteknologi dan Sains Biomolekul

Kemajuan industri yang pesat telah menyebabkan peningkatan kepekatan logam berat dalam alam sekitar, yang menimbulkan kebimbangan. Di antara logam-logam ini, ion Hg^{2+} adalah salah satu daripada ion logam yang membahayakan kesihatan manusia akibat sifat neurotoksik dan nefrotoksiknya. Sehingga kini, kaedah untuk menentukan kehadiran ion Hg^{2+} memerlukan peralatan mahal dan operasi yang canggih. Ini tidak praktikal untuk pemantauan berterusan Hg^{2+} dalam sumber air. Oleh itu, pembangunkan kaedah yang mudah tetapi berkesan adalah perlu untuk memantau tahap logam berat di persekitaran kita. Baru-baru ini, emas-nanopartikel (AuNPs) telah menarik banyak perhatian kerana sifat resonans plasmoniknya yang kuat yang boleh dieksploitasi untuk pengujian warna yang mudah dan cepat. Apabila jarak antara nanopartikel berkurangan, kumpulan plasmon emas-nanopartikel akan beralih dan seterusnya mengubah warna AuNPs dari merah ke biru. Fenomena ini boleh digunakan sebagai mekanisme penderiaan dengan cara mengawal pengagregatan AuNPs melalui pendekatan kimia permukaan. Di sini dilaporkan ujian kolorimetric yang sensitif dan praktikal untuk pengesanan in-situ ion Hg^{2+} dalam air menggunakan cystiene-nanopartikel emas (Cys-AuNPs) bersama dendrimer polyamidoamine (PAMAM). Dalam kajian ini, dendrimer Cys-AuNPs dan PAMAM secara khusus akan menangkap Hg^{2+} dan mendorong perubahan warna koloid. Mekanisme interaksi Hg^{2+} dengan sistem ini disiasat menggunakan spektrofotometer UV-vis, penyebaran cahaya dinamik (DLS) dan mikroskop pengaliran elektron (TEM). Untuk meningkatkan nilai praktikal sistem, kotak kedap cahaya telah dibina untuk disepadukan dengan telefon bimbit bagi tujuan analisis in-situ. Oleh itu, perubahan warna boleh diiktiraf menggunakan aplikasi mudah alih. Sistem pengesanan yang baru dibangunkan dapat mengesan Hg^{2+} serendah 0.5 bahagian per bilion (ppb) dalam sampel yang disediakan atau dalam sampel air sungai sebenar dan disahkan menggunakan penganalisis merkuri penguraian haba langsung. Pendekatan kolorimetric yang kos efektif adalah praktikal untuk menentukan kehadiran Hg^{2+} dalam sumber air.

ACKNOWLEDGEMENTS

In the name of Allah, the most gracious, the most merciful, all praises and glorification are due to Him.

First and foremost, I would like to extend my utmost gratitude to my Advisor, Dr. Amir Syahir Bin Amir Hamzah for his thoughtful insights and mentorship throughout the completion of my study. My heartiest thanks also go to my co-supervisors Prof. Dato' Dr. Abu Bakar bin Salleh and Dr. Asilah binti Ahmad Tajudin for their constant assistance and tireless effort in helping me throughout my Master journey.

My appreciation to Dr. Mohd Rashdan bin Saad and his team from Universiti Pertahanan Nasional Malaysia (UPNM) for their assistance for making part of this research smoother. My thanks also go to all my EMTECH fellow and friends who being a wonderful comrade throughout this time.

Last but definitely not least, my deepest appreciation goes to both of my parents, my lovely husband and family for their never-ending support, prayers, and encouragement throughout the duration of my study in UPM. May Allah repay the kindness of all the living souls who keep me motivated and have helped me in accomplishing my Master Degree

“ . And put thy trust in Allah and enough is Allah as a disposer of affairs.” [33:3]

I certify that a Thesis Examination Committee has met on 25 April 2018 to conduct the final examination of Nurfatini Idayu binti Busairi on her thesis entitled "A Colorimetric Assay to Detect Mercury (II) Ions in Water Sources Using Conjugated Gold-Nanoparticles" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Tan Wen Siang, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Nor Azah binti Yusof, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Ruslinda A. Rahim, PhD

Senior Lecturer
Universiti Malaysia Perlis
Malaysia
(External Examiner)



RUSLI HAJI ABDULLAH, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 30 August 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Amir Syahir Bin Amir Hamzah, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Science
Universiti Putra Malaysia
(Chairman)

Abu Bakar Bin Salleh, PhD

Professor
Enzyme Technology, Faculty of Biotechnology and Biomolecular Science
Universiti Putra Malaysia
(Member)

Asilah Binti Ahmad Tajudin, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Science
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of the thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No: Nurfatini Idayu Binti Busairi, GS45716

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of Chairman of
Supervisory
Committee: _____

Signature: _____

Name of Member of
Supervisory
Committee: _____

Signature: _____

Name of Member of
Supervisory
Committee: _____

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	ii
ACKNOWLEDGEMENTS	iii
APPROVAL	iv
DECLARATION	vi
LIST OF TABLES	x
LIST OF FIGURES	xi
LIST OF ABBREVIATIONS	xiii
CHAPTER	
1 INTRODUCTION	1
1.1 Background of Study	1
1.2 Problem Statement	2
1.3 Objectives	4
2 LITERATURE REVIEW	5
2.1 Occurrence of Heavy Metals	5
2.1.1 The Occurrence and Mobility of Mercury, Hg ²⁺	6
2.2 Development of Different Types of Sensor to Detect Hg ²⁺	10
2.2.1 Current Methods	10
2.2.2 Bio-sensing Methods	11
2.2.3 Ligand of Chelating Agent	13
2.3 Colorimetric Assay as a Biosensor to Detect Heavy Metals	15
2.4 Conjugated Gold-Nanoparticles (AuNPs)	16
2.5 Capturing Agent for Mercuric Ions, Hg ²⁺	17
2.6 Polyamioamide G2 as a Capturing Agent for Mercury, Hg ²⁺	17
2.7 Conclusion	18
3 MATERIALS AND METHODS / METHODOLOGY	19
3.1 Material	19
3.1.1 All Chemicals	19
3.1.2 Equipment	20
3.2 Overall work	20
3.3 Methods	22
3.3.1 General Preparation	22
3.3.2 Chemical Preparation for Bio-Conjugation	22
3.3.3 Chemical Preparation for Sensitivity Improvement	23

3.3.4	Preparation of Bio-Conjugated Gold-Nanoparticles	23
3.3.5	Colorimetric Assay of Hg ²⁺ Ions	24
3.3.6	Characterization of Gold-Nanoparticles	25
3.3.7	Specificity Study	25
3.3.8	Mobile Chamber Fabrication, Optimization and Printing	25
3.3.9	In-situ Analysis using sensor System	29
3.3.10	Data Analysis	29
4	RESULTS AND DISCUSSION	31
4.1	Colorimetric Assay of Hg ²⁺	31
4.1.1	The Determination of Hg ²⁺ Specific Capturing Agent	31
4.1.2	Cysteine-AuNPs-PAMAM, for Detection of Hg ²⁺	33
4.2	Selectivity Study	43
4.2.1	TEM and DLS data on Selectivity study	46
4.3	Color Analysis Apps for Detection of Hg ²⁺	46
4.4	Validation Study (<i>In-Situ Analysis</i>)	49
5	SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	52
5.1	Conclusion	52
5.2	Recommendation for Future Research	54
	REFERENCES/BIBLIOGRAPHY	56
	BIO DATA OF STUDENT	70

LIST OF TABLES

Table		Page
2.1	Mercury concentration in several rivers around Malaysia.	14
2.2	Reported concentration of mercury using bio monitors in Malaysia.	15
2.3	Nanomaterial based-sensor for rapid detection of Hg ²⁺ .	19
2.4	Capturing agents that have been used to capture mercury (II) ions.	22
3.1	List of chemicals.	29
3.2	List of equipment.	30
4.1	Capturing agents that have been used to capture mercury (II) ions.	47
4.2	Comparison of Color Analysis Apps data with data captured with 12.1-megapixel camera.	66
4.3	Data on pH value and concentration of heavy metals and salt in three different samples.	71
5.1	Nanomaterial based-sensor for rapid detection of mercury ions, Hg ²⁺ and a comparison with a recent work.	74

LIST OF FIGURES

Figures		Page
1.1	Mercury emission percentage based on different sources of emission.	4
21	Mercury cycle schematic diagram.	12
3.1	Overall workflow of this study.	31
3.2	First chamber design using SketchUp: (a) Top right view; (b) Bottom view; (c) Top left view and (d) Top view	39
3.3	3D Ponterface printing software set up before printing process.	41
3.4	Final product of mobile chamber: (a) After 3D printing; (b) Complete chamber with LED, switch and battery; and (c) Chamber with phone.	42
3.5	(a) Formula figure of merit; and (b) Method to calculate figure of merit using absorbance graph.	43
4.1	Schematic diagram of Hg^{2+} colorimetric assay.	45
4.2	Schematic diagram of Hg^{2+} colorimetric assay using cysteine conjugated gold-nanoparticles, enhanced with PAMAM G2.	49
4.3	Colorimetric result for two stages of this sensor in increasing concentration of Hg^{2+} (1000ppb to 0.1 ppb).	50
4.4	Corresponding UV-vis data for naked eye detection of Hg^{2+} with varies concentration (0.1ppb to 1000ppb): (a) First stage (b) Enhancement stage.	51
4.5	Figure of merit data for both stages with increasing Hg^{2+} concentration (0.5 ppb-1000ppb).	52
4.6	Dynamic light scattering (DLS) graph for first stage of the reaction (size in radius).	54
4.7	Dynamic light scattering (DLS) graph for the enhancement stage of the reaction (size in radius).	55
4.8	Dynamic Light Scattering data for average size of particles for both stages.	56
4.9	20 nm Cys-AuNP system (first stage) image under transmission electron microscope at 200kV with; 1 ppb Hg^{2+} (a) 10 ppb Hg^{2+} (b) 100 ppb Hg^{2+} (c).	57
4.10	20 nm Cys-AuNPs-PAMAM system (enhancement stage) image under transmission electron microscope at 200kV with; 0.1 ppb Hg^{2+} (a) 0.5 ppb Hg^{2+} (b) 1 ppb Hg^{2+} (c).	58
4.11	TEM image of (a) Cys-AuNPs with 1 ppb Hg^{2+} , (b) Cys-AuNPs-PAMAM with 1 ppb of Hg^{2+} . Figure (c) and (d) is the DLS graph for (a) and (b) respectively.	60
4.12	UV-vis absorption spectra (a) and the corresponding photo image (b) of Cys-AuNPs-PAMAM containing 100 ppb of different metal ions.	62
4.13	Figure of merit graph with different heavy metal ions at different concentration ranging from 0.1 to 1000 ppb.	63
4.14	TEM image of; (a) Cys-AuNPs-PAMAM (control); (b) Cys-AuNPs-PAMAM with Hg^{2+} at 1 ppb; (c) Cys-AuNPs-PAMAM with other heavy metal ions at 1 ppb. Picture (d) (e)	64

	(f) is the respective DLS data on average size of sample for (a) (b) and (c) respectively.	
4.15	RGB value of Cys-AuNPs-PAMAM with different concentration of mercury (II) ions.	68
4.16	HSB value of Cys-AuNPs-PAMAM with different concentration of mercury (II) ions.	68
4.17	UV-vis graph for naked eye detection of Hg^{2+} in three different rivers. (b) Naked eye data for each sample taken from three different rivers.	70
4.18	The saturation value of each sample against standard curve for HSV data.	70



LIST OF ABBREVIATIONS

%	Percent
°C	Degree Celcius
<i>et al.,</i>	And friends
Ag	Silver
As	Arsenic
AuNPs	Gold-nanoparticles
Cd	Cadmium
Co	Cobalt
Cr	Chromium
Cu	Copper
Cys-AuNPs	Cysteine-gold nanoparticles
DLS	Dynamic light scattering
g	Gram
Hg	Mercury
M	Molar
mg	Milligram
mL	Milliliter
μL	Microlitre
mM	Millimolar
μM	Micromolar
Ni	Nickel
PAMAM	Polyamidoamine
Pb	Lead
ppb	Part per billion
Ppm	Part per million
POC	Point of care
Ps	Picoseconds
TEM	Transmission electron microscope
Zn	Zinc

CHAPTER 1

INTRODUCTION

1.1 Background of Study

Mercury is classified as heavy metals and regarded as one of the most toxic metals by many international agencies due to its persistence, bioaccumulation, and toxicity (PBT). It is a non-transition metal and exists in three main form; elemental mercury (Hg^0), organic mercury (methylmercury, CH_3Hg & ethyl mercury, $\text{C}_2\text{H}_5\text{Hg}$) and inorganic mercury (Hg^+ , Hg^{2+}). These three forms possess different toxicity and health's implication. Generally, inorganic mercury has a high toxicity level compared to the other two forms. Mercury is known for its neurotoxic and nephrotoxic properties. It can cause neurological and behavioural disorder (Liu et al., 2013) and permanent damage to the kidney and digestive system. Exposure to mercury as low as 0.01 ppm either through breast milk or through water sources, can cause early childhood neurocognitive effects towards babies and toddlers (Karagas et al., 2012). Inorganic mercury exposure can also impaired children's immune system which eventually makes the children susceptible to infections. (Mutter and Yeter, 2008).

Elemental mercury (Hg^0), exist in a form of volatile liquid which can lead to a prolong anthropogenic release since it can remain suspended in the atmosphere for a one or two year long (Eckley et al., 2012; Eckley et al., 2015). Hg^0 can be oxidized to its inorganic form Hg^+ or Hg^{2+} and will be deposited into soil or water sources through the rain. Industrial release such as waste from incinerators, coal-burning power plant and gold mining industry can also be the sources of the inorganic mercury. In the water sources, inorganic mercury will be methylated by aquatic anaerobic sulfate-reducing bacteria to become methylmercury (MeHg). This methylmercury (MeHg) will then be ingested by fish, shellfish and other aquatic life and accumulated as it goes up the food chain.

Due to increasing industrial and mining activity globally, we are now facing a silent threat from mercury contamination. Human exposure to mercury can occur via eating MeHg-contaminated food (fish, shellfish, and other aquatic life), dental amalgam procedure, usage of inorganic mercury products (medication, germicidal soap and skin cream), occupational exposure (mining or mercury-related industry) and usage of any other mercury-based products (thermometer, sphygmomanometer, fluorescent light bulbs and batteries).

The United Nations Environmental Programme (UNEP) and the Arctic Monitoring and Assessment Programme (AMAP), estimated anthropogenic mercury emissions globally in 2010 to about 1960 tonnes·year⁻¹ with an uncertainty range of 1010–4070 tonnes·year⁻¹. Artisanal and small-scale gold mining (ASGM) were identified as the major anthropogenic sources (37%) and the dental amalgam emitted through cremation

were identified as the minor anthropogenic sources (0.2%) (United Nations Environment Programme, Global Mercury Assessment, 2013). Figure below shows a mercury emission percentage based on different sources of emission.




Figure 1.1: Mercury Emission Percentage Based on Different Sources of Emission.

Artisanal and small-scale gold mining (ASGM) were identified as the major anthropogenic sources (37%) and the dental amalgam emitted through cremation were identified as the minor anthropogenic sources (0.2%).

There are several cases worth mentioning to show that contaminated water is one of the main sources where mercury contamination could affect most lives. Water sources are the reservoir where all the inorganic mercury would reside before it will be methylated and would be magnified in the higher organism hierarchy as it passes up a food chain. In general population, dietary intake is the most common pathway for mercury contamination. The classic case of Minamata Bay Japan, in 1953 witnessed 2252 victims poisoned with MeHg (5.61ppm to 35.7 ppm) in their marine product (Harada, 1995). In 1964, the similar case has happened in Niigata with approximately 700 victims through the same exposure of the contaminated marine products (Maruyama et al., 2012). In 2006, China faced a threat from mercury pollution due to their mining and coal-burning industry. National Bureau of Oceanography (NBO) of China stated that there is approximately 77 ton of Hg were released into the coastal area through river annually (Jiang et al., 2006; Li et al., 2009).

1.2 Problem Statement

Due to the increasing amount of mercury release to the environment through anthropogenic activities and the alarming health risk of mercury consumption, monitoring it becoming an important yet challenging task to the authorities. In most

cases, mercury contamination in fish can be contained or intervened if authorities are able to constantly monitor the mercury level in water sources, for fast prevention or remediation action to be taken.

There are several detection techniques that have been developed and commercially available to detect mercury in water sources including, atomic absorption spectrometry (AAS) (Bannon and Chisolm, 2001; Fang et al., 1984; Maquieira and Elmahadi, 1994), Synchrotron X-ray fluorescence spectrometry (SXRF) spectrometry (Arai et al., 2003), microwave plasma atomic emission spectroscopy (MP-AES) (Poirier et al., 2017; Rios et al., 2017), inductively coupled plasma mass spectrometry (ICP-MS) (Shih et al., 2016; Yamakawa et al., 2017), cold vapor atomic fluorescence spectrometry (CVASF) (Ebdon et al., 2002) and thermal decomposition mercury analyser (TDMA) (Sasamoto et al., 2017). These techniques provide sensitive and accurate result in determining the level of heavy metal in the environment. However, these techniques require expensive instruments, sophisticated method and time-consuming procedure which make them less mobile and certainly non-practical to be used as *in-situ* analytical tools for frequent monitoring (Zhong et.al, 2015 and Xia et.al, 2012). These limitations lead to the needs of developing a simple yet effective technique as a first line screening to detect mercury ions in water sources.

The goal of this research is to overcome limitations of the current methods through biosensor technique. Biosensor is a device that uses biologically based element including cells, proteins, enzymes, nucleic acids or small molecules to detect the presence of targeted compound; in this case, Hg^{2+} . Biosensor can be divided into several types such as electrochemical, thermometric, potentiometric and colorimetric assay. A simple colorimetric assay that can be detected via naked eye is beneficial for *in-situ* analysis since it will provide rapid measurement, generally simple and most importantly, it can be portable (Liu and Lu, 2003; Zhao et al., 2008; Wang et al., 2012).

In a colorimetric assay, there are several colouring agents that are commonly used which are, enzymes (Kaur et.al, 2014), 4-aminoantipyrine (Kim *et.al*, 2015), Prussian Blue (Zhai *et.al*, 2013), redox enzyme horseradish peroxide (HRP) (Han *et.al*, 2001) and gold-nanoparticles. Recently, gold nanoparticles as colorimetric assay agent have been widely used for heavy metal detection in aqueous solution (Wang & Ma, 2009). Gold nanoparticles (AuNPs) possess an exploitable plasmonic property that can be manipulated via surface functionalization (Knech & Sethi, 2009). It has high extinction coefficient which makes it suitable to act as a sensing platform (Aragay et al., 2011; Guo et al., 2011) As the distance between particles decreases below 4 nm, plasmon coupling will decrease the plasmon-band energy level results in a change of absorption maxima of AuNP solution that consequently can change the colour of colution from red to blue colour; namely a red shift (Ghosh & Pal, 2007). This is the phenomenon where the absorption peak of the AuNPs, shifts towards the infrared region. Using a bare AuNP will allow the detection of approximately 5 ppm for almost all metal ion. Hence, it requires some surface modifications for it to be more selective and more sensitive towards targeted compound. This is often accomplished using aptamer, DNA or protein (Aragay et al., 2011). Hence, it is a promising approach to enhance the selectivity and sensitivity of the AuNP.

Mercury contamination in Malaysia is a raising concern among authorities. Even though there is no big mercury spillage case Malaysia, but according to Praveena et al (2013), the concentration of mercury in Malaysia water sources specifically in Terengganu, Johor, and Penang is above the permissible limit set by WHO and Department of Environmental of Malaysia (DOE). The current trends demand a development of a cheap, simple, yet sensitive bio sensor method to be used for a constant monitoring of mercury in more than 1000 monitoring points of the Malaysia rivers. These monitoring points are currently being observed by the Department of Environmental of Malaysia quarterly.

To the best of our knowledge, there are only a few studies on using colorimetric assay through bio-conjugated gold nanoparticles to detect inorganic mercury in water sources. There are still problems related to sensitivity as they cannot achieve the sensitivity limit set by World Health Organization (WHO) which is at 0.001 (mg/L) or ppm level. Some study did not provide field study data for *in-situ* analysis of real samples and make it hard for us to see if it can be practical for monitoring mercury (II) ions (Hg^{2+}) in water sources. Therefore, this research comes to fill in the gap and limitations stated above.

1.3 Objectives

The main objective of this study is to build the sensor system where it can detect Hg^{2+} down to at least 1 $\mu\text{g/L}$ (ppb) level to match the WHO permissible limit of mercury in water sources. This objective can be achieved through sub-objective listed below:

1. To conjugate gold-nanoparticles with several groups of compounds to search for the best capturing agent for Hg^{2+} .
2. To fabricate and build a mobile detection system for a practical *in-situ* analysis.

REFERENCES

- Akter, S., Salam, A., & Khan, A. H. (2015). Efficiency of Natural Bio-Adsorbents To Remove Heavy Metal Ions from Artificially Contaminated Water. *Bangladesh J*, 37(1), 57-63.
- Ali, H., Khan, E., & Sajad, M. A. (2013). Phytoremediation of heavy metals— concepts and applications. *Chemosphere*, 91(7), 869-881.
- Alkarkhi, F. A., Ismail, N., & Easa, A. M. (2008). Assessment of arsenic and heavy metal contents in cockles (*Anadara granosa*) using multivariate statistical techniques. *Journal of hazardous materials*, 150(3), 783-789.
- Aragay, G., Pons, J., & Merkoçi, A. (2011). Recent trends in macro-, micro-, and nanomaterial-based tools and strategies for heavy-metal detection. *Chemical reviews*, 111(5), 3433-3458.
- Aragay, G., Pons, J., & Merkoçi, A. (2011). Enhanced electrochemical detection of heavy metals at heated graphite nanoparticle-based screen-printed electrodes. *Journal of Materials Chemistry*, 21(12), 4326-4331.
- Arai, Y., Lanzirrotti, A., Sutton, S., Davis, J. A., & Sparks, D. L. (2003). Arsenic speciation and reactivity in poultry litter. *Environmental Science & Technology*, 37(18), 4083-4090.
- Atafar, Z., Mesdaghinia, A., Nouri, J., Homaei, M., Yunesian, M., Ahmadi Moghaddam, M., & Mahvi, A. H. (2010). Effect of fertilizer application on soil heavy metal concentration. *Environmental monitoring and assessment*, 160(1), 83-89.
- Balian, H. G., & Eddy, N. W. (1977). Figure-of-merit (FOM), an improved criterion over the normalized chi-squared test for assessing goodness-of-fit of gamma-ray spectral peaks. *Nuclear Instruments and Methods*, 145(2), 389-395.
- Basta, N. T., Ryan, J. A., & Chaney, R. L. (2005). Trace element chemistry in residual-treated soil. *Journal of environmental quality*, 34(1), 49-63.
- Bannon, D. I., & Chisolm, J. J. (2001). Anodic stripping voltammetry compared with graphite furnace atomic absorption spectrophotometry for blood lead analysis. *Clinical chemistry*, 47(9), 1703-1704.
- Batool, Z., Yousafzai, N. A., Murad, M. S., Shahid, S., & Iqbal, A. (2017). Lead Toxicity and Evaluation of Oxidative Stress in Humans. *PSM Biological Research*, 2(2), 79-82.
- Bellinger, D. C., Chen, A., & Lanphear, B. P. (2017). Establishing and Achieving National Goals for Preventing Lead Toxicity and Exposure in Children. *JAMA pediatrics*. 171(7), 616 -618.

- Bhalla, V., Tejpal, R., & Kumar, M. (2010). Rhodamine appended terphenyl: a reversible “off-on” fluorescent chemosensor for mercury ions. *Sensors and Actuators B: Chemical*, 151(1), 180-185.
- Bhalla, V., Tejpal, R., Kumar, M., & Sethi, A. (2009). Terphenyl derivatives as “turn on” fluorescent sensors for mercury. *Inorganic chemistry*, 48(24), 11677-11684.
- Carocci A., Rovito N., Sinicropi M.S., Genchi G. (2014) Mercury Toxicity and Neurodegenerative Effects. In: Whitacre D. (eds) *Reviews of Environmental Contamination and Toxicology. Reviews of Environmental Contamination and Toxicology (Continuation of Residue Reviews)*, 229, 1-18.
- Chai, F., Wang, C., Wang, T., Ma, Z., & Su, Z. (2009). L-cysteine functionalized gold nanoparticles for the colorimetric detection of Hg²⁺ induced by ultraviolet light. *Nanotechnology*, 21(2), 025501.
- Chandurvelan, R., Marsden, I. D., Gaw, S., & Glover, C. N. (2017). Acute and sub-chronic effects of sub-lethal cadmium exposure on energy metabolism in the freshwater shrimp, *Paratya curvirostris*. *Ecotoxicology and environmental safety*, 135, 60-67.
- Chansuvarn, W., & Imyim, A. (2012). Visual and colorimetric detection of mercury (II) ion using gold nanoparticles stabilized with a dithia-diaza ligand. *Microchimica Acta*, 176(1-2), 57-64.
- Chen, Z., Meng, H., Xing, G., Chen, C., Zhao, Y., Jia, G., & Chai, Z. (2006). Acute toxicological effects of copper nanoparticles in vivo. *Toxicology letters*, 163(2), 109-120.
- Cózar, A., Echevarría, F., González-Gordillo, J. I., Irigoien, X., Úbeda, B., Hernández-León, S., & Fernández-de-Puelles, M. L. (2014). Plastic debris in the open ocean. *Proceedings of the National Academy of Sciences*, 111(28), 10239-10244.
- Darbha, G. K., Singh, A. K., Rai, U. S., Yu, E., Yu, H., & Chandra Ray, P. (2008). Selective detection of mercury (II) ion using nonlinear optical properties of gold nanoparticles. *Journal of the American Chemical Society*, 130(25), 8038-8043.
- De Almeida Lopes, A. C. B., Silbergeld, E. K., Navas-Acien, A., Zamoiski, R., da Cunha Martins Jr, A., Camargo, A. E. I., & Paoliello, M. M. B. (2017). Association between blood lead and blood pressure: a population-based study in Brazilian adults. *Environmental Health*, 16(1), 27.
- De Roma, A., Esposito, M., Chiaravalle, E., Miedico, O., De Filippis, S. P., & Brambilla, G. (2017). Occurrence of cadmium, lead, mercury, and arsenic in prepared meals in Italy: Potential relevance for intake assessment. *Journal of Food Composition and Analysis*, 63, 28-33.

- Denizli, A., Senel, S., Alsancak, G., Tüzmen, N., & Say, R. (2003). Mercury removal from synthetic solutions using poly (2-hydroxyethylmethacrylate) gel beads modified with poly (ethyleneimine). *Reactive and Functional Polymers*, 55(2), 121-130.
- Di Felice, R., & Selloni, A. (2004). Adsorption modes of cysteine on Au (111): Thiolate, amino-thiolate, disulfide. *The Journal of chemical physics*, 120(10), 4906-4914.
- Diallo, M. S., Balogh, L., Christie, S., Swaminathan, P., Shi, X., Goddard III, W. A., & Johnson Jr, J. H. (2005). Dendritic nanoscale chelating agents: synthesis, characterization, and environmental applications. *Nanotechnology and the environment*, 89(32), 238-247.
- Ding, S. Y., Dong, M., Wang, Y. W., Chen, Y. T., Wang, H. Z., Su, C. Y., & Wang, W. (2016). Thioether-based fluorescent covalent organic framework for selective detection and facile removal of mercury (II). *Journal of the American Chemical Society*, 138(9), 3031-3037.
- Duffus, J. H. (2002). "Heavy metals" a meaningless term? (IUPAC Technical Report). *Pure and applied chemistry*, 74(5), 793-807.
- Ebdon, L., Foulkes, M. E., Le Roux, S., & Muñoz-Olivas, R. (2002). Cold vapour atomic fluorescence spectrometry and gas chromatography-pyrolysis-atomic fluorescence spectrometry for routine determination of total and organometallic mercury in food samples. *Analyst*, 127(8), 1108-1114.
- Eckley, C. S., Parsons, M. T., Mintz, R., Lapalme, M., Mazur, M., Tordon, R., & St Louis, V. (2013). Impact of closing Canada's largest point-source of mercury emissions on local atmospheric mercury concentrations. *Environmental science & technology*, 47(18), 10339-10348.
- Eckley, C. S., Blanchard, P., McLennan, D., Mintz, R., & Sekela, M. (2015). Soil-air mercury flux near a large industrial emission source before and after Closure (Flin Flon, Manitoba, Canada). *Environmental science & technology*, 49(16), 9750-9757.
- Esfand, R., & Tomalia, D. A. (2001). Poly (amidoamine)(PAMAM) dendrimers: from biomimicry to drug delivery and biomedical applications. *Drug discovery today*, 6(8), 427-436.
- Fang, Y., Sun, X., Yang, W., Ma, N., Xin, Z., Fu, J., & Hu, Q. (2014). Concentrations and health risks of lead, cadmium, arsenic, and mercury in rice and edible mushrooms in China. *Food chemistry*, 147, 147-151.
- Fang, Y., & Liu, Y. (2013). Ultrasensitive and selective sensing of heavy metal ions with modified graphene. *Chem. Communication*, 49, 6492-6494.

- Fang, Z., Růžička, J., & Hansen, E. H. (1984). An efficient flow-injection system with on-line ion-exchange pre-concentration for the determination of trace amounts of heavy metals by atomic absorption spectrometry. *Analytica chimica acta*, 164, 23-39.
- Flora, G., Gupta, D., & Tiwari, A. (2012). Toxicity of lead: a review with recent updates. *Interdisciplinary Toxicol*, 5(2), 47-58.
- Foti, C., Giuffrè, O., Lando, G., & Sammartano, S. (2009). Interaction of inorganic mercury (II) with polyamines, polycarboxylates, and amino acids. *Journal of Chemical & Engineering Data*, 54(3), 893-903.
- García-Ordiales, E., Esbrí, J. M., Covelli, S., López-Berdonces, M. A., Higuera, P. L., & Loredó, J. (2016). Heavy metal contamination in sediments of an artificial reservoir impacted by long-term mining activity in the Almadén mercury district (Spain). *Environmental Science and Pollution Research*, 23(7), 6024-6038.
- García-Sánchez, A., & Álvarez-Ayuso, E. (2003). Arsenic in soils and waters and its relation to geology and mining activities (Salamanca Province, Spain). *Journal of Geochemical Exploration*, 80(1), 69-79.
- Genchi, G., Sinicropi, M. S., Carocci, A., Lauria, G., & Catalano, A. (2017). Mercury Exposure and Heart Diseases. *International journal of environmental research and public health*, 14(1), 74.
- Ghosh, S. K., & Pal, T. (2007). Interparticle coupling effect on the surface plasmon resonance of gold nanoparticles: from theory to applications. *Chemical reviews*, 107(11), 4797-4862.
- Gong, J., Zhou, T., Song, D., Zhang, L., & Hu, X. (2009). Stripping voltammetric detection of mercury (II) based on a bimetallic Au–Pt inorganic–organic hybrid nanocomposite modified glassy carbon electrode. *Analytical chemistry*, 82(2), 567-573.
- Gong, J., Zhou, T., Song, D., & Zhang, L. (2010). Monodispersed Au nanoparticles decorated graphene as an enhanced sensing platform for ultrasensitive stripping voltammetric detection of mercury (II). *Sensors and actuators B: Chemical*, 150(2), 491-497.
- Guariglia, S. R., Stansfield, K. H., McGlothlan, J., & Guilarte, T. R. (2016). Chronic early life lead (Pb 2+) exposure alters presynaptic vesicle pools in hippocampal synapses. *BMC Pharmacology and Toxicology*, 17(1), 56.
- Güney, O., & Cebeci, F. Ç. (2010). Molecularly imprinted fluorescent polymers as chemosensors for the detection of mercury ions in aqueous media. *Journal of applied polymer science*, 117(4), 2373-2379.

- Guo, Y., Wang, Z., Qu, W., Shao, H., & Jiang, X. (2011). Colorimetric detection of mercury, lead and copper ions simultaneously using protein-functionalized gold nanoparticles. *Biosensors and Bioelectronics*, 26(10), 4064-4069.
- Gupta, V. K., Chandra, S., & Agarwal, S. (2003). Mercury selective electrochemical sensor based on a double armed crown ether as ionophore. *Indian Journal of Chemistry*, 42, 813-818.
- Hajeb, P., Jinap, S., Ismail, A., Fatimah, A. B., Jamilah, B., & Rahim, M. A. (2009). Assessment of mercury level in commonly consumed marine fishes in Malaysia. *Food Control*, 20(1), 79-84.
- Han, S., Zhu, M., Yuan, Z., & Li, X. (2001). A methylene blue-mediated enzyme electrode for the determination of trace mercury(II), mercury(I), methylmercury, and mercury-glutathione complex. *Biosensors and Bioelectronics*, 16, 9-16.
- Harada, M. (1995). Minamata disease: methylmercury poisoning in Japan caused by environmental pollution. *Critical reviews in toxicology*, 25(1), 1-24.
- Hilson, G. (2006). Abatement of mercury pollution in the small-scale gold mining Industry: restructuring the policy and research agendas. *Science of the Total Environment*, 362(1), 1-14.
- Hsieh, N. H., Chung, S. H., Chen, S. C., Chen, W. Y., Cheng, Y. H., Lin, Y. J., ... & Liao, C. M. (2017). Anemia risk in relation to lead exposure in lead-related manufacturing. *BMC public health*, 17(1), 389.
- Huang, C. C., Yang, Z., Lee, K. H., & Chang, H. T. (2007). Synthesis of highly fluorescent gold nanoparticles for sensing mercury (II). *Angewandte Chemie*, 119(36), 6948-6952.
- Hung, Y. L., Hsiung, T. M., Chen, Y. Y., Huang, Y. F., & Huang, C. C. (2010). Colorimetric detection of heavy metal ions using label-free gold nanoparticles and alkanethiols. *The Journal of Physical Chemistry C*, 114(39), 16329-16334.
- Illustrations of Chemistry. *Sci Am*. 1846; 1 (22):2-27.
- Islam, M. M., Halim, M. A., Safiullah, S., Hoque, S. W., & Islam, M. S. (2009). Heavy metal (Pb, Cd, Zn, Cu, Cr, Fe and Mn) content in textile sludge in Gazipur, Bangladesh. *Research Journal of Environmental Sciences*, 3(3), 311-315.
- Jaishankar, M., Tseten, T., Anbalagan, N., Mathew, B. B., & Beeregowda, K. N. (2014). Toxicity, mechanism and health effects of some heavy metals. *Interdisciplinary toxicology*, 7(2), 60-72.
- Jiang, G. B., Shi, J. B., & Feng, X. B. (2006). Mercury pollution in China. *Environmental science and technology*. 40 (12), 3673-3578.

- Jo, H. Y., Park, G. J., Bok, K. H., Park, K.-M., Chang, P.-S., & Kim, C. (2015). An asymmetric naked-eye chemo-sensor for Cu²⁺ in aqueous solution. *Inorganic Chemistry Communications*, 51, 90–94.
- Jonsson, S., Andersson, A., Nilsson, M. B., Skyllberg, U., Lundberg, E., Schaefer, J. K., & Björn, E. (2017). Terrestrial discharges mediate trophic shifts and enhance methylmercury accumulation in estuarine biota. *Science Advances*, 3(1), e1601239.
- Kamaruzzaman, B. Y., Rina, Z., John, B. A., & Jalal, K. C. A. (2011). Heavy metal accumulation in commercially important fishes of South West Malaysian coast. *Research journal of environmental sciences*, 5(6), 595.
- Karagas, M. R., Choi, A. L., Oken, E., Horvat, M., Schoeny, R., Kamai, E., & Korrick, S. (2012). Evidence on the human health effects of low-level methylmercury exposure. *Environmental Health Perspectives*, 120(6), 799.
- Kaur, H., Kumar, S., & Verma, N. (2014). Enzyme-based colorimetric and potentiometric biosensor for detecting Pb (II) ions in milk. *Brazilian Archives of Biology and Technology*, 57(4), 613-619.
- Kim, H. Y., Lee, H. J., & Chang, S.-K. (2015). Reaction-based colorimetric signaling of Cu²⁺ ions by oxidative coupling of phenols with 4-aminoantipyrine. *Talanta*, 132, 625–629.
- Kim, H. N., Lee, M. H., Kim, H. J., Kim, J. S., & Yoon, J. (2008). A new trend in rhodamine-based chemosensors: application of spirolactam ring-opening to sensing ions. *Chemical society reviews*, 37(8), 1465-1472.
- King, J. K., Kostka, J. E., Frischer, M. E., & Saunders, F. M. (2000). Sulfate-reducing bacteria methylate mercury at variable rates in pure culture and in marine sediments. *Applied and environmental microbiology*, 66(6), 2430-2437.
- Knecht, M. R., & Sethi, M. (2009) Bio-inspired colorimetric detection of Hg²⁺ and Pb²⁺ heavy metal ions using Au nanoparticles. *Analytical and bioanalytical chemistry*, 394(1), 33-46.
- Ko, S. K., Yang, Y. K., Tae, J., & Shin, I. (2006). In vivo monitoring of mercury ions using a rhodamine-based molecular probe. *Journal of the American Chemical Society*, 128(43), 14150-14155.
- Lamborg, C. H., Hammerschmidt, C. R., Bowman, K. L., Swarr, G. J., Munson, K. M., Ohnemus, D. C., & Saito, M. A. (2014). A global ocean inventory of anthropogenic mercury based on water column measurements. *Nature*, 512(7512), 65-68.
- Law, A. T., & Singh, A. (1991). Relationships between heavy metal content and body weight of fish from the Kelang estuary, Malaysia. *Marine Pollution Bulletin*, 22(2), 86-89.

- Li, H., Toh, P. Z., Tan, J. Y., Zin, M. T., Lee, C. Y., Li, B., & Kang, L. (2016). Selected Biomarkers Revealed Potential Skin Toxicity Caused by Certain Copper Compounds. *Scientific reports*, 6, 37664.
- Li, P., Feng, X. B., Qiu, G. L., Shang, L. H., & Li, Z. G. (2009). Mercury pollution in Asia: a review of the contaminated sites. *Journal of hazardous materials*, 168(2), 591-601.
- Lidsky, T. I., & Schneider, J. S. (2003). Lead neurotoxicity in children: basic mechanisms and clinical correlates. *Brain*, 126(1), 5-19.
- Liu, H. W., Jiang, S. J., & Liu, S. H. (1999). Determination of cadmium, mercury and lead in seawater by electrothermal vaporization isotope dilution inductively coupled plasma mass spectrometry. *Spectrochimica Acta Part B: Atomic Spectroscopy*, 54(9), 1367-1375.
- Liu, J., & Lu, Y. (2003). A colorimetric lead biosensor using DNAzyme-directed assembly of gold nanoparticles. *Journal of the American Chemical Society*, 125(22), 6642-6643.
- Liu, D., Qu, W., Chen, W., Zhang, W., Wang, Z., & Jiang, X. (2010). Highly sensitive, colorimetric detection of mercury (II) in aqueous media by quaternary ammonium group-capped gold nanoparticles at room temperature. *Analytical chemistry*, 82(23), 9606-9610.
- Liu, J., Wu, D., Yan, X., & Guan, Y. (2013). Naked-eye sensor for rapid determination of mercury ion. *Talanta*, 116, 563-8.
- Liu, Q., Cai, H., Xu, Y., Xiao, L., Yang, M., & Wang, P. (2007). Detection of heavy metal toxicity using cardiac cell-based biosensor. *Biosensors and Bioelectronics*, 22(12), 3224-3229.
- Liu, Y., Chang, X., Yang, D., Guo, Y., & Meng, S. (2005). Highly selective determination of inorganic mercury (II) after preconcentration with Hg (II)-imprinted diazoaminobenzene-vinylpyridine copolymers. *Analytica Chimica Acta*, 538(1-2), 85-91.
- Long, F., Zhu, A., Shi, H., Wang, H., & Liu, J. (2013). Rapid on-site/in-situ detection of heavy metal ions in environmental water using a structure-switching DNA optical biosensor. *Scientific reports*, 3, 2308.
- López, F. A., Alguacil, F. J., Roman, C. P., Tayibi, H., & López-Delgado, A. (2008). Disposal of elemental mercury via sulphur reaction by milling. In *Proceeding of the First International Conference on "Hazardous Waste Management"*, Crete, Greece (pp. 479-80).
- Malik, R. N., Hashmi, M. Z., & Huma, Y. (2014). Heavy metal accumulation in edible fish species from Rawal Lake Reservoir, Pakistan. *Environmental Science and Pollution Research*, 21(2), 1188-1196.

- Maquieira, A., Elmahadi, H. A., & Puchades, R. (1994). Immobilized cyanobacteria for online trace metal enrichment by flow injection atomic absorption spectrometry. *Analytical Chemistry*, 66(21), 3632-3638.
- Marcus, S. A., Sidiropoulos, S. W., Steinberg, H., & Talaat, A. M. (2016). CsoR is essential for maintaining copper homeostasis in Mycobacterium tuberculosis. *PloS one*, 11(3).
- Maruyama, K., Yorifuji, T., Tsuda, T., Sekikawa, T., Nakadaira, H., & Saito, H. (2012). Methyl mercury exposure at Niigata, Japan: results of neurological examinations of 103 adults. *BioMed Research International*, 2012.
- Mehmood, T., Chaudhry, M. M., Tufail, M., & Irfan, N. (2009). Heavy metal pollution from phosphate rock used for the production of fertilizer in Pakistan. *Microchemical Journal*, 91(1), 94-99.
- Melnick, J. G., & Parkin, G. (2007). Cleaving mercury-alkyl bonds: a functional model for mercury detoxification by MerB. *Science*, 317(5835), 225-227.
- Mishra, A. K. (Ed.). (2016). Dendrimers and mesoporous materials for heavy metal removal from aqueous solution. *Nanomaterials for Water Remediation: Carbon-Based Materials*, 1, 115-116.
- Mortvedt, J. J. (1996). Heavy metal contaminants in inorganic and organic fertilizers. In *Fertilizers and Environment* (pp. 5-11).
- Moyo, M. (2014). Horseradish Peroxidase Biosensor to Detect Zinc Ions in Aqueous Solutions. *Open Journal of Applied Biosensor*, 3(2), 1-7.
- Mutter, J., & Yeter, D. (2008). Kawasaki's disease, acrodynia, and mercury. *Current medicinal chemistry*, 15(28), 3000-3010.
- Nagajyoti, P. C., Lee, K. D., & Sreekanth, T. V. M. (2010). Heavy metals, occurrence and toxicity for plants: a review. *Environmental Chemistry Letters*, 8(3), 199-216.
- Nakata, H., Nakayama, S. M., Oroszlany, B., Ikenaka, Y., Mizukawa, H., Tanaka, K., & Saengtienchai, A. (2017). Monitoring lead (Pb) pollution and identifying Pb pollution sources in Japan using stable Pb isotope analysis with kidneys of wild rats. *International journal of environmental research and public health*, 14(1), 56.
- Navarro, R. R., Sumi, K., Fujii, N., & Matsumura, M. (1996). Mercury removal from wastewater using porous cellulose carrier modified with polyethyleneimine. *Water Research*, 30(10), 2488-2494.
- Niu, Y., Qu, R., Chen, H., Mu, L., Liu, X., Wang, T., & Sun, C. (2014). Synthesis of silica gel supported salicylaldehyde modified PAMAM dendrimers for the effective removal of Hg (II) from aqueous solution. *Journal of hazardous materials*, 278, 267-278.

- Oh, E., Delehanty, J. B., Sapsford, K. E., Susumu, K., Goswami, R., Blanco-Canosa, J. B., & Goering, P. L. (2011). Cellular uptake and fate of PEGylated gold nanoparticles is dependent on both cell-penetration peptides and particle size. *Acs Nano*, 5(8), 6434-6448.
- Omichinski, J. G. (2007). Toward methylmercury bioremediation. *Science*, 317(5835), 205-206.
- Orisakwe, O. E., Asomugha, R., Afonne, O. J., Anisi, C. N., Obi, E., & Dioka, C. E. (2004). Impact of effluents from a car battery manufacturing plant in Nigeria on water, soil, and food qualities. *Archives of Environmental Health: An International Journal*, 59(1), 31-36.
- Othman, A. B., Lee, J. W., Wu, J. S., Kim, J. S., Abidi, R., Thuéry, P., ... & Vicens, J. (2007). Calix [4] arene-based, Hg²⁺-induced intramolecular fluorescence resonance energy transfer chemosensor. *The Journal of organic chemistry*, 72(20), 7634-7640.
- Pacyna, J. M. (1987). Atmospheric Emissions of Arsenic, Cadmium, Lead and Mercury from High Temperature Processes in Power Generation and Industry. *John. Lead, Mercury, Cadmium and Arsenic in Environment*, 7, 69-8.
- Poirier, L., Nelson, J., Gilleland, G., Wall, S., Berhane, L., & Lopez-Linares, F. (2017). Comparison of Preparation Methods for the Determination of Metals in Petroleum Fractions (1000° F+) by Microwave Plasma Atomic Emission Spectroscopy. *Energy & Fuels*, 31(8), 7809-7815.
- Pirrone, N., Cinnirella, S., Feng, X., Finkelman, R. B., Friedli, H. R., Leaner, J., & Telmer, K. (2010). Global mercury emissions to the atmosphere from anthropogenic and natural sources. *Atmospheric Chemistry and Physics*, 10(13), 5951-5964.
- Prashanth, L., Kattapagari, K. K., Chitturi, R. T., Baddam, V. R. R., & Prasad, L. K. (2015). A review on role of essential trace elements in health and disease. *Journal of Dr. NTR University of Health Sciences*, 4(2), 75.
- Praveena, S. M., de Burbure, C., Aris, A. Z., & Hashim, Z. (2013). Mini review of mercury contamination in environment and human with an emphasis on Malaysia: status and needs. *Reviews on environmental health*, 28(4), 195-202.
- Priyadarshini, E., & Pradhan, N. (2017). Gold nanoparticles as efficient sensors in colorimetric detection of toxic metal ions: A review. *Sensors and Actuators B: Chemical*, 238, 888-902.
- Refaie, F. M., Esmat, A. Y., Mohamed, A. F., & Nour, W. H. A. (2009). Effect of chromium supplementation on the diabetes induced-oxidative stress in liver and brain of adult rats. *Biometals*, 22(6), 1075.

- Ratnarathorn, N., Chailapakul, O., & Dungchai, W. (2015). Highly sensitive colorimetric detection of lead using maleic acid functionalized gold nanoparticles. *Talanta*, 132, 613-618.
- Ravindran, A., Mani, V., Chandrasekaran, N., & Mukherjee, A. (2011). Selective colorimetric sensing of cysteine in aqueous solutions using silver nanoparticles in the presence of Cr³⁺. *Talanta*, 85(1), 533-540.
- Rether, A., & Schuster, M. (2003). Selective separation and recovery of heavy metal ions using water-soluble N-benzoylthiourea modified PAMAM polymers. *Reactive and Functional Polymers*, 57(1), 13-21.
- Rezai, H., Yusoff, F. M., & Yap, C. K. (2003). Mercury in zooplankton from the Malacca Straits, 32(3), 240-243
- Ripp, S., DiClaudio, M. L., & Sayler, G. S. (2010). Biosensors as environmental monitors. *Environmental Microbiology*, 2, 213-233.
- Ríos, S. E. G., Peñuela, G. A., & Botero, C. M. R. (2017). Method Validation for the Determination of Mercury, Cadmium, Lead, Arsenic, Copper, Iron, and Zinc in Fish Through Microwave-Induced Plasma Optical Emission Spectrometry (MIP OES). *Food Analytical Methods*, 10(10), 1-8.
- Sakamoto, H., Ichikawa, T., Tomiyasu, T., & SATO, M. (2004). Mercury concentration in environmental samples of Malaysia. *鹿児島大学理学部紀要* 37, 83-90.
- Saha, A., Karnik, A., Sathawara, N., Kulkarni, P., & Singh, V. (2008). Ceruloplasmin as a marker of occupational copper exposure. *Journal of Exposure Science and Environmental Epidemiology*, 18(3), 332-337.
- Sasamoto, T., Otani, H., Hirayama, I., Hayashi, M., Baba, I., Iida, K., & Osugi, N. (2016). Dietary Exposure to Dioxins, Polychlorinated Biphenyls, and Heavy Metals in the Tokyo Metropolitan Area from 1999 to 2014. In *Persistent Organic Chemicals in the Environment: Status and Trends in the Pacific Basin Countries I Contamination Status*, 4, 85-106.
- Selin, N. E. (2009). Global biogeochemical cycling of mercury: a review. *Annual Review of Environment and Resources*, 34, 43-63.
- Shallari, S., Schwartz, C., Hasko, A., & Morel, J. L. (1998). Heavy metals in soils and plants of serpentine and industrial sites of Albania. *Science of the total environment*, 209(2-3), 133-142.
- Sharaf, E. F., & Alharbi, E. (2013). Removal of heavy metals from waste water of tanning leather industry by fungal species isolated from polluted soil, 12(27), 4351-4355.

- Shellaiah, M., Simon, T., Sun, K. W., & Ko, F. H. (2016). Simple bare gold nanoparticles for rapid colorimetric detection of Cr³⁺ ions in aqueous medium with real sample applications. *Sensors and Actuators B: Chemical*, 226, 44-51.
- Shih, T. T., Hsieh, C. C., Luo, Y. T., Su, Y. A., Chen, P. H., Chuang, Y. C., & Sun, Y. C. (2016). A high-throughput solid-phase extraction microchip combined with inductively coupled plasma-mass spectrometry for rapid determination of trace heavy metals in natural water. *Analytica chimica acta*, 916, 24-32.
- Shu, Q., Liu, M., Ouyang, H., & Fu, Z. (2017). Label-free fluorescent immunoassay for Cu²⁺ ion detection based on UV degradation of immunocomplex and metal ion chelates. *Nanoscale*, 9(34), 12302-12306.
- Siddiqui, M. A., Alhadlaq, H. A., Ahmad, J., Al-Khedhairi, A. A., Musarrat, J., & Ahamed, M. (2013). Copper oxide nanoparticles induced mitochondria mediated apoptosis in human hepatocarcinoma cells. *PLoS One*, 8(8).
- Simões, M. R., Preti, S. C., Azevedo, B. F., Fiorim, J., Freire, D. D., Covre, E. P., ... & dos Santos, L. (2017). Low-level chronic lead exposure impairs neural control of blood pressure and heart rate in rats. *Cardiovascular toxicology*, 17(2), 190-199.
- Spiegel, S. J., & Veiga, M. M. (2010). International guidelines on mercury management in small-scale gold mining. *Journal of Cleaner Production*, 18(4), 375-385.
- Srivastava, N. K., & Majumder, C. B. (2008). Novel biofiltration methods for the treatment of heavy metals from industrial wastewater. *Journal of Hazardous Materials*, 151(1), 1-8.
- Strydom, C., Robinson, C., Pretorius, E., Whitcutt, J. M., Marx, J., & Bornman, M. S. (2006). The effect of selected metals on the central metabolic pathways in biology: a review. *Water SA*, 32(4), 543-554.
- Su, H., Fan, H., Ai, S., Wu, N., Fan, H., Bian, P., & Liu, J. (2011). Selective determination of melamine in milk samples using 3-mercapto-1-propanesulfonate-modified gold nanoparticles as colorimetric probe. *Talanta*, 85(3), 1338-1343.
- Sudibya, H. G., He, Q., Zhang, H., & Chen, P. (2011). Electrical detection of metal ions using field-effect transistors based on micropatterned reduced graphene oxide films. *ACS nano*, 5(3), 1990.
- Sugunan, A., Thanachayanont, C., Dutta, J., & Hilborn, J. G. (2005). Heavy-metal ion sensors using chitosan-capped gold nanoparticles. *Science and Technology of Advanced Materials*, 6(3), 335-340.
- Sungur, Ş., & Gülmez, F. (2015). Determination of metal contents of various fibers used in textile industry by MP-AES. *Journal of Spectroscopy*, 2015.

- Tan, J., Liu, R., Wang, W., Liu, W., Tian, Y., Wu, M., & Huang, Y. (2009). Controllable aggregation and reversible pH sensitivity of AuNPs regulated by carboxymethyl cellulose. *Langmuir*, 26(3), 2093-2098.
- Tanaka, Y., Oda, S., Yamaguchi, H., Kondo, Y., Kojima, C., & Ono, A. (2007). ¹⁵N-¹⁵N J-coupling across HgII: Direct observation of HgII-mediated TT base pairs in a DNA duplex. *Journal of the American Chemical Society*, 129(2), 244-245.
- Toscano, C. M., Simões, M. R., Alonso, M. J., Salices, M., Vassallo, D. V., & Fioresi, M. (2017). Sub-chronic lead exposure produces β 1-adrenoceptor downregulation decreasing arterial pressure reactivity in rats. *Life Sciences*, 180, 93-101.
- Tomalia, D. A., Baker, H., Dewald, J., Hall, M., Kallos, G., Martin, S., & Smith, P. (1985). A new class of polymers: starburst-dendritic macromolecules. *Polymer Journal*, 17(1), 117.
- Tomalia, D.A. Dendrimer Research. *Science* 1991, 252, 1231.
- Turner, A. P. (2013). Biosensors: sense and sensibility. *Chemical Society Reviews*, 42(8), 3184-3196.
- Wang, G. L., Zhu, X. Y., Jiao, H. J., Dong, Y. M., & Li, Z. J. (2012). Ultrasensitive and dual functional colorimetric sensors for mercury (II) ions and hydrogen peroxide based on catalytic reduction property of silver nanoparticles. *Biosensors and Bioelectronics*, 31(1), 337-342.
- Wang, Z., & Ma, L. (2009). Gold nanoparticle probes. *Coordination Chemistry Reviews*, 253(11), 1607-1618.
- Wang, Guang-Li, et al. "Ultrasensitive and dual functional colorimetric sensors for mercury (II) ions and hydrogen peroxide based on catalytic reduction property of silver nanoparticles." *Biosensors and Bioelectronics* 31.1 (2012): 337-342.
- Wei, H., Li, B., Li, J., Dong, S., & Wang, E. (2008). DNAzyme-based colorimetric sensing of lead (Pb²⁺) using unmodified gold nanoparticle probes. *Nanotechnology*, 19(9), 095501.
- Wei, Q., Nagi, R., Sadeghi, K., Feng, S., Yan, E., Ki, S. J., & Ozcan, A. (2014). Detection and spatial mapping of mercury contamination in water samples using a smart-phone. *ACS nano*, 8(2), 1121-1129.
- Wu, X., Cobbina, S. J., Mao, G., Xu, H., Zhang, Z., & Yang, L. (2016). A review of toxicity and mechanisms of individual and mixtures of heavy metals in the environment. *Environmental Science and Pollution Research*, 23(9), 8244-8259.

- Wuana, R. A., & Okieimen, F. E. (2011). Heavy metals in contaminated soils: a review of sources, chemistry, risks and best available strategies for remediation. *Isrn Ecology*, 2011.
- Wylie, D. E., Lu, D., Carlson, L. D., Carlson, R., Babacan, K. F., Schuster, S. M., & Wagner, F. W. (1992). Monoclonal antibodies specific for mercuric ions. *Proceedings of the National Academy of Sciences*, 89(9), 4104-4108.
- Xia, N., Shi, Y., Zhang, R., Zhao, F., Liu, F., & Liu, L. (2012). Simple, rapid and label-free colorimetric assay for arsenic based on unmodified gold nanoparticles and a phytochelatin-like peptide. *Analytical Methods*, 4(12), 3937-3941.
- Xu, H., Zeng, L., Xing, S., Shi, G., Xian, Y., & Jin, L. (2008). Microwave-radiated synthesis of gold nanoparticles/carbon nanotubes composites and its application to voltammetric detection of trace mercury (II). *Electrochemistry Communications*, 12(10), 1839-1843.
- Xu, R. X., Yu, X. Y., Gao, C., Jiang, Y. J., Han, D. D., Liu, J. H., & Huang, X. J. (2013). Non-conductive nanomaterial enhanced electrochemical response in stripping voltammetry: the use of nanostructured magnesium silicate hollow spheres for heavy metal ions detection. *Analytica chimica acta*, 790, 31-38.
- Yamakawa, A., Moriya, K., & Yoshinaga, J. (2017). Determination of isotopic composition of atmospheric mercury in urban-industrial and coastal regions of Chiba, Japan, using cold vapor multicollector inductively coupled plasma mass spectrometry. *Chemical Geology*, 448, 84-92
- Yang, H., Zhou, Z., Huang, K., Yu, M., Li, F., Yi, T., & Huang, C. (2007). Multisignaling optical-electrochemical sensor for Hg²⁺ based on a rhodamine derivative with a ferrocene unit. *Organic letters*, 9(23), 4729-4732.
- Yang, Z., Liu, Z. W., Allaker, R. P., Reip, P., Oxford, J., Ahmad, Z., & Reng, G. (2013). A review of nanoparticle functionality and toxicity on the central nervous system. In *Nanotechnology, the Brain, and the Future* (pp. 313-332). Springer Netherlands.
- Yoon, S., Albers, A. E., Wong, A. P., & Chang, C. J. (2005). Screening mercury levels in fish with a selective fluorescent chemosensor. *Journal of the American Chemical Society*, 127(46), 16030-16031.
- Yu, C. J., & Tseng, W. L. (2008). Colorimetric detection of mercury (II) in a high-salinity solution using gold nanoparticles capped with 3-mercaptopropionate acid and adenosine monophosphate. *Langmuir*, 24(21), 12717-12722.
- Yuan, S., Peng, D., Song, D., & Gong, J. (2013). Layered titanate nanosheets as an enhanced sensing platform for ultrasensitive stripping voltammetric detection of mercury (II). *Sensors and Actuators B: Chemical*, 181, 432-438.

- Zhai, J., Yong, D., Li, J., & Dong, S. (2013). A novel colorimetric biosensor for monitoring and detecting acute toxicity in water. *Analyst*, 138(2), 702-707.
- Zhang, X., Yang, L., Li, Y., Li, H., Wang, W., & Ye, B. (2012). Impacts of lead/zinc mining and smelting on the environment and human health in China. *Environmental monitoring and assessment*, 184(4), 2261-2273.
- Zhao, Q., Wang, Y., Cao, Y., Chen, A., Ren, M., Ge, Y., & Ruan, L. (2014). Potential health risks of heavy metals in cultivated topsoil and grain, including correlations with human primary liver, lung and gastric cancer, in Anhui province, Eastern China. *Science of the Total Environment*, 470, 340-347.
- Zhao, W., Brook, M. A., & Li, Y. (2008). Design of gold nanoparticle-based colorimetric biosensing assays. *ChemBioChem*, 9(15), 2363-2371.
- Zhong, G., Liu, J., & Liu, X. (2015). A Fast Colourimetric Assay for Lead Detection Using Label-Free Gold Nanoparticles (AuNPs). *Micromachines*, 6(4), 462-472.
- Zhou, Y., Wang, S., Zhang, K., & Jiang, X. (2008). Visual Detection of Copper (II) by Azide-and Alkyne-Functionalized Gold Nanoparticles Using Click Chemistry. *Angewandte Chemie*, 120(39), 7564-7566.
- Zhu, X., & Alexandratos, S. D. (2005). Polystyrene-supported amines: affinity for mercury (II) as a function of the pendant groups and the Hg (II) counterion. *Industrial & engineering chemistry research*, 44(23), 8605-8610.