

1 **Potential use of composts and vermicomposts as low-cost adsorbents**
2 **for dye removal: an overlooked application**

3 Remigio Paradelo¹, Xanel Vecino², Ana Belén Moldes³, María Teresa Barral¹

4 ¹ Universidade de Santiago de Compostela, Departamento de Edafoloxía e Química
5 Agrícola, Facultade de Farmacia, Praza Seminario de Estudos Galegos s/n, 15782
6 Santiago de Compostela (Spain)

7 ² Polytechnic University of Catalonia (UPC)-Barcelona TECH, Chemical Engineering
8 Department, Barcelona East School of Engineering (EEBE); Barcelona Research Center
9 for Multiscale Science and Engineering, Campus Diagonal-Besòs, 08930 Barcelona
10 (Spain)

11 ³ University of Vigo, Chemical Engineering Department, School of Industrial
12 Engineering – Módulo Tecnológico Industrial (MTI), Campus As Lagoas-Marcosende,
13 36310 Vigo (Spain)

14
15 **Corresponding author:** Remigio Paradelo Núñez. Phone: +34 881 815 042. E-mail:
16 remigio.paradelo.nunez@usc.es

17 ORCID: 0000-0002-4165-177X

18

19

20 **Acknowledgements**

21 Dr R. Paradelo and Dr X. Vecino are grateful to the Spanish Ministry of Economy and
22 Competitiveness (MINECO) for award of a *Ramón y Cajal* fellowship (RYC-2016-
23 19286) and a *Juan de la Cierva* contract (ref. IJCI-2016-27445), respectively.

24

25

1 **Summary**

2 The use of composts and vermicomposts as adsorbents is an important topic of study in
3 the field of environmental remediation. These materials are rich in organic matter and
4 have functional groups that can interact with organic and inorganic compounds. They
5 also contain microorganisms that can promote biodegradation of organic substances.
6 Composts that cannot be used for agronomic purposes (owing to e.g. low nutrient levels
7 or phytotoxicity) may be valuable for soil remediation or pollutant removal. In this
8 review paper, we discuss other papers on this topic, with the objective of drawing
9 attention to the potential use of composts and vermicomposts and to recommend further
10 investigation on this subject. Few published studies have investigated the use of
11 vermicomposts to remove dyes and other coloured compounds. However, preliminary
12 results show that these materials are potentially good adsorbents, particularly for basic
13 dyes. However, there remain several uncertainties regarding this application. For
14 example, very few dyes have been studied so far, and little is known about the influence
15 of the properties of composts/vermicomposts on the dye removal process. Moreover, the
16 possible use of vermicompost to enhance biodegradation processes has not been
17 explored. All of these questions should be addressed in future research.

18 **Keywords:** Organic waste; Waste management; Waste water; Coloured compounds;
19 Adsorption.

20

21

22 **1. Need to treat dye-contaminated effluents**

23 The extensive use of dyes in many industries, such as the dyestuffs, textile, paper and
24 plastic industries, has led to the production of huge amounts of coloured waste water
25 (Crini 2006). In general, dyes are not readily degraded under the aerobic conditions of

1 biological treatment plants, and the effluents discharged from these plants are usually
2 coloured. The presence of dyes in waste water, even in small amounts, is a major
3 problem for these industries, as colours and dyes are inherently highly visible, which
4 greatly affects the perception of water quality by the public (Slokar and Majcen Le
5 Marechal 1997; Crini 2006). In addition, dye compounds can interfere with the growth
6 of aquatic organisms through absorption and reflection of sunlight (Slokar and Majcen
7 Le Marechal 1997), and some azo dyes are also suspected of being
8 mutagenic/carcinogenic and toxic to aquatic life (Gottlieb et al. 2003). The removal of
9 dyes from waste water is a challenging task because of the inherent properties of these
10 molecules, including colour fastness, stability and resistance to degradation (Sun and
11 Yang 2003). Among the different types of dyes, reactive and acid dyes are the most
12 problematical, as they usually pass unaffected through conventional treatment systems
13 (Willmott et al. 1998; McKay et al. 2011).

14 Dye-containing effluents are currently treated in two ways: by chemical or physical
15 methods of dye removal, or by biodegradation (Slokar and Majcen Le Marechal 1997).
16 Among the physical and chemical techniques used for dye removal, adsorption is the
17 most common procedure and has been shown to produce the best results for different
18 types of dyes (Ho and McKay 2003; Jain et al. 2003). Activated carbon adsorption is
19 one of the best available control technologies in this respect, according to the US
20 Environmental Protection Agency (Derbyshire et al. 2001), although its use for the
21 treatment of high volumes of waste water is restricted by its high cost. Alternative
22 adsorbents must be found in order to reduce the cost of the treatments, and there is an
23 abundance of literature on the use of inexpensive materials for the removal of dyes from
24 waste water, with several review articles dealing specifically with this subject (Table 1).
25 An adsorbent can be considered low-cost if it requires little processing, is abundant in

1 nature or is a by-product or waste material from another industry (Bailey et al. 1999). In
2 addition, it should ideally be efficient for the removal of a wide variety of dyes, have a
3 high adsorption capacity and be tolerant to a wide range of physicochemical parameters
4 of waste water (Crini 2006). Extensive reviews of low-cost adsorbents proposed for dye
5 removal have been provided, for numerous materials, including agricultural waste,
6 industrial waste products, biomolecules such as chitosan, peats, and more (e.g. Crini
7 2006) (Table 1). In contrast to the large amount of research on the use of other types of
8 waste material as low-cost adsorbents for dye removal, existing research on the use of
9 composts and vermicomposts for this application is relatively scarce, despite the
10 potential advantages of using these materials.

11

12 **2. Properties of composts and vermicomposts**

13 Composts and vermicomposts are produced as a result of the transformation of different
14 types of organic waste. Composting is the biological decomposition of the organic
15 matter contained in wastes by mesophilic and thermophilic microorganisms, leading to
16 stabilized organic amendments with fertilization potential (de Bertoldi et al. 1983).
17 Vermicomposting is used to valorize organic waste, and the process relies on the action
18 of epigeic earthworm species to accelerate biodegradation and stabilization processes
19 (Gómez-Brandón and Domínguez 2014). Urban waste was initially the most common
20 type of feedstock used for composting, including municipal solid waste and sewage
21 sludge, in response to the needs of waste management in growing cities. Extension of
22 composting and vermicomposting to the treatment of materials of agricultural and
23 industrial origin has resulted in the worldwide use of these processes. Composting and
24 vermicomposting are now routinely used in the treatment of municipal solid waste
25 (Farrell and Jones 2009) and sewage sludge (U.S. EPA 2002), animal residues such as

1 solid and liquid manures (Larney et al. 2006), agricultural waste (Mortier et al. 2016),
2 food-processing waste such as winery or olive oil mill waste (Cegarra and Paredes
3 2008), and industrial waste, including textile, papermill and tannery waste (Bhat et al.
4 2018).

5 In addition to their use as waste management treatments, composting and
6 vermicomposting also yield valuable products. Composts and vermicomposts are
7 typically stable, safe products, with neutral or slightly alkaline pH. They are rich in
8 organic matter (often more than 50% of weight) and, as a consequence, they are highly
9 porous and have a low bulk density and a high water-holding capacity. They are also
10 characterized by high concentrations of plant nutrients (N, P, K) in different forms and
11 by large amounts and varieties of microbial communities. Thus, the composts and
12 vermicomposts produced by these processes have many properties that make them
13 valuable resources in several fields.

14

15 **3. The use of composts and vermicomposts as adsorbents**

16 Composts and vermicomposts have several applications, including agronomic uses as
17 organic amendments in agricultural soils (Hargreaves et al. 2008; Diacono and
18 Montemurro 2010), as components in soil-less horticultural substrates (Carmona and
19 Abad 2008; Paradelo et al. 2012, 2019), and for the remediation of degraded or polluted
20 soils (Semple et al. 2001; Paradelo et al. 2007, 2009a,b, 2011; Park et al. 2011; Huang
21 et al. 2016). In addition, the use of composts and vermicomposts as adsorbents for the
22 treating polluted waters is a promising field of study. Composts and vermicomposts can
23 be produced inexpensively from by-products and waste materials, and their elaboration
24 usually requires little processing. Compared with the cost of activated carbon (> 300 \$

1 m⁻³), composts can typically be produced from urban waste at much lower cost (around
2 20 \$ m⁻³).

3 Composts and vermicomposts are rich in organic matter with functional groups
4 (polar ionisable and non-ionisable groups, non-polar aromatic and aliphatic groups) that
5 can interact with neutral, cationic and anionic compounds and provide a high adsorption
6 capacity for a wide range of organic and inorganic substances. They are also rich in
7 microorganisms that can promote the biodegradation of organic compounds. These
8 characteristics make composts and vermicomposts of potential value for the treatment
9 of waters polluted with a range of substances, including metallic elements and
10 pesticides and organic compounds such as dyes. This represents an excellent alternative
11 use for several types of composts that are not suitable for agronomic purposes because
12 of e.g. phytotoxicity or low nutrient contents.

13 As a result of these characteristics, composted and vermicomposted materials have
14 frequently been studied as potential biosorbents for the removal of pesticides and
15 inorganic pollutants from water (Boni and Scaffoni 2009; Kocasoy and Guvener 2009;
16 Paradelo and Barral 2012; Carrillo-Zenteno et al. 2013; Barral et al. 2014; Singh and
17 Kaur 2015; Cancelo-González et al. 2017; He et al. 2017), and many works have
18 addressed the topic (Table 2). By contrast, few studies have considered the use of these
19 products for dye removal, and research is still at a preliminary stage.

20

21 **4. Studies on the use of compost and vermicomposts for dye removal**

22 After running a Scopus search with the terms “dye” and “adsorption” or “removal” and
23 “compost” or “vermicompost”, we found the 13 papers listed in Table 3. Most existing
24 studies concern pure dye solutions (20 different molecules), and among these,
25 methylene blue and malachite green (also known as Basic Green 4) are the most

1 commonly studied dyes, with almost all studied only once. In addition, some studies
2 have investigated coloured effluents from the wine industry (Paradelo et al. 2009;
3 Pérez-Ameneiro et al. 2014, 2015), which are distinctly red and similar to the colour of
4 commercial amaranth dye (Pérez-Ameneiro et al. 2014). Figure 1 shows the chemical
5 structures of the dyes studied. They are all ionisable organic molecules with aromatic
6 structures, so they can interact with the functional groups present in composts and
7 vermicomposts either through electrostatic interactions or through non polar dispersion
8 forces.

9 Regarding the chemical nature of the molecules in the research involving pure dye
10 solutions, most studies have investigated basic dyes, which are cationic molecules at
11 most pH values. Only three studies included more than one type of dye (Tsui et al. 2003;
12 Jozwiak et al. 2013; Toptas et al. 2014), of which only one compared four types of dyes
13 (Tsui et al. 2003). Anionic dyes, including acid dyes, direct dyes (molecules with high
14 affinity for fibre) and reactive dyes (molecules that react with fibre) have been much
15 less well studied than basic dyes. This precedence is probably due to the fact that
16 composted materials have higher retention capacity for cations than for anions, which
17 should result in a better performance for the removal of basic dyes relative to anionic
18 dyes.

19 Approaches to the treatment of dye-contaminated waters also vary widely: some
20 studies have included detailed adsorption studies (Kyziol-Komosińska et al. 2010;
21 McKay et al. 2011; Jozwiak et al. 2013; Toptas et al. 2014; Bhagavathi et al. 2015;
22 Anastopoulos et al. 2018), whereas others have only investigated dye removal without
23 considering the adsorption process (Tsui et al. 2003; Bhagavathi et al. 2016a,b;
24 Anastopoulos et al. 2018). In addition to studies focused on eliminating dyes from water

1 by contact with an adsorbent, one of the studies addresses bioremediation during
2 composting (Dey et al. 2017).

3

4 4.1 Dye removal capacity and nature of the removal process

5 Although simultaneous comparison of the removal of different types of dyes by
6 composts and vermicomposts is not common, basic dyes are consistently more
7 efficiently removed than direct, reactive or acid dyes. This has been observed by Tsui et
8 al. (2003), Jozwiak et al. (2013) and Toptas et al. (2014) after comparison of the
9 adsorption of several dyes on the same composts. Overall, the findings of all the
10 research reviewed here sustains this observation (Table 4, Figure 1). As composts are
11 mainly negatively charged, due to the presence of functional groups –COOH and –OH,
12 stronger interaction will take place with positively charged compounds (cationic dyes),
13 as a result of electrostatic attraction, than with anionic compounds, as a consequence of
14 electrostatic repulsion.

15 Regarding the nature of the process, some studies have shown that removal takes
16 place by adsorption mechanisms. In most studies, the adsorption process follows the
17 Langmuir model (Kyziol-Komosińska et al. 2010; McKay et al. 2011; Jozwiak et al.
18 2013; Toptas et al. 2014; Bhagavathi et al. 2015; Anastopoulos et al. 2018), which
19 assumes that adsorption takes place on a homogeneous surface by monolayer adsorption,
20 with no significant interaction between the adsorbed molecules. This model implies that
21 retention reaches a maximum due to the limited number of sites available for adsorption.

22 Regarding kinetics, there is also some agreement among these studies that adsorption
23 of dyes follows a pseudo-second order model (McKay et al. 2011; Toptas et al. 2014;
24 Bhagavathi et al. 2015). This assumes a chemisorption mechanism in which the rate-
25 limiting step is the surface adsorption; it has commonly been observed that sorbent-

1 mediated removal of pollutants follows this model (Ho and MacKay 1999). There is
2 evidence that the removal process occurs rapidly: high removal percentages are
3 generally reached within short contact times, in some cases of minutes. De Godoi
4 Pereira et al. (2009) observed quantitative retention of crystal violet onto a
5 vermicompost after 10 minutes and of methylene blue after only one minute.
6 Anastopoulos et al (2018) also observed maximum adsorption of methylene blue after
7 one minute of contact. For other molecules, equilibrium times of 3-5 hours were
8 observed (Tsui et al. 2009; Jozwiak et al. 2013; Toptas et al. 2014). The faster
9 adsorption kinetics of methylene blue may be explained by steric hindrance, as
10 methylene blue molecules are smaller than in other dyes. This is important for the
11 treatment of waste water in continuous flow systems, in which high reaction rates imply
12 low contact times and higher flows, thereby increasing the capacity of water treatment
13 systems.

14 Regarding biodegradation as an alternative to adsorption for dye removal, Dey et al.
15 (2017) studied the removal of methylene blue from sugarcane bagasse during
16 vermicomposting with *Eisenia foetida*. These researchers observed that the combined
17 activities of earthworms and microbes led to the removal of 61% and 98% of methylene
18 blue after 30 and 60 days, respectively.

19

20 4.2 Factors in dye removal

21 Among the conditions that affect the process of dye removal, the pH of the solution
22 influences both the charge of the adsorbent and the ionic forms of the dye in solution.
23 As expected, the effect of pH has been shown to depend on the chemical nature of the
24 dye: the adsorption of anionic dyes increases at low pH values, with optimal values
25 around 2-3 for acid and reactive dyes (McKay et al. 2011; Jozwiak et al. 2013; Toptas et

1 al. 2014). In turn, adsorption of basic dyes increases with pH, with variable optimal
2 values always over 5 (Jozwiak et al. 2013; Toptas et al. 2014; Bhagavathi et al. 2015).
3 Basic dyes are generally best removed at neutral or slightly alkaline pH, whereas
4 anionic dyes are optimally removed at acid pH (Jozwiak et al. 2013; Toptas et al. 2014).
5 This occurs because anionic dyes generally have functional acid groups that are
6 negatively charged at pH values above the pKa, and neutral at lower pH. As composts
7 are also negatively charged, decreasing the pH will lead to a reduction in anion
8 repulsion and therefore to potentially higher adsorption.

9 The properties and composition of the compost or vermicompost will obviously also
10 have a strong influence on their capacity to adsorb dyes. Factors related to composition
11 include pH, cation and anion exchange capacity, organic matter content and nature, and
12 the specific surface area. In this respect, Kyziol-Komosińska et al. (2010) reported the
13 compost had a lower adsorption capacity than several peats, probably due to the lower
14 organic matter content and specific surface area and higher pH of the compost, as the
15 combination of both factors will reduce the capacity to adsorb for anionic dyes.
16 Bhagavathi et al. (2016a) compared the adsorption of crystal violet on four composts,
17 but sound conclusions cannot be reached regarding the influence of these factors on
18 adsorption, because of the narrow ranges of pH and OM content considered.

19 Regarding the role of the properties of compost or vermicomposts on their efficiency
20 as adsorbents for dye removal, the properties of the final composts are not the only
21 important factors. Physicochemical properties are modified during composting of
22 organic wastes, and the process of composting/vermicomposting itself may affect the
23 capacity of materials to remove dyes. Only one of the studies reviewed here compared
24 dye removal with composted and non-composted waste. Thus, Anastopoulos et al.
25 (2018) examined the effect of composting on the capacity of olive tree pruning waste to

1 adsorb methylene blue. These researchers found that the maximum adsorption capacity
2 was 250 mg g⁻¹ for composted material and 130 mg g⁻¹ for non-composted material,
3 indicating that composting greatly improved the adsorptive properties of pruning waste.
4 Similar findings may be obtained with other materials and dyes, as some researchers
5 have observed that composting can increase the capacity of waste to adsorb other
6 pollutants (Liu et al. 2018). The maturity of the compost may also affect its capacity for
7 dye removal, as the nature and composition of organic matter evolves during
8 composting. In this respect, Lashermes et al. (2010) observed that the maturity of
9 compost modifies its capacity to adsorb organic pollutants, and the same may be true for
10 organic dyes.

11

12 4.3. Application to coloured effluents

13 In addition to pure dye solutions, some studies have addressed the treatment of coloured
14 effluents from the winery industry, which are rich in natural colorants. The compounds
15 producing the colour of red vinasses include polyphenols, melanoidins produced by the
16 reaction of sugars and proteins with furfurals (Pant and Adhleya 2007). Pérez-Ameneiro
17 et al. (2014, 2015) used an adsorbent based on composted grape marc, immobilized in
18 calcium alginate beads, to remove pigments from vinasses. The immobilized composted
19 grape marc was able to eliminate between 95% and 100% of the pigments present in
20 winery effluents. As with pure dye solutions, the adsorption process followed a pseudo-
21 second order kinetic model (Pérez-Ameneiro et al. 2014); the adsorption equilibrium
22 process was described by the Freundlich isotherm (Pérez-Ameneiro et al. 2015).
23 Paradelo et al. (2009c) used several non-conventional low-cost natural adsorbents to
24 remove the colour from red wine vinasses. Among the materials assayed, grape marc
25 vermicompost produced the best results (80% colour removal), comparable to those

1 achieved with activated carbon. The good results obtained by the vermicompost in this
2 study may be partly related to its high organic matter concentration (91%). Nevertheless,
3 this was not the only factor, as composted pine bark also has a high organic matter
4 content (98%), but it did not yield a very high level of colour removal (below 15%).
5 The nature of organic matter is therefore expected to play an important role in the
6 performance of the adsorbent.

7

8 4.4 Possible modifications of biosorbents

9 Composts and vermicomposts may also represent a source of more efficient biosorbents
10 via modifications that increase the adsorption capacity of the original material. Washing
11 some composts before use has been recommended (Toptas et al. 2014) in order to
12 remove excessive amounts of soluble C, which can affect the dye removal process. In a
13 study with winery effluents, Paradelo et al. (2009c) observed that grinding or boiling the
14 biosorbents before treatment improved colour removal, as a consequence of the
15 increased surface area, in the first case, and temperature-mediate activation in the
16 second. On the other hand, there is some evidence that activated carbon or char
17 produced from compost could also be used for dye removal (Qian et al. 2008; Yang et al.
18 2016). Qian et al. (2008) studied the adsorption of methylene blue on activated carbon
19 prepared from composted cattle manure and found maximum adsorption capacity of
20 around 500 mg g⁻¹ (higher than those reported for other composts), due to the high
21 surface area and large mesopore volume. Yang et al. (2016) observed that increasing the
22 temperature of carbonization had contrasting effects on the adsorption capacity of a
23 vermicompost, increasing the adsorption of Congo red (a direct dye) and decreasing that
24 of methylene blue. Carbonisation of composts will probably increase the dye removal
25 capacity, at least for some molecules; however, unfortunately, neither of these studies

1 compared the performance of the original compost/vermicompost with that of the
2 charred products. Therefore, the available direct evidence is not sufficient, and further
3 studies should compare the adsorption capacity of charred and non-charred materials for
4 dyes, as done for other organic pollutants (Tsui and Juang 2010).

5

6 4.5 Limitations and future development

7 Overall, this review of existing studies show that there remain some uncertainties
8 regarding the potential application of composts and vermicomposts for dye removal.
9 Further research is therefore necessary in order to overcome these limitations. The
10 small number of studies, in addition to the variable experimental conditions (common in
11 adsorption studies), make it difficult to reach sound conclusions about this issue.
12 Varying conditions of ionic strength, pH or solid/liquid ratios during adsorption
13 experiments are problematical as regards the valid comparison of different studies.
14 Moreover, the study results are not always reported in the same way: although in most
15 cases complete adsorption parameters are provided, some researchers only report
16 removal percentages. Another drawback is the scarcity of results regarding the removal
17 of each dye, with the exception of methylene blue. The studies reviewed here involve a
18 total of 20 dyes (Table 3), and although some researchers report the removal of more
19 than one dye by the same adsorbent and under comparable conditions, additional studies
20 are necessary before strong conclusions can be reached about each compound, in
21 particular direct dyes.

22 In most of the studies reviewed here, details are not given about the feedstock used to
23 produce the composts or the characteristics of the composting process, despite the
24 potential influence of these factors on the performance of the adsorbents. Future studies
25 should include more detailed information about the composting process, the feedstock

1 used and the maturity of the materials used for dye removal. Additional studies should
2 be conducted to compare composted/vermicomposted and untreated waste.

3 However, studies comparing the performance of several composts with different
4 properties on the removal of one or more dyes remain scarce. In the absence of direct
5 comparison of dye removal by several composts, indirect inference by comparison of
6 the results from different studies may also be helpful. Unfortunately, these studies do
7 not always provide details of the characteristics of the composts used, so comparison is
8 difficult. There is therefore a need for further studies that compare the removal of one or
9 more dyes by several composts under comparable experimental conditions.

10 Finally, most of the studies reviewed consider adsorption as the removal method,
11 although this is not the only possible mechanism of removal during contact between dye
12 solutions and compost. The main processes regulating dye removal are adsorption and
13 biodegradation. Although the factors that affect adsorption are well known, because
14 they have been clearly established for organic pesticides, including the nature of the
15 molecule (ionizable or non-ionizable, pKa, solubility, etc.), biodegradation is also
16 important and has been less well studied. Indeed, among the studies reviewed here, only
17 one considered the biodegradation process. The contribution of biodegradative
18 mechanisms to dye removal should also be explored further.

19

20 **5. Conclusions**

21 Composts and vermicomposts have properties that make them potentially valuable for
22 use as adsorbents of a range of substances, including metallic ions and pesticides and
23 organic compounds such as dyes. However, the last application has received little
24 attention, especially relative to the numerous studies concerning other types of
25 biosorbents, and it represent a promising line of research. The results of the studies

1 reviewed here highlight both the unexplored potential of the process of dye removal by
2 adsorption onto composts and vermicomposts and the need for further investigation of
3 the processes leading to colour removal. In particular, review of the existing literature
4 suggests significant research gaps: very few dyes have been studied, and additional
5 research on dyes of all classes is required to determine which compounds or groups of
6 compounds are most suitable for this treatment. Further studies evaluating the removal
7 of one or more dyes by several composts under comparable experimental conditions are
8 also required in order to obtain further information about the role of the properties of
9 composts on the process of dye removal. In this respect, a more detailed
10 characterization of the composts and vermicomposts used in dye removal studies is also
11 necessary, as well as more complete description of the feedstocks and the characteristics
12 of the process of composting or vermicomposting in each case. Finally, studies on the
13 use of biodegradation mechanisms involved in removal are also necessary. Future
14 research on the use of compost and vermicomposts as low-cost sorbents for dye removal
15 should focus on resolving these shortcomings.

16

17 **References**

- 18 Adegoke KA, Bello OS (2015) Dye sequestration using agricultural wastes as
19 adsorbents. *Water Resour Industry* 12:8-24.
- 20 Ahmad T, Danish M (2018) Prospects of banana waste utilization in wastewater
21 treatment: A review. *J Environ Manage* 206:330-348.
- 22 Ahmad T, Danish M, Rafatullah M, Ghazali A, Sulaiman O, Hashim R, Ibrahim MNM
23 (2012) The use of date palm as a potential adsorbent for wastewater treatment: A
24 review. *Env Sci Pollut Res* 19:1464-1484.

1 Ahmad T, Rafatullah M, Ghazali A, Sulaiman O, Hashim R (2011) Oil palm biomass-
2 based adsorbents for the removal of water pollutants-a review. *J Environ Sci Heal C*
3 29:177-222.

4 Ahmaruzzaman M (2010) A review on the utilization of fly ash. *Prog Energ Combust*
5 36:327-363.

6 Anastopoulos I, Margiotoudis I, Massas I (2018) The use of olive tree pruning waste
7 compost to sequester methylene blue dye from aqueous solution. *Int J*
8 *Phytoremediat* 20:831-838.

9 Bailey SE, Olin TJ, Bricka RM, Adrian DD (1999) A review of potentially low-cost
10 sorbents for heavy metals. *Water Res* 33:2469-2479.

11 Barral MT, Paradelo R, Liste A, Cancelo-González J, Balufo A, Prieto DM (2014)
12 Reutilization of granite powder as a component of permeable reactive barriers for
13 the treatment of Cr(VI)-contaminated waters. *Span J Soil Sci* 4:179-191.

14 Bello OS, Bello IA, Adegoke KA (2013) Adsorption of dyes using different types of
15 sand: A review. *S Afr J Chem* 66:117-129.

16 Bhagavathi Pushpa T, Vijayaraghavan J, Sardhar Basha SJ, Sekaran V, Vijayaraghavan
17 K, Jegan J (2015) Investigation on removal of malachite green using EM based
18 compost as adsorbent. *Ecotox Environ Safe* 118:177–182.

19 Bhagavathi Pushpa T, Sekaran V, Sardhar Basha SJ, Jegan J (2016a) Investigation on
20 preparation, characterization and application of effective microorganisms (EM)
21 based composts - An Ecofriendly Solution. *Nature Environ Pollut Technol* 15:153-
22 158.

23 Bhagavathi Pushpa T, Vijayaraghavan J, Vijayaraghavan K, Jegan J (2016b) Utilization
24 of Effective Microorganisms based water hyacinth compost as biosorbent for the
25 removal of basic dyes. *Desalin Water Treat* 57: 24368-24377.

- 1 Bhat SA, Singh S, Singh J, Kumar S, Bhawana, Pal Vig A, (2018) Bioremediation and
2 detoxification of industrial wastes by earthworms: Vermicompost as powerful crop
3 nutrient in sustainable agriculture. *Bioresource Technol* 252:172-179.
- 4 Boni MR, Sbaffoni S (2009) The potential of compost-based biobarriers for Cr(VI)
5 removal from contaminated groundwater: Column test. *J Hazard Mater* 166:1087-
6 1095.
- 7 Cancelo-González J, Martiñá-Prieto D, Hernández-Huerta D, Barral MT (2017) Metal
8 removal by pine bark compost using a permeable reactive barrier device at
9 laboratory scale. *Environ Chem* 14:310.
- 10 Carmona E, Abad M (2008) Aplicación del compost en viveros y semilleros. In:
11 Moreno J, Moral R (eds) *Compostaje*. Mundi-Prensa, Madrid, pp 397-424.
- 12 Carrillo Zenteno MD, De Freitas RCA, Fernandes RBA, Fontes MPF, Jordão CP (2013)
13 Sorption of cadmium in some soil amendments for in situ recovery of contaminated
14 soils. *Water Air Soil Pollut* 224:14-18.
- 15 Cegarra J, Paredes C (2008) Residuos agroindustriales. In: Moreno J, Moral R (eds)
16 *Compostaje*. Mundi-Prensa, Madrid, pp 519-551.
- 17 Crini G (2005) Recent developments in polysaccharide-based materials used as
18 adsorbents in wastewater treatment. *Prog Polym Sci* 30:38-70.
- 19 Crini G (2006) Non-conventional low-cost adsorbents for dye removal: A review.
20 *Bioresource Technol* 97:1061-1085.
- 21 de Bertoldi M, Vallini G, Pera A (1983) The biology of composting: A review waste
22 *Manage Res* 1:157-176.
- 23 De Gisi S, Lofrano G, Grassi M, Notarnicola M (2016) Characteristics and adsorption
24 capacities of low-cost sorbents for wastewater treatment: A review. *Sustainable*
25 *Materials and Technologies* 9:10-40.

1 De Godoi Pereira M, Arruda MAZ (2004) Preconcentration of Cd(II) and Pb(II) using
2 humic substances and flow systems coupled to flame atomic absorption
3 spectrometry. *Microchim Acta* 146:215-222.

4 De Godoi Pereira M, Korn M, Barros Santos B, Guia Ramos M (2009) Vermicompost
5 for tinted organic cationic dyes retention. *Water Air Soil Pollut* 200:227-235.

6 Demirbas A (2009) Agricultural based activated carbons for the removal of dyes from
7 aqueous solutions: A review. *J Hazard Mater* 167:1-9.

8 Derbyshire F, Andrews R, Jacques D, Jagloyen M, Kimber G, Rantelle T (2001)
9 Synthesis of isotropic carbon fibers and activated carbon fibers from pitch
10 precursors. *Fuel* 80:345-356.

11 Dey MD, Das S, Kumar R, Doley R, Bhattacharya SS, Mukhopadhyay R (2017)
12 Vermiremoval of methylene blue using *Eisenia fetida*: A potential strategy for
13 bioremediation of synthetic dye-containing effluents. *Ecol Eng* 106:200–208.

14 Diacono M, Montemurro F (2010) Long-term effects of organic amendments on soil
15 fertility. A review. *Agron Sustain Dev* 30:410-422.

16 Doke KM, Khan EM (2013) Adsorption thermodynamics to clean up wastewater;
17 critical review. *Rev Environ Sci Bio* 12:25-44.

18 Farrell M, Jones DL (2009) Critical evaluation of municipal solid waste composting and
19 potential compost markets. *Bioresource Technol* 100:4301-4310.

20 Geetha K, Velmani N (2015) Diverse technology and methods for dye treatment. *Asian*
21 *J Chem* 27:1177-1184.

22 Gómez-Brandón M, Domínguez J (2014) Recycling of solid organic wastes through
23 vermicomposting: microbial community changes throughout the process and use of
24 vermicompost as a soil amendment. *Crit Rev Env Sci Tech* 44:1289-1312.

- 1 Gottlieb A, Shaw C, Smith C, Wheatle A, Forsyth S (2003) Toxicity of textile reactive
2 azo dyes after hydrolysis and decolorization. *J Biotec* 101:49–56.
- 3 Gupta VK, Suhas (2009) Application of low-cost adsorbents for dye removal - A review.
4 *J Environ Manage* 90:2313-2342.
- 5 Hargreaves JC, Adl MS, Warman PR (2008) A review of the use of composted
6 municipal solid waste in agriculture. *Agric Ecosyst Environ* 123:1-14.
- 7 He X, Zhang Y, Shen M, Tian Y, Zheng K, Zeng G (2017) Vermicompost as a natural
8 adsorbent: evaluation of simultaneous metals (Pb, Cd) and tetracycline adsorption
9 by sewage sludge-derived vermicompost. *Environ Sci Pollut Res* 24:8375-8384.
- 10 Ho YS, McKay G (1999) Pseudo-second order model for sorption processes. *Process*
11 *Biochem* 34:451-465.
- 12 Ho YS, McKay G (2003) Sorption of dyes and copper ions onto biosorbents. *Process*
13 *Biochem* 38:1047–1061.
- 14 Huang M, Zhu Y, Li Zh, Huang B, Luo N, Liu Ch, Zeng G (2016) Compost as a soil
15 amendment to remediate heavy metal-contaminated agricultural soil: Mechanisms,
16 efficacy, problems, and strategies. *Water Air Soil Pollut* 227:359.
- 17 Jain AK, Gupta VK, Bhatnagar A, Suhas (2003) Utilization of industrial waste products
18 as adsorbents for the removal of dyes. *J Hazard Mater B*101:31–42.
- 19 Jordão CP, Fernandes RBA, de Lima Ribeiro K, de Souza Nascimento B, de Barros PM
20 (2009) Zn(II) adsorption from synthetic solution and kaolin wastewater onto
21 vermicompost. *J Hazard Mater* 162:804-811.
- 22 Jordão CP, Pereira WL, Carari DM, Fernandes RBA, de Almeida RM, Fontes MPF
23 (2011) Adsorption from Brazilian soils of Cu(II) and Cd(II) using cattle manure
24 vermicompost. *Int J Environ Stud* 68:719-736.

- 1 Józwiak T, Filipkowska U, Rodziewicz J, Mielcarek A, Owczarkowska D (2013)
2 Application of compost as a cheap sorbent for dyes removal from aqueous solutions.
3 *Rocznik Ochrona Srodowiska* 15:2398–2411.
- 4 Kharat DS (2015) Preparing agricultural residue based adsorbents for removal of dyes
5 from effluents - A review. *Braz J Chem Eng* 32:1-12.
- 6 Koay YS, Ahamad IS, Nourouzi MM, Abdullah LC, Choong TSY (2014) Development
7 of novel low-cost quaternized adsorbent from palm oil agriculture waste for reactive
8 dye removal. *BioResources* 9:66-85.
- 9 Kocasoy G, Güvener Z (2009) Efficiency of compost in the removal of heavy metals
10 from the industrial wastewater. *Environ Geol* 57:291-296.
- 11 Krishna SK, Sivaprakash S (2015) Removal of dyes by using various adsorbents: A
12 review. *Int J Appl Chem* 11:195-202.
- 13 Kyziol-Komosińska J, Rosik-Dulewska C, Pająk M (2010) The potential of metal-
14 complex dyes removal from wastewater the sorption method onto organic-matter
15 rich substances. In: Pawlowski L, Dudzinska MR, Pawlowski A (eds),
16 *Environmental Engineering III*. CRC Press, pp 197-202.
- 17 Larney FJ, Sullivan DM, Buckley KE, Eghball B (2006) The role of composting in
18 recycling manure nutrients. *Can J Soil Sci* 86:597-611.
- 19 Lashermes G, Houot S, Barriuso E (2010) Sorption and mineralization of organic
20 pollutants during different stages of composting, *Chemosphere* 79:455-462.
- 21 Liu L, Guo X, Zhang Ch, Luo Ch, Xiao Ch, Li R (2018) Adsorption behaviours and
22 mechanisms of heavy metal ions' impact on municipal waste composts with
23 different degree of maturity. *Environ Technol* (in press).
- 24 Matos GD, Arruda MAZ (2003) Vermicompost as an adsorbent for removing metal ions
25 from laboratory effluents. *Process Biochem* 39:81–88.

- 1 McKay G, Hadi M, Samadi MT, Rahmani AR, Aminabad MS, Nazemi F (2011)
2 Adsorption of reactive dye from aqueous solutions by compost. *Desalin Water Treat*
3 28:164-173.
- 4 Mendes CB, Lima GDF, Alves VN, Coelho NMM, Dragunski DC, Tarley CRT (2012)
5 Evaluation of vermicompost as a raw natural adsorbent for adsorption of pesticide
6 methylparathion. *Environ Technol* 33:167-172.
- 7 Momina, Shahadat M, Isamil S (2018) Regeneration performance of clay-based
8 adsorbents for the removal of industrial dyes: A review. *RSC Adv* 8:24571-24587.
- 9 Mortier N, Velghe F, Verstichel S (2016) Organic recycling of agricultural waste today:
10 composting and anaerobic digestion. In: D'Urso OF (ed), *Biotransformation of*
11 *Agricultural Waste and By-products*, 1st edn. Elsevier, pp 69-124.
- 12 Ong ST, Keng PS, Lee SL, Hung YT (2014) Low cost adsorbents for sustainable dye
13 containing-wastewater treatment. *Asian J Chem* 26:1873-1881.
- 14 Pandey S (2017) A comprehensive review on recent developments in bentonite-based
15 materials used as adsorbents for wastewater treatment. *J Mol Liq* 241:1091-1113.
- 16 Pant D, Adholeya A (2007) Biological approaches for treatment of distillery wastewater:
17 A review. *Bioresource Technol* 98:2321-2334.
- 18 Paradelo R, Barral MT (2012) Evaluation of the potential capacity as metal biosorbents
19 of two MSW composts with different Cu, Pb and Zn content. *Bioresource Technol*
20 104:810–813.
- 21 Paradelo R, Cendón Y, Moldes AB, Barral MT (2007) A pot experiment with mixtures
22 of slate processing fines and compost. *Geoderma* 141:363-369.
- 23 Paradelo R, Moldes AB, Barral MT (2009a) Properties of slate mining wastes incubated
24 with grape marc compost under laboratory conditions. *Waste Manage* 29:579-584.

- 1 Paradelo R, Moldes AB, Barral MT (2009b) Amelioration of the physical properties of
2 slate processing fines using grape marc compost and vermicompost. *Soil Sci Soc*
3 *Amer J* 73:1251-1260.
- 4 Paradelo R, Moldes AB, Barral MT (2009c) Treatment of red wine vinasses with non-
5 conventional substrates for removing coloured compounds. *Water Sci Technol*
6 59:1585-1592.
- 7 Paradelo R, Villada A, Barral MT (2011) Reduction of the short-term availability of
8 copper, lead and zinc in a contaminated soil amended with MSW compost. *J Hazard*
9 *Mater* 188:96-104.
- 10 Paradelo R, Moldes AB, González D, Barral MT (2012) Plant tests for determining the
11 suitability of grape marc composts as components of plant growth media. *Waste*
12 *Manage Res* 30:1059-1065.
- 13 Paradelo R, Basanta R, Barral MT (2019) Water-holding capacity and plant growth in
14 compost-based substrates modified with polyacrylamide, guar gum or bentonite. *Sci*
15 *Hortic-Amsterdam* 243:344-349.
- 16 Park JH, Lamb D, Paneerselvam P, Choppala G, Bolan N, Chung J-W (2011) Role of
17 organic amendments on enhanced bioremediation of heavy metal(loid) contaminated
18 soils. *J Hazard Mater* 185:549-574.
- 19 Pereira MG, Arruda MAZ (2003) Vermicompost as a natural adsorbent material:
20 Characterization and potentialities for cadmium adsorption. *J Brazil Chem Soc*
21 14:39-47.
- 22 Perez-Ameneiro M, Vecino X, Barbosa-Pereira L, Cruz JM, Moldes AB (2014)
23 Removal of pigments from aqueous solution by a calcium alginate-grape marc
24 biopolymer: A kinetic study. *Carbohydr Polym* 101:954-960.

- 1 Perez-Ameneiro M, Vecino X, Cruz JM, Moldes AB (2015) Physicochemical study of a
2 bio-based adsorbent made from grape marc. *Ecol Eng* 84:190-193.
- 3 Pradhan AK, Beura K, Das I, De N (2014) Cadmium sorption by vermicompost
4 produced from recycled organic wastes. *Ecol Environ Conserv* 20:S271-S276.
- 5 Qian QR, Machida M, Tatsumoto H (2008) Textural and surface chemical
6 characteristics of activated carbons prepared from cattle manure compost. *Waste*
7 *Manage* 28:1064–1071.
- 8 Rafatullah M, Sulaiman O, Hashim R, Ahmad A (2010) Adsorption of methylene blue
9 on low-cost adsorbents: A review. *J Hazard Mater* 177:70-80.
- 10 Ramesh A, Lee DJ, Wong JWC (2005) Adsorption equilibrium of heavy metals and
11 dyes from wastewater with low-cost adsorbents: A review. *J Chin Inst Chem Eng*
12 36:203-222.
- 13 Semple KT, Reid BJ, Fermor TR (2001) Impact of composting strategies on the
14 treatment of soils contaminated with organic pollutants. *Environ Pollut* 112:269-83.
- 15 Sharma P, Kaur H, Sharma M, Sahore V (2011) A review on applicability of naturally
16 available adsorbents for the removal of hazardous dyes from aqueous waste.
17 *Environ Monitor Assess* 183:151-195.
- 18 Singh J, Kaur A (2015) Vermicompost as a strong buffer and natural adsorbent for
19 reducing transition metals, BOD, COD from industrial effluent. *Ecol Eng* 74:13-19.
- 20 Slokar YM, Majcen Le Marechal A (1997) Methods of decoloration of textile
21 wastewaters. *Dyes Pigments* 37:335-356.
- 22 Sulyman M, Namiesnik J, Gierak A (2017) Low-cost adsorbents derived from
23 agricultural by-products/wastes for enhancing contaminant uptakes from wastewater:
24 A review. *Pol J Environ Stud* 26:479-510.

- 1 Sun Q, Yang L (2003) The adsorption of basic dyes from aqueous solution on modified
2 peat-resin particle. *Water Res* 37:1535–1544.
- 3 Toptas A, Demierege S, Ayan EM, Yanik J (2014) Spent mushroom compost as
4 biosorbent for dye biosorption. *Clean – Soil Air Water* 42:1721–1728.
- 5 Tsui LS, Roy WR, Cole MA (2003) Removal of dissolved textile dyes from wastewater
6 by a compost sorbent. *Coloration Technology* 119:14-18.
- 7 Tsui L, Juang MA (2010) Effects of composting on sorption capacity of bagasse-based
8 chars. *Waste Management* 30:995-999.
- 9 Urdaneta C, Parra LMM, Matute S, Garaboto MA, Barros H, Vázquez C (2008)
10 Evaluation of vermicompost as bioadsorbent substrate of Pb, Ni, V and Cr for waste
11 waters remediation using Total Reflection X-ray Fluorescence. *Spectrochim Acta B*
12 63:1455-1460.
- 13 U.S. EPA (2002) Biosolids technology fact sheet: Use of composting for biosolids
14 management. EPA/832-F-02-024. Office of Solid Waste and Emergency Response,
15 U.S. EPA, Washington, D.C.
- 16 Wang S, Wu H (2006) Environmental-benign utilisation of fly ash as low-cost
17 adsorbents. *J Hazard Mater* 136:482-501.
- 18 Wang Z, Guo H, Shen F, Duan D (2015) Production of biochar by vermicompost
19 carbonization and its adsorption to Rhodamine-B. *Huanjing Kexue Xuebao*
20 35:3170-3177.
- 21 Willmott N, Guthrie J, Nelson G (1998) The biotechnology approach to colour removal
22 from textile effluents. *Dyes Colour* 114:38–41.
- 23 Wu L, Liu Y, Li Y, Shen F, Yang G, Wu J (2016) Removal of 17 β -estradiol from
24 aqueous solutions by vermicompost-derived biochars adsorption. *Res Environ Sci*
25 29:1537-1545.

- 1 Yagub MT, Sen TK, Afroze S, Ang HM (2014) Dye and its removal from aqueous
2 solution by adsorption: A review. *Adv Colloid Interfac* 209:172-184.
- 3 Yang G, Wang Z, Xian Q, Shen F, Sun C, Zhang Y, Wu J (2015) Effects of pyrolysis
4 temperature on the physicochemical properties of biochar derived from
5 vermicompost and its potential use as an environmental amendment. *RSC Adv*
6 5:40117-40125.
- 7 Yang G, Wu L, Xian Q, Shen F, Wu J, Zhang Y (2016) Removal of congo red and
8 methylene blue from aqueous solutions by vermicompost-derived biochars. *PLoS*
9 *ONE* 11(5):e0154562.
- 10

1 FIGURE CAPTIONS

2

3 Figure 1. Molecular structures of dyes used in the studies reviewed here.

4

5 Figure 2. Maximum adsorption capacities (Q_m , from the Langmuir model) reported for
6 basic, acid, reactive and direct dyes on composts and vermicomposts.

1 Table 1. Review papers published between 2005 and 2018 dealing with the use of low-cost adsorbents for dye or colored compounds removal.

Reference	Adsorbent	Dye classes
Ahmad and Danish (2018)	Banana waste	<i>Acid dyes:</i> orange 52 (Methyl orange), Egacid Orange II, Brilliant blue <i>Basic dyes:</i> blue 9 (Methylene blue), green 4 (Malachite green), violet 10 (Rhodamine B) <i>Direct dyes:</i> red <i>Other dyes:</i> Safranin, Novacron blue FN-R, Methyl violet, Methyl red
Momina et al. (2018)	Clays	<i>Acid dyes:</i> blue 193, orange 52 (Methyl orange) <i>Basic dyes:</i> red 2, green 4 (Malachite green), blue 9 (Methylene blue) <i>Reactive dyes:</i> black 5
Pandey (2017)	Clays	<i>Other dyes:</i> Amido black 10B
Sulyman et al. (2017)	Agricultural wastes	<i>Acid dyes:</i> blue 25, blue 92, orange 52 (Methyl orange) <i>Basic dyes:</i> red 29, blue 9 (Methylene blue), violet 3 (Crystal violet), green 4 (Malachite green), violet 10 (Rhodamine B) <i>Direct dyes:</i> red 28 (Congo red) <i>Reactive dyes:</i> Remazol brilliant yellow <i>Other dyes:</i> Methyl violet, Methyl red, Red MX 3B, Cibacron yellow
De Gisi et al. (2016)	Agricultural and household wastes, industrial waste, soil and ore materials, metal oxides and hydroxides	<i>Acid dyes:</i> yellow 36, blue 92, blue 25, blue 80, blue 264, red 14, red 114, orange 10, orange 52 (Methyl orange), violet <i>Basic dyes:</i> blue 9 (Methylene blue), blue 69, red 22, green 4 (Malachite green), violet 10 (Rhodamine B), red 9 <i>Direct dyes:</i> red 28 (Congo red), red 80, red 81, blue 71 <i>Reactive dyes:</i> Brilliant green <i>Disperse dyes:</i> orange 25
Adegoke and Bello (2015)	Agricultural wastes	<i>Acid dyes:</i> violet 17, red 119, blue, blue 15 <i>Basic dyes:</i> blue 9 (Methylene blue), yellow 21, violet 10 (Rhodamine B), green 4 (Malachite green) <i>Direct dyes:</i> F. Scarlet, red 23, red 28 (Congo red) <i>Reactive dyes:</i> red 23, blue 19 <i>Other dyes:</i> Red brown C4R

Kharat (2015)	Agricultural wastes	<p><i>Acid dyes:</i> green 20, orange 7, blue 29</p> <p><i>Basic dyes:</i> blue 3, blue 9 (Methylene blue), violet 1, violet 3 (Crystal violet), violet 10 (Rhodamine B), Rhodamine 6G, green 4 (Malachite green)</p> <p><i>Direct dyes:</i> red 28 (Congo red), orange 26</p> <p><i>Reactive dyes:</i> red 3 BS, red 2, red 120, red 198, red 228, orange, orange 16, black 5, Turquoise blue QG, Remazol black B, Remazol brilliant blue R, Remazol brilliant red</p> <p><i>Disperse dyes:</i> red 1</p> <p><i>Other dyes:</i> Erichrome black T, Dark green PLS, Coomassie brilliant, Methylene red, Ethylene blue, Acridine orange, Aniline blue, Safranin O, Alpacide yellow, Tartrazine</p>
Geetha and Velmani (2015)	Agricultural wastes, industrial waste	<p><i>Basic dyes:</i> yellow 21</p> <p><i>Direct dyes:</i> red 28 (Congo red)</p> <p><i>Reactive dyes:</i> orange 107, Remazol yellow</p>
Krishna and Sivaprakash (2015)	Agricultural and household wastes, industrial waste	
Koay et al. (2014)	Agricultural wastes	<p><i>Reactive dyes:</i> black 5</p> <p><i>Acid dyes:</i> Egacid Orange II, blue 74 (Indigo carmine), orange 52 (Methyl orange)</p> <p><i>Basic dyes:</i> blue 3, blue 9 (Methylene blue), red 46, yellow 21, violet 3 (Crystal violet), violet 10 (Rhodamine B), green 4 (Malachite green)</p> <p><i>Direct dyes:</i> navy blue 106, blue 86, yellow 12, red 28 (Congo red)</p> <p><i>Reactive dyes:</i> blue 19</p> <p><i>Other dyes:</i> Lanaset grey G, Astrazon yellow</p>
Ong et al. (2014)	Agricultural wastes, industrial waste, clays	<p><i>Acid dyes:</i> yellow 23, yellow 36, yellow 132, blue, blue 25, blue 29, blue 80, blue 193, blue 256, blue 264, red 18, red 27, red 73, violet, violet 17, orange 7, orange 10, orange 52 (Methyl orange), green 25</p> <p><i>Basic dyes:</i> blue 3G, blue 9 (Methylene blue), blue 69, blue 86, yellow 2, red 13, red 18, red 46, green 4 (Malachite green), violet 3 (Crystal violet), violet 10 (Rhodamine B)</p> <p><i>Direct dyes:</i> brown 2, red 23, red 28 (Congo red)</p> <p><i>Reactive dyes:</i> black 5, orange 16, blue 19, red 4, red 5, Brilliant green, Brilliant red HE-3B, Remazol black, Brilliant blue</p> <p><i>Disperse dyes:</i> red 1</p> <p><i>Other dyes:</i> Eriochrome black T, Astrazone black, Naphthol green</p>
Yagub et al. (2014)	Agricultural wastes, industrial waste, clays	

Bello et al. (2013)	Various sand types	<p>B, Indosol black, Methyl violet, Yellow X-GL, Vat red 10, Vat orange 11</p> <p><i>Basic dyes:</i> blue 9 (Methylene blue), green 4 (Malachite green)</p> <p><i>Other dyes:</i> Neutral red dye, Coomassie blue, Safranin orange</p> <p><i>Acid dyes:</i> violet 54, orange 52 (Methyl orange), Egacid Orange II</p> <p><i>Basic dyes:</i> orange, blue 3, blue 9 (Methylene blue), yellow 28, violet 3 (Crystal violet), green 4 (Malachite green)</p> <p><i>Direct dyes:</i> red 28 (Congo red)</p>
Doke and Khan (2013)	Agricultural wastes, industrial waste	<p><i>Reactive dyes:</i> black 5, Brilliant red HE-3B, Remazol brilliant orange 3R</p> <p><i>Other dyes:</i> Methyl violet, Eriochrome black T, Neutral red, Tartrazina, Amaranth</p> <p><i>Acid dyes:</i> green 25, black 26, blue 7</p> <p><i>Basic dyes:</i> blue 9 (Methylene blue)</p> <p><i>Reactive dyes:</i> Remazol</p>
Ahmad et al. (2012)	Agricultural wastes	<p><i>Acid dyes:</i> yellow 23, yellow 36, yellow 99, yellow 117, blue, blue 9, blue 25, blue 29, blue 74 (Indigo carmine), blue 80, blue 113, blue 264, orange 7, orange 10, orange 51, blue 80, red 91, red 114, brown 283, violet, Brilliant blue, Egacid orange II, Egacid red G, Egacid yellow G</p> <p><i>Basic dyes:</i> blue, blue 4, blue 9 (Methylene blue), blue 41, blue 69, violet 1, violet 3 (Crystal violet), violet 10 (Rhodamine B), green 4 (Malachite green), yellow, yellow 28, red, red 18, red 22, red 46, brown 1</p> <p><i>Direct dyes:</i> yellow 11, yellow 12, yellow 28, yellow 50, red 12b, red 28 (Congo red), red 89, black, black 19, brown</p> <p><i>Reactive dyes:</i> blue 2, yellow 2, yellow 23, red, red 2, red 4, red 120, red 141, Ramazol yellow, Ramazol back, Ramazol red</p> <p><i>Disperse dyes:</i> blue 79, red 1</p> <p><i>Other dyes:</i> α-picoline, Safranin, Midlon black VL, Brilliant green, Polar yellow, Polar blue RAWL, Methylene yellow, Methyl violet, Methyl red</p>
Sharma et al. (2011)	Agricultural wastes, industrial waste, clays	<p><i>Basic dyes:</i> blue 9 (Methylene blue), green 4 (Malachite green)</p> <p><i>Direct dyes:</i> blue 71, red 28 (Congo red)</p> <p><i>Reactive dyes:</i> black 5, red E</p> <p><i>Disperse dyes:</i> blue, red</p>
Ahmad et al. (2011)	Agricultural wastes	<p><i>Basic dyes:</i> blue 9 (Methylene blue), green 4 (Malachite green)</p> <p><i>Direct dyes:</i> blue 71, red 28 (Congo red)</p> <p><i>Reactive dyes:</i> black 5, red E</p> <p><i>Disperse dyes:</i> blue, red</p>

Ahmaruzzaman (2010)	Industrial wastes	<p><i>Acid dyes:</i> red 91, blue 9, blue 29, Egacid orange II, Egacid red G, Egacid yellow G</p> <p><i>Basic dyes:</i> blue 9 (Methylene blue), violet 3 (Crystal violet), violet 10 (Rhodamine B), green 4 (Malachite green)</p> <p><i>Direct dyes:</i> red 28 (Congo red)</p> <p><i>Other dyes:</i> Rosaniline hydrochloride, Midlon black VL, Orange-G</p>
Rafatullah et al. (2010)	Agricultural wastes, industrial waste, clays	<p><i>Basic dyes:</i> blue 9 (Methylene blue)</p> <p><i>Acid dyes:</i> blue 15, blue 74 (Indigo carmine), red 119, violet 17, violet 49, orange 52 (Methyl orange)</p>
Demirbas (2009)	Agricultural wastes	<p><i>Basic dyes:</i> blue 9 (Methylene blue)</p> <p><i>Direct dyes:</i> blue 86</p> <p><i>Other dyes:</i> Erythrosine, Quinoline yellow</p>
Gupta and Suhas (2009)	Agricultural wastes, industrial waste, clays	<p><i>Acid dyes:</i> brilliant blue, blue, blue 9, blue 25, blue 29, blue 40, blue 74 (Indigo carmine), blue 80, red 88, blue 92, blue 113, blue 193, blue 256, blue 264, red 1, red 14, red 18, red 51, red 73, red 88, red 114, yellow, yellow 11, yellow 17, yellow 36, yellow 99, yellow 117, yellow 132, brown, brown 283, black 26, green 25, orange 7, orange 10, orange 12, orange 52 (Methyl orange), violet, Ethyl orange</p> <p><i>Basic dyes:</i> blue 3, blue 6 (Meldolás blue), blue 9 (Methylene blue), blue 47, blue 54, blue 69 (Astrazone blue), green 4 (Malachite green), red 2, red 13, red 18, red 22, red 29, red 46, yellow 21, yellow 24, orange 2 (Chrysoidine G), violet 3 (Crystal violet), violet 10 (Rhodamine B), violet 14 (basic fuchsin)</p> <p><i>Direct dyes:</i> black 168, brown, brown 1, red, red 12b, red 23, red 28 (Congo red), red 79, red 80, red 81, red 89, blue, blue 86, yellow 12, green 26</p> <p><i>Reactive dyes:</i> black B, black 5, blue 2, blue 19, blue 114, yellow 2, yellow 23, yellow 64, yellow 86, yellow 176, Levafix, green 12, orange 16, red X6BN Sandoz, red 2 red 120, red 124, red 141, red 189, red 222, red 239, Remazol golden yellow, Remazol red BB, Remazol black B, Remazol blue</p> <p><i>Disperse dyes:</i> blue, red, red 1, orange 25</p> <p><i>Other dyes:</i> Metomega chrome orange, Sella fast brown H, Brilliant Red E-4BA, Vat blue 4</p>
Wang and Wu (2006)	Industrial wastes	<p><i>Acid dyes:</i> blue 9, blue 29, blue 40, red 1, red 88, red 91, Egacid orange II, Egacid Red G, Egacid yellow G</p> <p><i>Basic dyes:</i> blue 9 (Methylene blue), violet 3 (Crystal violet), violet</p>

Crini (2006)	Agricultural wastes, industrial waste, clays, soil materials, peat	<p>10 (Rhodamine B) <i>Direct dyes:</i> red 28 (Congo red) <i>Other dyes:</i> Rosaniline hydrochloride, Midlon Black VL</p> <p><i>Acid dyes:</i> yellow, yellow 17, yellow 36, yellow 99, yellow 117, yellow 132, blue, blue 9, blue 25, blue 29, blue 40, blue 80, blue 113, blue 193, blue 256, blue 264, red 4, red 18, red 73, red 88, red 114, violet, violet 17, orange 10, orange 12, orange 52 (Methyl orange), Ethyl orange, green 25</p> <p><i>Basic dyes:</i> yellow, yellow 21, yellow 24, red 2, red 13, red 18, red 22, red 29, red 46, blue 9 (Methylene blue), blue 47, blue 69, green 4 (Malachite green), violet 3 (Crystal violet), violet 10 (Rhodamine B), violet 14 (basic fuchsin)</p> <p><i>Direct dyes:</i> red, red 28 (Congo red), red 81, brown 1, yellow 12</p> <p><i>Reactive dyes:</i> red, red 2, red 4, red 5, red 120, red 124, red 141, red 189, red 222, red 239, E-4BA, yellow 2, yellow 23, yellow 64, yellow 86, yellow 176, yellow 208, blue 2, blue 19, blue 114, black 5, orange 16, orange 107, Remazol yellow, Remazol BB, Remazol blue</p> <p><i>Disperse dyes:</i> red 1</p> <p><i>Other dyes:</i> Alizarin sulfonic, Sella fast brown H, Methyl violet</p>
Ramesh et al. (2005)	Agricultural wastes, industrial waste	<p><i>Acid dyes:</i> blue 113, yellow 36, red 114</p>
Crini (2005)	Synthetic biopolymers	<p><i>Basic dyes:</i> violet 3 (Crystal violet) <i>Direct dyes:</i> red 28 (Congo red) <i>Reactive dyes:</i> red 2, red 189, red 141, red 189, blue 2</p>

1 Table 2. Papers on compost and vermicompost use as adsorbents for pollutant removal.

Pretreatment	Composition of compost	Contaminant	Reference
<i>Eisenia fetida</i> earthworm (40 days)	-Sewage sludge from the second municipal wastewater treatment plant -Different additive materials (soil, straw, fly ash, and sawdust) were mixed with sludge	Heavy metals (Pb (II) and Cd (II)) and tetracycline (TC)	He et al. (2017)
Pyrolysis at 300°C, 500°C and 700°C	Vermicompost biochars	17β-estradiol	Wu et al. (2016)
Slow pyrolysis	Vermicompost biochar	Rhodamine B	Wang et al. (2015)
<i>E. fetida</i> earthworm	Cattle dung vermicompost	Cu, Mn, Fe, Zn	Singh and Kaur (2015)
Pyrolysis at 400-700°C	Vermicompost biochar	Heavy metal ions, dyes and organic contaminants	Yang et al. (2015)
	Temple wastes: vegetable crop residues, grass residues, dry mango leaf litter, regular farmyard manure and cow dung vermicompost	Cd	Pradhan et al. (2014)
	Commercial cattle manure vermicompost	Cd	Carrillo Zenteno et al. (2013)
Dried in an oven at 60°C, and sieved less than 150 μm	Commercial vermicompost as gardening humus	Methylparathion	Mendes et al. (2012)
Air-dried for 72 h and sieved a 2 mm	Cattle manure vermicompost	Cu (II), Cd (II)	Jordão et al. (2011)
<i>E. fetida</i> earthworm	Vermicompost	Cr, Pb, Ni	Parra et al. (2010)
Sun-dried and sieved to 300 μm			
Air-dried for 72 h	Cattle manure vermicompost	Al (III), Fe (II)	Jordão et al. (2010)
Dried at 70°C for 4h	Cattle manure vermicompost	Zn (II)	Jordão et al. (2009)
	Local vermicompost	Pb, Ni, V, Cr	Urdaneta et al. (2008)
Dried and sieved between 75-150 μm	Local vermicompost	Cd (II), Pb (II)	De Godoi Pereira et al. (2004)
Dried and sieved size of ≤ 150, ≤ 355 or ≤ 600 μm	Local vermicompost	Cd (II), Cu (II), Pb (II), Zn (II)	Matos and Arruda (2003)
Dried at 60°C for 24 h	Vermicompost samples were obtained from different regions of Minas Gerais and São Paulo States (Brazil)	Cd (II)	Pereira and Arruda (2003)

2

1 Table 3. Papers on compost and vermicompost use as adsorbents for dye removal.

Reference	Adsorbent	Colorant(s) studied				
		Basic dyes	Acid dyes	Reactive dyes	Direct dyes	Other dyes
Tsui et al. (2003)	Compost	Blue 9, Green 4	Black 24, Orange 74	Orange 16, Red 2	Blue 71, Orange 39	
de Godoi Pereira et al. (2009)	Vermicompost	Crystal violet, Methylene blue				
Paradelo et al. (2009c)	Urban waste compost, grape marc vermicompost, pine bark compost					Winery wastewater
Kyziol-Komosińska et al. (2010)	Green waste compost		Blue 193, Black 194			
McKay et al. (2011)	Urban waste compost			Red 234		
Jozwiak et al. (2013)	Sewage sludge green waste compost	Green 4, Violet 10		Yellow 84, Black 5		
Toptas et al. (2014)	Spent mushroom compost	Red 18	Red 111	Brown 37		
Pérez-Ameneiro et al. (2014)	Grape marc compost					Winery wastewater
Bhagavathi et al. (2015)	Kitchen waste compost	Malachite green				
Bhagavathi et al. (2016a)	Kitchen waste compost, leaf waste compost, paper waste compost, water hyacinth compost	Crystal violet				
Bhagavathi et al. (2016b)	Water hyacinth compost	Methylene blue, Malachite green, Blue 41				
Dey et al. (2017)	Sugarcane bagasse vermicompost	Methylene blue				
Anastopoulos et al. (2018)	Pruning waste compost	Methylene blue				

2

- 1 Table 4. Summary of adsorption parameters of dyes on composts/vermicomposts. Q_m : maximum adsorption capacity at the Langmuir equation;
- 2 t_{eq} : contact time for maximum adsorption.

Dye	Adsorption	Kinetics	Optimal pH	Reference
	Q_m (mg g ⁻¹)	t_{eq} (min)		
Methylene blue	5.47	500		de Godoi Pereira et al. (2009)
Methylene blue	296	1		Bhagavathi et al. (2016b)
Methylene blue	250	1	no optimal pH	Anastopoulos et al. (2018)
Malachite green	151	300	8	Bhagavathi et al. (2015)
Basic green 4 (Malachite green oxalate)	26.41	300	5	Jozwiak et al. (2013)
Malachite green	153	240		Bhagavathi et al. (2016b)
Crystal violet	0.78			de Godoi Pereira et al. (2009)
Basic blue 41	158	10		Bhagavathi et al. (2016b)
Basic red 18	400		6	Toptas et al. (2014)
Basic violet 10/Rhodamine B	27,2	180	5	Jozwiak et al. (2013)
Basic blue 9	0.08	180		Tsui et al. (2003)
Acid orange 74	0.005	180		Tsui et al. (2003)
Acid red 111	141		3	Toptas et al. (2014)
Acid black 24	0.014	360		Tsui et al. (2003)
Acid blue 193	9.3			Kyziol-Komosińska et al. (2010)
Acid black 194	15.9			Kyziol-Komosińska et al. (2010)
Reactive yellow 84	2.15		3	Jozwiak et al. (2013)
Reactive black 5	4.79	180	3	Jozwiak et al. (2013)
Reactive red 234	0.718	24 h	2.3	McKay et al. (2011)
Reactive red 2	0.003	180		Tsui et al. (2003)
Levafix braun (reactive brown 37)	169.5		2	Toptas et al. (2014)
Direct orange 39	0.002	180		Tsui et al. (2003)

3