

PAPER • OPEN ACCESS

Looping and clustering model for the organization of protein-DNA complexes on the bacterial genome

To cite this article: Jean-Charles Walter *et al* 2018 *New J. Phys.* **20** 035002

View the [article online](#) for updates and enhancements.

Related content

- [DNA sequence-dependent mechanics and protein-assisted bending in repressor-mediated loop formation](#)
James Q Boedicker, Hernan G Garcia, Stephanie Johnson *et al.*
- [Physical descriptions of the bacterial nucleoid at large scales, and their biological implications](#)
Vincenzo G Benza, Bruno Bassetti, Kevin D Dorfman *et al.*
- [Review Article](#)
Joel Stavans and Amos Oppenheim

Recent citations

- [Focus on bacterial mechanics](#)
Stefan Klumpp *et al*



PAPER

Looping and clustering model for the organization of protein-DNA complexes on the bacterial genome

OPEN ACCESS

RECEIVED

27 June 2017

REVISED

30 January 2018

ACCEPTED FOR PUBLICATION

6 February 2018

PUBLISHED

26 March 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Jean-Charles Walter¹, Nils-Ole Walliser¹, Gabriel David¹, Jérôme Dornignac¹, Frédéric Geniet¹, John Palmeri¹, Andrea Parmeggiani^{1,2}, Ned S Wingreen³ and Chase P Broedersz⁴ 

¹ L2C, Université de Montpellier, CNRS, Montpellier, France

² DIMNP, Université de Montpellier, CNRS, Montpellier, France

³ Department of Molecular Biology and Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton NJ 08544, United States of America

⁴ Arnold-Sommerfeld-Center for Theoretical Physics and Center for NanoScience, Ludwig-Maximilians-Universität München, D-80333 München, Germany

E-mail: c.broedersz@lmu.de


Keywords: Chromosome organization, ParABS, protein-DNA cluster

Abstract

The bacterial genome is organized by a variety of associated proteins inside a structure called the nucleoid. These proteins can form complexes on DNA that play a central role in various biological processes, including chromosome segregation. A prominent example is the large ParB-DNA complex, which forms an essential component of the segregation machinery in many bacteria. ChIP-Seq experiments show that ParB proteins localize around centromere-like *parS* sites on the DNA to which ParB binds specifically, and spreads from there over large sections of the chromosome. Recent theoretical and experimental studies suggest that DNA-bound ParB proteins can interact with each other to condense into a coherent 3D complex on the DNA. However, the structural organization of this protein-DNA complex remains unclear, and a predictive quantitative theory for the distribution of ParB proteins on DNA is lacking. Here, we propose the looping and clustering model, which employs a statistical physics approach to describe protein-DNA complexes. The looping and clustering model accounts for the extrusion of DNA loops from a cluster of interacting DNA-bound proteins that is organized around a single high-affinity binding site. Conceptually, the structure of the protein-DNA complex is determined by a competition between attractive protein interactions and loop closure entropy of this protein-DNA cluster on the one hand, and the positional entropy for placing loops within the cluster on the other. Indeed, we show that the protein interaction strength determines the ‘tightness’ of the loopy protein-DNA complex. Thus, our model provides a theoretical framework for quantitatively computing the binding profiles of ParB-like proteins around a cognate (*parS*) binding site.

1. Introduction

Understanding the biophysical principles that govern chromosome structure in both eukaryotic and prokaryotic cells remains an outstanding challenge [1–7]. Many bacteria have a single chromosome with a length three orders of magnitude longer than the cell itself, posing a daunting organizational problem. Owing to recent technological advances in live-cell imaging and chromosome conformation capture based approaches, it is becoming increasingly clear that the DNA is not coiled like a simple amorphous polymer inside the cell [8–10], but rather exhibits a high degree of organization over a broad range of lengthscales [11]. It remains unclear, however, how this spatial and dynamic organization of the chromosome is established and maintained inside living bacteria [12]. A host of nucleoid-associated proteins (NAPs) have been shown to play a central role in the spatial organization of the bacterial chromosome [12–14]. Such NAPs bind to the DNA in large numbers, and by interacting with each other and with DNA in both sequence-dependent and sequence-independent manners



they can collectively structure the DNA polymer and control chromosome organization, e.g. by DNA looping. More generally, the formation of loops appears to play a fundamental role in structuring DNA in both bacterial DNA and eukaryotic chromatin and further theoretical and simulation studies are needed to clarify the mechanisms at play [15, 16].

In many bacterial species, the broadly conserved ParABS system is responsible for chromosome and plasmid segregation [12, 17]. A central component of this system is the partitioning module, which is formed by a large protein–DNA complex of ParB proteins that assembles around centromere-like *parS* sites, frequently located near the origin of replication. The ParBS complexes can subsequently interact with ParA ATPases, leading to the segregation of replicated origins [18–25]. How is this ParBS partitioning module physically organized on the DNA? ParB is known to bind specifically to *parS*, triggering the formation of a large protein–DNA cluster, which is visible as a tight focus in microscopy images of fluorescently labeled ParB [17, 21, 26, 27]. The propensity of ParB to form foci around *parS* has been exploited in recent studies, which used exogenous expression of fluorescently labeled ParB along with *parS* insertion to label DNA loci for live-cell imaging [28, 29]. In the F-plasmid of *Escherichia coli* cells, each ParB focus contains roughly 300 proteins, together representing 90% of all ParB present in the cell [27]. High-precision ChIP-Seq experiments on this system provide quantitative ParB binding profiles along the DNA, which are strongly peaked around *parS* with a broad decay over a distance of up to 13 kilobasepairs (kb), consistent with earlier observations [26, 30].

Various models have been introduced to explain the distribution of ParB along DNA around *parS* sites. An early study of the distribution of ParB proposed that ParB proteins spread from the *parS* sequence by nearest-neighbor interactions, forming a continuous filament-like structure along the DNA [30]. This model was termed the spreading model. However, this is effectively a 1D model with short range interactions. On general statistical physical grounds, such a 1D model cannot be expected to account for the formation of a large coherent protein–DNA complex, given physiological protein interaction strengths [31]. Furthermore, the number of ParB proteins available in the cell is not sufficient to allow enrichment by simple 1D polymerization of ParB along DNA at genomic distances from *parS* as large as observed experimentally [27]. To resolve the puzzle of how ParB proteins organize around a *parS* site, we recently introduced a novel theoretical framework to study the collective behavior of interacting proteins that can bind to a DNA polymer [31]. This model suggested that ParB assembles into a three-dimensional complex on the DNA, as illustrated in figures 1(a) and (b). Single molecule experiments provided direct evidence for the presence of 3D bridging interactions between two ParB proteins on DNA [32, 33]. We showed that a combination of such a 3D bridging bond and 1D spreading bonds between ParB proteins constitutes a minimal model for the condensation of ParB proteins on DNA into a coherent complex [31], consistent with the observation that ParB-GFP fusion proteins form a tight fluorescent focus on the DNA [17, 21, 26, 27].

The statistical properties of the 3D structure of ParB-DNA complexes determines the binding profile of ParB on DNA, which can be accurately measured in ChIP-Seq experiments. However, it is computationally demanding to simulate these binding profiles with the spreading and bridging model. The protein binding profiles can be easily calculated analytically in the limit of strong protein-protein interactions, where the cluster of ParB on the DNA becomes compact with a corresponding triangular distribution of ParB along DNA. The protein binding profiles can also be estimated in the limit of weak protein-protein interactions with the so-called stochastic binding model, where a sphere of high ParB concentration is assumed to exist within which a DNA polymer freely fluctuates [27] (see figure 1(c)). The description of the average protein binding profile is thus similar to the return statistics of the polymer into the ParB sphere [34], suggesting a long range (power-law) distribution of ParB proteins along DNA. Importantly, however, neither of these two existing approaches provide a simple way of computing ParB binding profiles around *parS* sites over the full relevant range of system parameters. In addition, it remains unclear how the spreading and bridging model and the stochastic binding model relate to each other.

Here, we propose a theoretical approach to describe the distribution of ParB proteins around *parS* sites on the DNA in terms of molecular interaction parameters and protein expression levels. To this end, we develop a simple model for protein-DNA clusters that explicitly accounts for the competition between the positional entropy associated with placing the loops on the cluster, which favors a looser cluster configuration, and both protein-protein interactions and loop closure entropy, which tend to favor a compact cluster. This looping and clustering (LC) model represents a reduced, approximate version of the full spreading and bridging model that incorporates the key physical ingredients needed to provide a clearer understanding and at the same time greatly facilitates calculations of the distribution profile of ParB (or other proteins that form protein-DNA clusters). Thus, our approach can be used to estimate molecular interactions between proteins from experimentally determined protein binding profiles.

2. The looping and clustering (LC) model

To theoretically describe the protein binding profiles of ParB on DNA, we first consider a DNA polymer of length L that can move in space on a 3D cubic lattice and with a finite number of proteins m . Since the number of ParB proteins in the protein-DNA cluster has been observed to include the vast majority of proteins in the cell [27], we employ a canonical ensemble with a fixed number of ParB proteins m in the ParB complex. These proteins are able to diffuse along the DNA. Importantly, in this model the DNA itself is also dynamic and fluctuates between different three-dimensional configurations, which are affected by the presence of interacting DNA bound proteins. When proteins are bound to the DNA, they are assumed to be able to interact attractively with each other by contact interactions in two distinct ways: (i) 1D spreading interactions with coupling strength J_S , defined as an interaction between proteins on nearest-neighbor sites along the polymer, and (ii) a 3D bridging interaction with strength J_B between two proteins bound to sites on non-nearest neighbor-sites on the DNA, but which are positioned at nearest neighbor-sites in 3D space (see figures 1(a) and (b)). Thus, these bridging interactions couple to the 3D configuration of the DNA, while the 1D spreading interactions do not. Single-molecule experiments provide evidence for bridging bonds [32], with the bridging valency of a ParB protein limited to one [35, 36]. Even in this case where each protein can form two spreading bonds and a single bridging bond, the system has been shown to exhibit a condensation transition where the majority of the proteins form a single large cluster that can be localized by a single *parS* site on the DNA [31].

While it is possible to perform Monte Carlo simulations of the spreading and bridging model for a lattice polymer, such simulations are computationally demanding. In this paper, we aim to provide a simple analytical description for the average binding profile of proteins along the DNA (see right panels in figure 1). With this aim in mind, we can simplify our description by realizing that the configurations of ParB proteins along the DNA are more sensitive to J_S than to J_B , for sufficiently large J_B . While both spreading and bridging bonds are necessary for the condensation of all proteins into a single cluster, loop extrusion from the cluster is controlled by J_S , and such loop extrusion strongly impacts the binding profile of proteins on the DNA. Indeed, a loop can be extruded from the protein-DNA cluster by breaking a spreading bond, but without effecting the internal configuration of the bridging bonds. Therefore, we will assume that J_B is sufficiently large to maintain a coherent 3D protein-DNA cluster, leaving J_S as the main adjustable parameter in the model.

A contiguous 3D cluster of proteins on DNA with loops can effectively be represented graphically by a disconnected 1D cluster along the DNA, where connections in 3D between the 1D subclusters are implied, and domains of protein-free DNA within the disconnected 1D cluster represent loops that emanate from the 3D cluster (see figures 1(b) and (c)). We can describe this system by a reduced model for the effective 1D cluster in which we account for the entropy of the loops that originate from the protein-DNA cluster. In this model, the spreading bond energy set by the parameter J_S combined with the cost in loop closure entropy, competes with

the positional entropy for placing loops on the cluster and will therefore play a crucial role in determining the binding profile of ParB on DNA around a *parS* site.

To capture these effects, we propose the reduced LC model, which offers a simplified description of 3D protein-DNA clusters with spreading and bridging bonds. In this model a loop is formed whenever there is a gap between 1D clusters. We can make the connection between the gaps in the 1D cluster and the number of loops extending from the 3D cluster explicit by writing down the partition function for this model. The effective 1D cluster corresponding to a 3D cluster with m proteins and n loops has a multiplicity:

$$\Omega_{\text{cluster}} = \frac{(m-1)!}{(m-n-1)!n!}, \quad (1)$$

which counts the number of ways in which one can partition m proteins into $n+1$ subclusters in 1D. This multiplicity leads to a positional entropy of mixing, $S_{\text{cluster}} = \ln \Omega_{\text{cluster}}$, for placing n loops at $m-1$ possible positions (in units of k_B). Note, we do not explicitly include the number of ways in which the bridging bonds can be formed, since loop formation is not expected to substantially affect the possible configurations of bridging bonds. However, creating n loops will require breaking n spreading bonds, and the probability at equilibrium for this to occur will include a Boltzmann factor $\sim \exp(-nJ_S)$, where the interaction energy is expressed in units of $k_B T$. Within our simple description, we do not consider how the formation of a loop affects the full internal entropy of the protein-DNA cluster, but this can be expected to be a fixed number per loop that can be absorbed into J_S . Furthermore, the loops that are formed are assumed to be independent, and thus contribute to the loop closure entropy (in units of k_B) as [34]:

$$S_{\text{loop}} = -d\nu \sum_{i=1}^n \ln(\ell_i), \quad (2)$$

where d is the spatial dimension, ν is the Flory exponent, and the loop length is measured in units of the lattice spacing of the polymer a , which we take to be equal to the footprint of a ParB protein, e.g. 16 bp for the exogenous ParABS system of *E. coli* [27].

This entropy is obtained by considering both the loops formed within the protein cluster and the protein-free segment of DNA outside the cluster. Indeed, the number of configurations associated with loop i for a Gaussian polymer is given by $z^{\ell_i} \ell_i^{-d\nu}$ [34, 37], where z is the lattice coordination number. Therefore, there is also an extensive contribution to the entropy given by $k_B \ell_i \ln(z)$. However, when a loop of length ℓ_i forms, the same length of polymer is removed from the DNA outside of the cluster, which also results in a reduction of the entropy by $k_B \ell_i \ln(z)$. Thus, there is a precise cancelation between the extensive contribution to the entropy associated with the loop inside the cluster and the extensive contribution due to effectively shortening the DNA outside the cluster⁵.


It is now straightforward to write down the partition function of the LC model:

$$\begin{aligned} Z_{\text{LC}} &= \sum_{n=0}^{m-1} \frac{(m-1)!}{(m-n-1)!n!} \exp(-nJ_S) \int_{\ell_0}^{\ell_{\max}} d\ell_1 \ell_1^{-d\nu} \dots \int_{\ell_0}^{\ell_{\max}} d\ell_n \ell_n^{-d\nu} \\ &\xrightarrow{\ell_{\max} \rightarrow \infty} [1 + \exp(-J'_S)]^{m-1}, \end{aligned} \quad (3)$$

where $J'_S = J_S + \ln[\ell_0^{d\nu-1}(d\nu-1)] > J_S$ is a renormalized loop activation energy that includes the cost in loop closure entropy. All lengths are measured in units of the protein's footprint a , ℓ_0 is the lower cutoff of loop sizes and approximately represents the persistence length of DNA, and the bond interactions are in units of $k_B T$. In the partition function, we conveniently set the upper boundary of integration, ℓ_{\max} , to infinity. Strictly speaking, the upper boundary for ℓ_j should be $L - (m + L_j)$, where $L_j = \sum_{i=1}^{j-1} \ell_i$ represents the total accumulated loop length before loop j . In practice, however, for chromosomes, but arguably also for plasmids, $L \gg m$ and the probability to have a large loop is very small. For instance, if we consider the F-plasmid of *E. coli* with a length of 60 kbp, we have $L = 3750$ in units of the ParB footprint of 16 bp [27, 38]. For this system, Monte Carlo (MC) simulations (see appendix A) of the LC model, with $m = 100$ reveal that the average cumulated loop size is ≈ 500 for small couplings ($J_S = 1$) down to ≈ 25 for large couplings ($J_S = 4$), which in both cases is much less than the DNA length. Thus, for biologically relevant cases it is reasonable to assume that the length of the DNA polymer is much larger than the footprint of the whole protein complex on the DNA.

The LC model constitutes a simple statistical mechanics approach to describe how proteins assemble into a protein-DNA cluster with multiple loops. Next, we will include a *parS* site on the DNA, to which ParB proteins bind with a higher affinity than the other non-specific binding sites on the DNA. Our central aim is to compute the binding profile of ParB around this *parS* site.

⁵ Although this reasoning is not strictly true for self-avoiding polymers, it does hold if we adopt the usual approximation used in the Poland-Scheraga model for DNA melting that self-avoidance acts only within individual loops.



3. Profile of Par B for fixed number and sizes of loops

With our approach, we aim to quantitatively describe average ParB binding profiles, which are directly measurable by ChIP-Seq experiments. By fitting our model to such ChIP-Seq data, it would be possible to extract microscopic parameters such as the number of proteins in the ParB clusters and the protein–protein interaction parameters such as J_s . In this section, we will describe how to compute the ParB binding profile around this $parS$ site given a fixed number of loops with specified loop lengths. Then we will use the statistical mechanics framework provided above, to perform a weighted average over all possible loop numbers and sizes to arrive at a simple predictive theory for the ParB binding profile.

3.1. 1-loop binding profile

It is instructive to start our analysis of ParB binding profiles by first calculating the probability of ParB occupancy as a function of distance from the $parS$ site for the case of a protein-DNA cluster with only one DNA loop ($n = 1$) with fixed loop length ℓ . We will assume a fixed number m of ParB proteins in this 1-loop protein-DNA cluster, and that one of these proteins is bound to the $parS$ site at any time, as illustrated in figure 2. Thus, to calculate the 1-loop ParB binding probability, $P_1(s, \ell)$, at a distance s from $parS$, we need to consider all possible configurations of proteins in the protein-DNA cluster subject to these constraints.

First, we note that $P_1(s, \ell) = 0$ for $s > m + \ell$, because the 1D cluster can maximally extend to a distance $m + \ell$, which occurs when the 1D cluster adopts a configuration that lies entirely on one side of the $parS$ site. For a binding site at a distance $s < m + \ell$, the ParB binding probability is reduced, either by configurations where this site is located on the DNA loop within the 1D cluster, or by states where the 1D cluster adopts a configuration around the $parS$ site that does not extend to the binding site at s , placing this site outside the 1D cluster. To capture these effects, it is helpful to express $P_1(s, \ell)$ in terms of conditional probabilities:

$$\begin{aligned} P_1(s, \ell) &= P_1(s, \ell | \text{loop}(s)) p_{\text{loop}}(s) + P_1(s, \ell | \overline{\text{loop}}(s)) p_{\overline{\text{loop}}}(s) \\ &= P_1(s, \ell | \overline{\text{loop}}(s)) p_{\overline{\text{loop}}}(s), \end{aligned} \quad (4)$$

where ‘loop(s)’ represents a condition with probability $p_{\text{loop}}(s)$ corresponding to site s being part of a loop extruding from the cluster, i.e. an unoccupied site on the DNA within the protein cluster, as depicted in figure 2. The overbar here represents the complementary condition, and the expression above simplifies because $P_1(s, \ell | \text{loop}(s)) = 0$ by construction.

We can proceed to calculate the conditional probability, $P_1(s, \ell | \overline{\text{loop}}(s))$, by decomposing this contributions as a sum of probabilities of mutually exclusive configurations, which are conditioned by the location s' of the right edge of the 1D ParB cluster denoted as ‘end(s')’ (see figure 2). Then, we will take a continuous limit for the binding profile assuming $m \gg 1$, and express the binding profile $P_1(s, \ell | \overline{\text{loop}}(s))$ in terms of probabilities, $p_{\text{end}}(s')$, for the condition describing the position of the right edge of the cluster. Thus, we first write the conditional probability $P_1(s, \ell | \overline{\text{loop}}(s))$ for $s \geq 0$ (the case $s < 0$ is obtained by symmetry) as

$$\begin{aligned} P_1(s, \ell | \overline{\text{loop}}(s)) &= \sum_{s'=0}^{m+\ell} P_1(s, \ell | \overline{\text{loop}}(s), \text{end}(s')) p_{\text{end}}(s'), \\ &\approx \int_0^{m+\ell} ds' \Theta(s' - s) p_{\text{end}}(s'). \end{aligned} \quad (5)$$

Clearly, $P_1(s, \ell | \overline{\text{loop}}(s), \text{end}(s')) = 1$ when $s < s'$ and zero otherwise, and thus we have replaced this term by the Heaviside step function $\Theta(s' - s)$ and approximated the sum by an integral in the second line above.

To calculate $p_{\text{end}}(s')$, it is convenient to introduce two subclusters, 1 and 2, with m_1 and $m - m_1$ proteins respectively ($0 < m_1 < m$), such that cluster 1 with m_1 proteins is overlapping with $parS$, as shown in figure 2. Given two such subclusters, two equally likely situations can occur: (i) the leftmost cluster overlaps with $parS$, i.e. $m - m_1 + \ell \leq s' < m + \ell$ or (ii) the rightmost cluster overlaps with $parS$, i.e. $0 \leq s' < m_1$. This directly

allows us to construct the conditional probability to find the right edge of the whole system, such that one of the m_1 proteins in the cluster overlaps with *parS*

$$p_{\text{end}}(s'|m_1) = \frac{1}{2m_1} [\Theta(s' - (m - m_1 + \ell))\Theta(m + \ell - s') + \Theta(m_1 - s')], \quad (6)$$

where the prefactor $1/2$ comes from the equal probabilities to find the system in one of the two cases (i) and (ii). The conditions (i) and (ii) are encoded with a product of two unit step functions for (i) and a single step function for (ii). Each single realization can be obtained by shifting the position of the site in cluster 1 overlapping with *parS* and is equally likely, giving rise to an overall prefactor $1/m_1$. From this, we can obtain the full probability $p_{\text{end}}(s')$ by integrating over m_1 :

$$\begin{aligned} p_{\text{end}}(s') &\approx \int_1^{m-1} dm_1 p_{\text{end}}(s'|m_1) p(m_1) \\ &= \frac{\Theta(m + \ell - s')}{m(m-2)} [\Theta(1 - (m + \ell - s'))(m-2) \\ &\quad + \Theta(m + \ell - s' - 1)\Theta(s' - \ell - 1)(s' - \ell - 1) \\ &\quad + \Theta(1 - s')(m-2) + \Theta(s' - 1)\Theta(m - 1 - s')(m - 1 - s)], \end{aligned} \quad (7)$$

where we used $p(m_1) = 2m_1/(m(m-2))$, since the number of configurations to place cluster 1 is $\propto m_1$ and $m_1 \in [1, m-1]$. Using this expression for the normalized probability distribution for the right edge of the 1D cluster to be positioned at s' , we can compute the conditional probability in equation (5):

$$\begin{aligned} P_1(s, \ell | \overline{\text{loop}}(s)) &\approx \frac{\Theta(\ell + m - s)}{(m-2)m} \left[\frac{(m-2)^2}{2} \Theta(\ell - s + 1) + (m-2)\Theta(\ell + m - s - 1) \right. \\ &\quad + (m-2)(m + \ell - s)\Theta(s - m - \ell + 1) \\ &\quad + (m + \ell - s - 1)m - \ell + s - 3)\Theta(s - \ell - 1) \\ &\quad + \Theta(1 - s) \left(\frac{(m-2)^2}{2} \Theta(m - s - 1) + (m-2)(1 - s) \right) \\ &\quad \left. + \frac{(s - m + 1)^2}{2} \Theta(s - 1)\Theta(m - s - 1) \right]. \end{aligned} \quad (8)$$

To obtain the full 1-loop protein distribution (equation (4)), we first need to compute the probability for a site to not be part of loop,

$$p_{\overline{\text{loop}}}(s) = 1 - p_{\text{loop}}(s). \quad (9)$$


If the loop density, ρ , were uniform, we would simply have $p_{\text{loop}}^{\text{uni}}(s) = \ell \rho^{\text{uni}}(m, \ell) = \frac{\ell}{m + \ell}$, since the 1D cluster has a total length of $m + \ell$ with a single loop of length ℓ . This uniform condition would only apply if we randomly choose ℓ sites to be part of the loop and ignore the requirement that all these loop sites need to be neighboring. In a real cluster, however, we expect the loop density $\rho_{\text{loop}}(s)$ to be higher in the bulk of the 1D cluster than close to the *parS* site or the edges, because fewer loops can be formed near the *parS* site or near the boundaries of the 1D cluster, at which a protein must be bound by construction. In particular, we expect the loop density, $\rho(s, m, \ell) \propto \min(s, \ell, m + \ell - s)$, which measures the number of ways a site at s can be part of a loop. This results in the normalized probability

$$\begin{aligned} p_{\text{loop}}(s) &= \ell \rho(s, m, \ell) \\ &= \frac{\ell \min(s, \ell, m + \ell - s)}{\Theta(m - \ell)[\ell^2 + \ell(m - \ell)] + \Theta(\ell - m) \left(\frac{m + \ell}{2} \right)^2}. \end{aligned} \quad (10)$$

In the normalization of this expression we distinguish the cases where the loop is either smaller or larger than the number of proteins in the cluster. With equations (8) and (10), we have all the elements to calculate the 1-loop protein binding profile $P_1(s, \ell)$ from equation (4).

We investigated the binding profiles $P_1(s, \ell)$ predicted by this model for a selected set of parameters, as shown in figure 3. We only show $s > 0$ because of the symmetry of the binding profile. It is instructive to contrast these profiles with the triangular profile (black curve) for a cluster with no loops. As expected, the addition of loops widens the profile, allowing the tail of the distribution to extend out to a distance $m + \ell$. The widening of the binding profile is accompanied by a faster decay of the profile in the vicinity of *parS*, which crosses over to a flatter profile at distances $s > \ell$ due to additional contributions from configurations where the loop lies between the *parS* site and site s .

Interestingly, for large loop size the profile can even become non-monotonic with a slight increase near the far edges of the domain. These features of the profile reflect the reduced loop density near *parS* and near the far



edges of the cluster. Note that the integral under this curve remains constant for varying ℓ to conserve the number of particles in the cluster. To verify the validity of the analytical approximations leading to $P_1(s, \ell)$, we used exact enumeration as a benchmark. Overall, the numerics (dashed lines) and the analytics (solid lines) are in good agreement for the 1-loop case, as shown in figure 3. In the next section, we employ the approximate analytical expressions obtained above, to efficiently calculate the full binding profile averaged over all configurations.

4. Protein binding profiles and statistics of the LC model

Above we defined the LC model and calculated the binding profile of proteins around a *parS* site for a cluster with 1 loop with fixed length. Real protein-DNA clusters, however, are expected to fluctuate with new loops forming and disappearing continuously. To capture such fluctuations, we will use the expressions for the binding profile of a static cluster with fixed loop length together with a statistical mechanics description of the LC model to obtain *average* binding profiles for dynamic clusters, including an ensemble average over both the number of loops and the loop lengths.

To obtain a full binding profile averaged over all realizations, it is useful to investigate the statistics of loops that extend from the protein-DNA cluster and how these statistics are determined by the underlying microscopic parameters of the model. We start by considering the number of loops that extend from the cluster. Using the partition function in equation (3), it is possible to calculate the basic features of the LC model. For instance, the moments of the distribution of the number of loops are given by

$$\langle n^\alpha \rangle = \frac{1}{Z_{LC}} \sum_{n=0}^{m-1} n^\alpha \frac{(m-1)!}{(m-n-1)!n!} e^{-nJ_S} \left[\frac{\ell_0^{1-d\nu}}{d\nu-1} \right]^n. \quad (11)$$

From this, we find the the average loop number is given by

$$\langle n \rangle = (m-1) \frac{1}{1 + e^{J'_S}} \underset{m, J'_S \gg 1}{\sim} m e^{-J_S}, \quad (12)$$

where J'_S is the renormalized loop activation energy introduced in equation (3). The average loop number $\langle n \rangle$ is depicted in figure 4(a), demonstrating the exponential dependence on the spreading energy J_S . In figure 4(b), we plot $\langle n \rangle$ as a function of the total number of proteins m in the protein-DNA cluster. Indeed, we observe the expected linear dependence of the average loop number $\langle n \rangle$ on m over a broad range of parameters. These results illustrate how the average number of loops is determined by the competition between the effective renormalized loop activation energy, J'_S (including the cost in loop closure entropy), and the gain in the positional entropy of mixing (see appendix B).

The linear dependence on m in equation (12) reflects that loops are assumed to be able to form anywhere in the cluster in the LC model. However, one would naively expect that loops can only form at the surface of a 3D cluster, resulting in a dependence $\langle n \rangle \sim m^{2/3}$ for a compact, spherical cluster. However, Monte Carlo simulations of the full S&B model have revealed that the protein-DNA clusters are not compact [31], but rather




Figure 4. (a) Average number of loops, $\langle n \rangle$, as a function of the spreading coupling strength J_S obtained from equation (12). The different curves correspond to protein number $m = 50$ (black), $m = 100$ (red), $m = 200$ (green), and $m = 400$ (blue), with loop-size cutoff $\ell_0 = 10$. We observe an exponential decrease $\langle n \rangle \propto e^{-J_S}$ in accordance with equation (12). Inset: same data replotted with the expected dependence of the average loop number on m scaled out. (b) Average number of loops $\langle n \rangle$ as a function of m for $J_S = 1, 2, 3$, and 4. The behavior is linear as expected from equation (12). The prefactor that determines the vertical shift between the different curves scales with e^{-J_S} , as demonstrated in the inset of panel (b). (c) Average loop probability as a function of the genomic coordinate with $m = 200$ and $L = 4000$ for protein-DNA clusters with fluctuating loop number and loop lengths. Different curves correspond to different spreading couplings $J_S = 1, 2, 3$, and 4. The analytic approximation (symbols) using equation (15) for the loop density, averaged over different loop configurations with the appropriate Boltzmann factor as in equation (14), is compared to MC simulations (dashed curves) of the LC model (see appendix A).

have a surface that scales almost linearly in m , close to the behavior of the simplified LC model presented here. The non-compact nature of the protein-DNA cluster is perhaps not surprising because each protein can form only one bridging bond.

A closely related statistic is the average accumulated loop length $\langle \ell \rangle$. From the LC partition function, we notice that the loop length is completely decoupled from the coupling constant J_S and depends only on the upper

cutoff ℓ_{\max} . Therefore, the cumulated average loop length becomes

$$\langle \ell \rangle_{\ell_{\max} \gg \ell_0} = \frac{d\nu - 1}{2 - d\nu} \frac{\ell_{\max}^{2-d\nu}}{\ell_0^{1-d\nu}} \langle n \rangle, \quad (13)$$

where the factor in front of $\langle n \rangle$ represents the average length per loop. This prefactor induces a small algebraic dependence on ℓ_{\max} , in contrast to $\langle n \rangle$ which depends only on the lower cutoff ℓ_0 .

The loop statistics of protein-DNA clusters are not easily accessible in experiments. Instead, the most relevant results for which this model can provide insight come from ChIP-Seq experiments. These experiments yield data for the enrichment of bound ParB as a function of genomic position on the DNA, providing a measure of the average protein binding profile of ParB on DNA [26, 27]. In the LC model, the ParB density profile along DNA can be calculated from


$$P_{\text{LC}}(s) = \frac{1}{Z_{\text{LC}}} \sum_{n=0}^{m-1} \frac{(m-1)!}{(m-n-1)!n!} \exp(-nJ_S) \times \int_{\ell_0}^{\infty} d\ell_1 \ell_1^{-d\nu} \dots \int_{\ell_0}^{\infty} d\ell_n \ell_n^{-d\nu} P_n(s, \{\ell_i\}), \quad (14)$$

where Z_{LC} is given in equation (3). Here, $P_n(s, \{\ell_i\})$ represents the multiloop ParB binding profile with n loops of length $\{\ell_i\} = \{\ell_1, \dots, \ell_n\}$. For simplicity, we approximate this multiloop profile by the analytical 1-loop conditional probability, $P_1(s, \ell | \text{loop}(s))$, with the loop length equal to the accumulated loop length, i.e. $\ell \rightarrow \sum_i \ell_i$, weighted by the loop probability $p_{\text{loop}}(s, \{\ell_i\}) \approx \sum_{i=1}^n \ell_i \rho(s, m, \ell_i, \ell)$. In the expression for the loop probability, $\rho(s, m, \ell_i, \ell)$ is defined as the contribution to the loop density of a loop of length ℓ_i in a cluster of m proteins with a total accumulated loop length ℓ , and we neglected correlations between contributions from different loops. Furthermore, we approximate $\rho(s, m, \ell_i, \ell)$ by using a generalization of the 1-loop expression in equation (10),

$$\rho(s, m, \ell_i, \ell) \approx \frac{\min(s, \ell_i, m + \ell - s)}{\Theta(m + \ell - 2\ell_i)[\ell_i^2 + \ell_i(m + \ell - 2\ell_i)] + \Theta(2\ell_i - (\ell + m))\left(\frac{m + \ell}{2}\right)^2}. \quad (15)$$

In the analysis above, we aimed to capture the effects of multiple loops in a simple way by assuming statistical independence of the loops, and by using the analytical 1-loop expressions to approximate the impact of loop formation on the loop density and the ParB binding profile of the protein-DNA complex. To test the validity of these approximations, we performed MC simulations of the complete LC model. We find that the numerically obtained average loop probability is in reasonable agreement with our approximate expression for the multiloop density, as shown in figure 4(c). Thus, despite the simplicity of our approach, the analytical model provided here captures the essential features of looping in protein-DNA clusters.


The protein binding profile $P_{\text{LC}}(s)$ around a *parS* site is calculated by averaging the static binding profile for different total loop numbers and loop lengths using the Boltzmann factor (see equation (3)) from the LC model as the appropriate weighting factor. The resulting expression in equation (14) for the protein binding profile of a protein-DNA cluster is the central result of this paper. We use this expression to compute binding profiles for the full LC model, which are shown in figure 5 as a function of the distance s to *parS* for $m = 100, 200$, and 400 . By construction, the site $s = 0$ corresponding to *parS* is always occupied, and thus $P(s = 0) = 1$ for all values of the spreading energy J_S . This feature of the LC model captures the assumed strong affinity of ParB for a *parS* binding site. For $J_S = 4$, the binding profile converges to a triangular profile, implying a very tight cluster of proteins on the DNA with almost no loops. The triangular profile in this case results from all the distinct configurations in which this tight cluster can bind to DNA such that one of the proteins in the cluster is bound to *parS*, and therefore the probability drops linearly to 0 at $s \approx m$. The same triangular binding profile was observed for the S&B model in the strong coupling limit $J_S \rightarrow \infty$ [31]. Interestingly, as J_S becomes weaker, we observe a faster decrease of the binding profile near *parS* together with a broadening of the tail of the distribution for distances far from *parS*. This behavior results from the increase in the number of loops that extrude from the ParB-DNA cluster with decreasing spreading bond strength J_S . The insertion of loops in the cluster allows binding of ParB to occur at larger distances from *parS*. Thus, the genomic range of the ParB binding profiles is set by $s_{\max} \approx m + \langle \ell \rangle$, where the average cumulated loop length $\langle \ell \rangle$ is controlled by J_S (see equation (13)) and m . These results illustrate how the full average binding profile is controlled by the spreading bond strength J_S : the weaker J_S , the looser the protein-DNA cluster becomes, which results in a much wider binding profile of proteins around *parS*. In the limit $J_S \rightarrow 0$, the LC model quantitatively reduces to the statistics of non-interacting loops, as shown in the inset of figure 5. In this case, the binding profiles exhibit asymptotic behavior $P_{\text{LC}}(s) \propto s^{-d\nu}$ for large s , as in the stochastic binding model [27]. Interestingly, we observe a weaker scaling $P_{\text{LC}}(s)$ with s at intermediate genomic distances, which we attribute to the reduced loop density near *parS* (see figure 4(c)).



To investigate how the functional shape of the binding profile is determined by the total number of proteins in the cluster, we plot the binding probability versus the scaled variable s/m for $m = 100, 200$, and 400 , as shown in figure 6. For fixed J_S , the data approximately collapse onto a single curve as a function of the scaled distance s/m . This implies that the functional shape of the ParB binding profile is largely determined by the spreading bond strength J_S , while the number of proteins in the cluster determines the width of the profile.

5. Discussion

The LC model introduced here allows us to access the average binding profile of proteins making up a large 3D protein-DNA complex. In our model, the formation of a coherent cluster of ParB proteins is ensured by a




combination of spreading and bridging bonds between DNA bound proteins, which together can drive a condensation transition in which all ParB proteins form a large protein-DNA complex localized around a $parS$ site [31]. We do not assume, however, that this protein-DNA cluster is compact. Indeed, loops of protein-free DNA may extend from the cluster, which strongly influences the average spatial configuration of proteins along the DNA. In the LC model, the formation of loops in the protein-DNA cluster is controlled by the strength of spreading bonds, i.e. the bond between proteins bound to nearest neighbor sites on the DNA. Specifically, for every protein-free loop of DNA that extends from the cluster, a single spreading bond between two proteins within the cluster must be broken. Thus, if the spreading interaction energy, J_S , is sufficiently small, thermal fluctuations will enable the transient formation and breaking of spreading bonds, thereby allowing multiple loops of DNA to emanate from the protein cluster (see figure 1).

Conceptually, the spreading bond interaction determines how ‘loose’ the protein-DNA cluster is, which directly impacts the ParB binding profiles. When J_S is large, loop formation is unlikely, resulting in a compact protein-DNA cluster with a corresponding triangular protein binding profile centered around $parS$ [31]. At intermediate J_S , the protein-DNA cluster becomes looser with the formation of loops, resulting in a binding profiles that are more strongly peaked around $parS$ but with far-reaching tails. Importantly, the LC model enables us to establish a link between the spreading and bridging model and the stochastic binding model [27]. The first used a microscopic approach based on the types of interactions between proteins on the DNA polymer, while the second employed a more macroscopic approach based on the polymer configurations around a dense sphere of proteins. In the limit $J_S \rightarrow 0$, the LC model is consistent with the stochastic binding model with a profile of the form [27] given by $P(s) \propto s^{-d\nu}$ (inset figure 5(a)). Thus, the LC model offers a description for a broad parameter regime, connecting two limits investigated in preceding studies [27, 31]. Therefore, our work also resolves the apparent contradiction between these previous models.

The LC model, which we introduce to calculate the binding profile of ParB-like proteins on the DNA, is a simple theoretical framework similar to the Poland–Scheraga model for DNA melting [39, 40]. An important difference in the LC model with respect to the homogeneous Poland–Scheraga model, is that translational symmetry is broken due to the presence of a $parS$ site at which a protein is bound with a high affinity. Thus, the protein-DNA cluster can adopt a wide range of configurations as long as one of the proteins is bound to the $parS$ site. As a result, loops are effectively excluded in the vicinity of $parS$. The central new result of this work is a simple way of computing the protein binding profiles around such a $parS$ site in terms of molecular interaction parameters. We show that the binding profiles predicted by this model are sensitive to both the expression level of proteins and the spreading interaction strength, which directly controls the formation of loops in the protein-DNA cluster. The LC model predicts a profile in good quantitative agreement with binding profiles measured with ChIP-Seq on the F-plasmid of *E. coli*, as shown in figure 5(c). Importantly, from this analysis we extract the spreading interaction strength $J_S \approx 1k_B T$ and the number of proteins in the cluster $m \approx 400$. The value of m is consistent with estimates based on quantitative Western blots [27]. The quantity J_S is a more challenging quantity to estimate, but gel-shift or similar assays could be used on short DNA oligomers to estimate this spreading bond strength.

Our results also have implications for experiments that employ fluorescent labeling of DNA loci by exogenous ParBs [28, 29]. Indeed, our model can be used to investigate how the protein interaction strengths



determine the 3D structure and mobility of the ParB-DNA cluster, as well as the tendency of multiple ParB foci to adhere to each other. This model thus provides an insightful quantitative tool that could be employed to analyze and interpret ChIP-Seq and fluorescence data of ParB-like proteins on chromosomes and plasmids.

Acknowledgments

This project was supported by the German Excellence Initiative via the program NanoSystems Initiative Munich (NIM) (CPB), the Deutsche Forschungsgemeinschaft (DFG) Grant TRR174 (CPB), and the National Science Foundation Grant PHY-1305525 (NSW). The authors acknowledge financial support from the *Agence Nationale de la Recherche* (IBM project ANR-14-CE09-0025-01) and from the CNRS Défi Inphyniti (*Projet Structurant* 2015-2016). This work is also part of the program ‘Investissements d’Avenir’ ANR-10-LABX-0020 and Labex NUMEV (AAP 2013-2-005, 2015-2-055, 2016-1-024). We also thank J-Y Bouet for helpful comments on the manuscript.

Appendix A. Monte Carlo simulations and numerical integration procedures

A.1. Monte Carlo procedure

Using the partition function, we can formulate an effective 1D Hamiltonian for the LC model, which explicitly accounts for the spreading bonds and loop entropy:

$$\mathcal{H}_{\text{LC}} = -J_S \sum_{i=1}^{L-1} \phi_i \phi_{i+1} + d\nu \sum_{i=1}^n \ln(\ell_i + \ell_0). \quad (\text{A1})$$

This effective Hamiltonian is useful to perform Monte Carlo simulations of the model as a benchmark for the approximations performed in the analytical approach (see figure 7). The proteins are modeled as particles that bind/unbind onto sites of a one-dimensional lattice with free boundary conditions. The lattice size $L = 4000$ is chosen to prevent finite size effect for the range of proteins considered. Note that, in these MC simulations, the total size of the loops is limited to $L - m$.

The simulations are performed with the standard Metropolis rules:

1. Propose a move of a particle randomly chosen to a random empty site of the lattice (conserved order parameter). A MC iteration step consists of m attempts of move.
2. Calculate the difference of energy $\Delta\mathcal{H} = \mathcal{H}_f - \mathcal{H}_i$ between final and initial configurations.
3. If $\Delta\mathcal{H} < 0$, the move is accepted with probability 1, otherwise it is accepted with probability $\exp(-\beta\Delta\mathcal{H})$.

The system is set initially with all particles in a single cluster ($J_S = \infty$), and then thermalized to the actual J_S of the simulations ranging from 1 to 4 (see figure 7). The sampling starts after thermalization of the system (40 000 MC iterations). A sampling of the systems configuration is performed every 100 MC iterations. All MC averages have been performed over 10^7 configurations, $\nu = 0.588$, $L = 4000$ and $\ell_0 = 10$. The numerical

results of this Monte Carlo simulation are in good agreement with our approximate analytic results, as shown in figure 7.

A.2. Numerical integration

To evaluate the binding profile $P_{LC}(s)$, we proceeded as follows. We carried out the evaluation of the simplified expression in equation (14) using numerical summation and integration. We truncated the summation at $n = 15$, instead of going up to $m - 1$, based on the corresponding average number of loops of figure 5. Finally, we introduced an upper cutoff for the loop-length, $\ell_{\max} = 100$, instead of going up to infinity. We confirmed that shape of the binding profiles does not change significantly for higher values of $\ell_{\max} = 100$.

The numerical evaluation of the multidimensional integrals in equation (14) have been performed with an accuracy and precision of respectively 2 and 3 effective digits in the final results. We have carried out convergence tests of the curve shapes in order to assess our parameter choice and rule out numerical instabilities. All computations have been performed by routines written in the Wolfram Language and executed by the Mathematica software suite (version 10 and 11).

Appendix B. Formal connection between the LC model and a Lattice Gas with renormalized coupling

For $m, n \gg 1$ (thermodynamic limit), we can formulate a saddle point approximation to evaluate the partition function and $\langle n \rangle$, by approximating the entropic (factorial) term in Z (equation (11)) using the standard entropy of mixing for placing n loops on $m - 1$ possible sites. This approach gives physical insight into how the loop entropy contributes to a renormalized protein–protein interaction and how the competition between this renormalized interaction and the entropy of mixing controls $\langle n \rangle$. Taking the thermodynamic limit leads to a partition function:

$$Z' = \int_0^1 d\rho_\ell \exp[-(m-1)F_{\text{eff}}(\rho_\ell)], \quad (\text{B1})$$

where $\rho_\ell = n/(m-1)$ is the concentration of loops ($0 \leq \rho_\ell \leq 1$) and

$$F_{\text{eff}}(\rho_\ell) = \rho_\ell J'_S + \{[1 - \rho_\ell] \ln[1 - \rho_\ell] + \rho_\ell \ln[\rho_\ell]\} \quad (\text{B2})$$

an effective free energy where $J'_S = J_S + \ln \alpha_0$ is a loop activation energy renormalized by the cost in loop entropy with $\alpha_0 = \ell_0^{d\nu-1}(d\nu-1)$. In the limit $m \rightarrow \infty$, the approximate partition function Z' becomes exact and can be evaluated exactly in the saddle point approximation by minimizing F_{eff} . The solution, ρ_{SP} , to the saddle point equation, $dF_{\text{eff}}(\rho_\ell)/d\rho_\ell = 0$, is

$$\rho_{\text{SP}} = \frac{1}{1 + e^{J'_S}}. \quad (\text{B3})$$

The entropic contribution to F_{eff} (second term) vanishes at $\rho_\ell = 0$ and 1, and reaches a minimum at $\rho_\ell = 1/2$, which is the exact result for ρ_{SP} at vanishing renormalized loop activation energy J'_S because the entropy of mixing is then maximized. For $J'_S > 0$, ρ_{SP} decreases from 1/2 to vanish in the limit $J'_S \rightarrow \infty$ as $\rho_{\text{SP}} \rightarrow e^{-J'_S}$. In this limit only the no and one loop states contribute and the asymptotic behavior can be simply obtained by a series expansion of the partition sum and the corresponding expression for $\langle n \rangle$. The saddle point result leads to

$$n_{\text{SP}} = \frac{m-1}{1 + e^{J'_S}}, \quad (\text{B4})$$

which turns out to be the exact result, thanks to compensating errors, for $\langle n \rangle$ for finite m (which can be obtained by differentiating the exact Z with respect to J_S , see equation (12)). For example, for $\ell_0 = 10$, $d = 3$, and $\nu = 0.588$, $\alpha_0 = 4.437$ and $\ln(\alpha_0) = 1.49$, which is not negligible if $J_S = \mathcal{O}(1)$.

ORCID iDs

Chase P Broedersz  <https://orcid.org/0000-0001-7283-3704>

References

- [1] Dekker J, Marti-Renom MA and Mirny LA 2013 Exploring the three-dimensional organization of genomes: interpreting chromatin interaction data *Nat. Rev. Genet.* **14** 390–403
- [2] Scolari V F and Lagomarsino M C 2015 Combined collapse by bridging and self-adhesion in a prototypical polymer model inspired by the bacterial nucleoid *Soft Matter* **11** 1677–87
- [3] Dame RT, Tark-Dame M and Schiessel H 2011 A physical approach to segregation and folding of the *Caulobacter crescentus* genome *Mol. Microbiol.* **82** 1311–5

- [4] Jun S and Mulder B 2006 Entropy-driven spatial organization of highly confined polymers: lessons for the bacterial chromosome *Proc. Natl. Acad. Sci.* **103** 12388–93
- [5] Emanuel M, Radja N H, Henriksson A and Schiessel H 2009 The physics behind the larger scale organization of DNA in eukaryotes *Phys. Biol.* **6** 025008
- [6] Marenduzzo D, Micheletti C and Cook P R 2006 Entropy-driven genome organization *Biophys. J.* **90** 3712–21
- [7] Mirny L A 2011 The fractal globule as a model of chromatin architecture in the cell *Chromosome Res.* **19** 37–51
- [8] Umbarger M A, Toro E, Wright M A, Porreca G J, Bau D, Hong S H and Shapiro L 2011 The three-dimensional architecture of a bacterial genome and its alteration by genetic perturbation *Mol. Cell* **44** 252–64
- [9] Le T B, Imakaev M V, Mirny L A and Laub M T 2013 High-resolution mapping of the spatial organization of a bacterial chromosome *Science* **342** 731–4
- [10] Viollier P H, Thanbichler M, McGrath P T, West L, Meewan M, McAdams H H and Shapiro L 2004 Rapid and sequential movement of individual chromosomal loci to specific subcellular locations during bacterial DNA replication *Proc. Natl. Acad. Sci. USA* **101** 9257–62
- [11] Lagomarsino M C, Espéli O and Junier I 2015 From structure to function of bacterial chromosomes: evolutionary perspectives and ideas for new experiments *FEBS Lett.* **589** 2996–3004
- [12] Wang X, Llopis P M and Rudner D Z 2013 Organization and segregation of bacterial chromosomes *Nat. Rev. Genet.* **14** 191–203
- [13] Dillon S C and Dorman C J 2010 Bacterial nucleoid-associated proteins, nucleoid structure and gene expression *Nat. Rev. Microbiol.* **8** 185–95
- [14] Dame R T 2005 The role of nucleoid associated proteins in the organization and compaction of bacterial chromatin *Mol. Microbiol.* **56** 858–70
- [15] Benza V G, Bassetti B, Dorfman K D, Scolari V F, Bromek K, Cicuta P and Lagomarsino M C 2012 Physical descriptions of the bacterial nucleoid at large scales, and their biological implications *Rep. Prog. Phys.* **75** 076602
- [16] Johnson J, Brackley C A, Cook P R and Marenduzzo D 2015 *J. Phys.: Condens Matter* **27** 064119
- [17] Mohl D A and Gober J W 1997 Cell cycle-dependent polar localization of chromosome partitioning proteins in *Caulobacter crescentus* *Cell* **88** 675–84
- [18] Lim H C, Surovtsev I V, Beltran B G, Huang F, Bewersdorf J and Jacobs-Wagner C 2014 Evidence for a DNA-relay mechanism in ParABS-mediated chromosome segregation *Elife* **3** e02758
- [19] Banigan E J, Gelbart M A, Gitai Z, Wingreen N S and Liu A J 2011 Filament depolymerization can explain chromosome pulling during bacterial mitosis *PLoS Comput. Biol.* **7** e1002145
- [20] Ptacin J L, Lee S F, Garner E C, Toro E, Eckart M, Comolli L R and Shapiro L 2010 A spindle-like apparatus guides bacterial chromosome segregation *Nat. Cell Biol.* **12** 791–8
- [21] Le Gall A, Cattoni D I, Guilhas B, Mathieu-Demazière C, Oudjedi L, Fiche J B and Nollmann M 2016 Bacterial partition complexes segregate within the volume of the nucleoid *Nat. Commun.* **7** 12107
- [22] Walter J C, Dorignac J, Lorman V, Rech J, Bouet J Y, Nollmann M, Palmeri J, Parmeggiani A and Geniet F 2017 Surfing on protein waves: proteophoresis as a mechanism for bacterial genome partitioning *Phys. Rev. Lett.* **119** 028101
- [23] Vecchiarelli A G, Neuman K C and Mizuuchi K 2014 A propagating ATPase gradient drives transport of surface-confined cellular cargo *Proc. Natl. Acad. Sci.* **111** 4880–5
- [24] Jindal L and Emberly E 2015 Operational principles for the dynamics of the *in vitro* ParA-ParB system *PLOS Comput. Biol.* **11** e1004651
- [25] Surovtsev I V, Campos M and Jacobs-Wagner C 2016 DNA-relay mechanism is sufficient to explain ParA-dependent intracellular transport and patterning of single and multiple cargos *Proc. Natl. Acad. Sci.* **113** E7268–76
- [26] Breier A M and Grossman A D 2007 Whole genome analysis of the chromosome partitioning and sporulation protein Spo0J (ParB) reveals spreading and origin distal sites on the *Bacillus subtilis* chromosome *Mol. Microbiol.* **64** 703–18
- [27] Sanchez A, Cattoni D I, Walter J C, Rech J, Parmeggiani A, Nollmann M and Bouet J Y 2015 Stochastic self-assembly of ParB proteins builds the bacterial DNA segregation apparatus *Cell Syst.* **1** 163–73
- [28] Chen B, Guan J and Huang B 2016 Imaging specific genomic DNA in living cells *Annu. Rev. Biophys.* **45** 1–23
- [29] Saad H, Gallardo F, Dalvai M, Tanguy-le-Gac N, Lane D and Bystricky K 2014 DNA dynamics during early double-strand break processing revealed by non-intrusive imaging of living cells *PLoS Genet.* **10** e1004187
- [30] Rodionov O, Lobočka M and Yarmolinsky M 1999 Silencing of genes flanking the P1 plasmid centromere *Science* **283** 546–9
- [31] Broedersz C P, Wang X, Meir Y, Loparo J J, Rudner D Z and Wingreen N S 2014 Condensation and localization of the partitioning protein ParB on the bacterial chromosome *Proc. Natl. Acad. Sci.* **111** 8809–14
- [32] Graham T G, Wang X, Song D, Etson C M, van Oijen A M, Rudner D Z and Loparo J J 2014 ParB spreading requires DNA bridging *Genes Dev.* **28** 1228–38
- [33] Taylor J A, Pastrana C L, Butterer A, Pernstich C, Gwynn E J, Sobott F and Dillingham M S 2015 Specific and non-specific interactions of ParB with DNA: implications for chromosome segregation *Nucleic Acids Res.* **43** 719–31
- [34] De Gennes P G 1979 *Scaling Concepts in Polymer Physics* (Ithaca, NY: Cornell University Press)
- [35] Leonard T A, Butler P J and Löwe J 2005 Bacterial chromosome segregation: structure and DNA binding of the Soj dimer a conserved biological switch *EMBO J.* **24** 270–82
- [36] Fisher G L, Pastrana C L, Higman V A, Koh A, Taylor J A, Butterer A and Moreno-Herrero F 2017 The C-terminal domain of ParB is critical for dynamic dna binding and bridging interactions which condense the bacterial centromere bioRxiv:122986
- [37] Hanke A and Metzler R 2003 Entropy loss in long-distance DNA looping *Biophys. J.* **85** 167–73
- [38] Bouet J Y and Lane D 2009 Molecular basis of the supercoil deficit induced by the mini-F plasmid partition complex *J. Biol. Chem.* **284** 165–73
- [39] Poland D and Scheraga H A 1970 *Theory of Helix-Coil Transitions in Biopolymers* (New York: Academic)
- [40] Everaers R, Kumar S and Simm C 2007 Unified description of poly- and oligonucleotide DNA melting: nearest-neighbor, Poland-Sheraga, and lattice models *Phys. Rev. E* **75** 041918