

Spiranoid Withanolides from *Jaborosa odonelliana* and *Jaborosa runcinata*

Adriana M. Cirigliano and Rosana I. Misico

Departamento de Química Orgánica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Peabellón 2, Ciudad Universitaria (C1428EGA) Buenos Aires, Argentina

Reprint requests to Dr. Adriana M. Cirigliano. Tel/Fax: (54-11) 4576-3346.

E-mail: adrianac@qo.fcen.uba.ar

Z. Naturforsch. **60b**, 867–869 (2005); received December 28, 2004

Two new spiranoid withanolides, (20*R*,22*R*,23*R*)-12 β ,17 β ,22-trihydroxy-1-oxo-12,23-cycloergosta-2,24-dien-26,23-olide (**3**) and (23*R*)-5 α ,6 β ,12 β ,21-tetrahydroxy-1,22-dioxo-12,23-cycloergosta-2,17,24-trien-26,23-olide (**4**) were isolated from plants of *Jaborosa odonelliana* and *Jaborosa runcinata*, respectively. Compounds **3** and **4** were characterized by a combination of spectroscopic methods (1D and 2D NMR, MS) and molecular modelling.

Key words: *Jaborosa*, Withanolides, Jaborosalactone

Introduction

The withanolides are a group of natural C₂₈-steroidal lactones built on an intact or arranged ergostane framework that occurs mainly in plants of certain genera of *Solanaceae*. The first member of this group of compounds, withafarin A, was isolated from the well-known Indian medicinal plant, *Withania somnifera* [1] and its structure was fully elucidated by Lavie and coworkers in 1965 [2]. The withanolides exhibit a variety of biological activities as antifeedant, immunosuppressive and cancer chemoprevention activity [3].

Jaborosa Miers is a South American genus belonging to the *Solanaceae* family that comprises about 23 species, 11 of which are almost exclusively distributed in Argentina [4]. Previous studies on populations of *J. odonelliana*, *J. runcinata* and *J. araucana*, gave a group of withanolides with a spiranoid lactone ring in the side chain. The compounds isolated from *J. odonelliana* presented a 17,22-diol functionality and a 23*S* stereochemistry at the spiranoid center (e.g. Jaborosalactone P, **1**) [5], while those isolated from *J. runcinata* and *J. araucana* showed a 17(20)-ene-22-keto system and a 23*R* stereochemistry (e.g. Jaborosalactone 2, **2**) [6], in agreement with that found in **1**.

Studies on cancer chemopreventive activity of withanolides as inducers of phase II detoxification enzymes indicated that the spiranoid withanolides jaborosalactone P and jaborosalactone 1 were promising agents in terms of inducing potency and low toxicity [7].

Results and Discussion

Jaborosalactone 24 (**3**) was isolated as a minor component from the aerial part of *J. odonelliana*. The HREIMS showed a molecular ion corresponding to the formula C₂₈H₃₆O₆, whereas the EIMS showed peaks at *m/z* 299 (42) and 168 (31) corresponding to the cleavage between C-20-C-17 and C-23-C-12, distinctive for this type of structure [8]. ¹H and ¹³C NMR chemical shift values in jaborosalactone 24 (**3**) were closely related to those reported for jaborosalactone P (**1**) [5]. In the low-field end of the ¹H NMR spectrum signals at $\delta = 5.81$, 6.73 and 5.56 were assigned to H-2, H-3 and H-6, respectively, of a 1-oxo-2,5-dienewithanolide. The typical pattern of the spiranoid arrangement was inferred from the resonances of carbons 23–28 (Table 1) and the low-field ¹H resonances of methyls 27 and 28, observed as quartets (*J* = 1.0 Hz) due to their mutual homoallylic coupling. Despite these similarities, detailed comparison of the NMR spectral data of **1** and **3** showed small but clear differences in the ¹³C chemical shifts of the α , β -unsaturated- γ -lactone ring, and a downfield shift of H-22. This indicated that the side chain of jaborosalactone 24 (**3**) had an arrangement different from that found in **1**, possibly due to an inverted stereochemistry of the spiranoid center at C-23. The configuration at C-23 was confirmed to be *R* by comparison of the NOESY spectra of **1** and **3**. Thus, the correlation observed for the pair H-28/H-22 in the NOESY spectrum

Table 1. ^{13}C NMR spectral data of compounds **3** (125.77 MHz) and **4** (50.32 MHz) in CDCl_3 .

C	3	4	C	3	4
1	203.8	203.3	2	127.6	128.8
3	145.0	141.9	4	33.3	34.3
5	135.4	79.7	6	124.9	73.8
7	34.5	31.9	8	32.9	29.5
9	39.9	38.3	10	50.0	51.6
11	30.0	35.4	12	78.9	75.2
13	50.8	48.9	14	43.6	46.4
15	23.3	23.3	16	32.7	25.3
17	83.5	165.6	18	10.7	14.8
19	17.9	15.1	20	40.7	127.3
21	11.8	58.2	22	69.0	193.5
23	94.5	91.0	24	163.1	161.9
25	126.7	128.1	26	171.6	173.4
27	8.6	18.9	28	15.2	12.1

C multiplicities were determined from DEPT data.





Fig. 1. AM1 calculated structures of a) jaborosalactone **24** (**3**) and b) jaborosalactone **P** (**1**), indicating relevant NOEs observed.


of **3** was only possible in the $23R$ stereoisomer, while a correlation for the pair H-28/H-20 was observed in the NOESY spectrum of **1** in agreement with the $23S$ stereochemistry assigned to this compound (Fig. 1).

Jaborosalactone **25** (**4**), was isolated as a minor component from the aerial part of *J. runcinata*. The ^1H and ^{13}C NMR spectral data of **4** were very similar to those of jaborosalactone **2** (**2**) [6], however the absence of a singlet for H-21 in the high-field end of the ^1H NMR spectrum and the appearance of an AB quartet at 4.21–4.33 ppm indicated the presence of an isolated C-21 hydroxymethylene group. This functionality has been previously found in jaborosalactones **4** and **5** isolated from *J. runcinata* [6]. The ^{13}C NMR spectrum (Table 1) showed only four methyl groups that were coincident with C-18, C-19, C-27 and C-28 in **2**. The methylene signal at 58.2 ppm confirmed the presence of a hydroxyl group at C-21. The molecular ion was absent in the EIMS, but a peak at m/z 480 (1%) corresponding to the ion $[\text{M}-\text{H}_2\text{O}]^+$ was observed. A significant fragment was at m/z 355 (4%), corresponding to the cleavage between C-23 and C-22 and the



1 $23S$

3 $23R$



2 R=H

4 R=OH

subsequent loss of the γ -lactone ring and CO. The FABMS (glycerol) of jaborosalactone **25** (**4**) showed a $[\text{M}+1]^+$ ion at m/z 499 consistent with the formula $\text{C}_{28}\text{H}_{34}\text{O}_8$.

Experimental Section

General experimental procedures

^1H and ^{13}C NMR spectra were recorded on a Bruker AC-200 NMR spectrometer at 200.13 and 50.32 MHz, respectively, or a Bruker AM-500 at 500.13 and 125.77 MHz, respectively. Multiplicity determinations (DEPT-135) and 2D spectra (COSY-45 and NOESY) were obtained using standard Bruker software. Chemical shifts are given in (δ) downfield from TMS as internal standard. EIMS were collected on a VG Trio-2 mass spectrometer at 70 eV by direct inlet; FABMS and HREIMS (70 eV) were measured on a VG ZAB-BEQQ mass spectrometer. IR and UV spectra were measured on a Nicolet Magna 550 FT IR and a Hewlett-Packard 8451A spectrophotometer, respectively. AM1 calculations were performed with the MOPAC module in Chem3D 8.0 (Cambridge Soft). Melting points were taken on a Fisher-Johns apparatus and are uncorrected. HPLC separations were carried out on a YMC-Pack ODS-AQ column (250×10 mm ID) and a mixture of MeOH– H_2O (70:30) as eluant, with UV detection at 245 nm. Vacuum liquid chromatography (VLC) and column flash chromatography were carried out on Kieselgel 60-G (Merck) and Kieselgel S 0.040–0.063 mm, respectively. TLC analysis was performed on silica gel 60 F254 (0.2 mm thick).

Plant material

Whole *J. runcinata* plants were collected in March 1995 in El Jagüel, departamento Paraná, Entre Ríos Province, Argentina. Aerial parts of *J. odonelliana* were collected in April and December 1996 in Salta Province, Argentina. Voucher specimens of both species are deposited at the Museo Botánico, Universidad Nacional de Córdoba under No. CORD 248 (*J. runcinata*) and CORD No. 25540 (*J. odonelliana*).

Extraction and isolation

Leaves and stems (1.15 g) of *J. odonelliana* were extracted as previously described [8]. The residue obtained after evaporation of the combined extracts was initially fractionated by vacuum liquid chromatography using hexane-EtOAc mixtures of increasing polarity (100:0-0:100) as eluant.

The fraction eluted with hexane-EtOAc (40:60) was further fractionated by flash chromatography and purified by HPLC yielding jaborosalactone P (**1**) (176 mg) and jaborosalactone 24 (**3**) (3.4 mg).

The dried and pulverized aerial parts of *J. runcinata* (935 mg) were extracted and fractionated as previously described [6]. The fraction eluted with hexane-EtOAc (20:80) was further purified by flash chromatography using mixtures of CH₂Cl₂–MeOH (100:5-100:10) yielding jaborosalactone 25 (**4**) (4.5 mg).

Jaborosalactone 24 (3)

Amorphous solid. – UV/vis (MeOH): $\lambda_{\max}(\log \epsilon) = 223$ nm (3.07). – IR (dry film): $\tilde{\nu} = 3458, 2966, 1747, 1676, 1390, 1270, 1006, 728$ cm⁻¹; ¹H NMR (500.13 MHz, CDCl₃, assignments based on ¹H-¹H-COSY): $\delta = 1.06$ (s,

3H, 18-H), 1.16 (d, $J = 6.4$ Hz, 1H, 21-H), 1.16 (s, 3H, 19-H), 1.48 (m, 1H, 11 β -H), 1.80 (m, 1H, 7 α -H), 1.99 (q, $J = 1.0$ Hz, 3H, 27-H), 2.02 (m, 1H, 7 β -H), 2.09 (m, 1H, H-20), 2.15 (q, $J = 1.0$ Hz, 3H, 28-H), 2.46 (dd, $J = 10.4, 2.6$ Hz, 1H, 11 α -H), 2.83 (dd, $J = 21.0, 5.0$ Hz, 1H, 4 α -H), 3.24 (dt, $J = 21.0, 2.5$ Hz, 1H, 4 β -H), 4.03 (d, $J = 11.1$ Hz, 1H, 22-H), 5.56 (dd, $J = 4.1, 2.0$ Hz, 1H, 6-H), 5.81 (dd, $J = 10.0, 2.5$ Hz, 1H, 2-H), 6.73 (ddd, $J = 10.0, 5.0, 2.5$ Hz, 1H, 3-H). – ¹³C NMR (125.77 MHz) see Table 1. – MS (EI, 70 eV): m/z (%) = 468 (4) [M⁺], 450 (2), 432 (2), 299 (42), 283 (10), 265 (4), 168 (31), 107 (18), 97 (21), 43 (100); HREIMS $m/z = 468.2512$ [M⁺] (C₂₈H₃₆O₆, requires 468.2516).

Jaborosalactone 25 (4)

White solid. – Mp 253–255 °C. – UV/vis (MeOH): $\lambda_{\max}(\log \epsilon) = 226$ nm (3.25). – IR (dry film): $\tilde{\nu} = 3450, 1742, 1673, 1381, 1254, 1018$ cm⁻¹. – ¹H NMR (200.13 MHz, CDCl₃, assignments based on ¹H-¹H-COSY): $\delta = 1.14$ (s, 3H, 18-H), 1.26 (s, 3H, 19-H), 2.03 (q, $J = 1.0$ Hz, 3H, 27-H), 2.10 (dd, $J = 19.2, 5.0$ Hz, 1H, H-4 α), 2.25 (q, $J = 1.0$ Hz, 3H, 28-H), 2.64 (m, 1H, 16-H), 3.25 (dt, $J = 19.2, 2.2$ Hz, 1H, 4 β -H), 3.66 (t, $J = 2.6$ Hz, 1H, 6-H), 4.21 (d, $J = 12.2$ Hz, 1H, 21b-H), 4.33 (d, $J = 12.3$ Hz, 1H, 21a-H), 5.82 (dd, $J = 10.1, 2.2$ Hz, 1H, 2-H), 6.60 (ddd, $J = 10.1, 5.0, 2.2$ Hz, 1H, 3-H). – ¹³C NMR (50.32 MHz) see Table 1. – MS (EI, 70 eV): m/z (%) = 480 (1) [M⁺ - H₂O] (1), 355 (4), 107 (10), 97 (20), 43 (100); FABMS (glycerol) m/z 499 [M + H]⁺.

Acknowledgements

We thank the late Prof. A. T. Hunziker, Universidad Nacional de Córdoba, for identification of the plants. Financial support by CONICET (Argentina) and Universidad de Buenos Aires is gratefully acknowledged.

- [1] P. A. Kurup, *Current Sci.* **25**, 57 (1956).
 [2] D. Lavie, E. Glotter, Y. Shvo, *J. Chem. Soc.* 7517 (1965).
 [3] A. S. Anjaneyulu, D. S. Rao, P. W. Lequesne, *Stud. Nat. Prod. Chem.* **20**, 135 (1998).
 [4] A. T. Hunziker, G. Barboza, *Flora, Fanerogámica Argentina* **54**, 1 (1998).
 [5] E. S. Monteagudo, J. C. Oberti, E. G. Gros, G. Burton, *Phytochemistry* **29**, 933 (1990).
 [6] A. M. Cirigliano, A. S. Veleiro, G. M. Bonetto, J. C. Oberti, G. Burton, *J. Nat. Prod.* **59**, 717.
 [7] R. I. Misico, L. L. Song, A. S. Veleiro, A. M. Cirigliano, M. C. Tettamanzi, G. Burton, G. M. Bonetto, V. E. Nicotra, G. Silva, R. R. Gil, J. C. Oberti, A. D. Kinghorn, J. M. Pezzuto, *J. Nat. Prod.* **65**, 677 (2002).
 [8] A. M. Cirigliano, A. S. Veleiro, J. C. Oberti, G. Burton, *J. Nat. Prod.* **65**, 1049 (2002).