

Technical University of Denmark



## Comparative analysis of the microbial diversity in liquid and foaming layer in biogas reactors

**Campanaro, Stefano; Treu, Laura; Kougias, Panagiotis; De Francisci, Davide; Angelidaki, Irini**

*Published in:*

Proceedings of the 2nd International Conference on Biogas Microbiology

*Publication date:*

2014

[Link back to DTU Orbit](#)

*Citation (APA):*

Campanaro, S., Treu, L., Kougias, P., De Francisci, D., & Angelidaki, I. (2014). Comparative analysis of the microbial diversity in liquid and foaming layer in biogas reactors. In Proceedings of the 2nd International Conference on Biogas Microbiology

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Comparative analysis of the microbial diversity in liquid and foaming layer in biogas reactors

Stefano Campanaro<sup>2</sup>, Laura Treu<sup>1</sup>, Panagiotis G. Kougias<sup>1</sup>, Davide De Francisci<sup>1</sup>, Irini Angelidaki<sup>1</sup>

<sup>1</sup>Department of Environmental Engineering, Technical University of Denmark, Kgs. Lyngby, DK-2800, Denmark

<sup>3</sup>Department of Biology, University of Padua, Via U. Bassi 58/b 35131 Padova, Italy

Foaming incidents have been recorded in many biogas plants causing severe operational, economical and environmental problems (Kougias et al., 2014). However, the foaming phenomenon in biogas reactors fed with agro-industrial wastes has not been extensively investigated, especially with respect to the microbial composition of the digesters (Moeller et al., 2012). In the cited literature, it has been reported that specific microorganisms, which are mainly filamentous (e.g. *Gordonia species*, *Microthrix parvicella*), are attached to biogas bubbles and transferred to the air/liquid interface of sludge reactors or wastewater treatment works (Ganidi et al., 2009). Once these microorganisms accumulate on the liquid surface, they initiate biosurfactants production due to their metabolic activity, leading to the decrease of the surface tension and thus generate foaming. The aim of the present study was to investigate the microbial diversity in the liquid versus the foaming layer in manure-based biogas reactors suffering by foaming incidents in order to elucidate potential role and contribution of the microorganisms in foam promotion.

The experimental work was carried out in three thermophilic continuous stirred tank reactors (CSTR) fed with manure and supplemental amounts of lipids, proteins and carbohydrates. Once foaming was formed in the reactors, samples from the liquid and foaming layer were obtained and screened using 16S rDNA sequencing.

The results of these analyses revealed that there are indeed some species that significantly vary their relative abundance in the foaming layer compared to the liquid one (e.g. *Methanoculleus sp.*, *Dialister sp.*). However, based on the cited literature and to the best of our knowledge there was not a direct correlation of these species with foaming. Further investigation is needed in order to define the properties of these species on foam generation. Finally, it was observed that particles of barley plant (that was contained in the raw manure as ingredient of animal nutrition) were accumulated in the foaming layer. It has been previously documented that barley contributes in stabilization of beer foam due to the activity of one of its proteins (Brey et al., 2003). For that reason, it could be hypothesised that the existence of barley particles in our reactors could contribute in foaming although their presence was also prior to the foaming incidents; their effect on foam formation or stabilization might be enhanced in correlation with other parameters (i.e. presence of specific microorganisms).

### References

1. Brey, S. E., de Costa, S., Rogers, P. J., Bryce, J. H., Morris, P. C., Mitchell, W. J., Stewart, G. G., 2003. The Effect of Proteinase A on Foam-Active Polypeptides During High and Low Gravity Fermentation. *Journal of the Institute of Brewing*. 109, 194-202.
2. Ganidi, N., Tyrrel, S., Cartmell, E., 2009. Anaerobic digestion foaming causes – A review. *Bioresource Technology*. 100, 5546-5554.
3. Kougias, P. G., Boe, K., Tsapekos, P., Angelidaki, I., 2014. Foam suppression in overloaded manure-based biogas reactors using antifoaming agents. *Bioresource Technology*. 153, 198-205.
4. Moeller, L., Goersch, K., Neuhaus, J., Zehnsdorf, A., Mueller, R., 2012. Comparative review of foam formation in biogas plants and ruminant bloat. *Energ Sustain Soc*. 2, 1-9.

P6

**Conference topic: - Microbial inhibition and adaptation to perturbations POSTER**  
**PRESENTATION**