

## Environmental controls on ozone fluxes in a poplar plantation in Western Europe

**Zona, D.; Gioli, B.; Fares, S.; De Groot, T.; Pilegaard, Kim; Ibrom, Andreas; Ceulemans, R.**

*Published in:*  
Environmental Pollution

*Link to article, DOI:*  
[10.1016/j.envpol.2013.08.032](https://doi.org/10.1016/j.envpol.2013.08.032)

*Publication date:*  
2014

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Zona, D., Gioli, B., Fares, S., De Groot, T., Pilegaard, K., Ibrom, A., & Ceulemans, R. (2014). Environmental controls on ozone fluxes in a poplar plantation in Western Europe. *Environmental Pollution*, 184(Sp. Iss. SI), 201-210. DOI: 10.1016/j.envpol.2013.08.032

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Environmental controls on ozone fluxes in a poplar plantation in Western Europe<sup>☆</sup>



D. Zona <sup>a, b, \*</sup>, B. Gioli <sup>c</sup>, S. Fares <sup>d</sup>, T. De Groote <sup>a, e</sup>, K. Pilegaard <sup>f</sup>, A. Ibrom <sup>f</sup>, R. Ceulemans <sup>a</sup>

<sup>a</sup> Department of Biology, Research Group of Plant and Vegetation Ecology, University of Antwerp, Wilrijk, Belgium

<sup>b</sup> Department of Animal and Plant Sciences, University of Sheffield, Sheffield, UK

<sup>c</sup> National Research Council (CNR), Biometeorology Institute, Florence, Italy

<sup>d</sup> Consiglio per la Ricerca e la sperimentazione in Agricoltura (CRA) – Centro di ricerca per lo studio delle Relazioni tra Pianta e Suolo, Rome, Italy

<sup>e</sup> Flemish Institute for Technological Research (VITO), Mol, Belgium

<sup>f</sup> Centre for Ecosystems and Environmental Sustainability, Department of Chemical and Biochemical Engineering, Technical University of Denmark, Roskilde, Denmark

## ARTICLE INFO

### Article history:

Received 24 March 2013

Received in revised form

6 August 2013

Accepted 9 August 2013

### Keywords:

Eddy covariance

AOT40

Stomatal O<sub>3</sub> uptake

Deposition velocity

## ABSTRACT

Tropospheric O<sub>3</sub> is a strong oxidant that may affect vegetation and human health. Here we report on the O<sub>3</sub> fluxes from a poplar plantation in Belgium during one year. Surprisingly, the winter and autumn O<sub>3</sub> fluxes were of similar magnitude to ones observed during most of the peak vegetation development. Largest O<sub>3</sub> uptakes were recorded at the beginning of the growing season in correspondence to a minimum stomatal uptake. Wind speed was the most important control and explained 44% of the variability in the nighttime O<sub>3</sub> fluxes, suggesting that turbulent mixing and the mechanical destruction of O<sub>3</sub> played a substantial role in the O<sub>3</sub> fluxes. The stomatal O<sub>3</sub> uptake accounted for a seasonal average of 59% of the total O<sub>3</sub> uptake. Multiple regression and partial correlation analyses showed that net ecosystem exchange was not affected by the stomatal O<sub>3</sub> uptake.

© 2013 The Authors. Published by Elsevier Ltd. All rights reserved.

## 1. Introduction

Tropospheric ozone is a very strong oxidant, with the potential to damage human health, and vegetation functioning (Fowler, 1992; Lefohn, 1992; Wittig et al., 2007; Royal Society, 2008). Over the last 100 years tropospheric ozone pollution has significantly increased (Vingarzan, 2004; Derwent et al., 2007), and [O<sub>3</sub>] is predicted to continue rising in the next decades (IPCC-DDC, 2004; Vingarzan, 2004; Royal Society, 2008). Importantly, because of a more stringent regulation in the emission limits of ozone precursors in both the USA and Western Europe, peak ozone concentrations are decreasing in these areas (Vingarzan, 2004; Solberg et al., 2005; Derwent et al., 2007; Jenkin, 2008; Lefohn et al., 2008). Background concentrations are however increasing due to the transport of precursors from the developing world (NEGTA, 2001; Vingarzan, 2004; Dentener et al., 2006; Davis et al., 2010).

These background concentrations are forecast to increase to more than 60 ppb over the next century (IPCC-DDC, 2004) making it extremely challenging to maintain clean air standards (e.g. [O<sub>3</sub>] < 80 ppb, IPCC, 2001) in densely populated areas (IPCC, 2001; Vingarzan, 2004). The observed increase in [O<sub>3</sub>] has already been proved to affect agricultural yield (Benton et al., 2000; Fumagalli et al., 2001), and to lead to biomass/yield reduction in crops in Europe (Mills et al., 2011), potentially reducing the global carbon sink of terrestrial vegetation (Sitch et al., 2007).

Within Europe, high levels of radiation and warmer temperatures, together with the significant anthropogenic sources of ozone precursors, could make the southern ecosystems among the most exposed to O<sub>3</sub> (Cieslik and Labatut, 1997; Nolle et al., 2002; Paoletti, 2006; Manes et al., 2007). However, as ozone impact on vegetation depends on the ozone entering the stomata and oxidizing plant tissues, stomatal ozone uptake better describes the possible effect to the vegetation than ambient ozone concentrations (Emberson et al., 2000; Uddling et al., 2004; Matyssek et al., 2007). Stomatal ozone uptake is the highest under high soil water availability, and consequently ozone can be more damaging in non-water limited ecosystems in Central and Northern Europe (Emberson et al., 2000; Cieslik, 2009; Karlsson et al., 2009; Marzuoli et al., 2009; Zapletal et al., 2011), even with lower [O<sub>3</sub>] (Emberson et al., 2000; Mills

<sup>☆</sup> This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-No Derivative Works License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

\* Corresponding author.

E-mail addresses: [D.zona@sheffield.ac.uk](mailto:D.zona@sheffield.ac.uk), [donatellazona@yahoo.it](mailto:donatellazona@yahoo.it) (D. Zona).

et al., 2011). However, long-term ozone flux measurements of vegetation are still not very frequent (Fowler et al., 2001; Mikkelsen et al., 2000, 2004; Cieslik, 2009; Mills et al., 2011; Rannik et al., 2012), leaving uncertainties about the impact of this pollutant on net ecosystem exchange (NEE) and on plant productivity. In general, the majority of the studies on ozone impacts on vegetation have been carried out in controlled environments where plants are subjected to ozone treatments (e.g. Bortier et al., 1999; Bortier et al., 2000; Novak et al., 2003, 2005; Karlsson et al., 2003a; Wittig et al., 2007; Bussotti et al., 2011). Ideally, ozone impact should be studied under natural conditions (Pilegaard et al., 1998; Kolb and Matyssek, 2001; Gerosa et al., 2003; Fares et al., 2010a). However, such conditions make it a challenge to separate the impact of ozone on carbon assimilation from other factors (such as drought and temperature stress).

The impact of ozone on vegetation has usually been quantified using the AOT40 index, which sums the differences between the hourly mean  $[O_3]$  above 40 ppb during daylight hours (Appendix A, Kärenlampi and Skärby, 1996). A more valid approach is to define the amount of ozone entering the leaves through the stomates (Fuhrer, 2000; Emberson et al., 2000; Matyssek et al., 2007; Tuovinen et al., 2007). This stomatal ozone uptake has been described with different criteria in the literature, such as the accumulated stomatal fluxes ( $AF_{ST}$ , Mills, 2004; Karlsson et al., 2004; currently indicated as  $POD_{Ygen}$ , ICP, 2004), leaf cumulative uptake (CUO) over a threshold (for example cumulative ozone uptake above  $1.6 \text{ nmol m}^{-2}\text{s}^{-1}$  ( $CUO > 1.6$ , Karlsson et al., 2003b, 2004), and accumulated ozone dose (AOD) (Pollastrini et al., 2010).

Among different plant species, fast-growing poplars are especially susceptible to ozone impacts, because of the high stomatal conductance and thus high ozone uptake (Pye, 1988; Bortier et al., 1999; Novak et al., 2003, 2005). Several studies reported that ozone damage affects the photosynthetic apparatus, leading to visible leaf injury, damage to photosystems and growth reduction in poplar (Bortier et al., 1999, 2000; Oksanen et al., 2001; Fares et al., 2006; Marzuoli et al., 2009; Pollastrini et al., 2010; Bussotti et al., 2011). The sensitivity of poplar to ozone makes the investigation of the environmental controls on ozone uptake particularly relevant. The aim of this study is to investigate the environmental controls on ozone fluxes in a high-density poplar plantation, to quantify the stomatal ozone uptake, and the effect of this ozone uptake on NEE during the growing season.

## 2. Materials and methods

### 2.1. Site description

An eddy covariance (EC) tower was installed in a poplar plantation in Lochristi, Belgium ( $51^{\circ}06'44'' \text{ N}$ ,  $3^{\circ}51'02'' \text{ E}$ ) in spring/summer 2010. The plantation (18.4 ha) was established in April 2010 with different poplar clones of *Populus deltoides*, *P. maximowiczii*, *P. nigra*, and *P. trichocarpa*, and interspecific hybrids. The climate is maritime with long-term average annual temperature of  $9.5^{\circ}\text{C}$ , and precipitation of 726 mm that is uniformly distributed over the year (Royal Meteorological Institute of Belgium, <http://www.meteo.be>). During the second growing season (2011) after the establishment of the plantation, the canopy height increased from  $2.4 \pm 0.07 \text{ m}$  in April to  $4.6 \pm 0.14 \text{ m}$  in October. The understory of the plantation was typically composed of bare ground with few weeds, mostly thistles. More details on the plantation establishment and development, and on the management of the site, etc. are provided in Broeckx et al. (2012).

### 2.2. Eddy covariance (EC) system

The EC system was composed of a 3-D sonic anemometer (CSAT 3-D, Campbell Sci., UT, USA) and of several fast-response analyzers, including a LOZ-3F  $O_3$  analyzer (Drummond Technology Inc., Ontario, Canada) and a  $CO_2/H_2O$  infrared analyzer (LI-7000, LI-COR, Lincoln, NE, USA). The sonic anemometer was used to calculate the momentum and the sensible heat fluxes. More details on these analyzers are presented in Zona et al. (2013a) and Zona et al. (2013b). The LOZ-3F  $O_3$  analyzer is based on chemiluminescence with Eosin-Y dye (Ray et al., 1986; Topham et al., 1992; Rannik et al., 2012) circulated continuously through a peristaltic pump in the

sample cell, and measures  $O_3$  mixing ratio at a 10 Hz sampling frequency. The instrument performs an automatic zero check every hour for about 10 s, then saves the signal acquired during this zeroing mode, and subtracts it from the signal during the following measuring mode. After the zeroing correction, the instrument applies the appropriate calibration factors to output the final corrected mixing ratio (in ppb). A slow response  $O_3$  analyzer (API 400E, Teledyne Instruments, CA, USA) was used continuously to monitor  $[O_3]$ , which was then compared with the concentrations measured by the LOZ-3F, to control whether any drifting occurred in the signal of the LOZ-3F. The API was calibrated every six months by the Flemish Environment Agency (VMM).

The LI-7000  $CO_2/H_2O$  analyzer was used to measure  $CO_2$  and  $H_2O$  mixing ratios, to calculate  $CO_2$  fluxes (NEE) and latent heat (LE) fluxes. Two sampling lines (Teflon tubing about 12 m long and 8 mm inner diameter) were used for the EC system, one for the LI-7000 and another separate line for the two  $O_3$  analyzers (API and LOZ-3F). The two inlets were positioned about 10–15 cm from the center of the sonic anemometer. A  $1 \mu\text{m}$  teflon filter (Gelman) was placed 10 cm after the inlet of the sampling line of the LI-7000. No filter was placed in front of the inlet line of the  $O_3$  analyzers. A coarser filter (20–30  $\mu\text{m}$ , Zitex) was placed in front of the inlet of the LOZ-3F analyzer, and of the API analyzer (5  $\mu\text{m}$ , Savillex), to avoid contamination of the cells. The flow of the main line of the two  $O_3$  analyzers was  $20\text{--}27 \text{ l min}^{-1}$ . As described in Pilegaard et al. (1998), the flow rate used was sufficiently high to maintain a turbulent flow needed to limit possible errors due to tube attenuation (e.g. Reynolds number  $Re \geq 3000$ , Lenschow and Raupach, 1991). A mass flow controller was used to regulate the flow through all these sampling lines. Prior to the inlets of the two instruments, a Teflon manifold was used to split the air sampled into three parts, as described in Pilegaard et al. (1998); one line was connected to the slow response API analyzer (with flow of about  $0.76 \text{ l min}^{-1}$ ), one to the LOZ-3F (with a flow of about  $1.9 \text{ l min}^{-1}$ ), and the last one was used as a bypass and regulated with a mass flow controller. After the splitting of the flow, the outlets from the instruments were connected again to the main flow outlet. The flow meter inside the LOZ-3F was continuously operated and monitored, assuring a constant and stable flow which, together with the mass flow controller, prevented pressure fluctuations. The LOZ-3F measures cell pressure and temperature, and provides actual mixing ratio thus not requiring any further temperature and vapor density corrections. A buffer volume was placed between the main flow outlet and the pump to dampen the fluctuation of the pump.

### 2.3. Data processing and filtering of the $[O_3]$ and $O_3$ fluxes

EdiRe (R. Clement, University of Edinburgh, U.K.) and EddyPro (LI-COR, Lincoln, NE, USA, Fratini et al., 2012) were used for the computation of the fluxes of  $CO_2$ , latent heat (LE), sensible heat (H), and momentum. EddyPro was used to calculate the  $O_3$  fluxes. The cross-comparison of LE and  $CO_2$  fluxes calculated using these two software packages (EdiRe and EddyPro) showed very similar results, within 1–2% (data not shown). Consequently,  $CO_2$  fluxes (NEE) calculated using EdiRe (Zona et al., 2012, 2013), and  $O_3$  fluxes calculated using EddyPro were used in this paper.


The mean lateral and vertical velocity components were set to zero by application of a two-component rotation. The maximum cross covariance function (within 30 min averaging time) was used to determine the lag time for  $O_3$ , which was about 2 s. A drop-out filter was applied to remove the zeroing periods (10 s every hour) from the raw 10 Hz data. These removed data accounted for a total of only one day per year, limiting biases to the flux computation. An additional pre-processing routine was applied to remove any drift in the  $[O_3]$  of the fast-response analyzer. The  $[O_3]$  measured by LOZ-3F was compared to the  $[O_3]$  measured by the API analyzer (both measured at 5.8 m above the surface from January to 31 August 2011, and at 6.6 m thereafter), and the appropriate offsets and multipliers were applied to the half-hourly averaged 10 Hz data. This calibrated 10 Hz dataset was then used for the final  $O_3$  flux calculation. As the reaction cell of the LOZ-3F contains a liquid solution of Eosin-Y, eliminating water vapor interference, and as the  $[O_3]$  were expressed as mixing ratios, i.e. related to dry air, no Webb, Pearman and Leuning (WPL, Webb et al., 1980) correction was needed, as described in Ibrom et al. (2007). Frequency response correction was performed according to Moncrieff et al. (1997), which is an analytical correction modified from the Kaimal formulation (Kaimal et al., 1972). A high-pass filtering correction, as described in Moncrieff et al. (2004), was applied to the  $O_3$  fluxes in the low frequency range (high-pass filtering effect). A low-pass filtering correction was applied as described in Ibrom et al. (2007), and a sensor separation correction according to Horst and Lenschow (2009).

Data were removed if they matched one of the following filtering criteria: during malfunctioning of the instrument (22% of the entire dataset), during calibration, replacement of the filters, and maintenance of the instrument (which together accounted for  $\sim 0.03\%$  of the data), when the half-hour average  $[O_3]$  presented more than four spikes (when the standard deviation was  $>8$ ,  $\sim 0.03\%$  of the data), when the  $[O_3]$  was more than 500 ppb (unrealistic values,  $\sim 2\%$  of the data, when the wind direction was outside of the footprint of interest (i.e.  $>250$  and  $<50^{\circ}$ , 35% of the data). As no standardized gapfilling methodology has been developed for eddy covariance  $O_3$  fluxes, the fluxes were gap-filled using a feed-forward back propagation Artificial Neural Network (ANN) (nntool on MATLAB, 2011; Mathworks, Natick, Massachusetts, USA) as described in Papale and Valentini (2003). The meteorological variables used as input for the ANN were: air temperature (air T),

relative humidity (RH), wind speed, rainfall, short-wave and long-wave radiation, vapor pressure deficit (VPD), and  $[O_3]$ . The ANN had one layer (and 10 neurons) and was trained with 60% of the data and validated with 20% of the data; another 20% of the data was used for independent testing. The training data were used to build the model, which were then used to gap-fill the entire dataset; the validation and independent test data were not used to build the model, but used to check the performance of the gap-filling procedure. The training and validation presented  $R^2 = 83\%$  and  $86\%$ , respectively. Because of good performance of the model, the chosen percentage of training and validation data were considered reasonable. The data cleaning and gap-filling of the slow response analyzer were performed

according to the following procedure: negative  $[O_3]$  (caused by malfunctioning of the instrument) were removed and if the resulting gaps were equal to or less than 3 h, the  $[O_3]$  were gap-filled by linear interpolation; for gaps longer than 3 h, the gaps were replaced by the  $[O_3]$  measured at a nearby station (Admiraaldruff, Destelbergen, Belgium, Flemish Environment Agency, VMM), 7.5 km from our research site and in a similar rural setting. The stomatal uptake (indicated as  $F_{O3ST}$  hereafter) estimation and data filtering are described in Appendix A.

Concentrations of NO and  $NO_2$  were measured in three stations in the proximity of Gent (i.e. Baudelstraat in Gent; Admiraaldruff in Destelbergen; Schuitstraat in Sint-Kruis-Winkel), using a TRACE Level  $NO_x$  Analyzer (Model 421-TL, Thermo



**Fig. 1.** Seasonal trend in daily averaged short-wave radiation ( $W m^{-2}$ ), daily averaged vapor pressure deficit (VPD, kPa), daily averaged air temperature (air T,  $^{\circ}C$ ) and daily total rainfall (mm), daily averaged  $NO_2$  and NO (average from three stations: Baudelstraat, Gent; Admiraaldruff, Destelbergen; Schuitstraat in Sint-Kruis-Winkel, Belgium), and daily averaged  $[O_3]$  (at the field site) in 2011.




Fig. 2. Average diurnal trends in ozone concentrations ( $[O_3]$  ppb) during each of the indicated months. Displayed are ensemble averages and their standard error of the mean.

Scientific Waltham, MA, USA), at about 3 m above the ground, by the Flemish Environment Agency (VMM). The Admiraalreef and Sint-Kruis-Winkel were in more rural settings than Baudelostraat. The concentrations of  $NO_x$  reported in this paper are averages of the measurements in these three stations.

#### 2.4. Statistical analysis

The non-gap-filled ozone fluxes and the deposition velocity (defined as  $-F_{O_3}/[O_3]$ ), were modeled as a function of several environmental variables (air T, RH, VPD, wind speed, short and long-wave radiation) using a general linear model, GLM (Systat version 13, Systat Software Inc., 2002) to test the importance of each variable as control on the fluxes and on the deposition velocity. We performed this analysis on the entire dataset (daytime and nighttime data together) and on the data from daytime and nighttime separately (when short wave radiation was either  $\geq 20 \text{ W m}^{-2}$  or  $< 20 \text{ W m}^{-2}$ ). These analyses were performed on the ozone fluxes and the deposition velocity averaged on a half-hour time scale. The impact of  $F_{O_3ST}$  was estimated using multiple regression and partial correlation analyses of the residuals from 14 days, derived from a non-linear light-response fits of NEE (see Appendix A for more details). For this analysis a Michaelis–Menten type light response model was used as described in Pilegaard et al. (2011).

### 3. Results

#### 3.1. Ozone concentration and environmental conditions

The annual course of the daily averages of short wave radiation, air temperature, precipitation, VPD, and concentrations of  $NO_x$  and  $O_3$  is shown in Fig. 1. The environmental conditions in April and May 2011 were abnormally dry, with a precipitation (9 mm and 16.8 mm in April and May, respectively) much lower than what usually observed in this area (e.g. the ten year mean of 32.1 mm and 42.1 mm in April and May, respectively). As low precipitation is usually related to high ozone production (Comrie, 1990; Davis et al., 2010), the abnormally dry weather conditions during spring contributed to the fairly high  $[O_3]$  (Fig. 1). A spring maximum in  $[O_3]$  is typical of background sites in the Northern Hemisphere (Scheel et al., 1997; Angle and Sandhu, 1989; Meagher et al., 1987), where the absence of a summer peak may indicate low influence

from the local photochemical ozone production from precursor emission (Hough and Derwent, 1990).

However, the difference between the mean diurnal minimum and maximum  $[O_3]$  in our field site was 30 ppb or larger during June–August (Fig. 2), much higher than the 10 ppb observed in conditions not affected by local combustion (Mikkelsen et al., 2004). A spring  $[O_3]$  peak has also been related to the photochemical reaction of  $NO_x$  originating from combustion processes, with increase in solar radiation during spring (Dibb et al., 2003). On the other hand, the long-term (ten years)  $[O_3]$  data from another monitoring site in Belgium (Brasschaat), about 60 km from this field site, showed a summer maximum in  $[O_3]$  (Neiryck et al., 2012). This suggests that abnormally dry conditions in spring 2011 may have led to unexpectedly high  $[O_3]$ . An early season  $[O_3]$  peak may





Fig. 3. Cospectra of the fluctuation in vertical wind velocity and in the sonic temperature ( $w'Ts'$ ) and in  $O_3$  ( $w'O_3'$ ) averaged for June–July 2011 (in logarithmic scale) as function of the sampling frequency (Hz).

potentially be very damaging to the vegetation, if it occurs when vegetation is not water limited and therefore stomatal  $O_3$  uptake is maximum (Novak et al., 2005). The maximum  $[O_3]$  recorded in our field site was 90.8 ppb on the 24 of April 2011; we investigated the leaves for visible symptoms of  $O_3$  damage. No leaf damage was observed during the measurement period. In November a sudden drop in  $[O_3]$  and minimum  $O_3$  fluxes were coinciding with a peak increase in  $[NO]$  (Fig. 1). This increase in  $[NO]$  was probably linked to the strong ground surface inversion, which resulted in a minimum boundary layer depth during intense local traffic (L. Verlinden, VMM, personal communication).

### 3.2. Total, stomatal $O_3$ fluxes, & deposition velocity

Cospectral analysis was used to assess the performance of the LOZ-F instrument for eddy covariance, which revealed some noise in the mid to high frequency part of the spectra, but reasonable comparison with the cospectra of  $w'Ts'$  (Fig. 3). The temporal data coverage of the  $O_3$  fluxes ( $F_{O_3}$ ) after filtering was 41%. The seasonal trend in the daily averaged gap-filled total  $O_3$  fluxes (with the percentage of data coverage) and deposition velocity are shown in Fig. 4. Generally, the  $O_3$  fluxes were the most negative from April to July and closely related to the  $[O_3]$  (Fig. 1). Substantial  $O_3$  fluxes



**Fig. 4.** Wind speed, deposition velocity (not gap-filled, averaged in daily and monthly time scales), daily averaged gap-filled  $O_3$  fluxes (with the percentage of data coverage), gap-filled  $O_3$  fluxes averaged during daytime and nighttime (solar radiation  $>20 W m^{-2}$  or  $<20 W m^{-2}$ , respectively), and their percentage difference (area plot). Note the substantial  $O_3$  sink when temperature was below freezing from 4 to 6 February (red arrow). Displayed are averages and standard errors of the means (monthly averaged deposition velocity  $n = 407, 821, 613, 567, 728, 715, 431, 494, 529, 514, 381, 63$  for each month from January to December). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

were observed when the vegetation was not active yet, with a peak flux in co-occurrence of high  $[O_3]$  (4–6 February) during a storm (Fig. 1) and with a wind speed  $> 7 \text{ m s}^{-1}$  (Fig. 4). On 4–6 February, because of the low temperature and radiation (below freezing air temperatures and short wave radiation below  $100 \text{ W m}^{-2}$ , Fig. 1), the high  $[O_3]$  was probably resulting from the intrusion of stratospheric  $O_3$  into the troposphere (Weber and Prevot, 2002; Cieslik, 2009). The  $O_3$  fluxes (Fig. 4) observed in January–March (when the vegetation was not active yet, and the stem area index (SAI) was about  $0.33 \pm 0.19 \text{ m}^2 \text{ m}^{-2}$  (Broeckx, personal communication) had comparable rates to the ones observed during the growing season with substantial canopy development (with maximum seasonal leaf area index (LAI) of about  $2.5 \pm 1 \text{ m}^2 \text{ m}^{-2}$ ). The relevance of the stomatal uptake varied considerably during the season (Fig. 5), accounting for about 39% of the total  $O_3$  flux in April, 42% in May, increasing to 65% in June, 72% in July, and 77% in September, decreasing again to 59% in October. The decrease in VPD from June onwards together with the increase in LAI increased the importance of the  $F_{O_3ST}$  later in the summer (Fig. 5). Generally, the  $F_{O_3ST}$  was at maximum around noon (Fig. 5), while the  $[O_3]$  was at maximum early in the afternoon (around 16:00, Fig. 2). Monthly averaged deposition velocities reached maximum values in February ( $4.9 \pm 0.1 \text{ mm s}^{-1}$ ), and in October and November

( $5.5 \pm 0.1 \text{ mm s}^{-1}$  and  $6.0 \pm 0.2$  respectively, mean  $\pm$  st. errors, see Fig. 4), before canopy development and after leaf loss.

The non-stomatal uptake increased during daytime (most evident in March, Fig. 6). The controls on the non-stomatal uptake were investigated by examining the environmental controls on  $O_3$  fluxes and on the deposition velocity, for the entire day and separately during daytime and nighttime, as suggested by Fowler et al. (2001). The most important environmental control on the nighttime half-hourly averaged  $O_3$  fluxes was wind speed (Table 1). Wind speed was also highly significant but explained only less than 1% of the variability in the half-hourly averaged nighttime deposition velocity. During daytime, the single most important control on total  $O_3$  uptake was VPD. Wind speed was still an important control, and increased the explanatory power of the model. VPD and wind speed were also significant but only explained very small percentage of the deposition velocity (Table 1).

The official AOT40 reported by VMM in Gent (about 10 km from the research field site) was about 3 ppm h in 2011. At the field site we estimated AOT40 to be 5.6 ppm h from April until the end of October. The cumulative stomatal uptake from April to November, estimated as a daily average uptake for each of the months was about  $25 \text{ mmol m}^{-2}$ . This value is an underestimation as the stomatal uptake during mid-July to mid-September could not be





Fig. 5. Diurnal trend of not gap-filled  $O_3$  fluxes (total:  $F_{O_3}$ ; stomatal:  $F_{O_3ST}$ ) for each of the indicated months. Displayed are ensemble averages and standard error of the mean. A monthly average leaf area index (LAI)  $\pm$  standard errors are displayed in each panel.



**Fig. 6.** Diurnal trend of not gap-filled total  $O_3$  fluxes ( $F_{O_3}$ ) for each of the indicated months. The vegetation was not active during these periods (with the exception of November when leaf fall was still occurring). Displayed are averages for the same time of the day and standard error of the mean.

modeled. If the stomatal uptake during this period is assumed to be 59% of the total fluxes (i.e. the seasonal average stomatal ozone uptake), the resulting cumulative stomatal uptake for the entire growing season would be  $27 \text{ mmol m}^{-2}$ . Multiple regression and partial correlation analysis of the  $CO_2$  flux data from this study showed no impact of  $O_3$  on photosynthesis (Appendix A).

#### 4. Discussion

The comparable  $O_3$  fluxes in January–March (when the vegetation was not active yet), and during most of the growing season (with substantial canopy development), highlights a dominant role of non-stomatal  $O_3$  uptake in this system. The importance of wind speed on  $O_3$  fluxes more than on deposition velocity suggests that the turbulent mixing and the mechanical destruction of  $O_3$  played a role on non-stomatal  $O_3$  fluxes, but not on the surface exchange. Deposition velocity was poorly explained by any models, which suggests that a half-hour time scale might not be appropriate for the investigation of the controls on deposition velocity. A mechanical destruction of  $O_3$  by contact with snowflakes under turbulent conditions (Cieslik, 2009), was probably responsible for the substantial  $O_3$  fluxes observed during the storm at the beginning of February in absence of active vegetation. Ozone deposition to snow has previously been reported (Helmig et al., 2007; Cieslik, 2009; Bocquet et al., 2011) and explained as chemical interaction in the

snowpack (Helmig et al., 2007). However, in this field site snow did not deposit on the ground during the peak  $O_3$  flux in February, we therefore assume the mechanical destruction (Cieslik, 2009) was more important than deposition to snow for the observed peak in  $O_3$  fluxes.

The difference between the daytime and nighttime  $O_3$  fluxes was higher from April to May when the  $F_{O_3ST}$  was the lowest, as the dry condition, particularly during May, limited  $CO_2$  uptake by the vegetation (Zona et al., 2013) and consequently the stomatal ozone uptake. This highlights the importance of non-stomatal uptake during the light hours. A daytime increase in the non-stomatal uptake also occurred when vegetation was not active (most evident in March). This increase has been linked to the daytime increase in  $[O_3]$  and the decrease in aerodynamic resistance (Fowler et al., 2001). The observed mismatch between the peak  $O_3$  uptake and maximum  $[O_3]$  has already been reported in several ecosystem (Kurpius and Goldstein, 2003; Mikkelsen et al., 2004; Fares et al., 2013), and it is related to delay in the ozone formation through the photochemical process.

Mechanisms leading to  $O_3$  deposition are connected to VOC and NO emission from vegetation and from the soil. The importance of NO as  $O_3$  sink has already been reported by several studies (Wesely et al., 1982; Pilegaard et al., 1998; Dorsey et al., 2004; Michou et al., 2005; Wang et al., 2006). Gas-phase reaction between ozone and BVOC showed to be the major non-stomatal sink in a multitude of crop and forest ecosystems (Kurpius and Goldstein, 2003; Fares et al., 2010b, 2012; Rannik et al., 2012). Certain classes of BVOC, in particular monoterpenes and sesquiterpenes, seem to be associated with the non-stomatal ozone fluxes (Holzinger et al., 2005; Fares et al., 2012; Rannik et al., 2012). On the other hand, emission of BVOC, and isoprene in particular, can also contribute to the  $O_3$  formation (Jacob and Winner, 2009; Beltman et al., 2013). Poplar is known to be a major emitter of isoprene, the most abundant BVOC emitted on Earth by plant ecosystems (Guenther et al., 2006; Jardine et al., 2012). Isoprene production is maximized by light and temperature (Guenther et al., 1993), the same environmental conditions which promote photochemical ozone formation. It is therefore plausible that isoprene might have contributed to

**Table 1**

Statistical results (GLM) of the half-hourly averaged not gap-filled total  $O_3$  fluxes and deposition velocity for the daytime and nighttime (solar radiation  $>20 \text{ W m}^{-2}$  or  $<20 \text{ W m}^{-2}$ , respectively) datasets. Displayed are the explanatory powers and the statistical significance ( $p$ -values) of the listed variables.

	$O_3$ fluxes		Deposition velocity	
	Daytime	Nighttimes	Daytime	Nighttimes
Wind speed		44%( $p<0.001$ )		$<1\%$ ( $p<0.001$ )
VPD	38%( $p<0.001$ )		2%( $p<0.001$ )	
VPD & wind speed	47%( $p<0.001$ )		8%( $p<0.001$ )	



enhance  $[O_3]$  and ultimately increase the  $O_3$  fluxes. However, additional measurements on BVOC emissions and their oxidation products are required to clarify the mechanisms responsible for the chemical degradation component of non-stomatal  $O_3$  sink.

Critical values (corresponding to a 2–4% biomass loss or growth reduction, ICP, 2004) have been set to 5 ppm h for forests (AOT40), similar to the ones observed at our site. Foliar injury to poplar has been observed for AOT40 equal to or higher than 10 ppm h and for a stomatal ozone uptake equal to or higher than about 30 mmol  $m^{-2}$  (Marzuoli et al., 2009). The cumulative  $O_3$  uptake estimated in this study (25–27 mmol  $m^{-2}$ ) is lower but similar to values linked to visible damage in poplar (between 27.85 and 30.40 mmol  $m^{-2}$ , Marzuoli et al., 2009). In agreement with this observation, we did not notice visible damage from  $O_3$  exposure neither on the leaves, nor on NEE. However, a reduction in primary productivity has been shown even without visible damage (Wang et al., 1986). Several poplar species including some of the hybrids used in this study are known to be sensitive to  $O_3$  such as *Populus nigra* (Novak et al., 2007; Bortier et al., 2000) and *Populus trichocarpa* (Street et al., 2011), *Populus maximowizii* × *trichocarpa* (Landry and Pell, 1993), *Populus tremuloides* (Oksanen et al., 2001; Berrang et al., 1991; Karnosky et al., 1992) but these studies usually reported significant impacts with  $[O_3]$  much higher (e.g.  $\geq 100$  ppb) than the ones observed in this study.

## 5. Conclusions

Over the entire year of measurements, the non-stomatal  $O_3$  sink was a substantial component of the total  $O_3$  fluxes at our research site. The relative importance of non-stomatal  $O_3$  uptake (as percentage of the total  $O_3$  fluxes) was at its maximum during April–May and during daytime. Deposition velocities were the highest during February and October–November, in absence of a developed canopy. Substantial  $O_3$  deposition was observed during a snow storm, with rates comparable to the ones observed when the vegetation was at peak development. Wind speed represented the most important control on nighttime  $O_3$  fluxes, and therefore on the non-stomatal  $O_3$  sink. Despite the dominance of the non-stomatal  $O_3$  flux, the vegetation still absorbed a considerable amount of  $O_3$ , with an average stomatal contribution of about 59% of the total  $O_3$  uptake during the growing season. This uptake, however, did not result neither in any visible damage to the poplar leaves nor caused a reduction in NEE. More studies are needed to explore the individual contribution of stomatal and different non-stomatal pathways and sinks to total ozone removal from the atmosphere in this and other ecosystems.

## Acknowledgments

This research was funded by the European Research Council (FP7/2007–2013) ERC Advanced Grant n° 233366 (POPFULL) and by the Flemish Hercules Foundation as a Research Infrastructure Grant (ZW09–06). D. Zona was supported by a Marie Curie Reintegration Grant (PIRG07–GA-2010–268257), T. De Groote is a Ph.D. fellow of the Research Foundation – Flanders (FWO) and the Flemish Institute for Technological Research (VITO). We thank J. Cools and K. Mouton for logistic support, F. Miglietta, P. Toscano, G. Alberti and A. Zaldei for help with the experimental set-up, G. Fratini for help with data processing, the Royal Meteorological Institute of Belgium and L. Verlinden (VMM) for supply of monitoring data, E. Paoletti for recommendations on the data interpretation, J. Drummond for the help with LOZ-3F instrument, C. Ammann, B. Loubet, and the participants of the COST Action ABBA/ECLAIRE workshop in Paris for useful discussions. The COST Action ABBA, ES0804, and the COST

Action FP0903 provided funding for Short Term Scientific Missions (STSM) to discuss about data analysis.

## Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.envpol.2013.08.032>.

## References

- Angle, R.P., Sandhu, H.S., 1989. Urban and rural ozone concentrations in Alberta, Canada. *Atmos. Environ.* 23, 215–221.
- Beltman, J.B., Hendriks, C., Tum, M., Schaap, M., 2013. The impact of large scale biomass production on ozone air pollution in Europe. *Atmos. Environ.* <http://dx.doi.org/10.1016/j.atmosenv.2013.02.019>.
- Benton, J., Fuhrer, J., Gimeno, B.S., Skarby, L., Palmer-Brown, D., Ball, G., Roadknight, C., Mills, G., 2000. An international cooperative programme indicates the widespread occurrence of ozone injury on crops. *Agri. Ecosyst. Environ.* 78, 19–30.
- Berrang, P., Karnosky, D.F., Bennett, J.P., 1991. Natural selection for ozone tolerance in *Populus tremuloides*: an evaluation of nationwide trends. *Can. J. For. Res.* 21, 1091.
- Bocquet, F., Helmig, D., Van Dam, B.A., Fairall, C.W., 2011. Evaluation of the flux gradient technique for measurement of ozone surface fluxes over snowpack at Summit, Greenland. *Atmos. Measur. Tech.* 4, 2305–2321.
- Bortier, K., Ceulemans, R., De Temmerman, L., 1999. Effects of tropospheric ozone on woody plants. In: Agrawal, S.B., Agrawal, M. (Eds.), *Environmental Pollution and Plant Responses*. CRC Press, New York, pp. 153–182.
- Bortier, K., De Temmerman, L., Ceulemans, R., 2000. Effects of ozone exposure in open-top chambers on poplar (*Populus nigra*) and beech (*Fagus sylvatica*): a comparison. *Environ. Pollut.* 109, 509–516.
- Broeckx, L.S., Verlinden, M.S., Ceulemans, R., 2012. Establishment and two-year growth of a bio-energy plantation with fast-growing *Populus* trees in Flanders Belgium, effects of genotype and former land use. *Biomass Bioenerg.* 42, 151–163.
- Bussotti, F., Desotgiu, R., Cascio, C., Pollastrini, M., Gravano, E., Gerosa, G., Marzuoli, R., Nali, C., Lorenzini, G., Salvatori, E., Manes, F., Schaub, M., Strasser, R.J., 2011. Ozone stress in woody plants assessed with chlorophyll a fluorescence. A critical reassessment of existing data. *Environ. Exp. Bot.* 73, 19–30.
- Cieslik, S., 2009. Ozone fluxes over various plant ecosystems in Italy: a review. *Environ. Pollut.* 157, 1487–1496.
- Cieslik, S., Labatut, A., 1997. Ozone and heat fluxes over a Mediterranean pseudosteppe. *Atmos. Environ.* 31, 177–184.
- Comrie, A.C., 1990. The climatology of surface ozone in rural-areas - a conceptual-model. *Prog. Phys. Geogr.* 14, 295–316.
- Davis, R.E., Normile, C.P., Sitka, L., Hondula, D.M., Knight, D.B., Gawtry, S.P., Stenger, P.J., 2010. A comparison of trajectory and air mass approaches to examine ozone variability. *Atmos. Environ.* 44, 64–74.
- Dentener, F., Stevenson, D., Ellingsen, K., van Noije, T., Schultz, M., Amann, J., Atherton, C., Bell, N., Bergmann, D., Bey, I., Bouwman, L., Butler, T., Cofala, J., Collins, B., Drevet, J., Doherty, J., Eickhout, B., Eskes, H., Fiore, A., Gauss, M., Hauglustaine, D., Horowitz, L., Isaksen, I.S.A., Josse, B., Lawrence, M., Krol, M., Lamarque, J.F., Montanaro, V., Müller, J.F., Peuch, V.H., Pitari, G., Pyle, J., Rast, S., Rodriguez, J., Sanderson, M., Savage, N.H., Shindell, D., Strahan, S., Szopa, S., Sudo, K., Van Dingenen, R., Wild, O., Zeng, G., 2006. The global atmospheric environment for the next generation. *Environ. Sci. Technol.* 40, 3586–3594.
- Derwent, R.G., Simmonds, P.G., Manning, A.J., Spain, T.G., 2007. Trends over a 20-year period from 1987 to 2007 in surface ozone at the atmospheric research station, Mace Head, Ireland. *Atmos. Environ.* 41, 9091–9098.
- Dibb, J.E., Talbot, R.W., Scheuer, E., Seid, G., Debell, L., Lefer, B., Ridley, B., 2003. Stratospheric influence on the northern North American free troposphere during TOPSE: be-7 as a stratospheric tracer. *J. Geophys. Res.-atmos.* 108. <http://dx.doi.org/10.1029/2001jd001347>.
- Dorsey, J.R., Duyzer, J.H., Gallagher, M.W., Coe, H., Pilegaard, K., Weststrate, J.H., Jensen, N.O., Walton, S., 2004. Oxidized nitrogen and ozone interaction with forests. I: experimental observations and analysis of exchange with Douglas fir. *Quart. J. Royal Meteorol. Soc.* 130, 1941–1955.
- Emberson, L.D., Ashmore, M.R., Cambridge, H.M., Simpson, D., Tuovinen, J.P., 2000. Modelling stomatal ozone flux across Europe. *Environ. Pollut.* 109, 403–413.
- Fares, S., Weber, R., Park, J.H., Gentner, D., Karlik, J., Goldstein, A.H., 2012. Ozone deposition to an orange orchard: partitioning between stomatal and non-stomatal sinks. *Environ. Pollut.* 169, 258–266.
- Fares, S., Barta, C., Ederli, L., Ferranti, F., Pasqualini, S., Reale, L., Brilli, F., Tricoli, D., Loreto, F., 2006. Impact of high ozone on isoprene emission and some anatomical and physiological parameters of developing *Populus alba* leaves directly or indirectly exposed to the pollutant. *Physiol. Plant.* 128, 456–465.
- Fares, S., Goldstein, A.H., Loreto, F., 2010b. Determinants of ozone fluxes and metrics for ozone risk assessment in plants. *J. Exp. Bot.* 61, 629–633.
- Fares, S., Matteucci, G., Scarascia Mugnozza, G., Morani, A., Calfapietra, C., Salvatori, E., Fusaro, L., Manes, F., Loreto, F., 2013. Testing of models of stomatal

- ozone fluxes with field measurements in a mixed Mediterranean forest. *Atmos. Environ.* 67, 242–251.
- Fares, S., McKay, M., Holzinger, R., Goldstein, A.H., 2010a. Ozone fluxes in a *Pinus ponderosa* ecosystem are dominated by non-stomatal processes: evidence from long-term continuous measurements. *Agric. For. Meteorol.* 150, 420–431.
- Fowler, D., 1992. Effects of acidic pollutants on terrestrial ecosystems. In: Radojevic, M., Harrison, R.M. (Eds.), *Atmospheric Acidity: Sources, Consequences and Abatement*. Elsevier, Amsterdam, pp. 341–361.
- Fowler, D., Flechard, C., Cape, J.N., Storeton-West, R.L., Coyle, M., 2001. Measurements of ozone deposition to vegetation quantifying the flux, the stomatal and nonstomatal components. *Water Air Soil Pollut.* 130, 63–74.
- Fratini, G., Ibrom, A., Arriga, N., Burba, G., Papale, D., 2012. Relative humidity effects on water vapour fluxes measured with closed-path eddy-covariance systems with short sampling lines. *Agric. For. Meteorol.* 165, 53–63.
- Fuhrer, J., 2000. Introduction to the special issue on ozone risk analysis for vegetation in Europe. *Environ. Pollut.* 109, 359–360.
- Fumagalli, I., Gimeno, B.S., Velissariou, D., De Temmerman, L., Mills, G., 2001. Evidence of ozone-induced adverse effects on crops in the Mediterranean region. *Atmos. Environ.* 35, 2583–2587.
- Gerosa, G., Marzuoli, R., Bussotti, F., Pancrazi, M., Ballarin-Denti, A., 2003. Ozone sensitivity of *Fagus sylvatica* and *Fraxinus excelsior* young trees in relation to leaf structure and foliar ozone uptake. *Environ. Pollut.* 125, 91–98.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P.I., Geron, C., 2006. Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature). *Atmos. Chem. Phys.* 6, 3181.
- Guenther, A., Zimmerman, P.R., Harley, P.C., Monson, R.K., Fall, R., 1993. Isoprene and monoterpene emission rate variability: model evaluations and sensitivity analyses. *J. Geophys. Res.* 98, 12609–12617.
- Helmig, D., Bocquet, F., Cohen, L., Oltmans, S.J., 2007. Ozone uptake to the polar snow at Summit, Greenland. *Atmos. Environ.* 41, 5061–5076.
- Holzinger, R., Lee, A., Paw, U.K.T., Goldstein, A.H., 2005. Observations of oxidation products above a forest imply biogenic emissions of very reactive compounds. *Atmos. Chem. Phys.* 5, 67–75. <http://dx.doi.org/10.5194/acp-5-67-2005>.
- Horst, T.W., Lenschow, D.H., 2009. Attenuation of scalar fluxes measured with spatially-displaced sensors. *Bound. Layer Meteorol.* 130, 275–300.
- Hough, A.M., Derwent, R.G., 1990. Changes in the global concentration of tropospheric ozone due to human activities. *Nature* 344, 645–648.
- Ibrom, A., Dellwik, E., Larsen, S.E., Pilegaard, K., 2007. On the use of the Webb – Pearman – Leuning – theory for closed-path eddy correlation measurements. *Tellus B* 59B, 937–946.
- ICP, 2004. Update (2010/2011) of “Mapping critical levels for Vegetation, manual on methodologies and criteria for mapping critical levels/loads and geographical areas where they are exceeded”, UBA Texte 71/96. ISSN: 0722-186X. Berlin 1996.
- IPCC, 2001. *Climate Change 2001: the Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK, ISBN 0521-80767.0, p. 881.
- IPCC-DDC (The Intergovernmental Panel on Climate Change Data Distribution Centre), 2004. <http://ipcc-ddc.cru.uea.ac.uk>.
- Jacob, D.J., Winner, D.A., 2009. Effect of climate change on air quality. *Atmos. Environ.* 43, 51–63.
- Jardine, K., Abrell, L., Jardine, A., Huxman, T., Saleska, S., Arnett, A., Monson, R., Karl, T., Fares, S., Loreto, F., Goldstein, A., 2012. Within-plant isoprene oxidation confirmed by direct emissions of oxidation products methyl vinyl ketone and methacrolein. *Glob. Change Biol.* 18, 973–984.
- Jenkin, M.E., 2008. Trends in ozone concentration distributions in the UK since 1990: local, regional and global influences. *Atmos. Environ.* 42, 5434–5445.
- Kaimal, J.C., Wyngard, J.C., Izumi, Y., Cote, O.R., 1972. Spectral characteristics of surface-layer turbulence. *Quart. J. Royal Meteorol. Soc.* 98, 563–589.
- Kärenlampi, L., Skärby, L., 1996. Critical Levels for Ozone in Europe. TEsting and Finalizing the Concepts. UN-ECE Workshop Report. Department of Ecology and Environmental Science, University of Kuopio, Finland.
- Karlsson, P.E., Uddling, J., Braun, S., Broadmeadow, M., Elvira, S., Gimeno, B.S., Le Thiec, D., Oksanen, E., Vandermeiren, K., Wilkinson, M., Emberson, L., 2004. New critical levels for ozone effects on young trees based on AOT40 and simulated cumulative leaf uptake of ozone. *Atmos. Environ.* 38, 2283–2294.
- Karlsson, P.E., Pleijel, H., Danielsson, H., Pihl Karlsson, G., Piikki, K., Uddling, J., 2009. Evidence for impacts of near-ambient ozone concentrations on vegetation in southern Sweden. *Ambio* 38, 425–431.
- Karlsson, P.E., Sellén, G., Pleijel, H., 2003b. Establishing Ozone Critical Levels II. UN-ECE Workshop Report, IVL Report B 1523. IVL Swedish Environmental Research Institute, Gothenburg, Sweden <http://www.ivl.se>.
- Karlsson, P.E., Uddling, J., Skärby, L., Wallin, G., Sellén, G., 2003a. Impact of ozone on the growth of birch (*Betula pendula*) saplings. *Environ. Pollut.* 124, 485–495.
- Karnosky, D.F., Gagnon, Z.E., Reed, D.D., Witter, J.A., 1992. Growth and biomass allocation of symptomatic and asymptomatic *Populus tremuloides* clones in response to seasonal ozone exposures. *Can. J. For. Res.* 22, 1785–1788.
- Kolb, T.E., Matyssek, R., 2001. Limitations and perspectives about scaling ozone impacts in trees. *Environ. Pollut.* 115, 373–392.
- Kurpius, M.R., Goldstein, A.H., 2003. Gas-phase chemistry dominates O<sub>3</sub> loss to a forest, implying a source of aerosols and hydroxyl radicals to the atmosphere. *Geophys. Res. Lett.* 30, 1371. <http://dx.doi.org/10.1029/2002GL016785>.
- Landry, L.G., Pell, E.J., 1993. Modification of Rubisco and altered proteolytic activity in O<sub>3</sub>-stressed hybrid poplar (*Populus maximowizii* × *trichocarpa*). *Plant Phys.* 101, 1355–1362.
- Lefohn, A.S., 1992. *Surface Level Ozone Exposures and Their Effects on Vegetation*. Lewis Publishers, Chelsea, Mich.
- Lefohn, A.S., Shadwick, D., Oltmans, S.J., 2008. Characterizing long-term changes in surface ozone levels in the United States (1980–2005). *Atmos. Environ.* 42, 8252–8262.
- Lenschow, D.H., Raupach, M.R., 1991. The attenuation of fluctuations in scalar concentrations through sampling tubes. *J. Geophys. Res.* 96, 15259–15268.
- Manes, F., Vitale, M., Maria Fabi, A., De Santis, F., Zona, D., 2007. Estimates of potential ozone stomatal uptake in mature trees of *Quercus ilex* in a Mediterranean climate. *Environ. Exp. Bot.* 59, 235–241.
- Marzuoli, R., Gerosa, G., Desotgiu, R., Bussotti, F., Ballarin-Denti, A., 2009. Ozone fluxes and foliar injury development in the ozone-sensitive poplar clone Oxford (*Populus maximowizii* × *Populus berolinensis*): a dose-response analysis. *Tree Physiol.* 29, 67–76.
- Matyssek, R., Bytnerowicz, A., Karlsson, P.E., Paoletti, E., Sanz, M., Schaub, M., Wieser, G., 2007. Promoting the O<sub>3</sub> flux concept for European forest trees. *Environ. Pollut.* 146, 587–607.
- Meagher, J.F., Lee, N.T., Valente, R.J., Parkhurst, W.J., 1987. Rural ozone in the southeastern United-states. *Atmos. Environ.* 21, 605–615.
- Michou, M., Laville, P., Serca, D., Fotiadi, A., Bouchou, P., Peuch, V.H., 2005. Measured and modeled dry deposition velocities over the ESCOMPTE area. *Atmos. Res.* 74, 89–116.
- Mikkelsen, T.N., Ro-Poulsen, H., Hovmand, M.F., Jensen, N.O., Pilegaard, K., Egelov, A.H., 2004. Five-year measurements of ozone fluxes to a Danish Norway spruce canopy. *Atmos. Environ.* 38, 2361–2371.
- Mikkelsen, T.N., Ro-Poulsen, H., Pilegaard, K., Hovmand, M.F., Jensen, N.O., Christensen, C.S., Hummelshøj, P., 2000. Ozone uptake by an evergreen forest canopy—temporal variation and possible mechanisms. *Environ. Pollut.* 109, 423–429.
- Mills, G., 2004. Mapping critical levels for vegetation. In: UBA (Ed.), *UNECE Convention on Long-range Transboundary Air Pollution. Manual on Methodologies and Criteria for Mapping Critical Loads and Levels and Air Pollution Effects, Risks and Trends*. Constantly updated version available at: [www.oekodata.com/icpmapping/](http://www.oekodata.com/icpmapping/).
- Mills, G., Hayes, F., Simpson, D., Emberson, L., Norris, D., Harmens, H., Büker, P., 2011. Evidence of widespread effects of ozone on crops and (semi-)natural vegetation in Europe (1990–2006) in relation to AOT40- and flux-based risk maps. *Glob. Change Biol.* 17, 592–613.
- Moncrieff, J.B., Clement, R., Finnigan, J., Meyers, T., 2004. Averaging, detrending and filtering of eddy covariance time series. In: Lee, X., Massman, W.J., Law, B.E. (Eds.), *Handbook of Micrometeorology: a Guide for Surface Flux Measurements*. Kluwer Academic, Dordrecht, pp. 7–31.
- Moncrieff, J.B., Massheder, J.M., de Bruin, H., Ebers, J., Friborg, T., Heusinkveld, B., Kabat, P., Scott, S., Soegaard, H., Verhoef, A., 1997. A system to measure surface fluxes of momentum, sensible heat, water vapour and carbon dioxide. *J. Hydrol.* 188–189, 589–611.
- NEG-TAP, 2001. *National Expert Group on Transboundary Air Pollution. Transboundary Air Pollution: Acidification, Eutrophication and Ground-level Ozone in the UK*. Department for Environment, Food and Rural Affairs, UK.
- Neiryck, J., Gielen, B., Janssens, I.A., Ceulemans, R., 2012. Insights into ozone deposition patterns from decade-long ozone flux measurements over a mixed temperate forest. *J. Environ. Monit.* 14, 1684–1695.
- Nolle, M., Ellul, R., Heinrich, G., Gusten, H., 2002. A long-term study of background ozone concentrations in the central Mediterranean - diurnal and seasonal variations on the island of Gozo. *Atmos. Environ.* 36, 1391–1402.
- Novak, K., Cherubini, P., Saurer, M., Fuhrer, J., Skelly, J.M., Krauchi, N., Schaub, M., 2007. Ozone air pollution effects on tree-ring growth, delta C-13, visible foliar injury and leaf gas exchange in three ozone-sensitive woody plant species. *Tree Physiol.* 27, 941–949.
- Novak, K., Schaub, M., Fuhrer, J., Skelly, J.M., Hug, C., Landolt, W., Bleuler, P., Krauchi, N., 2005. Seasonal trends in reduced leaf gas exchange and ozone-induced foliar injury in three ozone sensitive woody plant species. *Environ. Pollut.* 136, 33–45.
- Novak, K., Skelly, J.M., Schaub, M., Krauchi, N., Hug, C., Landolt, W., Bleuler, P., 2003. Ozone air pollution and foliar injury development on native plants of Switzerland. *Environ. Pollut.* 125, 41–52.
- Oksanen, E., Amores, G., Kokko, H., Santamaria, J.M., Kärenlampi, L., 2001. Genotypic variation in growth and physiological responses of Finnish hybrid aspen (*Populus tremuloides* × *P. tremula*) to elevated tropospheric ozone concentration. *Tree Physiol.* 21, 1171–1181.
- Paoletti, E., 2006. Impact of ozone on Mediterranean forests: a review. *Environ. Pollut.* 144, 463–474.
- Papale, D., Valentini, R., 2003. A new assessment of European forests carbon exchanges by eddy fluxes and artificial neural network spatialization. *Glob. Change Biol.* 9, 525–535.
- Pilegaard, K., Hummelshøj, P., Jensen, N.O., 1998. Fluxes of ozone and nitrogen dioxide measured by Eddy correlation over a harvested wheat field. *Atmos. Environ.* 32, 1167–1177.
- Pilegaard, K., Ibrom, A., Courtney, M.S., Hummelshøj, P., Jensen, N.O., 2011. Increasing net CO<sub>2</sub> uptake by a Danish beech forest during the period from 1996 to 2009. *Agric. For. Meteorol.* 151, 934–946.
- Pollastrini, M., Desotgiu, R., Cascio, C., Bussotti, F., Cherubini, P., Saurer, M., Gerosa, G., Marzuoli, R., 2010. Growth and physiological responses to ozone and

- mild drought stress of tree species with different ecological requirements. *Trees – Struct. Funct.* 24, 695–704.
- Pye, J.M., 1988. Impact of ozone on the growth and yield of trees: a review. *J. Environ. Qual.* 17, 347–360.
- Rannik, Ü., Altimir, N., Mammarella, I., Bäck, J., Rinne, J., Ruuskanen, T.M., Hari, P., Vesala, T., Kulmala, M., 2012. Ozone deposition into a boreal forest over a decade of observations: evaluating deposition partitioning and driving variables. *Atmos. Chem. Phys.* 12, 12165–12182.
- Ray, J.D., Stedman, D.H., Wendel, G.J., 1986. Fast chemiluminescent method for measurement of ambient ozone. *Anal. Chem.* 58, 598–600.
- Royal Society, 2008. Ground-level Ozone in the 21st Century: Future Trends, Impacts and Policy Implications. Science Policy Report 15/08.
- Scheel, H.E., Areskoug, H., Geiss, H., Gomiscek, B., Granby, K., Haszpra, L., Klasinc, L., Kley, D., Laurila, T., Lindskog, A., Roemer, M., Schmitt, R., Simmonds, P., Solberg, S., Toupance, G., 1997. On the spatial distribution and seasonal variation of lower-troposphere ozone over Europe. *J. Atmos. Chem.* 28, 11–28.
- Sitch, S., Cox, P.M., Collins, W.J., Huntingford, C., 2007. Indirect radiative forcing of climate change through ozone effects on the land-carbon sink. *Nature* 448, 791–794.
- Solberg, S., Bergstrom, R., Langner, J., Laurila, T., Lindskog, A., 2005. Changes in Nordic surface ozone episodes due to European emission reductions in the 1990s. *Atmos. Environ.* 39, 179–192.
- Street, N.R., Tallis, M.J., Tucker, J., Brosche, M., Kangasjarvi, J., Broadmeadow, M., Taylor, G., 2011. The physiological, transcriptional and genetic responses of an ozone-sensitive and an ozone-tolerant poplar and selected extremes of their F-2 progeny. *Environ. Pollut.* 159, 45–54.
- Topham, L.A., MacKay, G.I., Schiff, H.I., 1992. Performance assessment of the portable and lightweight LOZ-3 chemiluminescence type ozone monitor. In: Proceedings of the International Symposium and Course on Measurement of Toxic and Related Air Pollutants. Air and Waste Management Association, Durham, North Carolina.
- Tuovinen, J.P., Simpson, D., Emberson, L., Ashmore, M., Gerosa, G., 2007. Robustness of modelled ozone exposures and doses. *Environ. Pollut.* 146, 578–586.
- Uddling, J., Günthardt-Goerg, M.S., Matyssek, R., Oksanen, E., Pleijel, H., Selldén, G., Karlsson, P.E., 2004. Biomass reduction of juvenile birch is more strongly related to stomatal uptake of ozone than to indices based on external exposure. *Atmos. Environ.* 38, 4709–4719.
- Vingarzan, R., 2004. A review of surface ozone background levels and trends. *Atmos. Environ.* 38, 3431–3442.
- Wang, D., Karnosky, D.F., Bormann, F.H., 1986. Effects of ambient ozone on the productivity of *Populus tremuloides* Michx. grown under field conditions. *Can. J. For. Res.* 16, 47–55.
- Wang, S., Ackermann, R., Stutz, J., 2006. Vertical profiles of O-3 and NO<sub>x</sub> chemistry in the polluted nocturnal boundary layer in Phoenix, Az: I. Field observations by long-path DOAS. *Atmos. Chem. Phys.* 6, 2671–2693.
- Webb, E.K., Pearman, G.I., Leuning, R., 1980. Correction of flux measurements for density effects due to heat and water-vapour transfer. *Quart. J. R. Meteorol. Soc.* 106, 85–100.
- Weber, R.O., Prevot, A.S.H., 2002. Climatology of ozone transport from the free troposphere into the boundary layer south of the Alps during North Foehn. *J. Geophys. Res.-atmos.* 107. <http://dx.doi.org/10.1029/2001JD000987>.
- Wesely, M.L., Eastman, J.A., Stedman, D.H., Yalvac, E.D., 1982. An eddy correlation measurement technique of NO<sub>2</sub> flux to vegetation and comparison to O<sub>3</sub> flux. *Atmos. Environ.* 16, 815–820.
- Wittig, V.E., Ainsworth, E.A., Long, S.P., 2007. To what extent do current and projected increases in surface ozone affect photosynthesis and stomatal conductance of trees? A meta-analytic review of the last 3 decades of experiments. *Plant Cell Environ.* 30, 1150–1162.
- Zapletal, M., et al., 2011. Ozone flux over a Norway spruce forest and correlation with net ecosystem production. *Environ. Pollut.* 159, 1024–1034.
- Zona, D., Janssens, I.A., Gioli, B., Jungkunst, H.F., Serrano, M.C., Ceulemans, R., 2013a. N<sub>2</sub>O fluxes of a bio-energy poplar plantation during a two years rotation period. *Glob. Change Biol. Bioenerg.* 5 (5), 536–547. <http://dx.doi.org/10.1111/gcbb.12019>.
- Zona, D., Janssens, I.A., Aubinet, M., Gioli, B., Vicca, S., Fichot, R., Ceulemans, R., 2013b. Fluxes of the greenhouse gases (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) above a short-rotation poplar plantation after conversion from agricultural land. *Agric. For. Meteorol.* 169, 100–110.