

Technical University of Denmark



Novel 4-component reaction for the synthesis of boron heterocyclic scaffolds

Petersen, Mette Terp; Flagstad, Thomas; Nielsen, Thomas Eiland

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Petersen, M. T., Flagstad, T., & Nielsen, T. E. (2014). Novel 4-component reaction for the synthesis of boron heterocyclic scaffolds. Abstract from 14th Belgian Organic Synthesis Symposium, Louvain-La-Neuve, Belgium.

DTU Library
Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


NOVEL 4-COMPONENT REACTION FOR THE SYNTHESIS OF BORON HETEROCYCLIC SCAFFOLDS

Mette T. Petersen (1), Thomas Flagstad (1), Thomas E. Nielsen (1,2)

1) Department of Chemistry, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark

2) Singapore Centre on Environmental Life Sciences Engineering, Nanyang Technological University, Singapore 637551, Singapore

Multi-component reactions allow for the generation of complex molecular scaffolds in a single synthetic operation from simple building blocks.¹ In this context, the Petasis 3-component reaction is particularly useful for the synthesis of *anti*-amino alcohols from amines, α -hydroxy aldehydes and boronic acids. We now wish to report a variant of this powerful reaction, where hydrazides **1**,² α -hydroxy aldehydes **2** and *two* boronic acids, typically an electron-rich boronic acid **3** and an electron-poor boronic acid **4**, are combined to give novel boron heterocyclic compounds **6** (see figure). This 4-component reaction is highly modular and proceeds with excellent diastereoselectivity, providing the desired products **6** in high yields. In recent years there has been profound interest in boron heterocycles, e.g. as protecting groups³ and organocatalysts,⁴ and we foresee many exciting applications of these scaffolds in the future.



References

- 1) de Graaff, C.; Ruijter, E.; Orru, R. V. A. *Chem. Soc. Rev.* 2012, 41, 3969-4009
- 2) Le Quement, S. T.; Flagstad, T.; Mikkelsen, R. J. T.; Hansen, M. R.; Givskov, M. C.; Nielsen, T. E. *Org. Lett.* 2012, 14, 640-643
- 3) For a recent report, see: Denis, J. D. St.; Scully, C. C. G.; Lee, C. F.; Yudin, A. K. *Org. Lett.* 2014, 16, 1338-1341
- 4) Dimitrijević, E.; Taylor, M. S. *ACS Catal.* 2013, 3, 945-962