

Pion condensation and phase diagram in the Polyakov-loop quark-meson model

Prabal Adhikari,^{1,*} Jens O. Andersen,^{2,†} and Patrick Kneschke^{3,‡}

¹*St. Olaf College, Physics Department, 1520 St. Olaf Avenue, Northfield, MN 55057, USA*

²*Department of Physics, Faculty of Natural Sciences, NTNU,*

Norwegian University of Science and Technology, Høgskoleringen 5, N-7491 Trondheim, Norway

³*Faculty of Science and Technology, University of Stavanger, N-4036 Stavanger, Norway*

(Dated: October 22, 2018)

We use the Polyakov-loop extended two-flavor quark-meson model as a low-energy effective model for QCD to study the phase diagram in the μ_I - T plane where μ_I is the isospin chemical potential. In particular, we focus on the Bose condensation of charged pions. At $T = 0$, the onset of pion condensation is at $\mu_I = \frac{1}{2}m_\pi$ in accordance with exact results. The phase transition to a Bose-condensed phase is of second order for all values of μ_I and in the $O(2)$ universality class. The chiral critical line joins the critical line for pion condensation at a point whose position depends on the Polyakov-loop potential and the sigma mass. For larger values of μ_I these curves are on top of each other. The deconfinement line enters smoothly the phase with the broken $O(2)$ symmetry. We compare our results with recent lattice simulations and find overall good agreement.

I. INTRODUCTION

The phases of QCD as functions of the baryon chemical potential μ_B or the quark chemical potential $\mu = \frac{1}{3}\mu_B$, and temperature T have been studied in detail since the first phase diagram was proposed more than forty years ago [1–3]. At vanishing baryon chemical potential, it is possible to perform lattice simulations to calculate the thermodynamic functions and the transition temperature associated with chiral symmetry restoration and deconfinement. For physical quark masses and two flavors, the transition is a crossover at a temperature of approximately 155 MeV [4–7].

At nonzero baryon chemical potential, however, Monte Carlo simulations are hampered by the so-called sign problem, namely that the fermion determinant becomes complex. Being complex, the usual interpretation of it as part of a probability distribution can no longer be upheld. The sign problem in QCD at finite baryon density has spurred the interest in QCD-like theories free of this problem. This includes QCD with quarks in the adjoint representation [8], two-color QCD [9], QCD at finite isospin μ_I [10], and QCD in a magnetic field B [11]. These theories are all interesting in their own right; QCD at finite isospin and QCD in a magnetic field are also relevant for compact stars. In addition, the application of Monte-Carlo methods allows a direct test of various model approaches in the cases mentioned above. Such a confrontation of model calculations with lattice simulations of QCD in a magnetic field has been very

fruitful in understanding their strengths and limitations [12, 13].

Lattice simulations of QCD at finite isospin have been performed in e.g. Refs. [14–18] with particular emphasis on Bose condensation of charged pions for isospin chemical potentials above the zero-temperature critical value $\mu_I^c = \frac{1}{2}m_\pi$. Chiral perturbation theory (ChPT) [10, 19–22], which is a model-independent low-energy theory for QCD valid at low densities has been used to study pion condensation. ChPT predicts a second-order transition, which is in agreement with lattice simulations. There have also been a number of other approaches and model calculations studying various aspects of the QCD phase diagram at finite isospin density, including the resonance gas model [23], random matrix models [24], the Nambu-Jona-Lasinio (NJL) model [25–37], the quark-meson (QM) model [38–41]¹, and effective theory at asymptotically high isospin [42].

Finally, we mention that one expects another phase transition at large isospin chemical potential. In perturbation theory, one-gluon exchange gives rise to an effective attractive interaction between u and \bar{d} quarks leading to the formation of Cooper pairs [10]. The transition from a Bose-Einstein condensate (BEC) to a Bardeen-Cooper-Schrieffer (BCS) state is expected to be an analytical crossover as the symmetry-breaking pattern is the same.

As pointed out in Ref. [38], there is a mapping of the quark-meson model at finite isospin and the corresponding two-color quark-meson-diquark model at finite baryon chemical potential. The neutral pion π_0 is replaced by an isovector triplet $\boldsymbol{\pi}$. The charged pions π^\pm

* adhika1@stolaf.edu

† andersen@tf.phys.ntnu.no

‡ patrick.kneschke@uis.no

¹ Or their Polyakov-loop extended versions (PNJL and PQM).

are replaced by a diquark-antidiquark pair Δ and Δ^* , which instead of being coupled to μ_I is now coupled to a baryon chemical potential μ_B .² Since the gauge groups $SU(2)$ and $SU(3)$ are fundamentally different, this mapping is valid for the matter sector; once we couple the QM model to the Polyakov loop, this identification is lost.

In the present paper, we study the QCD phase diagram at finite temperature and isospin density using the PQM model. The main conclusions of our work are

1. The second order transition to a BEC state. The transition is in the $O(2)$ universality class. At $T = 0$, the transition is exactly at $\mu_I = \frac{1}{2}m_\pi$.
2. The BEC and chiral transition lines meet at a point $(\mu_I^{\text{meet}}, T^{\text{meet}})$ and coincide for larger isospin chemical potentials μ_I .
3. The deconfinement and chiral transition lines coincide in the non-condensed phase for a logarithmic Polyakov-loop potential and a sufficiently low sigma mass.
4. The deconfinement line penetrates smoothly into the symmetry-broken phase.

These results are in agreement with the recent lattice simulations of Refs. [16–18].

The paper is organized as follows. In Sec. II, we briefly discuss the quark-meson model and in Sec. III we calculate the effective potential in the mean-field approximation. In Sec. IV, we discuss the coupling to the Polyakov loop, while in Sec. V, we present the phase diagram in the μ_I - T plane and compare it to recent lattice results. In Appendix A, we list a few integrals needed in the calculations, while Appendix B provides the reader with some details of how the parameters of the quark-meson model are determined.

II. QUARK-MESON MODEL

The Lagrangian of the two-flavor quark-meson model in Minkowski space is

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} [(\partial_\mu \sigma)(\partial^\mu \sigma) + (\partial_\mu \pi_3)(\partial^\mu \pi_3)] \\ & + (\partial_\mu + 2i\mu_I \delta_\mu^0) \pi^+ (\partial^\mu - 2i\mu_I \delta_0^\mu) \pi^- \\ & - \frac{1}{2} m^2 (\sigma^2 + \pi_3^2 + 2\pi^+ \pi^-) - \frac{\lambda}{24} (\sigma^2 + \pi_3^2 + 2\pi^+ \pi^-)^2 \\ & + h\sigma + \bar{\psi} [i\cancel{\partial} + \mu_f \gamma^0 - g(\sigma + i\gamma^5 \boldsymbol{\tau} \cdot \boldsymbol{\pi})] \psi, \end{aligned} \quad (1)$$

where ψ is a color N_c -plet, a four-component Dirac spinor as well as a flavor doublet

$$\psi = \begin{pmatrix} u \\ d \end{pmatrix}, \quad (2)$$

and $\mu_f = \text{diag}(\mu_u, \mu_d)$, where μ_u and μ_d , are the quark chemical potentials, μ_I is the isospin chemical potential, τ_i ($i = 1, 2, 3$) are the Pauli matrices in flavor space, $\boldsymbol{\pi} = (\pi_1, \pi_2, \pi_3)$, and $\pi^\pm = \frac{1}{\sqrt{2}}(\pi_1 \pm i\pi_2)$.

Apart from the global $SU(N_c)$ symmetry, the Lagrangian (1) has a $U(1)_B \times SU(2)_L \times SU(2)_R$ symmetry for $h = 0$ and a $U(1)_B \times SU(2)_V$ symmetry for $h \neq 0$. When $\mu_u \neq \mu_d$, this symmetry is reduced to $U(1)_B \times U_{I_3L}(1) \times U_{I_3R}(1)$ for $h = 0$ and $U(1)_B \times U_{I_3}(1)$ for $h \neq 0$.

The number density associated with a chemical potential μ_A is

$$n_A = -\frac{\partial V}{\partial \mu_A}, \quad (3)$$

where V is the effective potential. The baryon and isospin densities can be expressed in terms of the quark densities n_u and n_d as

$$n_B = \frac{1}{3}(n_u + n_d), \quad (4)$$

$$n_I = n_u - n_d. \quad (5)$$

Eqs. (4)–(5) together with the chain rule can be used to derive relations among the baryon and isospin chemical potentials and the quark chemical potentials. We have

$$\begin{aligned} n_I &= -\frac{\partial V}{\partial \mu_I} \\ &= -\left(\frac{\partial V}{\partial \mu_u} - \frac{\partial V}{\partial \mu_d} \right) \\ &= -\left(\frac{\partial \mu_u}{\partial \mu_I} \frac{\partial V}{\partial \mu_u} + \frac{\partial \mu_d}{\partial \mu_I} \frac{\partial V}{\partial \mu_d} \right). \end{aligned} \quad (6)$$

This yields

$$\frac{\partial \mu_u}{\partial \mu_I} = -\frac{\partial \mu_d}{\partial \mu_I} = 1. \quad (7)$$

Similarly, we find $\frac{\partial \mu_u}{\partial \mu_B} = \frac{\partial \mu_d}{\partial \mu_B} = \frac{1}{3}$. From this, we find the following relations among the chemical potentials

$$\mu_u = \frac{1}{3}\mu_B + \mu_I, \quad (8)$$

$$\mu_d = \frac{1}{3}\mu_B - \mu_I. \quad (9)$$

Introducing the quark chemical potential $\mu = \frac{1}{3}\mu_B$ and inverting the relations (8)–(9), we find

$$\mu = \frac{1}{2}(\mu_u + \mu_d), \quad (10)$$

$$\mu_I = \frac{1}{2}(\mu_u - \mu_d). \quad (11)$$

² The diquarks are the baryons of two-color QCD.

III. EFFECTIVE POTENTIAL

The expectation values of the fields are written as

$$\sigma = \phi_0, \quad \pi_1 = \pi_0, \quad (12)$$

where ϕ_0 and π_0 are constant in space. The former is the usual chiral condensate, while the latter represents a homogeneous pion condensate. A pion condensate breaks the $U_{I_3L}(1) \times U_{I_3R}(1)$ symmetry to $U_{I_3V}(1)$ or the $U_{I_3}(1)$ symmetry. Introducing $\Delta = g\phi_0$ and $\rho = g\pi_0$, the tree-level potential in Euclidean space can be written as

$$V_0 = \frac{1}{2} \frac{m^2}{g^2} \Delta^2 + \frac{1}{2} \frac{m^2 - 4\mu_I^2}{g^2} \rho^2 + \frac{\lambda}{24g^4} (\Delta^2 + \rho^2)^2 - \frac{h}{g} \Delta. \quad (13)$$

Expressing the parameters in Eq. (1) terms of the sigma mass m_σ , the pion mass m_π , the pion decay constant f_π , and the constituent quark mass m_q , we find

$$m^2 = -\frac{1}{2} (m_\sigma^2 - 3m_\pi^2), \quad \lambda = 3 \frac{(m_\sigma^2 - m_\pi^2)}{f_\pi^2}, \quad (14)$$

$$g^2 = \frac{m_q^2}{f_\pi^2}, \quad h = m_\pi^2 f_\pi. \quad (15)$$

Inserting these relations, we can write the tree-level potential as

$$V_0 = -\frac{1}{4} f_\pi^2 (m_\sigma^2 - 3m_\pi^2) \frac{\Delta^2 + \rho^2}{m_q^2} - 2\mu_I^2 f_\pi^2 \frac{\rho^2}{m_q^2} + \frac{1}{8} f_\pi^2 (m_\sigma^2 - m_\pi^2) \frac{(\Delta^2 + \rho^2)^2}{m_q^4} - m_\pi^2 f_\pi^2 \frac{\Delta}{m_q}. \quad (16)$$

$$V_{\text{div}} = -4N_c \int_p \left[\sqrt{p^2 + \Delta^2 + \rho^2} + \frac{\mu_I^2 \rho^2}{2(p^2 + \Delta^2 + \rho^2)^{\frac{3}{2}}} \right] = \frac{4N_c}{(4\pi)^2} \left(\frac{e^{\gamma_E} \Lambda^2}{\Delta^2 + \rho^2} \right)^\epsilon \left[(\Delta^2 + \rho^2)^2 \Gamma(-2 + \epsilon) - 2\mu_I^2 \rho^2 \Gamma(\epsilon) \right]. \quad (21)$$

The remainder V_{fin} is finite and reads

$$V_{\text{fin}} = V_1 - V_{\text{div}}. \quad (22)$$

Note that V_{fin} can be evaluated directly in $d = 3$ dimensions. In the present case, V_{fin} must be evaluated numerically. Using the expressions for the integrals listed in Appendix A, we can write the unrenormalized one-loop effective potential $V = V_0 + V_1$ as

$$V = \frac{1}{2} \frac{m^2}{g^2} \Delta^2 + \frac{1}{2} \frac{m^2 - 4\mu_I^2}{g^2} \rho^2 + \frac{\lambda}{24g^4} (\Delta^2 + \rho^2)^2 - \frac{h}{g} \Delta + \frac{2N_c}{(4\pi)^2} \left(\frac{\Lambda^2}{\Delta^2 + \rho^2} \right)^\epsilon \left[(\Delta^2 + \rho^2)^2 \left(\frac{1}{\epsilon} + \frac{3}{2} \right) - 4\mu_I^2 \rho^2 \frac{1}{\epsilon} \right] + V_{\text{fin}} + \mathcal{O}(\epsilon), \quad (23)$$

which contains poles in ϵ . These poles are removed by mass and coupling constant renormalization. In the $\overline{\text{MS}}$

The quark energies can be read off from the zeros of the determinant of the Dirac operator. One finds

$$E_u = E(-\mu_I), \quad E_d = E(\mu_I), \quad (17)$$

$$E_{\bar{u}} = E(\mu_I), \quad E_{\bar{d}} = E(-\mu_I). \quad (18)$$

where we have defined

$$E(\mu_I) = \left[\left(\sqrt{p^2 + \Delta^2} + \mu_I \right)^2 + \rho^2 \right]^{\frac{1}{2}}. \quad (19)$$

Note that the quark energies explicitly depend on μ_I . In the following we choose $\mu_I > 0$, but similar results are obtained for $\mu_I < 0$.

The one-loop contribution to the effective potential at $T = \mu_B = 0$ is

$$V_1 = -N_c \int_p (E_u + E_d + E_{\bar{u}} + E_{\bar{d}}), \quad (20)$$

where the integral is in $d = 3 - 2\epsilon$ dimensions (See Appendix A). The integral in Eq. (20) is ultraviolet divergent and in order to isolate the divergences, we need to expand the energies in powers of μ_I to the appropriate order. This yields

scheme this is achieved by making the substitutions $m^2 \rightarrow Z_{m^2} m^2$, $\lambda \rightarrow Z_\lambda \lambda$, $g^2 \rightarrow Z_{g^2} g^2$, and $h \rightarrow Z_h h$, where

$$Z_{m^2} = 1 + \frac{4N_c g^2}{(4\pi)^2 \epsilon}, \quad Z_\lambda = 1 + \frac{8N_c}{(4\pi)^2 \epsilon} \left[g^2 - 6 \frac{g^4}{\lambda} \right], \quad Z_{g^2} = 1 + \frac{4N_c g^2}{(4\pi)^2 \epsilon}, \quad Z_h = 1 + \frac{2N_c g^2}{(4\pi)^2 \epsilon}, \quad (24)$$

The renormalized one-loop effective potential then reads

$$V_{1\text{-loop}} = \frac{1}{2} \frac{m_{\overline{\text{MS}}}^2}{g_{\overline{\text{MS}}}^2} \Delta^2 + \frac{1}{2} \frac{m_{\overline{\text{MS}}}^2 - 4\mu_I^2}{g_{\overline{\text{MS}}}^2} \rho^2 + \frac{\lambda_{\overline{\text{MS}}}}{24g_{\overline{\text{MS}}}^4} (\Delta^2 + \rho^2)^2 - \frac{h_{\overline{\text{MS}}}}{g_{\overline{\text{MS}}}} \Delta + \frac{2N_c}{(4\pi)^2} \left\{ [(\Delta^2 + \rho^2)^2 - 4\mu_I^2 \rho^2] \log \left(\frac{\Lambda^2}{\Delta^2 + \rho^2} \right) + \frac{3}{2} (\Delta^2 + \rho^2)^2 \right\} + V_{\text{fin}}, \quad (25)$$

where the subscript $\overline{\text{MS}}$ indicates that the parameters are running with the renormalization scale Λ . Using Z_{g^2} in Eq. (24) and the wavefunction renormalization factor $Z_\phi = 1 - \frac{4N_c g^2}{(4\pi)^2 \epsilon}$, it is seen that the fields Δ and ρ do not run. In Appendix B, we discuss how one can express the parameters in the $\overline{\text{MS}}$ scheme in terms of physical masses and couplings. Using Eqs. (B14)–(B17), the final expression for the one-loop effective potential in the large- N_c limit becomes

$$V_{1\text{-loop}} = \frac{3}{4} m_\pi^2 f_\pi^2 \left\{ 1 - \frac{4m_q^2 N_c}{(4\pi)^2 f_\pi^2} m_\pi^2 F'(m_\pi^2) \right\} \frac{\Delta^2 + \rho^2}{m_q^2} - \frac{1}{4} m_\sigma^2 f_\pi^2 \left\{ 1 + \frac{4m_q^2 N_c}{(4\pi)^2 f_\pi^2} \left[\left(1 - \frac{4m_q^2}{m_\sigma^2} \right) F(m_\sigma^2) + \frac{4m_q^2}{m_\sigma^2} - F(m_\pi^2) - m_\pi^2 F'(m_\pi^2) \right] \right\} \frac{\Delta^2 + \rho^2}{m_q^2} - 2\mu_I^2 f_\pi^2 \left\{ 1 - \frac{4m_q^2 N_c}{(4\pi)^2 f_\pi^2} \left[\log \frac{\Delta^2 + \rho^2}{m_q^2} + F(m_\pi^2) + m_\pi^2 F'(m_\pi^2) \right] \right\} \frac{\rho^2}{m_q^2} + \frac{1}{8} m_\sigma^2 f_\pi^2 \left\{ 1 - \frac{4m_q^2 N_c}{(4\pi)^2 f_\pi^2} \left[\frac{4m_q^2}{m_\sigma^2} \left(\log \frac{\Delta^2 + \rho^2}{m_q^2} - \frac{3}{2} \right) - \left(1 - \frac{4m_q^2}{m_\sigma^2} \right) F(m_\sigma^2) + F(m_\pi^2) + m_\pi^2 F'(m_\pi^2) \right] \right\} \frac{(\Delta^2 + \rho^2)^2}{m_q^4} - \frac{1}{8} m_\pi^2 f_\pi^2 \left[1 - \frac{4m_q^2 N_c}{(4\pi)^2 f_\pi^2} m_\pi^2 F'(m_\pi^2) \right] \frac{(\Delta^2 + \rho^2)^2}{m_q^4} - m_\pi^2 f_\pi^2 \left[1 - \frac{4m_q^2 N_c}{(4\pi)^2 f_\pi^2} m_\pi^2 F'(m_\pi^2) \right] \frac{\Delta}{m_q} + V_{\text{fin}}. \quad (26)$$

The finite-temperature part of the one-loop effective potential at $\mu_B = 0$ is

$$V_T = -2N_c T \int_p \left\{ \log [1 + e^{-\beta E_u}] + \log [1 + e^{-\beta E_d}] + \log [1 + e^{-\beta E_{\bar{u}}}] + \log [1 + e^{-\beta E_{\bar{d}}}] \right\}. \quad (27)$$

The complete one-loop effective potential in the QM model in the large- N_c limit is then the sum of Eqs. (26) and (27). Note that Eq. (27) vanishes at $T = 0$ and that the only μ_I -dependence of $V_{1\text{-loop}}$ is line three of Eq. (26).

IV. COUPLING TO THE POLYAKOV LOOP

In a pure gauge theory, the Polyakov loop is an order parameter for deconfinement, as first discussed in Refs. [43, 44]. In QCD with dynamical quarks, it is an approximate order parameter. This is analogous to the quark condensate which is an exact order parameter for chiral

symmetry for massless quark but only an approximate order parameter for massive quarks. The Polyakov loop is defined as the trace of the thermal Wilson line, where the thermal Wilson line L is given by

$$L(\mathbf{x}) = \mathcal{P} \exp \left[i \int_0^\beta d\tau A_4(\mathbf{x}, \tau) \right], \quad (28)$$

where $A_4 = iA_0$ is the temporal component of the gauge field in Euclidean space, $A_0 = t_a A_0^a$, $t_a = \frac{1}{2} \lambda^a$ are the generators of $SU(3)_c$ gauge group, λ^a are the Gell-Mann matrices, and \mathcal{P} denotes path ordering. The background field A_4 in the Polyakov gauge is

$$A_4 = t_3 A_4^3 + t_8 A_4^8, \quad (29)$$

where A_4^3 and A_4^8 are time independent fields. Substituting Eq. (29) into Eq. (28), the Wilson line becomes

$$L = \begin{pmatrix} e^{i(\phi_1+\phi_2)} & 0 & 0 \\ 0 & e^{i(-\phi_1+\phi_2)} & 0 \\ 0 & 0 & e^{-2i\phi_2} \end{pmatrix}, \quad (30)$$

where we have defined $\phi_1 = \frac{1}{2}\beta A_4^3$ and $\phi_2 = \frac{1}{2\sqrt{3}}\beta A_4^8$. Introducing the Polyakov loop variables³

$$\Phi = \frac{1}{N_c} \text{Tr} L, \quad \bar{\Phi} = \frac{1}{N_c} \text{Tr} L^\dagger, \quad (31)$$

$$V_T = -2T \int \frac{d^3p}{(2\pi)^3} \left\{ \text{Tr} \log \left[1 + 3(\Phi + \bar{\Phi} e^{-\beta E_u}) e^{-\beta E_u} + e^{-3\beta E_u} \right] + \text{Tr} \log \left[1 + 3(\bar{\Phi} + \Phi e^{-\beta E_{\bar{u}}}) e^{-\beta E_{\bar{u}}} + e^{-3\beta E_{\bar{u}}} \right] \right. \\ \left. + \text{Tr} \log \left[1 + 3(\Phi + \bar{\Phi} e^{-\beta E_d}) e^{-\beta E_d} + e^{-3\beta E_d} \right] + \text{Tr} \log \left[1 + 3(\bar{\Phi} + \Phi e^{-\beta E_{\bar{d}}}) e^{-\beta E_{\bar{d}}} + e^{-3\beta E_{\bar{d}}} \right] \right\}. \quad (32)$$

Eq. (32) reduces to Eq. (27) upon setting $\Phi = \bar{\Phi} = 1$, i.e. we obtain the finite-temperature part of the effective potential in the quark-meson model.

The Polyakov loop has now been coupled to the quark sector of the model; we next need to include the contribution to the free energy density from the gauge sector. This is a phenomenological potential, which is a function of Φ and $\bar{\Phi}$, and is required to reproduce the pressure for pure-gluon QCD calculated on the lattice for temperatures around the transition temperature. There are several potentials on the market [45–48] with similar properties. We will first be using the polynomial potential of Ref. [45]

$$\frac{\mathcal{U}}{T^4} = -\frac{1}{2} b_2 \Phi \bar{\Phi} - \frac{1}{6} b_3 (\Phi^3 + \bar{\Phi}^3) + \frac{1}{4} b_4 (\Phi \bar{\Phi})^2, \quad (33)$$

where the constants are

$$b_2 = a_0 + a_1 \left(\frac{T_0}{T} \right) + a_2 \left(\frac{T_0}{T} \right)^2 + a_3 \left(\frac{T_0}{T} \right)^3, \quad (34)$$

$$b_3 = \frac{3}{4}, \quad (35)$$

$$b_4 = \frac{30}{4}, \quad (36)$$

with $a_0 = 6.75$, $a_1 = -1.95$, $a_2 = 2.625$, and, $a_3 = -7.44$.

the finite-temperature fermion contribution can then be written as

We will also use the logarithmic Polyakov-loop potential of Ref. [46]

$$\frac{\mathcal{U}}{T^4} = -\frac{1}{2} a \Phi \bar{\Phi} + b \log \left[1 - 6\Phi \bar{\Phi} + 4(\Phi^3 + \bar{\Phi}^3) - 3(\Phi \bar{\Phi})^2 \right], \quad (37)$$

with

$$a = 3.51 - 2.47 \left(\frac{T_0}{T} \right) + 15.2 \left(\frac{T_0}{T} \right)^2, \quad (38)$$

$$b = -1.75 \left(\frac{T_0}{T} \right)^3. \quad (39)$$

The temperature T_0 is defined by

$$T_0(N_f, \mu_I) = T_\tau e^{-1/(\alpha_0 b(\mu_I))}, \quad (40)$$

where we have modeled the μ_I -dependence in the same way as the μ_B -dependence in [47]

$$b(\mu_I) = \frac{1}{6\pi} (11N_c - 2N_f) - b_{\mu_I} \frac{\mu_I^2}{T_\tau^2}. \quad (41)$$

The parameter $T_\tau = 1.77$ GeV and $\alpha_0 = 0.304$ are determined such that the transition temperature for pure gluon at $\mu_I = 0$ is $T_0 = 270$ MeV [49]. The curvature of the deconfinement transition in μ_I direction is governed by b_{μ_I} , which is chosen as

$$b_{\mu_I} = \frac{16}{\pi} N_f. \quad (42)$$

The full thermodynamic potential is now given by the sum of Eqs. (26), (32), and (33) or (37) respectively. From Eqs. (17)–(18), it is easy to see that Eq. (32) is

³ We express the various contributions to the effective potential in terms of Φ and $\bar{\Phi}$, although they are equal in the present case.

real, thus there is no sign problem at $\mu_B = 0$. We also note that Eqs. (32), (33) and (37) vanish in the limit $T \rightarrow 0$ and the PQM model therefore reduces to the QM model.




FIG. 1. Normalized chiral condensate $\frac{\Delta}{m_q}$ (blue lines) and Polyakov-loop Φ (red lines) as functions of the temperature T for $\mu_B = \mu_I = 0$. See main text for details.

In Fig. 1, we show the normalized chiral condensate $\frac{\Delta}{m_q}$ (blue lines) and the expectation value of the Polyakov loop Φ as functions of the temperature T at $\mu_B = \mu_I = 0$ using the polynomial potential (33). The blue dashed line is the chiral condensate obtained in QM model while the blue solid line is obtained in the PQM model, i.e. with the coupling between the order parameters. Similarly, the red dashed line is obtained using the pure-gluon potential for Φ (with the N_f dependent $T_0 = 208$ MeV), while the red solid line is obtained in the PQM model. We notice that the critical temperature for the chiral transition moves to the right, i.e. to higher temperatures while the transition temperature for deconfinement moves to the left. They are now within a few MeV of each other, with the deconfinement transition occurring at slightly lower temperature than the chiral transition.

V. PHASE DIAGRAM

In this section, we discuss the phase diagram in the μ_I - T plane. In the numerical work below, we set $N_c = 3$, $m_\pi = 140$ MeV, and $f_\pi = 93$ MeV. We vary m_σ between 500 and 600 MeV.

In Fig. 2, we show the chiral (blue line) and pion condensates (red line) as functions of μ_I at zero temperature. We notice the onset of pion condensation which takes place at exactly $\mu_I = \frac{1}{2}m_\pi$ as we will discuss in some detail below. Moreover, the quark condensate de-

creases with μ_I once the pion condensate is nonzero. Finally, all physical quantities, are independent of μ_I from $\mu_I = 0$ all the way up to $\mu_I = \frac{1}{2}m_\pi$. For example, the effective potential is independent of μ_I , implying via Eq. (3) that the isospin density vanishes. This is an example of the Silver Blaze property [50] and was discussed in detail in the context of pion condensation in Refs. [38, 41]. We refer to this region as the vacuum phase.




FIG. 2. Chiral (blue line) and pion condensates (red line) Δ and ρ as functions of the isospin chemical potential μ_I at $T = 0$.

In Fig. 3, we show the phase diagram in the μ_I - T plane for $\mu_B = 0$ without the Polyakov loop, i.e. for the quark-meson model. The blue line is the transition line for the chiral transition and the green line is the transition line for condensation of π^+ , The blue line is defined by the inflection point of the order parameter Δ as functions of T for fixed μ_I . and the black dotted line indicates the crossover from a pion condensate to a BCS state with Cooper pairs.

The onset of pion condensation at $T = 0$ is for $\mu_I = \frac{1}{2}m_\pi$, which is guaranteed by the way we have determined the parameters in the Lagrangian. This was explicitly demonstrated in Ref. [41]. We can understand this result by considering the energy of a zero-momentum pion in the vacuum phase is $m_\pi - 2\mu_I$. If condensations of pions is a second order transition, it must take place exactly at a point where the (medium-dependent) mass of the pion drops to zero, because in the condensed phase there is a massless Nambu-Goldstone mode associated with the breaking of a $U(1)$ symmetry. If one uses matching at tree level, there will be finite corrections to this relation. Likewise, if one uses the effective potential itself to define the pion mass, one uses the pion self-energy at zero external momentum and so the pole of the propagator is not at the physical mass. Again there will be finite corrections and in some cases,

the deviation from the exact result can be significant [38].




FIG. 3. Phase diagram in the μ_I - T plane for $\mu_B = 0$ without Polyakov loop. See main text for details.

The condensation of pions is always a second-order transition with mean-field critical exponents. The order of the transition to a BEC is in agreement with the functional renormalization group application to the QM model in Ref. [38]. The critical isospin chemical potential is fairly constant for temperatures up to approximately $T = 100$ MeV, after which it rapidly increases. For large values of μ_I the critical temperature for pion condensation stays at $T_\rho \approx 187$ MeV. We also notice that the chiral transition temperature line T_χ meets the critical temperature line for pion condensation T_ρ at $(\mu_I^{\text{meet}}, T^{\text{meet}}) \approx (105, 159)$ MeV, and coincide for larger values of μ_I .

As we have seen, we enter the BEC phase when μ_I exceeds $\frac{1}{2}m_\pi$. As μ_I increases the quark mass Δ decreases as shown in Fig. 2. Once $\mu_I > \Delta$, the u -quark and \bar{d} -quark energies, Eqs. (17) and (18), are no longer minimized for $p = 0$, but for $p = \sqrt{\mu_I^2 - \Delta^2}$. This change is a signal of the BEC-BCS crossover. Although the BEC-BCS crossover is not particularly sharp, it is typically defined by the condition $\mu_I > \Delta$ [33, 51]. The crossover starts at $\mu_I = 113$ MeV for $T = 0$ and is almost independent of the temperature, as can be seen from the Figure.

In Fig. 4, we show the phase diagram in the μ_I - T plane at zero baryon chemical potential with the Polyakov loop and $\frac{U}{T^4}$ given by (33). The green line is the critical line for Bose-Einstein condensation of charged pions, the red line is the critical line for deconfinement, and the blue line is the critical line for the chiral transition. Finally, the black dotted line indicates the BEC-BCS transition line. The blue and red lines are defined by the inflection point of the order parameters Δ and Φ as functions of T for fixed μ_I . As




FIG. 4. Phase diagram in the μ_I - T plane for $\mu_B = 0$ with the Polyakov-loop potential Eq. (33). See main text for details.

in the QM model, the transition temperature line T_χ joins the critical temperature for pion condensation at $(\mu_I^{\text{meet}}, T^{\text{meet}}) \approx (116, 187)$ MeV. The transition line for deconfinement lies approximately 15 MeV below the chiral transition line for $\mu_I = 0$ increasing somewhat for large values of μ_I .




FIG. 5. Phase diagram in the μ_I - T plane for $\mu_B = 0$ with the Polyakov-loop potential Eq. (37). See main text for details.

The gap between the chiral and deconfinement line can be reduced by using a logarithmic Polyakov potential (37) instead of Eq. (33) and decreasing the sigma mass. For $m_\sigma = 500$ MeV the two lines basically coincide at $T = 0$ as seen in Fig. 5. The chiral and deconfinement transition line also meet the pion-condensed line at a point for a smaller value of μ_I as compared to Fig. 4, $(\mu_I^{\text{meet}}, T^{\text{meet}}) \approx (75, 164)$ MeV.

For completeness, we show in Fig. 6 the phase diagram in the standard mean-field approximation (sMFA),

which is a common approximation used in the literature, where one ignores the loop corrections to the vacuum potential, i.e. uses Eq. (16) instead of Eq. (26). We find the critical temperature for pion condensation to be smaller than for the one-loop potential in Fig. 4. We also find a first-order transition of the pion condensate above a critical isospin chemical potential $\mu_I \approx 86$ MeV, indicated by the black dot in the figure. This critical point is absent once we go beyond the sMFA, at least in the region of μ_I considered here.




FIG. 6. Phase diagram in the $\mu_I - T$ plane for $\mu_B = 0$ in the standard mean field approximation with the Polyakov-loop potential Eq. (33). See main text for details.

Our phase diagram is in qualitative good agreement with that obtained by Brandt, Endrődi, Schmalzbauer using lattice simulations [16–18], in particular if we use a logarithmic Polyakov loop potential and choose a lower sigma mass of 500 MeV. We believe that the quantitative differences (essentially the temperature dependence of the various transition lines) can mainly be attributed

to the fact that we have two light flavors, while they consider 2+1 flavors; for example the deconfinement transition temperature decreases with the number of quarks and our transition line is consistently higher.⁴ They find that chiral and BEC transition lines meet at a triple point, beyond which they coincide. The latter transition is again found to be second order for all values of μ_I and the scaling analysis is consistent with the $O(2)$ universality class. They computed contour lines of the expectation values of the renormalized Polyakov loop Φ for values 0.2, 0.4, 0.6, 0.8, and 1.0. Given their renormalization prescription for the Polyakov loop, developed in [52], a possible choice for T_{dec} is $\Phi = 1$, which implies that it coincides with T_χ within errors [18]. Finally, we mention that the deconfinement line penetrates smoothly into the BEC phase and that they identify this line with the BEC-BCS transition.

ACKNOWLEDGMENTS

The authors would like to thank G. Endrődi for useful discussions.

Appendix A: Integrals

With dimensional regularization, the momentum integral is generalized to $d = 3 - 2\epsilon$ spatial dimensions. We define the dimensionally regularized integral by

$$\int_p = \left(\frac{e^{\gamma_E} \Lambda^2}{4\pi} \right)^\epsilon \int \frac{d^d p}{(2\pi)^d}, \quad (\text{A1})$$

where Λ is the renormalization scale in the modified minimal subtraction scheme $\overline{\text{MS}}$.

In order to calculate the vacuum part of the effective potential, we need the vacuum integrals

$$\int_p \sqrt{p^2 + M^2} = -\frac{M^4}{(4\pi)^2} \left(\frac{e^{\gamma_E} \Lambda^2}{M^2} \right)^\epsilon \Gamma(-2 + \epsilon) = -\frac{M^4}{2(4\pi)^2} \left(\frac{\Lambda^2}{M^2} \right)^\epsilon \left[\frac{1}{\epsilon} + \frac{3}{2} + \mathcal{O}(\epsilon) \right], \quad (\text{A2})$$

$$\int_p \frac{1}{(p^2 + M^2)^{\frac{3}{2}}} = \frac{4}{(4\pi)^2} \left(\frac{e^{\gamma_E} \Lambda^2}{M^2} \right)^\epsilon \Gamma(\epsilon) = \frac{4}{(4\pi)^2} \left(\frac{\Lambda^2}{M^2} \right)^\epsilon \left[\frac{1}{\epsilon} + \mathcal{O}(\epsilon) \right]. \quad (\text{A3})$$

Appendix B: Parameter fixing

In this Appendix, we briefly discuss the fixing of the model parameters. At tree level, the relations between these parameters and the physical quantities are given by Eqs. (14)–(15). In the on-shell scheme, the counterterms

⁴ By using a smaller value of T_0 , we can bring down the transition line.

are chosen such that they exactly cancel the loop corrections to the self-energies and couplings evaluated on the mass shell, and such that the residues evaluated on shell are unity. Consequently, the renormalized parameters are independent of the renormalization scale and satisfy the tree-level relations [54–56]. In the $\overline{\text{MS}}$ scheme, the counterterms are chosen so that they cancel only the poles in ϵ of the loop corrections. The bare parameters are the same in the two schemes, which means that we can relate the corresponding renormalized parameters. The running parameters in the $\overline{\text{MS}}$ scheme can therefore be expressed in terms of the physical masses m_σ , m_π , and m_q as well as the pion decay constant f_π . In Ref. [53] we found

$$\begin{aligned} m_{\overline{\text{MS}}}^2 &= m^2 + 8ig^2 N_c \left[A(m_q^2) + \frac{1}{4}(m_\sigma^2 - 4m_q^2)B(m_\sigma^2) - \frac{3}{4}m_\pi^2 B(m_\pi^2) \right] - \delta m_{\overline{\text{MS}}}^2 \\ &= m^2 + \frac{4g^2 N_c}{(4\pi)^2} \left[m^2 \log \frac{\Lambda^2}{m_q^2} - 2m_q^2 - \frac{1}{2}(m_\sigma^2 - 4m_q^2) F(m_\sigma^2) + \frac{3}{2}m_\pi^2 F(m_\pi^2) \right], \end{aligned} \quad (\text{B1})$$

$$\begin{aligned} \lambda_{\overline{\text{MS}}} &= \lambda - \frac{12ig^2 N_c}{f_\pi^2} (m_\sigma^2 - 4m_q^2) B(m_\sigma^2) + \frac{12ig^2 N_c}{f_\pi^2} m_\pi^2 B(m_\pi^2) - 4i\lambda g^2 N_c [B(m_\pi^2) + m_\pi^2 B'(m_\pi^2)] - \delta \lambda_{\overline{\text{MS}}} \\ &= \lambda + \left\{ \frac{12g^2 N_c}{(4\pi)^2 f_\pi^2} \left[(m_\sigma^2 - 4m_q^2) \left(\log \frac{\Lambda^2}{m_q^2} + F(m_\sigma^2) \right) + m_\sigma^2 \left(\log \frac{\Lambda^2}{m_q^2} + F(m_\pi^2) + m_\pi^2 F'(m_\pi^2) \right) \right. \right. \\ &\quad \left. \left. - m_\pi^2 \left(2 \log \frac{\Lambda^2}{m_q^2} + 2F(m_\pi^2) + F'(m_\pi^2) \right) \right] \right\}, \end{aligned} \quad (\text{B2})$$

$$g_{\overline{\text{MS}}}^2 = g^2 - 4ig^4 N_c [B(m_\pi^2) + m_\pi^2 B'(m_\pi^2)] - \delta g_{\overline{\text{MS}}}^2 = \frac{m_q^2}{f_\pi^2} \left\{ 1 + \frac{4g^2 N_c}{(4\pi)^2} \left[\log \frac{\Lambda^2}{m_q^2} + F(m_\pi^2) + m_\pi^2 F'(m_\pi^2) \right] \right\}, \quad (\text{B3})$$

$$h_{\overline{\text{MS}}} = h - 2ig^2 N_c m_\pi^2 f_\pi [B(m_\pi^2) - m_\pi^2 B'(m_\pi^2)] - \delta h_{\overline{\text{MS}}} = h \left\{ 1 + \frac{2g^2 N_c}{(4\pi)^2} \left[\log \frac{\Lambda^2}{m_q^2} + F(m_\pi^2) - m_\pi^2 F'(m_\pi^2) \right] \right\} \quad (\text{B4})$$

where $A(m_q^2)$, $B(p^2)$, and $B'(p^2)$ are integrals in $d = 4 - 2\epsilon$ dimensions in Minkowski space. Going to Euclidean space, they can be straightforwardly computed and read

$$\begin{aligned} A(m_q^2) &= \int_k \frac{1}{k^2 - m_q^2} \\ &= \frac{im_q^2}{(4\pi)^2} \left(\frac{\Lambda^2}{m_q^2} \right) \left[\frac{1}{\epsilon} + 1 + \mathcal{O}(\epsilon) \right], \end{aligned} \quad (\text{B5})$$

$$\begin{aligned} B(p^2) &= \int_k \frac{1}{(k^2 - m_q^2)[(k+p)^2 - m_q^2]} \\ &= \frac{i}{(4\pi)^2} \left(\frac{\Lambda^2}{m_q^2} \right) \left[\frac{1}{\epsilon} + F(p^2) + \mathcal{O}(\epsilon) \right], \end{aligned} \quad (\text{B6})$$

$$B'(p^2) = \frac{i}{(4\pi)^2} F'(p^2), \quad (\text{B7})$$

where we have defined

$$F(p^2) = 2 - 2r \arctan \left(\frac{1}{r} \right), \quad (\text{B8})$$

$$F'(p^2) = \frac{4m_q^2 r}{p^2(4m_q^2 - r^2)} \arctan \left(\frac{1}{r} \right) - \frac{1}{p^2}, \quad (\text{B9})$$

with $r = \sqrt{\frac{4m_q^2}{p^2} - 1}$.

The running parameters satisfy the following renor-

malization group equations

$$\Lambda \frac{dm_{\overline{\text{MS}}}^2(\Lambda)}{d\Lambda} = \frac{8N_c m_{\overline{\text{MS}}}^2(\Lambda) g_{\overline{\text{MS}}}^2(\Lambda)}{(4\pi)^2}, \quad (\text{B10})$$

$$\Lambda \frac{dg_{\overline{\text{MS}}}^2(\Lambda)}{d\Lambda} = \frac{8N_c g_{\overline{\text{MS}}}^4(\Lambda)}{(4\pi)^2}, \quad (\text{B11})$$

$$\Lambda \frac{d\lambda_{\overline{\text{MS}}}(\Lambda)}{d\Lambda} = \frac{16N_c}{(4\pi)^2} [\lambda_{\overline{\text{MS}}}(\Lambda) g_{\overline{\text{MS}}}^2(\Lambda) - 6g_{\overline{\text{MS}}}^4(\Lambda)] \quad (\text{B12})$$

$$\Lambda \frac{dh_{\overline{\text{MS}}}(\Lambda)}{d\Lambda} = \frac{4N_c g_{\overline{\text{MS}}}^2(\Lambda) h_{\overline{\text{MS}}}(\Lambda)}{(4\pi)^2}. \quad (\text{B13})$$

The solutions to Eqs. (B10)–(B13) are

$$m_{\overline{\text{MS}}}^2(\Lambda) = \frac{m_0^2}{1 - \frac{4g_0^2 N_c}{(4\pi)^2} \log \frac{\Lambda^2}{\Lambda_0^2}}. \quad (\text{B14})$$

$$g_{\overline{\text{MS}}}^2(\Lambda) = \frac{g_0^2}{1 - \frac{4g_0^2 N_c}{(4\pi)^2} \log \frac{\Lambda^2}{\Lambda_0^2}}, \quad (\text{B15})$$

$$\lambda_{\overline{\text{MS}}}(\Lambda) = \frac{\lambda_0 - \frac{48g_0^4 N_c}{(4\pi)^2} \log \frac{\Lambda^2}{\Lambda_0^2}}{\left(1 - \frac{4g_0^2 N_c}{(4\pi)^2} \log \frac{\Lambda^2}{\Lambda_0^2} \right)^2}, \quad (\text{B16})$$

$$h_{\overline{\text{MS}}}(\Lambda) = \frac{h_0}{1 - \frac{2g_0^2 N_c}{(4\pi)^2} \log \frac{\Lambda^2}{\Lambda_0^2}}, \quad (\text{B17})$$

where m_0^2 , g_0^2 , λ_0 and h_0 , are the values of the running

parameters at the scale Λ_0 . We choose Λ_0 to satisfy

$$\log \frac{\Lambda_0^2}{m_q^2} + F(m_\pi^2) + m_\pi^2 F'(m_\pi^2) = 0. \quad (\text{B18})$$

One can now evaluate Eqs. (B1)–(B4) at the scale $\Lambda =$

Λ_0 to find m_0^2 , λ_0 , g_0^2 , and h_0 . Inserting Eqs. (B14)–(B17) into Eq. (25) using the results for m_0^2 , λ_0 , g_0^2 , and h_0 , we obtain the final result Eq. (26).

-
- [1] K. Rajagopal and F. Wilczek, *At the frontier of particle physics*, Vol. 3 (World Scientific, Singapore, p 2061) (2001).
- [2] M. G. Alford, A. Schmitt, K. Rajagopal, and T. Schäfer, *Rev. Mod. Phys.* **80**, 1455 (2008).
- [3] K. Fukushima and T. Hatsuda, *Rept. Prog. Phys.* **74**, 014001 (2011).
- [4] Y. Aoki, Z. Fodor, S. Katz, and K. Szabo, *Phys. Lett. B* **643**, 46 (2006).
- [5] Y. Aoki, S. Borsanyi, S. Dürr, Z. Fodor, S.D. Katz et al., *JHEP* **09** 06, 088 (2009),
- [6] S. Borsanyi et al. (Wuppertal-Budapest Collaboration), *JHEP* **10** **09**, 073 (2010).
- [7] A. Bazavov, T. Bhattacharya, M. Cheng, C. DeTar, H. Ding et al., *Phys. Rev. D* **85**, 054503 (2012).
- [8] J. B. Kogut, M. A. Stephanov, D. Toublan, and J. J. M. Verbaarschot, *Nucl. Phys. B* **582**, 477 (2000).
- [9] J. B. Kogut (Illinois U., Urbana), M. A. Stephanov, and D. Toublan, *Phys. Lett. B* **464**, 183 (1999).
- [10] D. T. Son and M. A. Stephanov, *Phys. Rev. Lett.* **86**, 592 (2001).
- [11] G. S. Bali, F. Bruckmann, G. Endrődi, Z. Fodor, S. D. Katz, S. Krieg, A. Schafer, and K. K. Szabo, *JHEP* **12** **02**, 044 (2012).
- [12] D. Kharzeev, K. Landsteiner, A. Schmitt, and H.-U. Yee (ed) *Lect. Notes Phys.* **871**, 1 (2013).
- [13] J. O. Andersen, W. R. Naylor, A. Tranberg, *Rev. Mod. Phys.* **88**, 025001 (2016).
- [14] J. B. Kogut and D. K. Sinclair, *Phys. Rev. D* **66**, 014508 (2002).
- [15] J. B. Kogut and D. K. Sinclair, *Phys. Rev. D* **66** 034505 (2002).
- [16] B. B. Brandt and G. Endrődi, *PoS LATTICE 2016*, 039 (2016).
- [17] B. B. Brandt, G. Endrődi, and S. Schmalzbauer, *EPJ Web Conf.* **175**, 07020 (2018).
- [18] B. B. Brandt, G. Endrődi, and S. Schmalzbauer, *Phys. Rev. D* **97**, 054514 (2018).
- [19] K. Splittorff, D. T. Son, M. A. Stephanov, *Phys. Rev. D* **64**, 016003 (2001).
- [20] M. Loewe and C. Villavicencio, *Phys. Rev. D* **67**, 074034 (2003).
- [21] E. S. Fraga, L. F. Palhares and C. Villavicencio, *Phys. Rev. D* **79**, 014021 (2009).
- [22] S. Carignano, L. Lepori, A. Mammarella, M. Mannarella and G. Pagliaroli, *Eur. Phys. J. A* **53**, 35 (2017).
- [23] D. Toublan and J. B. Kogut, *Phys. Lett. B* **605**, 129 (2005).
- [24] B. Klein, D. Toublan and J. J. M. Verbaarschot, *Phys. Rev. D* **68**, 014009 (2003).
- [25] M. Frank, M. Buballa, and M. Oertel, *Phys. Lett. B* **562**, 221 (2003).
- [26] D. Toublan, and J. B. Kogut, *Phys. Lett. B* **564**, 212 (2003).
- [27] A. Barducci, R. Casalbuoni, G. Pettini, and L. Ravagli *Phys. Rev. D* **69**, 096004 (2004).
- [28] L. He, and P.-F. Zhuang, *Phys. Lett. B* **615**, 93 (2005).
- [29] L. He, M. Jin and P.-F. Zhuang, *Phys. Rev. D* **71**, 116001, (2005).
- [30] L. He, M. Jin, and P.-F. Zhuang, *Phys. Rev. D* **74**, 036005 (2006).
- [31] D. Ebert and K. G. Klimenko, *J. Phys. G* **32**, 599 (2006).
- [32] D. Ebert and K. G. Klimenko, *Eur. Phys. J. C* **46**, 771 (2006).
- [33] G.-F. Sun, L. He, and P.-F. Zhuang, *Phys. Rev. D* **75**, 096004 (2007).
- [34] J. O. Andersen and L. Kyllingstad, *J. Phys. G* **37**, 015003 (2009).
- [35] H. Abuki, R. Anglani, R. Gatto, M. Pellicoro, and M. Ruggieri, *Phys. Rev. D* **79**, 034032 (2009).
- [36] C.-F. Mu, L. He, and Y. Liu, *Phys. Rev. D* **82**, 056006 (2010).
- [37] T. Xia, L. He and P. Zhuang, *Phys. Rev. D* **88**, 056013 (2013).
- [38] K. Kamikado, N. Strodthoff, L. von Smekal, and J. Wambach *Phys. Lett. B* **718**, 1044 (2013).
- [39] H. Ueda, T. Z. Nakano, A. Ohnishi, M. Ruggieri, and K. Sumiyoshi, *Phys. Rev. D* **88**, 074006 (2013).
- [40] R. Stiele, E. S. Fraga and J. Schaffner-Bielich, *Phys. Lett. B* **729**, 72 (2014).
- [41] J. O. Andersen and P. Kneschke, *Phys. Rev. D* **97**, 076005 (2018).
- [42] T. D. Cohen and S. Sen, *Nucl. Phys. A* **942**, 39 (2015).
- [43] L. G. Yaffe and B. Svetitsky, *Phys. Rev. D* **26**, 963 (1982).
- [44] B. Svetitsky and L. G. Yaffe, *Nucl. Phys. B* **210**, 423 (1982).
- [45] C. Ratti, M. A. Thaler, and W. Weise, *Phys. Rev. D* **73**, 014019 (2006).
- [46] S. Roessner, C. Ratti and W. Weise, *Phys. Rev. D* **75**, 034007 (2007).
- [47] B.-J. Schaefer, J. M. Pawłowski, and J. Wambach, *Phys. Rev. D* **76**, 074023 (2007).
- [48] K. Fukushima, *Phys. Rev. D* **77**, 114028. (2008).
- [49] Karsch, F., E. Laermann, and A. Peikert, *Nucl. Phys. B* **605**, 579 (2001).
- [50] T. D. Cohen, *Phys. Rev. Lett.* **91**, 222001 (2003).
- [51] T. Brauner, K. Fukushima, and Y. Hidaka, *Phys. Rev.*

- D **80**, 074035 (2009); *ibid* D **81**, 119904 (2010).
- [52] S. Borsanyi et al, JHEP 12 **08**, 126 (2012).
- [53] P. Adhikari, J. O. Andersen, and P. Kneschke, Phys. Rev. D **96**, 016013 (2017).
- [54] A. Sirlin, Phys. Rev. D **22**, 971 (1980).
- [55] A. Sirlin, Phys. Rev. D **29**, 89 (1984).
- [56] M. Bohm, H. Spiesberger, and W. Hollik, Fortsch. Phys. **34**, 687 (1986).