

"This is the peer reviewed version of the following article: Rørvik, K. A., Dessen, J. E., Åsli, M., Thomassen, M. S., Hoås, K. G., & Mørkøre, T. (2018). Low body fat content prior to declining day length in the autumn significantly increased growth and reduced weight dispersion in farmed Atlantic salmon *Salmo salar* L. *Aquaculture Research*, 49(5), 1944-1956., which has been published in final form at <https://doi.org/10.1111/are.13650> This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions."

1 **Low body fat content prior to declining day length in the autumn significantly increased**
2 **growth and reduced weight dispersion in farmed Atlantic salmon *Salmo salar* L.**

3

4 Kjell-Arne Rørvik^{1,2}, Jens-Erik Dessen^{1,2*}, Magnus Åsli^{1,2†}, Magny S. Thomassen², Kjellrun
5 G. Hoås¹ & Turid Mørkøre^{1,2}

6

7 ¹Nofima, NO-1432 Ås, Norway

8 ²Department of Animal and Aquaculture Sciences, Norwegian University of Life Sciences,
9 NO-1432 Ås, Norway

10

11 *Corresponding author: Jens-Erik Dessen; Nofima, 1432 Ås, Norway; Tel: +47 979 52 768;

12 Email: jens-erik.dessen@nofima.no

13 †Current address: Cermaq Group As, NO-0102 Oslo, Norway

14

15

16

17

18

19

20

21

22 **Running head:** Body fat and growth in Atlantic salmon

23 **Key words:** Salmon, growth response, body lipids, seasonal cues

24 **ABSTRACT:**

25

26 Based on the regulatory effects of body fat on appetite and seasonal variations in fat
27 deposition and growth of Atlantic salmon, the present study tested the hypothesis that body fat
28 content prior to declining day length in the autumn can significantly modulate growth rate.
29 The growth rate of salmon (mean initial body weight, BW=2.3 kg) with different muscle fat
30 content prior to autumn, subjected to natural photoperiod and temperature, during a 7-months
31 period (mean final BW=6.6 kg) was studied. In August, three fish groups (HF, LF and 0.5LF
32 group) with significantly different muscle fat content (HF=16.4%, LF=13.2% and
33 0.5LF=11.3%), individually marked with PIT-tag, were mixed into the four net pens and fed a
34 standard high-energy diet until March the following year. The muscle fat content prior to the
35 autumn had a highly significant ($P < 0.0001$) effect on growth during the seven month main-
36 dietary period, even after identical fat stores among the groups were restored, indicating a
37 more complex explanation than just a lipostatic regulation mechanism. Mean thermal growth
38 coefficients were HF=2.9, LF=3.4 and 0.5 LF=3.9, resulting in increased final weight gain for
39 LF and 0.5LF of 590 g. and 980 g., respectively, compared to the HF group. The LF groups
40 obtained a significantly higher homogeneity in BW and shape than HF fed fish in March,
41 optimizing automatic gutting and filleting at slaughter. The improved growth response among
42 the LF groups by reducing lipid levels can potentially be utilized in closed and semi-closed
43 production units where photoperiod can be manipulated.

44

45

46

47

48

49 **INTRODUCTION:**

50

51 Fish that encounter setbacks induced by nutritional deficit, feed deprivation or sub-optimal
52 conditions often display increased feed consumption (hyperphagia) and compensatory growth
53 (CG) when circumstances are normalized (Ali, Nicieza, & Wootton, 2003; Foss & Imsland,
54 2002; Metcalfe & Monaghan, 2001). The degree of CG in fish vary and is often categorized
55 based on the growth catch-up ability (Ali et al., 2003). Feed restriction or deprivation induce
56 changes in body energy by depleting lipid stores, and during the course of CG and
57 hyperphagia, body weight and lipid reserves are gradually restored (Ali et al., 2003; Bull &
58 Metcalfe, 1997; Jobling & Miglavs, 1993; Metcalfe & Thorpe, 1992). The lipostatic model is
59 often discussed within the circumstances of CG responses in fish (Jobling & Johansen, 1999;
60 Johansen, Ekli, Stangnes, & Jobling, 2001). The lipostatic regulation hypothesis identifies
61 adipose tissue and stored lipids to have an important role in governing appetite (Jobling &
62 Johansen, 1999; Keesey & Corbett, 1984; Kennedy, 1953). The model implies that the
63 amounts of stored fat has a negative feedback control on feed intake and is important for the
64 regulation of energy homeostasis. Hence, CG is not only a response to recover body weight,
65 but also a strong response to restore lipid levels and thereof CG will cease once this is
66 achieved (Ali et al., 2003; Jobling & Johansen, 1999; Johansen, Ekli, & Jobling, 2002).
67 Johansen et al., (2002) showed that altering body lipids of juvenile salmon by dietary
68 administration of low-fat feeds yield similar growth responses as deprivation or feed
69 restriction *per se*.

70

71 In modern high-fat diets for salmonids, lipids of marine and vegetable origin are the main
72 sources of energy and support growth efficiently if essential fatty acids requirements are met
73 (Bell et al., 2001; Thomassen & Røsjø, 1989; Torstensen, Lie, & Frøyland, 2000). Because

74 salmonids have a high ability to utilize large amount of lipids efficiently for growth, high-fat
75 diets with up to 380 g kg⁻¹ of fat are commonly used in intensive salmon farming (Torrissen et
76 al., 2011). However, salmonids also have the capacity to store large amounts of excess fat as
77 triacylglycerols mainly in the muscle and visceral cavity (Aursand, Bleivik, Rainuzzo, Leif, &
78 Mohr, 1994). Body lipid content of farmed salmonids correlates with fish size, dietary fat
79 level and feed intake (Aksnes, 1995; Hemre & Sandnes, 1999; Torstensen, Lie, & Hamre,
80 2001). Like other anadromous species, Atlantic salmon display seasonal changes in feed
81 intake, growth and lipid deposition during the seawater phase (Mørkøre & Rørvik, 2001).
82 Farmed Atlantic salmon display elevated deposition of lipids in muscle and increased
83 retention of lipids in whole body during declining day length in autumn, with a concomitant
84 increase in feed intake, somatic growth and condition factor (CF) (Alne, Oehme, Thomassen,
85 Terjesen, & Rørvik, 2011; Dessen, Weihe, Hatlen, Thomassen, & Rørvik, 2017; Mørkøre &
86 Rørvik, 2001; Rørvik et al., 2010). This is particularly pronounced for salmon reared at high
87 latitudes that experience long winters and late spring, which results in reduced lipid levels and
88 CF prior to summer and autumn.

89

90 The recent increase in automation of fish processing at slaughter requires uniform body
91 weight (BW) and shape among the salmon for optimal efficiency and quality of products such
92 as gutted fish and fillets. Increased uniformity of BW and CF reduces the need for manual
93 gutting/filleting of very small or large individuals. Due to this, the homogeneity in body shape
94 and mass of salmonids are important parameters in salmon farming industry and low
95 dispersion in BW and CF are beneficial at time of harvest. The homogeneity of BW may be
96 strongly influenced by events occurring during the production cycle, i.e. disease outbreaks,
97 handling stress, reduced seawater tolerance or competition of feed (McLoughlin, Nelson,
98 McCormick, Rowley, & Bryson, 2002; Ryer & Olla, 1996; Taksdal et al., 2007; Usher,

99 Talbot, & Eddy, 1991). The dispersion in the distribution of BW, length and CF are often
100 assessed by calculating the coefficient of variation (CV). The CV of BW for farmed salmon
101 grown from 70 until 300 g. and from 60 until 500 g. fed either in excess or restrictively for
102 period followed by unrestricted feeding, are reported to vary from 9 to 13% and 16 to 21%,
103 respectively (Johansen et al., 2001). In the latter study, no significant differences was
104 observed in the CV of BW between fish fed in excess and fish fed restrictively.

105

106 The majority of studies regarding growth responses related to lipid content are based on in-
107 house laboratory experiments with small juvenile salmonids under constant conditions. To our
108 knowledge, few have investigated grow out salmon with different lipid content subjected to
109 seasonal environmental changes in photoperiod and temperature. Due to the regulatory effects
110 of body fat on appetite and the observed fat storage in salmon linked to the seasonal cues, the
111 present study tested the hypothesis that lipid status prior to declining day length in the autumn
112 functions as a significant growth regulator. Accordingly, the growth rate for three groups of
113 salmon with different muscle fat content prior to autumn, subjected to natural photoperiod and
114 temperature, was studied throughout a 7-months period. About each second month, weight
115 samplings and analysis of muscle fat content was conducted to investigate any relationship
116 between fat accumulation and periodic growth rate, and to identify the duration of a potential
117 lipostatic regulatory effect. Changes in visceral fat, CF, length, and the dispersion in BW and
118 CF were further assessed.

119

120 **MATERIAL & METHODS:**

121

122 This experiment was conducted in accordance with laws and regulations that control
123 experiments and procedures in live animals in Norway, as overseen by the Norwegian Animal

124 Research Authority. Stunning and sampling of fish were performed in accordance with the
125 Norwegian Animal Welfare act. Fish were treated as production fish up to the point of tissue
126 sampling which was only conducted after the fish were put to death.

127

128 The experiment was conducted in seawater on the Norwegian west coast (Ekkilsøy, Norway
129 3° 03' N, 7° 35' E) at Nofima research center from August 2011 to March 2012. In July 2010,
130 the fish were transferred to seawater as S1 smolt, at which time the BW was 62 g. From the
131 10 to 12 of May 2011, the post-smolt were re-stocked into three net-pens (343 m³) with 650
132 fish per pen. Prior to this, all individual fish were measured for weight and length, and tagged
133 using passive integrated transmitter tags (PIT-tags) placed in the body cavity just posterior to
134 the gut. The average BW per pen was 1085 g. (SD = 79 g.) and each pen received different
135 dietary treatments: a high-fat diet (HF), a low-fat high-protein diet (LF) or half the ration of
136 the low-fat high-protein diet (0.5LF). The 0.5LF-group were given half the amount of the feed
137 provided to fish administrated the LF-diet the day before. Skretting (Averøy, Norway)
138 produced the feeds and the composition of the HF diet was (wet weight, as is basis): dry
139 matter 93.4%, crude protein 33.5%, crude lipid 34.1%, nitrogen-free extract (NFE) 21.2%,
140 ash 4.6% and gross energy of 25.1 MJ kg⁻¹. The composition of the LF diet was (wet weight,
141 as is basis): dry matter 91.7%, crude protein 49.9%, crude lipid 17.5%, NFE 17.1%, ash 7.2%
142 and gross energy 21.7 MJ kg⁻¹. The three dietary treatments were fed from 12 of May until 9
143 of August (pre-dietary phase). May 12th, the fish were sampled for analysis of initial muscle
144 fat content and biometric data. The analysis showed the following (mean ± SE, n = 30): BW:
145 1087 ± 97g, initial muscle fat: 12.2 ± 1.1% and initial CF: 1.10 ± 0.06. After ending the pre-
146 dietary phase, the PIT-tag, BW and length of all individual fish in the three pens were
147 recorded. In addition, fish from each pen were sampled for analysis of muscle and visceral fat
148 content. The pre-dietary feeding phase generated three fish groups with significantly different

149 ($P < 0.05$) muscle fat content, visceral fat and visceral mass (Table 1). During the pre-dietary
150 phase, 2.5%, 0.6% and 0.3% fish died in the HF, LF and 0.5LF group, respectively. The
151 majority of mortality occurred from May until mid-June and was not related to any disease
152 outbreak (non-specific mortality).

153

154 **(Table 1).**

155

156 At the 10 to 11 of August, the fish were restocked from the three original pens used in pre-
157 dietary phase into four new pens (125 m³). Each of the four pens contained 50 fish from each
158 of the three pre-dietary treatments (HF, LF and 0.5LF), 150 fish in total (Fig 1). During the
159 period from 11 of August until termination at 20 of March 2012 (main-dietary phase), the
160 pens were fed isonitrogenous and isoenergetic diets produced by Ewos (Bergneset, Norway)
161 (Table 2). The current experiment was an integrated part of a large study where potential
162 effects of dietary oil source were investigated. Therefore, two pens in the main-dietary phase
163 were fed a diet with a marine oil profile (MO), whereas the two other pens were fed a diet
164 with a rapeseed oil profile (RO). The marine oil diet (MO) had an inclusion of 70 % South
165 American fish oil and 30 % of rapeseed oil. The rapeseed oil diet (RO) had an inclusion of 70
166 % rapeseed oil and 30 % South American fish oil. During the main-dietary phase, the pellet
167 size was changed from 7 to 9 mm in December due to the increase in BW of the fish.

168

169 **(Fig. 1 and Table 2)**

170

171 In both periods, feed was administrated using automatic feeders (Betten Maskinstasjon AS,
172 Vågland, Norway) and uneaten feed was collected as described in Einen, Mørkøre, Røra &
173 Thomassen (1999) and corrected for the recovery of dry matter as described by Helland,

174 Grisdale-Helland & Nerland (1996). The fish groups (except the 0.5LF group during the pre-
175 dietary phase) were fed to satiation and the feed ration was set at 5-10 % in excess (*ad libitum*
176 feeding). The fish were fed four times a day until October 2011, after this, the fish were fed
177 three times a day until termination in March 2012. Adjustments of the feed ration was done
178 according to the daily amount of uneaten feed collected. Due to the stocking of 50 fish from
179 each of the pre-dietary treatments into each net pen, it was not possible to determine the feed
180 intake or feed utilization of the different pre-dietary groups during the main-dietary phase.
181 The pens were checked for mortalities daily and the dead fish were collected and weighed.
182 The fish were exposed to natural variations in photoperiod and sea temperature during the
183 experiment (Fig. 2).

184

185 **(Fig. 2)**

186

187 Three samplings were performed during the main-dietary phase; from 9 to 11 October 2011,
188 from 6 to 9 December 2011 and the final sampling and termination of the experiment was
189 conducted from 20 to 22 March 2012. At each sampling, all fish were anaesthetized (MS-222
190 metacaine 0.1 g L⁻¹, Alparma, Animal Health, Hampshire, UK) and the PIT-tag, fork length
191 and weight of each individual fish were recorded. All fish were starved two days prior to the
192 samplings in August and October, and three days prior to the samplings in December and
193 March to avoid feed matter in the gastrointestinal system. At each sampling, 10 fish from each
194 pre-dietary group in all the pens were sampled. The sampled fish at each sampling point were
195 selected so that the mean weight corresponded to the mean weight of all the fish in the
196 respective fish group within each pen (as all possible fish were weighted and PIT-tag read).
197 After anesthetization, a blow to the head was used to kill fish sampled for analysis. Then the
198 gill arches were cut and the fish were bled out in ice seawater. Length and weight of each

199 individual fish sampled for analysis were recorded after bleeding and the fish visually tagged.
200 The fish were then gutted and filleted by hand during the pre-rigor state. Norwegian Quality
201 Cut, NQC (NS9401, 1994) from the left fillet was photographed and the fat content was
202 predicted by digital image analyses (PhotoFish, AKVAgroup, Bryne, Norway), as described
203 by Folkestad et al. (2008). The visceral mass of the sampled fish were pooled on group level,
204 homogenised and frozen at - 20°C for later analyses of total lipid content as described by
205 Folch, Less & Stanley (1957). The proximate composition of crude protein, lipid (acidic-
206 hydrolysis method), starch and moisture of the diets were analysed according to the methods
207 described by Oehme et al. (2010). To determine the fatty acid (FA) composition of the diets,
208 lipids were first extracted according to Foch et al. (1957), and a sample of 2 ml from the
209 chloroform–methanol phase was dried under N₂ gas, then the residual lipid extract was trans-
210 methylated overnight with 2',2'-dimethoxypropane, methanolic HCl and benzene at room
211 temperature according to Mason & Waller (1964). Finally, the methyl esters were separated
212 by gas chromatography and individual FA were identified as described in Røsjø et al. (1994).

213

214 The growth rate of the fish are presented as the thermal growth coefficient (TGC), and were
215 calculated as described by Iwama & Tautz (1981): $TGC = [(M_1^{1/3} - M_0^{1/3}) \times (\Sigma T)^{-1}] \times 1000$,
216 where M_0 and M_1 are the initial and final BW, respectively, and ΣT is the sum of day degrees
217 during the period (feeding days x average temperature, °C). The mean TGC for the total
218 main-dietary phase was calculated as the weighted arithmetic mean of the periodical TGC to
219 balance these values in relation to their relative contribution to the weight gain.

220

221 All fish sampled and killed for analysis were starved and bled. The calculation of visceral-
222 somatic index is therefore based on BW with minimal blood content and no feed material in
223 the gastrointestinal system. Visceral-somatic index (VSI), was calculated as: $Y (g) \times BW (g)^{-1}$

224 $\times 100$, where Y is the measured visceral mass. The visceral mass was defined as all mass in
225 the abdominal cavity except liver, heart, kidney and swim bladder. The CF was calculated as:
226 $100 \times BW \text{ (g)} \times \text{fork length (cm)}^{-3}$. The dispersion in the distribution of BW, length and CF
227 were assessed by calculating the CV: $(\text{standard deviation} \times \text{mean value}^{-1}) \times 100$.

228

229 The results were analysed by the General Linear Model (GLM) procedure in the SAS 9.4
230 computer software (SAS Institute Inc., Cary, NC, USA). Mean results per fish group in each
231 pen were initially subjected to a two-way analysis of variance (ANOVA) to evaluate the
232 effects of muscle fat content due to the pre-dietary phase (0.5LF, LF and HF), main-dietary
233 treatment (oil source; MO and RO-diet) and their interaction (pre-diet \times main-diet). As the
234 statistical analysis showed that neither oil source nor the interaction term has significant
235 effects on the traits studied, the data was analysed using pre-dietary treatment as the only
236 experimental factor (one-way ANOVA). Significant differences among experimental groups
237 within treatments were indicated by Duncan's multiple range test. Least-square means
238 (lsmeans) comparison were also used to identify differences among variables within
239 treatments. The Pearson product-moment correlation coefficient was used to describe the
240 association between two variables. Linear regression analysis were conducted using Microsoft
241 excel. The proportion of total variance explained by the model was expressed by R^2 and the
242 level of significance was chosen at $P \leq 0.05$. Tendencies was identified at $P = 0.05 - 0.1$. The
243 results are presented as means \pm SEM, if not otherwise stated.

244

245 No significant effects of the main-dietary treatment (RO and MO-diet) or interaction term
246 (main \times pre-diet) *per se* were detected on the traits examined during the main-dietary phase.
247 Thus, only the effects of body fat content due to the pre-dietary treatment are presented in the

248 results. No significant differences in mortality among the pre-dietary groups were observed
249 during the main-dietary phase (24 out of 650 fish, 3.6%).

250

251 **RESULTS:**

252

253 The muscle fat content increased by 8.1% for 0.5LF fish, 5.6% for the LF group and 3.6% for
254 HF group from August to October (Fig 4A1). Thus, during an 8-week period of declining day
255 length, the initial significant differences in muscle fat content was equilibrated. TGC was
256 highest for the 0.5LF group, intermediate for the LF group and lowest for the HF group (Fig
257 5A). The growth rate and the increase in muscle fat content from August to October showed a
258 significant positive linear relationship, and the increase in muscle fat explained 81% of the
259 variation in growth (Fig 3). From August to October, the growth rates were therefore highly
260 affected by the pre-dietary treatment (ANOVA: $R^2 = 0.97$, $P < 0.001$). The muscle fat did not
261 differ significantly between the pre-dietary treatments in October or December (Fig. 4A1), but
262 pre-diet still significantly influenced the growth rates (ANOVA: $P < 0.05$, $R^2 = 0.51$) and the
263 TGCs were similar, relatively, to the period from August to October (0.5LF > LF > HF),
264 although no significant differences was found between LF and HF group. In the period
265 December to March, the TGC for the 0.5LF and LF group were significantly higher ($P <$
266 0.05) than the HF group (Fig 5A). At the end of the main-dietary phase, the muscle fat content
267 of the LF group was significantly lower ($P < 0.05$) than the 0.5LF group, and tended to be
268 lower ($P < 0.1$) than the HF group (Fig 4A2).

269

270 **(Fig. 3 and 4)**

271

272 The BW of the LF group reached a similar BW as the HF fish in October, whereas the 0.5LF
273 group reached a similar BW as the HF group in December (Fig 4B1). At the end of the trial in
274 March, the LF group (6.87 ± 0.07 kg.) had a significantly higher ($P < 0.05$) BW than the HF
275 group (6.40 ± 0.16 kg.) (Fig 4B2). The 0.5LF group (6.62 ± 0.12 kg.) had numerical higher
276 BW than the HF group, however, no statistically significant difference was detected. From
277 August 2011 to March 2012, the 0.5LF group gained 980 g. and the LF group gained 590 g.
278 more relative to the BW of the HF group (Fig 5B). The overall weighted mean TGC during
279 the main-dietary phase were 3.9 for the 0.5LF group, 3.4 for the LF group and 2.9 for the HF
280 group. Hence, the pre-dietary treatment and consequently the fat status in August 2011 had a
281 clear and significant effect on growth, weight gain and the changes in BW throughout the
282 whole main-dietary phase, with a total duration of seven months.

283

284 **(Fig. 5 and 6)**

285

286 No significant differences in length between LF and HF group were detected during the trial
287 (Fig 6B1). The strong growth spurt of the 0.5 LF group resulted in no significant differences
288 in length between the 0.5 LF (75.9 ± 0.2 cm.) and HF group (76.4 ± 0.8 cm.) at the trial
289 termination in March. However, the LF (77.9 ± 0.1 cm.) group was significantly longer ($P <$
290 0.05) than the 0.5LF group (Fig 6B2). The 0.5LF group that had the lowest CF in August,
291 ended up having the significantly highest CF at termination (Fig. 6A1 and A2). The overall
292 development in CF correlated well with the changes in muscle fat during the study ($r = 0.98$,
293 $P < 0.01$). Significant positive overall correlations were also observed between the final CF
294 and mean TGC ($r = 0.88$; $P < 0.001$), and between the final CF and total weight gain ($r =$
295 0.88 ; $P < 0.001$).

296

297 The visceral fat content of the HF group was consistently highest, although only significant in
298 October (Fig 7). The VSI of the LF group (8.5 ± 0.1) was significantly lower (ANOVA: $P <$
299 0.01) than the HF group (9.0 ± 0.1) in October, whereas the VSI of the 0.5LF group ($8.7 \pm$
300 0.1) was not different from the LF or HF group. No significant differences in VSI were
301 detected in December (overall mean; VSI: 8.8 ± 0.1) or March (overall mean; VSI: 9.8 ± 0.2).

302

303 **(Fig. 7)**

304

305 The 0.5LF group had the highest CV_{BW} at the end of the pre-dietary phase (Fig 8A). From
306 August to October, the CV_{BW} of the 0.5LF group decreased and no significant difference in
307 CV_{BW} was observed at the samplings in October and December. However, at termination in
308 March, the HF group had a significantly ($P < 0.05$) higher CV_{BW} compared to both LF and
309 0.5LF group. The CV_{CF} was lowest for the LF group and similar for the HF and 0.5LF group
310 at the end of the pre-dietary phase (Fig 8B). At the sampling in October, after the large
311 increase in fat deposition, growth and CF, the 0.5LF group had the highest CV_{CF} . The
312 variation within the CV of CF for the 0.5LF group was at this time very high and no
313 significant differences between the groups was detected. The CV_{CF} for the HF group
314 increased gradually from October to March. In line with the CV_{BW} , the HF group had a
315 significantly ($P < 0.05$) higher CV_{CF} compared to the 0.5LF and LF group at termination. No
316 significant differences in the CV_{LENGTH} was detected during the experiment (results not
317 shown).

318

319 **(Fig. 8)**

320

321

DISCUSSION:

323

324 The coinciding increase in fat and improved growth shown by the 0.5LF and LF group
325 compared to the HF group in the beginning of main-dietary phase (August and September),
326 seem to reflect a growth response similar to CG and lipostatic regulation observed in previous
327 studies in the field and laboratory (Ali et al., 2003; Jobling & Johansen, 1999; Johansen et al.,
328 2002, 2001). The obtained growth rates, fat increase and weight gain from August to October,
329 together with the high feed intake (on pen basis), indicate that the 0.5LF and LF group had
330 increased feed consumption and hyperphagic behaviour. In addition to the high growth rate of
331 the 0.5LF and LF groups, the increase in muscle and visceral fat content during August and
332 September were substantial for these two groups. However, the muscle fat of the HF group
333 also increased during this period (16.4% → 20.0%). The TGC of the HF group had an average
334 of 3.0, which is regarded as a normal and sufficient growth rate (Austreng, Storebakken, &
335 Åsgård, 1987; Thorarensen & Farrell, 2011). Thus, improved growth in the LF groups from
336 August to October, compared to the HF group, is not a result of impaired growth due to
337 adiposity in the latter group, but rather a stronger response among the fish in the LF and
338 0.5LF group. The growth responses from August to October differ from the observations of
339 Johansen, Sveier, & Jobling (2003), where Atlantic salmon fed a high fat diet during both the
340 build-up and main phase, maintained their body fat levels after the build-up phase, at the same
341 time as feed intake was down-regulated and growth impaired. In the present study, the salmon
342 were exposed to natural photoperiod, as opposed to the study by Johansen et al. (2003), where
343 the salmon were held under continuous light (24L:0D). It has been suggested that reduction in
344 day length is an important environmental factor that trigger the salmon to assess its current
345 mass during this time of the year (Maclean & Metcalfe, 2001). It may also apply for energy
346 status and body condition (Kadri, Mitchell, Metcalfe, Huntingford, & Thorpe, 1996). In

347 addition, high retention of dietary lipid, elevated fat deposition, increased CF and rapid
348 growth are observed during the autumn period (Alne et al., 2011; Dessen et al., 2017; Kadri et
349 al., 1996; Mørkøre & Rørvik, 2001). Hence, the influence of natural seasonal cues might be
350 the main reason for the observed differences in growth between the present study and the one
351 of Johansen et al. (2003).

352

353 In October, two months after the start of the main-dietary phase, muscle fat and CF were
354 restored in both the LF and 0.5LF group compared to the HF group. This observation shows
355 that Atlantic salmon is able to rapidly replenish lipid stores and body condition during the
356 autumn following a feeding period of a low-fat diet or restricted ration of this diet. In contrast,
357 the visceral fat content among the groups maintained about the same pattern throughout the
358 study. The level or severity of restricting lipid deposition during pre-dietary phase was highly
359 negatively related with the magnitude of the subsequent growth response from August to
360 October. This was particularly linked to the relative muscle fat content at termination of the
361 pre-dietary phase prior to autumn. The degree of CG response seem also related to the level of
362 deviance in body condition, length and mass in the restricted or deprived fish groups
363 compared to their non-treated counter-specifics (Alvarez & Nicieza, 2005; Johansen et al.,
364 2001; Johnsson & Bohlin, 2005; Johnsson & Bohlin, 2006). Although the deviance in mass
365 and length may have contributed to the growth response in the present study, the small
366 difference between the LF and HF groups in August and the strong correlation between
367 muscle fat and growth, indicate that fat/energy status seem to be the most important growth
368 regulator during August and September. The increased growth and rapid replenishment of
369 lipid stores suggest a robust mechanism for the regulation of body fat, and are in line with the
370 observation of Silverstein, Shearer, Dickhoff & Plisetskaya (1999).

371

372 Several studies have indicated that animals displaying CG prioritise the restoration of body
373 condition and fat stores before more resources are allocated to support structural and skeletal
374 growth (Broekhuizen, Gurney, Jones, & Bryant, 1994; Johnsson & Bohlin, 2006). In part, the
375 results of the present study support these observations, as both the relative muscle fat content
376 and CF were quickly restored in the 0.5LF group, but not that quickly restored for BW and
377 length. Some studies have also suggested that structural restoration can be delayed due to the
378 effects of food deprivation or restriction on the endocrine system, involved in the regulation
379 of growth (Björnsson, 1997; Johnsson, Jönsson, & Björnsson, 1996). There is evidence that
380 skeletal and muscle growth are independent processes and that the relationship between
381 length and weight is approximately cubic (Einen, Waagan, & Thomassen, 1998; Jobling,
382 2002; Mørkøre & Rørvik, 2001). Thus, changes in weight are relatively greater than in length,
383 and the rapid increase in BW and fat content observed among the 0.5LF group in the autumn,
384 may be a factor explaining why length are restored later than body shape and fat content.

385

386 The stabilisation of the muscle fat in October coincides with the study of Mørkøre & Rørvik
387 (2001). This may suggest that the capacity of muscle fat deposition has reached an upper limit
388 at this time point. There is documentation that CG responses will cease as lipid stores and
389 body condition are restored to similar levels as the non-affected conspecifics (Johansen *et al.*,
390 2001; Ali *et al.*, 2003; Alvarez & Nicieza, 2005; Johnsson & Bohlin, 2005). In the present
391 study, the LF and 0.5LF groups continued to grow faster than the HF group both during the
392 periods October to December and December to March. The improved growth of the LF
393 groups from December to March was evident although the relative muscle fat content, CF and
394 BW were restored prior to this period and not significantly different from the HF group.
395 Hence, the observed growth response in this period is not directly related to restoration of fat
396 or BW. The sexual maturation process in Atlantic salmon requires, in addition to photoperiod,

397 sufficient fat and energy reserves (Kadri et al., 1996; Rowe & Thorpe, 1990; Taranger et al.,
398 2010). The production of gonads are energetically expensive and acquire high-energy
399 investment (Fleming, 1996; Jonsson, Jonsson, & Hansen, 1997). Appropriate and available
400 energy and fat reserves during the spring period seem to be a major factor controlling
401 initiation and proceeding of the maturation process (Thorpe, 1994; Thorpe, Mangel, Metcalfe,
402 & Huntingford, 1998; Wright, 2007). Too low energy and fat levels may arrest the maturation
403 process and postpone reproduction (Duston & Saunders, 1999; Rowe & Thorpe, 1990; Rowe,
404 Thorpe, & Shanks, 1991; Thorpe, 1994; Thorpe, Talbot, Miles, & Keay, 1990). Hence, well
405 growing salmon with a high and stable fat content are more likely to adopt the development
406 pathway of becoming sexual mature (Thorpe, 1994). Following this line of arguments, the
407 stronger growth response observed in both LF groups compared to the HF group prior to the
408 spring period in the present study, may have been triggered by the salmon reproductive life
409 strategy. However, to verify this, the groups of salmon must be studied for a longer period
410 (during late spring, summer and autumn) and measurements of relevant plasma hormones,
411 gonad-somatic index and gene expression of e.g. myosin should be conducted. Unfortunately,
412 this was not possible in the present study. Anyhow, observation of a long-term improved
413 growth response is important for a further development of a dynamic seasonal feeding
414 concept in salmon farming. Not only for traditional sea cage farming, but also in closed and
415 semi-closed production units where photoperiod may be manipulated. Taken into
416 consideration that the initial BW of the 0.5LF group was 738 g. less than the HF group, a
417 relative increase in weight gain of 950 g. more than the HF group is impressive.

418

419 When feed availability is restricted, competition for the feed often increase and dominant
420 individuals may try to monopolize the feeding area to obtain larger amounts of feed that is
421 supplied (Maclean & Metcalfe, 2001; Ryer & Olla, 1996). High competition for feed may

422 therefore lead to increased disparities in feed intake and growth that consequently will give
423 higher variation in BW. To minimize such effects, the 0.5LF group was administrated all
424 daily feed in only one ration during the pre-dietary phase. The high dispersion in BW and CF
425 among the HF group at termination of the main-dietary phase indicates that the 0.5LF and LF
426 group had an increase in weight and CF that was more homogeneous than the HF group. This
427 was probably due to the increased growth of LF groups in latter stages of the trial. The
428 possibility that fish among the LF groups displayed aggressive behaviour and tried to
429 monopolize food in this period seem unlikely due to three main factors: i) the HF group
430 showed a normal and satisfying growth with mean TGC of 3.2, ii) feed was administered in
431 excess during the main-dietary phase to ensure *ad libitum* feeding and iii) no or little fin
432 damage were observed at termination.

433

434 In summary, salmon with low body fat levels (LF groups) prior to declining day lengths in the
435 autumn displayed significantly higher growth rate and weight gain compared to the control
436 fish (HF group). The initial differences in muscle fat and CF were restored after only two
437 months, displaying rapid replenishment of lipid stores and body condition. Differences in
438 body fat content prior to autumn had significant effect on growth throughout the whole seven-
439 month main-dietary phase, even after similar body fat stores among the groups were obtained,
440 indicating a more complex explanation than just a lipostatic regulation mechanism. The LF
441 and 0.5LF fed fish obtained a significantly lower variation in BW and CF than the HF fed fish
442 at trial termination. This increased uniformity of BW and CF may reduce the amount of
443 manual gutting and filleting of large and small individuals, which optimizes the efficiency of
444 automatic gutting and filleting of salmon at the time of slaughter.

445

446 **ACKNOWLEDGMENTS:**

447

448 The authors appreciate the excellent technical assistance and rearing of fish that was provided
449 by the staff from the former Nofima research station at Averøy (now Marine Harvest research
450 station), with special thanks to Sissel Nergaard. This work was supported by a project grant
451 from the Norwegian Seafood Research Fund (FHF, grant number 900653). The authors
452 declare that there are no potential sources of conflict of interest with this work.

453

454 **REFERENCES:**

455

456 Aksnes, A. (1995). Growth, feed efficiency and slaughter quality of salmon, *Salmo salar* L.,
457 given feeds with different ratios of carbohydrate and protein. *Aquaculture Nutrition*, **1**,
458 241–248. <https://doi.org/10.1111/j.1365-2095.1995.tb00050.x>

459 Ali, M., Nicieza, A., & Wootton, R. J. (2003). Compensatory growth in fishes: A response to
460 growth depression. *Fish and Fisheries*, **4**, 147–190. [https://doi.org/10.1046/j.1467-](https://doi.org/10.1046/j.1467-2979.2003.00120.x)
461 [2979.2003.00120.x](https://doi.org/10.1046/j.1467-2979.2003.00120.x)

462 Alne, H., Oehme, M., Thomassen, M., Terjesen, B., & Rørvik, K. A. (2011). Reduced growth,
463 condition factor and body energy levels in Atlantic salmon *Salmo salar* L. during their
464 first spring in the sea. *Aquaculture Research*, **42**, 248–259.

465 <https://doi.org/10.1111/j.1365-2109.2010.02618.x>

466 Alvarez, D., & Nicieza, A. G. (2005). Compensatory response “defends” energy levels but not
467 growth trajectories in brown trout, *Salmo trutta* L. *Proceedings of the Royal Society B:*
468 *Biological Sciences*, **272**, 601–7. <https://doi.org/10.1098/rspb.2004.2991>

469 Aursand, M., Bleivik, B., Rainuzzo, J. R., Leif, J., & Mohr, V. (1994). Lipid distribution and
470 composition of commercially farmed Atlantic salmon (*salmo salar*). *Journal of the*
471 *Science of Food and Agriculture*, **64**, 239–248. <https://doi.org/10.1002/jsfa.2740640214>

- 472 Austreng, E., Storebakken, T., & Åsgård, T. (1987). Growth rate estimates for cultured
473 Atlantic salmon and rainbow trout. *Aquaculture*, **60**, 157–160.
474 [https://doi.org/10.1016/0044-8486\(87\)90307-3](https://doi.org/10.1016/0044-8486(87)90307-3)
- 475 Bell, J. G., McEvoy, J., Tocher, D. R., McGhee, F., Campbell, P. J., & Sargent, J. R. (2001).
476 Replacement of fish oil with rapeseed oil in diets of Atlantic salmon (*Salmo salar*)
477 affects tissue lipid compositions and hepatocyte fatty acid metabolism. *The Journal of*
478 *Nutrition*, **131**, 1535–43. Retrieved from
479 <http://www.ncbi.nlm.nih.gov/pubmed/11340112>
- 480 Björnsson, B. T. (1997). The biology of salmon growth hormone: from daylight to
481 dominance. *Fish Physiology and Biochemistry*, **17**, 9–24.
482 <https://doi.org/10.1023/A:1007712413908>
- 483 Broekhuizen, N., Gurney, W. S. C., Jones, A., & Bryant, A.D. (1994). Modeling
484 Compensatory Growth. *Functional Ecology*, **8**, 770–782. Retrieved from
485 [isi:A1994PW40300013](http://www.isis.org.uk/isi:A1994PW40300013)
- 486 Bull, C. D., & Metcalfe, N. B. (1997). Regulation of hyperphagia in response to varying
487 energy deficits in overwintering juvenile Atlantic salmon. *Journal of Fish Biology*, **50**,
488 498–510. <https://doi.org/10.1111/j.1095-8649.1997.tb01945.x>
- 489 Dessen, J. E., Weihe, R., Hatlen, B., Thomassen, M. S., & Rørvik, K. A. (2017). Different
490 growth performance, lipid deposition, and nutrient utilization in in-season (S1) Atlantic
491 salmon post-smolt fed isoenergetic diets differing in protein-to-lipid ratio. *Aquaculture*,
492 **473**, 345–354. <https://doi.org/10.1016/j.aquaculture.2017.02.006>
- 493 Duston, J., & Saunders, R. L. (1999). Effect of winter food deprivation on growth and sexual
494 maturity of Atlantic salmon (*Salmo salar*) in sea water. *Canadian Journal of Fisheries*
495 *and Aquatic Sciences*, **56**, 201–207. <https://doi.org/10.1139/f98-165>
- 496 Einen, O., Mørkøre, T., Røra, A. M. B., & Thomassen, M. S. (1999). Feed ration prior to

- 497 slaughter - a potential tool for managing product quality of Atlantic salmon (*Salmo*
498 *salar*). *Aquaculture*, **178**(1–2), 149–169. [https://doi.org/10.1016/S0044-8486\(99\)00126-](https://doi.org/10.1016/S0044-8486(99)00126-)
499 X
- 500 Einen, O., Waagan, B., & Thomassen, M. S. (1998). Starvation prior to slaughter in Atlantic
501 salmon (*Salmo salar*): I. Effects on weight loss, body shape, slaughter- and fillet-yield,
502 proximate and fatty acid composition. *Aquaculture*, **166**, 85–104.
503 [https://doi.org/10.1016/S0044-8486\(98\)00279-8](https://doi.org/10.1016/S0044-8486(98)00279-8)
- 504 Fleming, I. a. (1996). Reproductive strategies of Atlantic salmon: ecology and evolution.
505 *Fisheries, Reviews in Fish Biology and Fisheries*, **6**, 349–416.
506 <https://doi.org/10.1007/BF00164323>
- 507 Folch, J., Lees, M., & Stanley, G. H. S. (1957). A simple method for the isolation and
508 purification of total lipides from animal tissues. *Journal of Biological Chemistry*, **226**,
509 497–509. <https://doi.org/10.1007/s10858-011-9570-9>
- 510 Folkestad, A., Wold, J. P., Rørvik, K. A., Tschudi, J., Haugholt, K. H., Kolstad, K., &
511 Mørkøre, T. (2008). Rapid and non-invasive measurements of fat and pigment
512 concentrations in live and slaughtered Atlantic salmon (*Salmo salar* L.). *Aquaculture*,
513 **280**, 129–135. <https://doi.org/10.1016/j.aquaculture.2008.04.037>
- 514 Foss, A., & Imsland, A. K. (2002). Compensatory growth in the spotted wolffish *Anarhichas*
515 *minor* (Olafsen) after a period of limited oxygen supply. *Aquaculture Research*, **33**,
516 1097–1101. <https://doi.org/10.1046/j.1365-2109.2002.00768.x>
- 517 Helland, S. J., Grisdale-Helland, B., & Nerland, S. (1996). A simple method for the
518 measurement of daily feed intake of groups of fish in tanks. *Aquaculture*, **139**, 157–163.
519 [https://doi.org/10.1016/0044-8486\(95\)01145-5](https://doi.org/10.1016/0044-8486(95)01145-5)
- 520 Hemre, G. I., & Sandnes, K. (1999). Effect of dietary lipid level on muscle composition in
521 Atlantic salmon *Salmo salar*. *Aquaculture Nutrition*, **5**, 9–16.

- 522 <https://doi.org/10.1046/j.1365-2095.1999.00081.x>
- 523 Iwama, G. K., & Tautz, A. F. (1981). A Simple Growth Model for Salmonids in Hatcheries.
524 *Canadian Journal of Fisheries and Aquatic Sciences*, **38**, 649–656.
525 <https://doi.org/10.1139/f81-087>
- 526 Jobling, M. (2002). Environmental factors and rates of development and growth. In *Handbook*
527 *of fish biology and fisheries. Volume 1: Fish biology* (pp. 97–122).
528 <https://doi.org/10.1002/9780470693803>
- 529 Jobling, M., & Johansen, S. J. S. (1999). The lipostat, hyperphagia and catch-up growth.
530 *Aquaculture Research*, **30**, 473–478. <https://doi.org/10.1046/j.1365-2109.1999.00358.x>
- 531 Jobling, M., & Miglav, I. (1993). The size of lipid depots - a factor contributing to the
532 control of food intake in Arctic charr, *Salvelinus alpinus*? *Journal of Fish Biology*, **43**,
533 487–489.
- 534 Johansen, S. J. S., Ekli, M., & Jobling, M. (2002). Is there lipostatic regulation of feed intake
535 in Atlantic salmon *Salmo salar* L.? *Aquaculture Research*, **33**, 515–524.
536 <https://doi.org/10.1046/j.1365-2109.2002.00736.x>
- 537 Johansen, S. J. S., Ekli, M., Stangnes, B., & Jobling, M. (2001). Weight gain and lipid
538 deposition in Atlantic salmon, *Salmo salar*, during compensatory growth: Evidence for
539 lipostatic regulation? *Aquaculture Research*, **32**, 963–974.
540 <https://doi.org/10.1046/j.1365-2109.2001.00632.x>
- 541 Johansen, S. J. S., Sveier, H., & Jobling, M. (2003). Lipostatic regulation of feed intake in
542 Atlantic salmon *Salmo salar* L. defending adiposity at the expense of growth?
543 *Aquaculture Research*, **34**, 317–331. <https://doi.org/10.1046/j.1365-2109.2003.00821.x>
- 544 Johnsson, J. I., & Bohlin, T. (2005). Compensatory growth for free? A field experiment on
545 brown trout, *Salmo trutta*. *Oikos*, **111**, 31–38. [https://doi.org/10.1111/j.0030-](https://doi.org/10.1111/j.0030-1299.2005.13972.x)
546 [1299.2005.13972.x](https://doi.org/10.1111/j.0030-1299.2005.13972.x)

- 547 Johnsson, J. I., & Bohlin, T. (2006). The cost of catching up: increased winter mortality
548 following structural growth compensation in the wild. *Proceedings. Biological Sciences /*
549 *The Royal Society*, **273**, 1281–6. <https://doi.org/10.1098/rspb.2005.3437>
- 550 Johnsson, J. I., Jönsson, E., & Björnsson, B. T. (1996). Dominance, nutritional state, and
551 growth hormone levels in rainbow trout (*Oncorhynchus mykiss*). *Hormones and*
552 *Behavior*, **30**, 13–21. <https://doi.org/10.1006/hbeh.1996.0003>
- 553 Jonsson, N., Jonsson, B., & Hansen, L. P. (1997). Changes in proximate composition and
554 estimates of energetic costs during upstream migration and spawning in Atlantic salmon
555 *Salmo salar*. *Journal of Animal Ecology*, **66**, 425–436. <https://doi.org/10.2307/5987>
- 556 Kadri, S., Mitchell, D. F., Metcalfe, N. B., Huntingford, F. A., & Thorpe, J. E. (1996).
557 Differential patterns of feeding and resource accumulation in maturing and immature
558 Atlantic salmon, *Salmo salar*. *Aquaculture*, **142**, 245–257. [https://doi.org/10.1016/0044-](https://doi.org/10.1016/0044-8486(96)01258-6)
559 [8486\(96\)01258-6](https://doi.org/10.1016/0044-8486(96)01258-6)
- 560 Keesey, R. E., & Corbett, S. W. (1984). Metabolic defense of the body weight set-point.
561 *Research Publications - Association for Research in Nervous and Mental Disease*, **62**,
562 87–96. Retrieved from <http://europepmc.org/abstract/MED/6695117>
- 563 Kennedy, G. C. (1953). The Role of Depot Fat in the Hypothalamic Control of Food Intake in
564 the Rat. *Proceedings of the Royal Society B: Biological Sciences*, **140**, 578–592.
565 <https://doi.org/10.1098/rspb.1953.0009>
- 566 Maclean, A., & Metcalfe, N. B. (2001). Social status, access to food, and compensatory
567 growth in juvenile Atlantic salmon. *Journal of Fish Biology*, **58**, 1331–1346.
568 <https://doi.org/10.1111/j.1095-8649.2001.tb02290.x>
- 569 Mason, M. E., & Waller, G. R. (1964). Dimethoxypropane Induced Transesterification of Fats
570 and Oils in Preparation of Methyl Esters for Gas Chromatographic Analysis. *Analytical*
571 *Chemistry*, **36**, 583–586. <https://doi.org/10.1021/ac60209a008>

- 572 McLoughlin, M. F., Nelson, R. N., McCormick, J. I., Rowley, H. M., & Bryson, D. B. (2002).
573 Clinical and histopathological features of naturally occurring pancreas disease in farmed
574 Atlantic salmon, *Salmo salar* L. *Journal of Fish Diseases*, **25**, 33–43.
575 <https://doi.org/10.1046/j.1365-2761.2002.00334.x>
- 576 Metcalfe, N. B., & Monaghan, P. (2001). Compensation for a bad start: Grow now, pay later?
577 *Trends in Ecology and Evolution*, **16**, 254–260. [https://doi.org/10.1016/S0169-](https://doi.org/10.1016/S0169-5347(01)02124-3)
578 [5347\(01\)02124-3](https://doi.org/10.1016/S0169-5347(01)02124-3)
- 579 Metcalfe, N. B., & Thorpe, J. E. (1992). Anorexia and defended energy levels in over-
580 wintering juvenile salmon. *Journal of Animal Ecology*, **61**, 175–181.
581 <https://doi.org/10.2307/5520>
- 582 Mørkøre, T., & Rørvik, K. -A. (2001). Seasonal variations in growth, feed utilisation and
583 product quality of farmed Atlantic salmon (*Salmo salar*) transferred to seawater as 0 +
584 smolts or 1 + smolts. *Aquaculture*, **199**, 145–157. [https://doi.org/10.1016/S0044-](https://doi.org/10.1016/S0044-8486(01)00524-5)
585 [8486\(01\)00524-5](https://doi.org/10.1016/S0044-8486(01)00524-5)
- 586 Norwegian Standards Association, 1994. Norwegian Quality Cut [NQC] NS 9401 (Oslo).
- 587 Oehme, M., Grammes, F., Takle, H., Zambonino-Infante, J. L., Refstie, S., Thomassen, M. S.,
588 Rørvik, K.-A & Terjesen, B. F. (2010). Dietary supplementation of glutamate and
589 arginine to Atlantic salmon (*Salmo salar* L.) increases growth during the first autumn in
590 sea. *Aquaculture*, **310**, 156–163. <https://doi.org/10.1016/j.aquaculture.2010.09.043>
- 591 Rowe, D. K., & Thorpe, J. E. (1990). Suppression of maturation in male Atlantic salmon
592 (*Salmo salar* L.) parr by reduction in feeding and growth during spring months.
593 *Aquaculture*, **86**, 291–313. [https://doi.org/10.1016/0044-8486\(90\)90121-3](https://doi.org/10.1016/0044-8486(90)90121-3)
- 594 Rowe, D. K., Thorpe, J. E., & Shanks, A. M. (1991). Role of Fat Stores in the Maturation of
595 Male Atlantic Salmon (*Salmo salar*) Parr. *Canadian Journal of Fisheries and Aquatic*
596 *Sciences*, **48**, 405–413. <https://doi.org/10.1139/f91-052>

- 597 Ryer, C. H., & Olla, B. L. (1996). Growth depensation and aggression in laboratory reared
598 coho salmon: the effect of food distribution and ration size. *Journal of Fish Biology*, **48**,
599 686–694. <https://doi.org/10.1111/j.1095-8649.1996.tb01464.x>
- 600 Rørvik, K.-A., Ytrestøyl, T., Lundberg, E., Jakobsen, F. A., Jakobsen, A. A., & Bjerkeng, B.
601 (2010). How Apparent Digestibility of Carotenoids, Macronutrients, and Minerals are
602 Differently Affected by Ration Level in Atlantic Salmon, *Salmo Salar*. *Journal of*
603 *Applied Aquaculture*, **22**, 123–139. <https://doi.org/10.1080/10454431003736227>
- 604 Røsjø, C., Berg, T., Manum, K., Gjøen, T., Magnusson, S., & Thomassen, M. S. (1994).
605 Effects of temperature and dietary n-3 and n-6 fatty acids on endocytic processes in
606 isolated rainbow trout (*Oncorhynchus mykiss*, Walbaum) hepatocytes. *Fish Physiology*
607 *and Biochemistry*, **13**, 119–32. <https://doi.org/10.1007/BF00004337>
- 608 Silverstein, J. T., Shearer, K. D., Dickhoff, W. W., & Plisetskaya, E. M. (1999). Regulation of
609 nutrient intake and energy balance in salmon. *Aquaculture*, **177**, 161-169.
610 [https://doi.org/10.1016/S0044-8486\(99\)00076-9](https://doi.org/10.1016/S0044-8486(99)00076-9)
- 611 Taksdal, T., Olsen, A. B., Bjerkås, I., Hjortaas, M. J., Dannevig, B. H., Graham, D. A., &
612 McLoughlin, M. F. (2007). Pancreas disease in farmed Atlantic salmon, *Salmo salar* L.,
613 and rainbow trout, *Oncorhynchus mykiss* (Walbaum), in Norway. *Journal of Fish*
614 *Diseases*, **30**, 545–558. <https://doi.org/10.1111/j.1365-2761.2007.00845.x>
- 615 Taranger, G. L., Carrillo, M., Schulz, R. W., Fontaine, P., Zanuy, S., Felip, A., Weltzien, F.-
616 A., Dufour, S., Karlsen, Ø., Norberg, B., Andersson, E., & Hansen, T. (2010). Control of
617 puberty in farmed fish. *General and Comparative Endocrinology*, **165**, 483–515.
618 <https://doi.org/10.1016/j.ygcen.2009.05.004>
- 619 Thomassen, M. S., & Røsjø, C. (1989). Different fats in feed for salmon: Influence on sensory
620 parameters, growth rate and fatty acids in muscle and heart. *Aquaculture*, **79**, 129–135.
621 [https://doi.org/10.1016/0044-8486\(89\)90453-5](https://doi.org/10.1016/0044-8486(89)90453-5)

- 622 Thorarensen, H., & Farrell, A. P. (2011). The biological requirements for post-smolt Atlantic
623 salmon in closed-containment systems. *Aquaculture*, **312**, 1-14.
624 <https://doi.org/10.1016/j.aquaculture.2010.11.043>
- 625 Thorpe, J. E. (1994). Reproductive strategies in Atlantic salmon, *Salmo salar* L. *Aquaculture*
626 *Research*, **25**, 77–87. <https://doi.org/10.1111/j.1365-2109.1994.tb00668.x>
- 627 Thorpe, J. E., Mangel, M., Metcalfe, N. B., & Huntingford, F. A. (1998). Modelling the
628 proximate basis of salmonid life-history variation, with application to Atlantic salmon,
629 *Salmo salar* L. *Evolutionary Ecology*, **12**, 581–599.
630 <https://doi.org/10.1023/A:1022351814644>
- 631 Thorpe, J. E., Talbot, C., Miles, M. S., & Keay, D. S. (1990). Control of maturation in
632 cultured Atlantic salmon, *Salmo salar*, in pumped seawater tanks, by restricting food
633 intake. *Aquaculture*, **86**, 315–326. [https://doi.org/10.1016/0044-8486\(90\)90122-4](https://doi.org/10.1016/0044-8486(90)90122-4)
- 634 Torrissen, O., Olsen, R. E., Toresen, R., Hemre, G. I., Tacon, A. G. J., Asche, F., Hardy, R.W.
635 & Lall, S. (2011). Atlantic Salmon (*Salmo salar*): The “Super-Chicken” of the Sea?
636 *Reviews in Fisheries Science*, **19**, 257–278.
637 <https://doi.org/10.1080/10641262.2011.597890>
- 638 Torstensen, B. E., Lie, Ø., & Frøyland, L. (2000). Lipid metabolism and tissue composition in
639 Atlantic salmon (*Salmo salar* L.)- Effects of capelin oil, palm oil, and oleic acid-enriched
640 sunflower oil as dietary lipid sources. *Lipids*, **35**, 653–664.
641 <https://doi.org/10.1007/s11745-000-0570-6>
- 642 Torstensen, B. E., Lie, & Hamre, K. (2001). A factorial experimental design for investigation
643 of effects of dietary lipid content and pro- and antioxidants on lipid composition in
644 Atlantic salmon (*Salmo salar*) tissues and lipoproteins. *Aquaculture Nutrition*, **7**, 265–
645 276. <https://doi.org/10.1046/j.1365-2095.2001.00184.x>
- 646 Usher, M. L., Talbot, C., & Eddy, F. B. (1991). Effects of transfer to seawater on growth and

647 feeding in Atlantic salmon smolts (*Salmo salar* L.). *Aquaculture*, **94**, 309–326.

648 [https://doi.org/10.1016/0044-8486\(91\)90176-8](https://doi.org/10.1016/0044-8486(91)90176-8)

649 Wright, P. J. (2007). Understanding the maturation process for field investigations of

650 fisheries-induced evolution. *Marine Ecology Progress Series*, **335**, 279-284.

651 <https://doi.org/10.3354/meps335279>

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667 **TABLES:**

668

669 **Table 1.** Biometrics and fat content of Atlantic salmon in August 2011 fed a diet high-fat diet
 670 (HF), low-fat high-protein diet (LF) or half ration of the low fat diet high-protein diet (0.5LF)
 671 from May until August 2011, referred to as pre-dietary feeding phase. Biometric parameters
 672 for all fish are presented as means \pm SD, whereas biometric parameters and fat content for
 673 sampled fish are presented as means \pm SEM together with indications of significant
 674 differences.

675

Dietary treatment	HF	LF	0.5LF
<i>Biometric parameters, all fish</i>			
Number of fish, n	584	584	602
Bodyweight, g	2651 \pm 335	2506 \pm 287	1865 \pm 253
Fork length, cm	59.1 \pm 2.3	59.1 \pm 2.1	55.8 \pm 2.3
CF	1.28 \pm 0.09	1.21 \pm 0.07	1.07 \pm 0.08
<i>Biometric parameters, sampled fish, n = 20</i>			
Bodyweight, g	2619 \pm 70 ^a	2515 \pm 63 ^a	1881 \pm 47 ^b
Fork length, cm	59.0 \pm 0.5 ^a	59.0 \pm 0.4 ^a	55.7 \pm 0.5 ^b
CF	1.22 \pm 0.02 ^a	1.18 \pm 0.02 ^a	1.03 \pm 0.01 ^b
VSI	11.3 \pm 0.4 ^a	9.6 \pm 0.2 ^b	8.5 \pm 0.1 ^c
<i>Fat content, sampled fish, n = 20</i>			
Muscle fat, %	16.4 \pm 0.3 ^a	13.1 \pm 0.2 ^b	11.3 \pm 0.3 ^c
Visceral fat[†], %	39.0	29.0	26.6

676 CF; condition factor, VSI; Visceral somatic index

677 [†]The analysis of visceral fat content was conducted on pooled samples in August 2011 (n=1)678 Values in the same row with different letters are significantly different ($P \leq 0.05$) determined by one-way

679 ANOVA followed by Duncan's multiple range test.

680

681

682

683

684 **Table 2.** Chemical compositions (wet weight, as is basis) and fatty acid composition (% of
685 total fatty acids) of the diets used in the main-dietary phase.

686

Diet code	7 mm pellet		9 mm pellet	
	MO	RO	MO	RO
<i>Chemical composition (wet weight, as is basis)</i>				
Dry matter, %	93.2	94.0	93.8	93.9
Crude protein (N x 6.25), %	41.2	41.7	34.4	34.6
Crude Lipid, %	31.2	31.4	37.0	35.7
Starch, %	6.2	6.1	6.7	6.8
Ash, %	4.8	4.8	5.1	5.1
NFE [†] , %	16.0	16.1	17.3	18.5
Crude protein/lipid ratio	1.32	1.33	0.93	0.97
<i>Calculated values[‡]</i>				
Gross energy, MJ kg ⁻¹	24.8	25.1	25.7	25.5
DP, g kg ⁻¹	354	359	296	298
DE, MJ kg ⁻¹	21.4	21.5	22.2	21.9
DP/DE ratio, g MJ kg ⁻¹	16.6	16.6	13.3	13.6
<i>Fatty acid composition (% of total fatty acids)</i>				
C 16:0	12.7	8.5	14.3	9.3
C 18:0	3.2	2.7	3.7	2.9
ΣSFA [§]	22.6	15.1	24.0	15.9
C 18:1 n-9	26.8	42.1	23.3	42.5
ΣMUFA [¶]	38.1	49.8	36.2	52.8
C 18:2 n-6	8.1	13.9	7.4	13.9
C 18:3 n-3	3.4	6.5	2.9	6.0
C 20:5 n-3	10.1	4.6	11.1	4.0
C 22:5 n-3	1.3	0.6	1.4	0.5
C 22:6 n-3	7.2	3.5	7.5	3.6
ΣPUFA [‡]	34.3	30.4	32.7	29.0
SUM EPA+DHA	17.4	8.1	18.6	7.5
n-6/n-3 ratio	0.4	0.9	0.4	1.0

687 MO; Marine oil profile, RO: Rapeseed oil profile, N; Nitrogen, NFE; Nitrogen-free extracts, DP; digestible
688 protein, DE; digestible energy, MJ; Mega joule, SFA; Saturated fatty acids, MUFA; monounsaturated fatty acids,
689 PUFA; polyunsaturated fatty acids.

690 [†]NFE was calculated as = 100 – (protein+lipids+ash+water)

691 [‡]Gross energy, DP and DE were estimated assuming 23.7, 39.5 and 17.2 MJ kg⁻¹ as the gross energy content of
692 protein, lipids and carbohydrates, respectively. The apparent digestibility coefficients (ADCs) for protein and
693 lipids used were 0.86 and 0.94, respectively (Einen & Roem 1997), whereas 0.50 was used for NFE (Arnesen &
694 Krogdahl 1993).

695 [§]SFA; C14:0, C15:0, C16:0, C18:0 and 22:0.

696 [¶]MUFA; C16:1n-9, C16:1n-7, C17:1n-7,C18:1n-7, C:18:1n-9, C20:1n-7, C20:1n-9,C20:1n-11, C22:1n-
697 9,C22:1n-11,C24:1n-9
698 [‡]PUFA; C16:2n-3, C16:3n-4, C18:2n-6,C18:3n-6, C18:3n-3, C18:4n-3, C20:4n-3, C20:2n-6, C20:3n-6, C20:4n-
699 6,C20:5n-3, C22:5-n-3, C22:6n-3.

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718 **FIGURE CAPTIONS:**

719

720 **Fig. 1** Schematic overview of the experimental design during the pre- and the main-dietary
 721 phase. The squares during the pre-dietary phase represent net-pens fed different diets; HF =
 722 high fat diet (black filled square), LF = low fat diet (grey filled square), 0.5LF = half ration of
 723 the low-fat diet (white filled square). The large squares in the main-dietary phase represents
 724 the net-pens and the squares within the net-pens are the pre-dietary groups.

725

726 **Fig. 2** Ambient daily sea temperature ($^{\circ}\text{C}$, y-axis) and hours of daylight (hours, z-axis) during
 727 the pre-dietary phase (May to August 2011) and the main-dietary phase (August 2011 to
 728 March 2012). The length of the different periods are indicate by the different grey colours
 729 (light grey = pre-dietary phase, dark grey = main-dietary phase).

730

731 **Fig. 3** Regression line between thermal growth coefficients (TGC) and the increase in muscle
 732 fat (%) from August to October in Atlantic salmon fed three different pre-dietary treatment
 733 from May to August 2011; high fat diet (black filled squares) = HF, low fat diet (grey filled
 734 triangles) = LF, half ration of the low-fat diet (white filled circles) = 0.5LF. Each point
 735 represents average per fish group/experimental unit ($n = 12$).

736

737 **Fig. 4** Muscle fat content (**A1**) and body weight (**B1**) development of Atlantic salmon fed
 738 three different pre-dietary treatment from May to August 2011. Values are means \pm SEM, $n =$
 739 4 ($n = 1$ at termination of the pre-dietary phase). Values not sharing common superscript
 740 letters within each sampling period are significantly different ($P \leq 0.05$). **A2** and **B2**, present
 741 the final muscle fat and BW of the groups, respectively. The values 11.3%, 13.2% and 16.4%

742 represent the obtained fat content at the beginning of the main-dietary phase (August 2011)
743 for the 0.5LF, LF and HF group, respectively. ns; not significant, *; trend ($P < 0.1$).

744

745

746 **Fig. 5** Thermal growth coefficients (TGC) (**A**) and weight gain (kg) (**B**) of Atlantic salmon
747 fed three different pre-dietary treatment from May to August 2011. Values are means \pm SEM,
748 $n = 4$. Values not sharing common superscript letters within each sampling period are
749 significantly different ($P \leq 0.05$). The values 11.3%, 13.2% and 16.4% represent the obtained
750 fat content at the beginning of the main-dietary phase (August 2011) for the 0.5LF, LF and
751 HF group, respectively.

752

753 **Fig. 6** Condition factor (CF) (**A1**) and fork length (cm) (**B1**) development of Atlantic salmon.
754 fed three different pre-dietary treatment from May to August 2011. Values are means \pm SEM,
755 $n = 4$ ($n = 1$ at termination of the pre-dietary phase). Values not sharing common superscript
756 letters within each sampling period are significantly different ($P \leq 0.05$). **A2** and **B2**, present
757 the final CF and fork length of the groups, respectively. The values 11.3%, 13.2% and 16.4%
758 represent the obtained fat content at the beginning of the main-dietary phase (August 2011)
759 for the 0.5LF, LF and HF group, respectively.

760

761 **Fig. 7** Visceral fat development of Atlantic salmon fed three different pre-dietary treatment
762 from May to August 2011. Values are means \pm SEM, $n = 4$ ($n = 1$ at termination of the pre-
763 dietary phase). Values not sharing common superscript letters within each sampling period are
764 significantly different ($P \leq 0.05$). The values 11.3%, 13.2% and 16.4% represent the obtained
765 fat content at the beginning of the main-dietary phase (August 2011) for the 0.5LF, LF and
766 HF group, respectively.

767

768 **Fig. 8** Variation in body weight (gram) (**A**) and condition factor (CF) (**B**) assessed using
769 coefficient of variation (CV; $\text{mean} \times \text{SD}^{-1}$) among Atlantic salmon fed three different pre-
770 dietary treatment from May to August 2011. Values are means \pm SEM, $n = 4$ ($n = 1$ at
771 termination of the pre-dietary phase). Values not sharing common superscript letters within
772 each sampling period are significantly different ($P \leq 0.05$). ns; not significant

773

774

775

776

777

778

779

780

781

782


783

784

785

786

787 FIGURE 1:



788

789

790

791

792


793

794

795

796

797 FIGURE 2:



798

799

800

801

802

803


804

805

806

807

808 FIGURE 3:



809

810

811

812

813

814

815


816

817

818

819 FIGURE 4:

820



821

822

823


824

825

826

827

828 FIGURE 5:



829

830

831


832

833

834

835 FIGURE 6:

836



837

838

839


840

841

842

843

844 FIGURE 7:



845

846

847

848

849

850

851


852

853

854

855 FIGURE 8:

856



857