

# Neurosteroid 3 $\alpha$ -Androstenediol Efficiently Counteracts Paclitaxel-Induced Peripheral Neuropathy and Painful Symptoms

Laurence Meyer<sup>✉</sup>, Christine Patte-Mensah<sup>✉</sup>, Omar Taleb, Ayikoe Guy Mensah-Nyagan<sup>\*</sup>

Biopathologie de la Myéline, Neuroprotection et Stratégies Thérapeutiques, INSERM U1119, Fédération de Médecine Translationnelle de Strasbourg (FMTS), Université de Strasbourg, Faculté de Médecine, Strasbourg, France

## Abstract

Painful peripheral neuropathy belongs to major side-effects limiting cancer chemotherapy. Paclitaxel, widely used to treat several cancers, induces neurological symptoms including burning pain, allodynia, hyperalgesia and numbness. Therefore, identification of drugs that may effectively counteract paclitaxel-induced neuropathic symptoms is crucial. Here, we combined histopathological, neurochemical, behavioral and electrophysiological methods to investigate the natural neurosteroid 3 $\alpha$ -androstenediol (3 $\alpha$ -DIOL) ability to counteract paclitaxel-evoked peripheral nerve tissue damages and neurological symptoms. Prophylactic or corrective 3 $\alpha$ -DIOL treatment (4 mg/kg/2days) prevented or suppressed PAC-evoked heat-thermal hyperalgesia, cold-allodynia and mechanical allodynia/hyperalgesia, by reversing to normal, decreased thermal and mechanical pain thresholds of PAC-treated rats. Electrophysiological studies demonstrated that 3 $\alpha$ -DIOL restored control values of nerve conduction velocity and action potential peak amplitude significantly altered by PAC-treatment. 3 $\alpha$ -DIOL also repaired PAC-induced nerve damages by restoring normal neurofilament-200 level in peripheral axons and control amount of 2',3'-cyclic-nucleotide-3'-phosphodiesterase in myelin sheaths. Decreased density of intraepidermal nerve fibers evoked by PAC-therapy was also counteracted by 3 $\alpha$ -DIOL treatment. More importantly, 3 $\alpha$ -DIOL beneficial effects were not sedation-dependent but resulted from its neuroprotective ability, nerve tissue repairing capacity and long-term analgesic action. Altogether, our results showing that 3 $\alpha$ -DIOL efficiently counteracted PAC-evoked painful symptoms, also offer interesting possibilities to develop neurosteroid-based strategies against chemotherapy-induced peripheral neuropathy. This article shows that the prophylactic or corrective treatment with 3 $\alpha$ -androstenediol prevents or suppresses PAC-evoked painful symptoms and peripheral nerve dysfunctions in rats. The data suggest that 3 $\alpha$ -androstenediol-based therapy may constitute an efficient strategy to explore in humans for the eradication of chemotherapy-induced peripheral neuropathy.

**Citation:** Meyer L, Patte-Mensah C, Taleb O, Mensah-Nyagan AG (2013) Neurosteroid 3 $\alpha$ -Androstenediol Efficiently Counteracts Paclitaxel-Induced Peripheral Neuropathy and Painful Symptoms. PLoS ONE 8(11): e80915. doi:10.1371/journal.pone.0080915

**Editor:** Mark L. Bacceti, University of Cincinnati, United States of America

**Received:** July 24, 2013; **Accepted:** October 15, 2013; **Published:** November 15, 2013

**Copyright:** © 2013 Meyer et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by grants from Association Ti'toine (Normandie, France, <http://www.titoin-normandie.org/>) and Université de Strasbourg (France). LM was a postdoctoral fellow supported by Association Ti'toine (Normandie, France). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

\* E-mail: [gmensah@unistra.fr](mailto:gmensah@unistra.fr)

✉ These authors contributed equally to this work.

## Introduction

Chemotherapy with antineoplastic drugs remains so far the widely used and unavoidable treatment of several cancers. However, the success of many cancer treatments is hampered by antineoplastic-induced painful peripheral neuropathy and neuropathological symptoms that constitute a major health concern. Neuropathic pain, histopathological, neurochemical and functional alterations of peripheral nerves evoked by anticancer drugs represent a leading cause of discontinuation of successful treatments [1,2]. These neuropathological

symptoms restrict therapeutic doses of antineoplastics to suboptimal concentrations unable to kill tumor cells and significantly affect the quality of life of millions of patients [3-5]. Painful neuropathological symptoms are generated by various anticancer drugs including paclitaxel (also called taxol®) that is widely used to treat breast cancer and several other malignancies such as ovarian, lung, head and bladder cancers and AIDS-related Kaposi's sarcoma [6-11]. Paclitaxel is known to induce sensory peripheral neuropathy with burning pain symptoms, allodynia, hyperalgesia tingling and numbness [1,3,12-18]. Several attempts have been made to treat or

prevent antineoplastic-induced painful neuropathy with various neuroprotective agents but the results are contradictory and most of the compounds used were ineffective or induced themselves adverse effects such as nausea, reflex dysfunctions, treatment-emergent nervousness, insomnia, tremor, anorexia or stomach burning [19-23]. Therefore, it is crucial to identify novel drugs that may efficiently prevent or suppress antineoplastic-induced painful and neuropathological symptoms without evoking themselves undesirable action or severe side-effects. In a recent study, we observed that progesterone-derived 3 $\alpha$ -5 $\alpha$ -reduced metabolites exerted potent neuroprotective effects against oxaliplatin- or vincristine-induced peripheral neuropathy in rats, suggesting that 3 $\alpha$ -5 $\alpha$ -neurosteroid-based strategies may offer serious opportunities to develop effective therapies against anticancer drug-evoked painful neuropathic symptoms [24,25]. Among the family of 3 $\alpha$ -5 $\alpha$ -neurosteroids are naturally-produced molecules such as allopregnanolone and 3 $\alpha$ -androstanediol (3 $\alpha$ -DIOL) which activate allosterically the GABA<sub>A</sub> receptor and potentiate the central inhibition [26-34]. While allopregnanolone has extensively been investigated in several experimental models and in humans as a neuroprotective, neurogenic, anaesthetic, anxiolytic or analgesic compound, the neurosteroid 3 $\alpha$ -DIOL has received little attention [26,35-42]. The present study was therefore designed to investigate whether 3 $\alpha$ -DIOL may also be a neuroprotective neurosteroid offering effective possibilities for the treatment of chemotherapy-induced painful neuropathy and neuropathological symptoms. Thus, we assessed the potential of 3 $\alpha$ -DIOL to prevent or suppress paclitaxel-evoked neurochemical and functional abnormalities in rat peripheral nerves. In particular, we used a multidisciplinary approach to determine the ability of 3 $\alpha$ -DIOL to prevent or counteract several neuropathological parameters evoked by paclitaxel therapy such as behavioral painful symptoms (mechanical and thermal allodynia or hyperalgesia), electrophysiological deficiency (decreased nerve conduction velocity) and histopathological alterations including the repression of neurofilament 200 kDa (NF200) in peripheral nerve axons, the decrease of intraepidermal nerve fiber (IENF) density in intraplantar skin and the reduction of peripheral expression of myelin enzyme 2',3'-cyclic-nucleotide-3'-phosphodiesterase (CNPase) that is pivotal for axonal survival [43,44].

## Methods

### Animals

Adult male Sprague-Dawley rats weighing 250-300 g were used. The experiments were performed with male animals in order to avoid eventual fluctuations of results that may happen in females through the different phases of the oestral cycle [45-47]. The animals were obtained from a commercial source (Janvier, Le Genest St Isle, France) and housed under standard laboratory conditions in a 14/10 h light/dark cycle with food and water *ad libitum*. Animals were allowed a 1 week acclimatization period before being used in experiments.

### Ethic statement

Animal care and manipulations were performed according to The European Community Council Directives (86/609/EC) and under the supervision of authorized investigators. All experiments were performed minimizing the number of animals used and their suffering in accordance with the Alsace Department of Veterinary Public Health Guide for the Care and Use of Laboratory Animals. The ethics committee of the Alsace Department of Veterinary Public Health Guide for the Care and Use of Laboratory Animals specifically approved the protocol used in the present study (Agreement number 67-186). The experiments also followed the International Association for the Study of Pain ethical guidelines [48].

### Drugs and treatments

Paclitaxel (PAC; Sequoia Research Products Ltd, UK) was dissolved in 10% Cremophor EL<sup>®</sup> in saline (Sigma-Aldrich, St Louis, MO, USA), used as vehicle (VEH<sub>crem</sub>) and stored at 4°C. PAC was intraperitoneally (i.p.) injected every-2 days during 1 week, at a concentration of 1 mg/kg/injection depending on the daily body weight [16]. The chemotherapy was therefore performed at D1, D3, D5 and D7. Thus, the animals received a total of 4 mg/kg of PAC. Control rats were injected with the vehicle '10% Cremophor EL<sup>®</sup> in saline' solution or VEH<sub>crem</sub> (1 ml/kg) according to a similar schedule. Before the onset of PAC (versus vehicle) treatment, behavioral tests were performed in all rats in order to determine the pre-injection values of the mechanical (Day 0 or D0), cold (D0) and heat thermal (Day 1 or D1) nociceptive thresholds. After PAC or vehicle treatment onset, behavioral analyses were realized once a week to assess alternately the mechanical or thermal sensitivity.

3 $\alpha$ -androstanediol or 3 $\alpha$ -DIOL (Steraloids, Newport, RI, USA) was diluted in hydroxypropylcellulose or HPC 0.3% (Sigma-Aldrich, St Louis, MO) in water used as vehicle (VEH<sub>hpc</sub>). Consequently, it should be noted that 4 different groups of control animals were used for the whole study: (i) naive non-treated rats, (ii) (VEH<sub>crem</sub>)-treated rats, (iii) (VEH<sub>hpc</sub>)-treated rats and (iv) (VEH<sub>crem</sub> + VEH<sub>hpc</sub>)-treated rats. 3 $\alpha$ -DIOL was i.p. administered immediately after the behavioral test session. Two types of treatments were performed using 3 $\alpha$ -DIOL. Corrective 3 $\alpha$ -DIOL treatment, which aimed at suppressing painful neuropathic symptoms induced by the 1 week PAC treatment (administered at D1, D3, D5, and D7), consisted in starting 3 $\alpha$ -DIOL i.p. administration 8 days after the end of PAC treatment (D15). Therapy with 3 $\alpha$ -DIOL lasted 2 weeks (D15, D17, D19, D21, D23, D25 and D27). Prophylactic 3 $\alpha$ -DIOL treatment, which aimed at preventing the development of PAC-induced neuropathic pain, consisted in 3 $\alpha$ -DIOL administration before the onset and during the treatment with PAC. Thus, for the prophylactic strategy (3 $\alpha$ -DIOL administration before PAC injections), 3 $\alpha$ -DIOL was administered at D1, D3, D5 and D7 before PAC treatment started at D8. Then, PAC was injected at D8, D10, D12 and D14 while 3 $\alpha$ -DIOL treatment was continued at D9, D11, D13, D15, D17, D19, D21, D23 and D25. Behavioral experiments, electrophysiological studies, immunohistochemical investigations, microscopic and cellular analyses were performed to determine 3 $\alpha$ -DIOL effects on PAC-induced

painful symptoms as well as peripheral nerve alterations. Experimenters were completely blind to the experimental conditions of animals. Corrective and prophylactic effects of 3 $\alpha$ -DIOL against PAC-induced painful neuropathy were investigated along the treatment period but also one week after the end of 3 $\alpha$ -DIOL administration.

### Nociceptive behavioral tests

The *mechanical nociceptive sensitivity* threshold was evaluated in individual rats placed on Plexiglas® boxes (30 x 30 x 25 cm) upon an elevated metal grid allowing access to the plantar surface of the hind paws. The presence of mechanical allodynia and hyperalgesia were assessed using a series of calibrated von Frey hairs (1, 2, 4, 6, 8, 10, 15, 26g; Stoelting, Wood Dale, IL), which were applied to the plantar surface of the hind paw with increasing force until the individual filament used just started to bend. The filament was applied for a period of 1-2 s and the procedure was repeated five times at 4-5 s intervals. The threshold for paw withdrawal was calculated by taking the average of ten (five per paw) repeated stimuli (in g) which induced a reflex paw withdrawal. Only robust and immediate withdrawal responses followed by a licking of the paw were considered as positive. Naive untreated rats never withdraw from stimulations less than 6 g but respond 15-20% of the time for 15 g stimulus and 35-40% for 26 g stimulus. Observation of responses for stimulations < 6 g after drug administrations is indicative of mechanical allodynia. Increased level of responding for 15 and 26 g after treatment is indicative of mechano-hyperalgesia. Decreased level of responding for 26 g after treatment is indicative of antinociceptive effect.

*Heat thermal nociceptive sensitivity* was assessed by using a Plantar test apparatus (Ugo Basile, Comerio, Italy) which measures the paw withdrawal latency in response to radiant heat [49]. The rats were first allowed to habituate to the apparatus for 10 min before testing. Each rat was placed individually in clear Plexiglas® boxes (23 x 18 x 14 cm) positioned on a clear plastic surface. The heat source was then positioned under the plantar surface of the hind paw and activated with an infra-red light beam. The heat source is connected to a timer which automatically switched off the heat when the paw was withdrawn. A cut-off time of 20 s was used to prevent tissue damage in absence of response. Three latencies were obtained alternately from each hind paw 5 min apart. No consistent left and right differences were observed. The mean paw withdrawal latency (in seconds) of hind paws was determined from an average of six separate measures (three per paw) at a given time point. Testing boxes were thoroughly cleaned between each test session.

*Cold thermal nociceptive sensitivity* was assessed by using the acetone test. Rats were placed on Plexiglas® boxes (30 x 30 x 25 cm) upon an elevated metal grid allowing access to the plantar surface of the hind paws. Fifty microliters of acetone (Sigma-Aldrich, St Louis, MO) was sprayed onto the plantar skin of each hind paw 3 times with a Hamilton syringe. Rats were observed 20 s from the start of the acetone spray. The latency is measured for the withdrawal followed by a licking of the hind paw. Normal rats ignore the stimulus or occasionally respond with a very small withdrawal taking place lately without

licking the hind paw. Neuropathic animals respond with a rapid and intense withdrawal followed by a licking of the hind paw.

Three latencies were obtained alternately from each hind paw 5 min apart. No consistent left and right differences were observed. The mean paw withdrawal latency (in seconds) of hind paws was determined from an average of six separate measures (three per paw) at a given time point. Testing boxes were thoroughly cleaned between each test session.

### Openfield test

The openfield (100x100x35 cm) has a 64-square grid floor (8x8 squares, 12.5 cm/side) with an overhead light for illumination. The number of squares entered and the number of rears on the wall were used as indexes of horizontal and vertical exploratory behavior, respectively. These numbers were recorded during a five-minute test period.

### Electrophysiological studies

Sciatic nerve action potential (NAP) recording was performed as previously described [24,25]. Briefly, sciatic nerves were rapidly dissected from decapitated rats under urethane anesthesia (25%, 0.5 ml/100g) and immersed in saline medium (NaCl 133, KCl 2, CaCl<sub>2</sub> 1, MgCl<sub>2</sub> 2, HEPES 10 and glucose 10 pH 7.4) 5 to 10 min before recording. The sciatic nerve was put on chlorinated silver electrodes mounted in a classical homemade electrodes holder chamber. Square-shaped stimulating pulses of 0.1ms duration generated by Clampex routine of the Pclamp software package (Axon Instruments, CA, USA) were applied to the distal end of the nerve through Digidata 1322A interface (Axon Instruments, CA, USA). Bipolar sciatic nerve action potentials (NAP) were recorded at the proximal end of the nerve ( $\Delta x=19$  mm distance from the stimulus point) using an ISO-DAM8A differential amplifier (World Precision Instruments, UK) with a bandwidth of 10 - 10,000 Hz. The signal was digitized (500 kHz) with the Digidata 1322A. Artifact of stimulation was obtained in isolation using a double pulse protocol allowing the second stimulation to occur in refractory period. This stimulus artifact was subtracted from recorded NAPs before analysis with Pclamp software.

Nerve conduction velocity (CV) was calculated using the latency between the beginning of stimulus artifact and the NAP onset ( $\Delta t_{\text{Latency}}$ ) or the NAP peak ( $\Delta t_{\text{Peak}}$ ) for the fastest and relatively slower fibers group respectively according to the equations:

$$CV_{\text{Latency}} = \Delta x / \Delta t_{\text{Latency}}$$

$$CV_{\text{Peak}} = \Delta x / \Delta t_{\text{Peak}}$$

Data obtained from the different conditions were compared using one way ANOVA statistical test followed by Bonferroni's Multiple Comparison Test. Statistical significance was fixed as usually at  $p<0.05$ .

### Immunohistochemical studies

At the end of the last behavioral test session, sciatic nerves and hind paw intra-plantar skins were removed to allow immunohistochemical studies. Thus, animals were deeply anesthetized with 25% urethane (0.5 ml/100g, i.p.) and perfused transcardially with 100 ml of 0.1 M phosphate buffered (PB; pH 7.4). The perfusion was carried out with 450

ml of fixative solution (4% formaldehyde and 0.2% picric acid in PB). Sciatic nerves and plantar skins were rapidly dissected and postfixed in the same fixative solution for 24h. Afterwards, the tissues were immersed in PB containing 15% sucrose for 24h and then transferred into 30% sucrose PB for 24h. Sciatic nerves and intraplantar skins were then placed in embedding medium (OCT, Tissue-Tek, Reichert-Jung, Nussloch, Germany) and immediately frozen at -80°C. Sagittal sections (10- $\mu$ m thick) were cut in a cryostat HM 560 (Microm, Francheville, France) and mounted on glass slides coated with gelatin and chromium potassium sulfate. Tissue sections were pre-incubated for 1 hour with 5% nonimmune donkey or goat serum in PB containing 0.3% Triton X-100 (PBT). Afterwards, the sciatic nerve sections were incubated overnight at 4°C with a mouse monoclonal antibody against NF200 (Clone N52; Sigma-Aldrich, St Louis, MO) diluted at 1:250 in PBT or a mouse monoclonal antibody against 2',3'-cyclic nucleotide 3'-phosphodiesterase (CNPase; Sigma-Aldrich, St Louis, MO) diluted at 1:100 in PBT. Hind paw intra-plantar skin sections were incubated overnight at 4°C with the rabbit polyclonal antibody against the protein gene product PGP9.5 (Cedarlane Laboratories, Ontario, Canada) diluted at 1:500 in PBT. The procedure was carried on by rinsing all of the tissue sections 3 times in PB (15 min/rinse) and transferring them for 2 hours into Alexa-488-conjugated donkey anti-mouse or FITC-conjugated goat anti-rabbit (Chemicon, Temecula, CA) diluted at 1:300 in PBT. Finally, the sections were rinsed for 1 hour in PB and mounted in Vectashield (Vector Laboratories, Burlingame, CA). Although specificity of the antibodies has previously been demonstrated [50-54], internal control experiments were performed in the present work as follows: (i) substitution of CNPase, NF200 or PGP9.5 antiserum with PBT, (ii) replacement of CNPase, NF200 or PGP9.5 antibody by non-immune mouse or rabbit serum and (iii) preincubation of CNPase, NF200 or PGP9.5 antibody with purified CNPase, NF200 or PGP9.5, respectively. The preparations were examined under a multichannel confocal laser-scanning microscope (Leica Confocal Systems, Paris, France) assisted by a pentium IV PC (Leica Microsystems). The number of CNPase or NF200 immunoreactive fibers was determined in a counting square of 200 x 200  $\mu$ m<sup>2</sup> on sagittal sections of sciatic nerves isolated from each experimental group. Quantification of NF200 immunoreactivity was performed on sciatic nerve sagittal sections by using ImageJ software. Results were expressed as % of control (+ SEM). The number of PGP9.5 positive-intraepidermal nerve fibers (IENF) was quantified as previously described [50,52]: all ascending nerve fibers that cross into the epidermis were counted, no minimum length was required and secondary branching into the epidermis was excluded from quantification. IENF counts are expressed as the number per mm of epidermal border. Experimenters were completely blind to the experimental conditions of animals.

### Statistical analyses

One or two-way repeated measures ANOVAs followed by Newman-Keuls *post hoc* comparisons were used. The data which did not exhibit a Gaussian distribution were analyzed by the non-parametric Mann-Whitney *U* test. The statistical

significance of differences between sciatic nerve conduction velocities or NAP amplitude peaks was assessed by one way ANOVA statistical test followed by Bonferroni's Multiple Comparison Test. Data were analyzed with Statistica Software 5.1 (Statsoft, Maisons-Alfort, France). A *p*-value of less than 0.05 was considered significant.

## Results

### Effects of PAC on the mechanical nociceptive threshold

Before the onset of PAC or vehicle (Cremophor 10%) treatment, the percentages of paw withdrawal for mechanical stimulations were 0% for von Frey filaments < 6 g, 10-15% for 15 g and 35-40% for 26 g. In animals receiving the vehicle, the percentages of withdrawal responses remained unchanged all treatment days (Figure 1A-C). In contrast, from D15, paw withdrawal responses in PAC-treated rats were observed (3-26%; *p*<0.005) for von Frey filaments < 6 g (mechanical allodynia) (Figure 1A) and increased levels of responding for 15g (30-71.5%; *p*<0.005) and for 26 g (60-87.5%; *p*<0.005) were detected (mechanical hyperalgesia) (Figure 1B-C). PAC-induced mechanical allodynia and hyperalgesia, which started one week after the withdrawal of PAC treatment, persisted, at least 3 weeks (Figure 1A-C).

### Effects of PAC on cold thermal nociceptive sensitivity


Prior to PAC or vehicle (10% Cremophor in saline) treatment, no paw withdrawal response was induced by acetone spray during the whole observation period (20 s), indicating that acetone-evoked cold stimulation is not noxious in naive rats. In animals receiving the vehicle, the withdrawal latency remained unchanged (no paw withdrawal response during 20 s) all treatment days (Figure 1D). In contrast, the withdrawal latency significantly decreased from 20 s to 15 s at D15 and to 10 s at D29 (last time point recorded) in PAC-treated animals (*p*<0.001), showing that PAC treatment caused cold thermal allodynia in rats (Figure 1D).

### Effects of PAC on heat thermal nociceptive sensitivity

The baseline withdrawal latency characterizing the heat thermal pain threshold was around 11.5  $\pm$  0.2 s on each paw of naive rats before the treatment onset. No significant changes of the thermal thresholds were observed in animal receiving the vehicle (Figure 1E). In contrast, a significant decrease of the heat thermal nociceptive threshold was observed in PAC-treated rats (from 11.4 to 9.2 s) at D16 (*p*<0.05) and this threshold remained significantly lower until D30 (*p*<0.001) (Figure 1E).


### Effects of PAC on sciatic nerve action potential and conduction velocity

The sciatic NAP recorded in our conditions concern the myelinated fibers that may essentially be A $\delta$  fibers owing to their conductance velocity range. As these fibers have different diameters, a large bandwidth of CVs is generated. The higher fiber CVs contribute to the NAP onset while the smallest CVs



**Figure 1. Effect of PAC treatment on the rat mechanical (A-C) and thermal (D, E) nociceptive threshold.** (A-C) Time-course of mechanical allodynia (A) and hyperalgesia (B,C) induced by PAC treatment. Graphs show the mean + SEM of the percentages of paw withdrawal responses to mechanical stimulation by von Frey filament 4 g (A), 15 g (B) or 26 g (C) (n=8 per group). (D,E) Effects of PAC treatment on rat cold (D) or heat (E) thermal nociceptive threshold assessed by acetone (D) or Plantar (E) tests. Each point represents the mean + SEM of 6 observations in each of 8 rats. Non-parametric Mann-Whitney *U* test was used for the analysis of the von Frey test results and one-way repeated measures ANOVAs followed by Newman-Keuls *post hoc* comparisons were used for acetone and Plantar tests. Statistical differences between control and paclitaxel group on each testing day are shown. \*  $p < 0.05$ , \*\*\*  $p < 0.005$ .

doi: 10.1371/journal.pone.0080915.g001



**Figure 2. PAC effects on rat sciatic nerve action potential (NPA) peak amplitude and conduction velocity.** (A) Mean traces of biphasic NAPs recorded from animals treated with vehicle (*black curve*) or PAC (*grey curve*). The nerve was stimulated with a supra-maximal pulse potential (1 V) as indicated by the protocol trace (*lower trace*). The arrows 1-3 indicate the beginning of the stimulation artifact (1) and the onset of NAP in control (2) and PAC (3) conditions giving a calculated NAP CV<sub>latency</sub> in this case of 39.6 and 23.2 m/s for control and PAC, respectively. Note the reduction of NAP peak amplitude in PAC condition (1.2 mV versus 2.9 mV in controls). (B,C) NAP conduction velocity (B) and peak amplitude (C) histograms of statistical data obtained for vehicle- and PAC-treated rats (n=8 for each condition). \*\*  $p < 0.01$ , \*\*\*  $p < 0.005$ .


doi: 10.1371/journal.pone.0080915.g002

are involved in the NAP peak. Figure 2A illustrates mean traces of sciatic NAPs recorded from vehicle- and PAC-treated rats. PAC affected both NAP peak amplitude and conduction velocity. In the case illustrated in Figure 2A, the CV of the fastest fibers (NAP onset) was reduced by 41.4% under PAC treatment (39.6 and 23.2 m/s for vehicle and PAC, respectively) while the slowest fibers (peak of NAP) was decreased about 33.6% (absolute values at NAP peak CV were 13.3 and 8.8 m/s in vehicle and PAC treated rats, respectively). Mean reduction values obtained for CV<sub>latency</sub> and CV<sub>peak</sub> were  $32.9 \pm 0.7$  and  $24.7 \pm 1.0\%$ , respectively (Figure 2B). The NAP peak amplitude was greatly reduced and the values for the NAP's first phase in the case illustrated were 2.9 and 1.2 mV for vehicle- and PAC-treated rats, respectively, giving a 58% inhibition of sciatic nerve function (Figure 2A). Mean values obtained in each condition were  $1.5 \pm 0.2$  and  $0.7$

$\pm 0.1$  mV respectively for vehicle- and PAC-treated rats, indicating that the mean decrease of NAP peak amplitude evoked by PAC was  $54 \pm 3\%$  (Figure 2C).

#### Effects of PAC on NF200 expression in peripheral nerve axons

To determine whether PAC treatment alters the structural organization of peripheral axons, we used immunohistochemical procedure to compare the expression of the key neuronal cytoskeleton protein NF200 [55] in PAC-treated and control rat sciatic nerve axons. Our results showed an intense NF200-like immunoreactivity in several processes in vehicle-treated or control rat sciatic nerves (Figure 3A). PAC-treatment drastically decreased NF200-immunofluorescence in sciatic nerve axons (Figure 3B). Quantitative analysis using the counting score method combined with ImageJ software



**Figure 3. Photomicrographs of sagittal sections of sciatic nerves (A,B,E,F) or hind paw intraplantar skins (C,D) dissected from vehicle (A,C,E)- and PAC (B,D,F)-treated rats.** Nerve sections were labeled with the monoclonal NF200 antibody (A,B) or with the monoclonal anti-CNPase (E,F) revealed with Alexa-488-conjugated donkey anti-mouse. (C,D) Intraplantar skin sections were labeled with the polyclonal anti-PGP9.5 revealed with FITC-conjugated goat anti-rabbit. White arrows indicated intraepidermal nerve fibers. Scale bar=50 $\mu$ m.

doi: 10.1371/journal.pone.0080915.g003

calculation revealed a 33% decrease of NF200-immunostaining in PAC-treated sciatic nerve axons compared to controls ( $p < 0.001$ ) (Figure 4A).


#### Effects of PAC on intraepidermal nerve fiber density in intraplantar skin

Intraepidermal nerve fibers (IENF) were firstly immunolabeled in rat hind paw intraplantar skin with a specific antibody against PGP9.5 (Figure 3C,D). Then, IENF density was quantified in vehicle- and PAC-treated rat hind paw intraplantar skins. PAC treatment decreased by 61% PGP9.5-

immunoreactivity or IENF density in intra-plantar epidermis ( $p < 0.001$ ) (Figure 4B).

#### Effects of PAC on CNPase expression in sciatic nerves

As previous studies demonstrated that the non-compact myelin protein CNPase, pivotal for axonal survival, is dysregulated in experimental peripheral neuropathies [24,43,44], we have determined the effects of PAC treatment on CNPase expression in Schwann cells surrounding axonal processes of the rat sciatic nerve. Intense immunoreactivity for CNPase was visualized on sagittal sections of naive or vehicle-treated rat sciatic nerves (Figure 3E). PAC-treatment



**Figure 4. Comparison of IENF density, NF-200 or CNPase expression in vehicle- and PAC-treated rat nerves.** (A) Comparative analysis of NF200-immunostaining density (actual counts for NF200-positive fibers) detected in sciatic nerve sections dissected from vehicle- and PAC-treated rats. (B) Comparative analysis of IENF density (counts for PGP9.5-positive terminals) measured in vehicle- and PAC-treated rat intraplantar skin sections. (C) Comparative analysis of the numbers of CNPase-positive Schwann cell bodies detected in sciatic nerve sections dissected from vehicle- and PAC-treated rats. Each value is expressed as percent (+ SEM) of CNPase-positive cells bodies detected in sciatic nerve sections of control (vehicle-treated) rats.  $n=8$  per group. \*\*\*  $p<0.001$ .

doi: 10.1371/journal.pone.0080915.g004

significantly reduced the intensity of CNPase-immunostaining in sciatic nerves (Figure 3F). Comparative analysis of the numbers of CNPase-positive Schwann cell bodies detected in counting squares ( $200 \times 200 \mu\text{m}^2$ ) showed that PAC treatment

caused a drastic decrease (-44%) of CNPase expression in sciatic nerves ( $p<0.001$ ) (Figure 4C).



### Action of 3 $\alpha$ -androstanediol against PAC-induced cold allodynia, heat hyperalgesia and mechanical allodynia or hyperalgesia

Although previous dose-response studies have identified 1 to 4 mg/kg as an effective dose range for 3 $\alpha$ -androstanediol neuroactivity *in vivo* [29,32,56,57], we investigated in the present work the dose- and injection frequency-dependent effects of 3 $\alpha$ -DIOL (by testing 2 or 4 mg/kg every 2 or 4 days) on the mechanical nociceptive thresholds of vehicle- and PAC-treated rats (Figure 5). In vehicle-treated animals, corrective 3 $\alpha$ -DIOL treatments at 2 or 4 mg/kg every 2 days (Figure 5A) or at 4 mg/kg every 4 days (Figure 5B) significantly decreased the percent of responses to 26 g ( $p < 0.05$ ), indicating an antinociceptive effect of 3 $\alpha$ -DIOL. In PAC-treated animals, the mechanical hyperalgesia ( $p < 0.05$ ) evoked by PAC was only partially reduced by 3 $\alpha$ -DIOL corrective treatment at the dose of 2 mg/kg every 2 or 4 days or at 4 mg/kg every 4 days ( $p < 0.05$ ; Figure 5). In contrast, at the dose of 4 mg/kg every 2 days, 3 $\alpha$ -DIOL corrective treatment was capable to abolish completely PAC-induced mechano-hyperalgesia and to restore a normal mechanical sensitivity threshold in PAC-treated rats ( $p < 0.05$ ; Figure 5A). Interestingly, we observed that the prophylactic treatment with 3 $\alpha$ -DIOL at the same dose (4 mg/kg/2 days) has also successfully prevented the occurrence of PAC-evoked abnormal mechanical perception (Figure 6). Consequently, 3 $\alpha$ -DIOL at 4 mg/kg injected every 2 days appeared as the best or optimal dose to be considered for the elaboration of an effective corrective or prophylactic strategy against PAC-induced neuropathic symptoms in rats. Indeed, 3 $\alpha$ -DIOL (4 mg/kg/2 days), which increased the mechanical and thermal pain thresholds ( $p < 0.05$ ), exerted a potent antinociceptive action in control rats (Figure 6C,E,H,J). When neuropathic pain symptoms are already installed in PAC-treated rats, corrective 3 $\alpha$ -DIOL treatment administered during two weeks (7 injections of 3 $\alpha$ -DIOL 4 mg/kg) totally abolished the cold allodynia, heat hyperalgesia as well as the mechanical allodynia and hyperalgesia ( $p < 0.005$ ) by restoring normal cold- and heat-thermal and mechanical pain thresholds in PAC-treated animals (Figure 6A-E). Prophylactic administration of 3 $\alpha$ -DIOL (4 mg/kg every 2 days), which started 8 days before PAC injections and was maintained during PAC treatment, completely prevented PAC-induced cold allodynia, heat hyperalgesia and mechanical allodynia or hyperalgesia ( $p < 0.005$ ) (Figure 6F-J). Neuropathic pain symptoms did not reappear in PAC-treated rats one week (D36 or D37) after the end of 3 $\alpha$ -DIOL corrective or prophylactic treatment (Figure 6). To verify that the beneficial action of 3 $\alpha$ -DIOL against PAC-induced painful neuropathy is not due to a persistent sedative effect causing weakness and reflex slowing, we have used the openfield test to assess the motor capacity of control and PAC-treated rats after 3 $\alpha$ -DIOL corrective or prophylactic treatment. As shown in Table 1, horizontal and vertical exploratory behaviors were similar in all tested groups indicating that 3 $\alpha$ -DIOL corrective or prophylactic treatment did not reduce the rat motor activity.

### Effects of 3 $\alpha$ -androstanediol against PAC-induced peripheral nerve functional deficit

Treatment of naive rats with 3 $\alpha$ -DIOL did not modify peripheral nerve activity even though an increase tendency of the mean values was observed in 3 $\alpha$ -DIOL-treated rats compared to vehicle treated animals (Figure 7). Most importantly, 3 $\alpha$ -DIOL reversed to normal values decreased CV<sub>latency</sub>, CV<sub>peak</sub> and NAP peak amplitude induced by PAC treatment (Figure 7A-C). CV<sub>latency</sub> ( $34.1 \pm 1.1$  m/s), CV<sub>peak</sub> ( $13.0 \pm 0.4$  m/s) and NAP peak amplitude ( $1.5 \pm 0.2$  mV) mean values of (PAC+3 $\alpha$ -DIOL)-treated rats were similar to the values of control animals ( $34.6 \pm 0.6$ ,  $13.0 \pm 0.4$  m/s and  $1.5 \pm 0.1$  mV, respectively).

### Effects of 3 $\alpha$ -androstanediol on PAC-induced NF200 repression in peripheral nerve axons

We observed that PAC treatment induced 30% decrease of NF200-immunoreactivity in rat sciatic nerves (Figure 8A,B and Figure 9A) ( $p < 0.001$ ). 3 $\alpha$ -DIOL treatment administered during two weeks totally reversed to normal decreased NF200 expression level induced by PAC in rat sciatic nerve axons ( $p < 0.001$ ) (Figure 8A-D and Figure 9A).

### Effects of 3 $\alpha$ -androstanediol on PAC-induced loss of intraepidermal nerve fibers


We detected 55% decrease of PGP9.5-immunoreactivity or IENF density in hind paw intra-plantar skins of PAC-treated rats compared to controls ( $p < 0.001$ ) (Figure 8E,F and Figure 9B). 3 $\alpha$ -DIOL treatment during two weeks restored PGP9.5 or IENF normal density in PAC-treated animals ( $p < 0.001$ ) (Figure 8E-H and Figure 9B).

### Effect of 3 $\alpha$ -androstanediol on CNPase expression in sciatic nerve

A strong increase of CNPase-immunoreactivity was visualized in sciatic nerves of 3 $\alpha$ -DIOL- compared to vehicle- and PAC-treated rats (Figure 8I-L). In particular, counting of CNPase-positive cells revealed that 3 $\alpha$ -DIOL induced a 26% increase ( $p < 0.001$ ) of CNPase expression in naive or healthy rat sciatic nerves (Figure 9C). Interestingly, the 40% decrease of CNPase-immunoreactivity evoked by PAC-treatment in rat sciatic nerves was totally abolished by 3 $\alpha$ -DIOL treatment administered during two weeks ( $p < 0.001$ ) (Figure 8I-L and Figure 9C).

## Discussion

The present study revealed that a 7 day-treatment schedule with PAC (1 mg/kg, daily body weight) induced painful peripheral neuropathy and associated neurological symptoms in rats. In particular, we observed that PAC treatment alters functional, histological and neurochemical parameters of the rat peripheral nerves as previously described in humans and various other experimental models [1,3,12-18,58,59]. Indeed, after one week treatment with PAC, the treated rats developed physiological and functional abnormalities such as mechanical allodynia and hyperalgesia, cold allodynia, heat-thermal




**Figure 5. Dose- and injection frequency-dependent effects of corrective 3 $\alpha$ -DIOL treatment on the mechanical nociceptive thresholds of control- and PAC-treated rats.** Corrective treatment every 2- (A) or 4-days (B) consisted in starting 3 $\alpha$ -DIOL (2 or 4 mg/kg) or VEH<sub>hpc</sub> i.p. administrations 2 days after the end of PAC treatment. Threshold values represent responses to 26 g von Frey filament (%  $\pm$  SEM). Non-parametric Mann-Whitney *U* test was used. Statistical differences between controls and each treatment group at each testing day are shown. *n*=6 per group; \* *p*<0.05. \* compared to VEH<sub>crem</sub>+VEH<sub>hpc</sub>; # compared to PAC +VEH<sub>hpc</sub>.

doi: 10.1371/journal.pone.0080915.g005

hyperalgesia, nerve conduction velocity decrease and NAP peak reduction. In addition, PAC also caused histopathological and neurochemical alterations including the repression of NF200 and CNPase in peripheral nerves and the decrease of IENF density in paw intraplantar skin. It is well known that NF200 is a constitutive protein of the cytoskeleton that pivotally controls neuronal cell morphology and axoplasmic transport. Therefore, our results showing PAC-evoked NF200 repression

in sciatic nerves suggest that PAC treatment may generate axonopathy in rats [1,3,60,61]. In agreement with this suggestion, neurophysiological investigation and peripheral nerve biopsy after docetaxel treatment in humans revealed an axonal neuropathy with a preferential loss of large myelinated fibers [62]. Our studies also show that the natural neurosteroid 3 $\alpha$ -DIOL is capable of preventing or correcting several major neuropathological symptoms evoked by PAC treatment.



**Figure 6. Effects of 3 $\alpha$ -DIOL (4 mg/kg/2 days) corrective (A-E) or prophylactic (F-J) treatment on PAC-induced neuropathic pain symptoms. (A-C,F-H)** Action of 3 $\alpha$ -DIOL against PAC-induced mechanical allodynia (A,F) and hyperalgesia (B,C,G,H). Chartbars show the mean + SEM of the percentages of paw withdrawal responses to mechanical stimulation by von Frey filament 4 g (A,F), 15 g (B,G) or 26 g (C,H) (n=8 per group). (D,I) Effect of 3 $\alpha$ -DIOL against PAC-evoked cold-allodynia. (E,J) 3 $\alpha$ -DIOL effects on the heat thermal nociceptive thresholds of vehicle- and PAC-treated rats. Each bar represents the mean + SEM of 6 observations in each of 8 rats. Non-parametric Mann-Whitney *U* test was used for the analysis of the von Frey test results and two-way repeated measures ANOVAs followed by Newman-Keuls *post hoc* comparisons were used for acetone and Plantar Tests. Statistical differences between controls and each treatment group at each testing day are shown. \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.005$ . \* vs (VEH<sub>crem</sub> + VEH<sub>hpc</sub>), # vs (PAC + VEH<sub>hpc</sub>).

doi: 10.1371/journal.pone.0080915.g006

**Table 1.** Effect of 3 $\alpha$ -DIOL treatment (4 mg/kg/every 2 days) on the exploratory behaviors of control- and PAC-treated rats.

3 $\alpha$ -DIOL Treatment Group		Total number of squares entered	Total number of rears on the wall
Corrective	VEH <sub>crem</sub> +VEH <sub>hpc</sub>	327.25 $\pm$ 6.41	32 $\pm$ 1.22
	VEH <sub>crem</sub> +3 $\alpha$ -DIOL	320.5 $\pm$ 5.55	30.25 $\pm$ 1.70
		ns1	ns1
	PAC+VEH <sub>hpc</sub>	317.5 $\pm$ 9.51	31.5 $\pm$ 1.66
		ns1	ns1
	PAC+3 $\alpha$ -DIOL	313 $\pm$ 8.18	30.5 $\pm$ 0.87
	ns1,2	ns1,2	
Prophylactic	VEH <sub>crem</sub> +VEH <sub>hpc</sub>	308 $\pm$ 13.28	29.75 $\pm$ 1.11
	VEH <sub>crem</sub> +3 $\alpha$ -DIOL	305.75 $\pm$ 7.87	30.75 $\pm$ 2.66
		ns1	ns1
	PAC+VEH <sub>hpc</sub>	295.75 $\pm$ 15.44	31.25 $\pm$ 1.65
		ns1	ns1
	PAC+3 $\alpha$ -DIOL	293 $\pm$ 6.31	31 $\pm$ 1.58
	ns1,2	ns1,2	

Total numbers of squares entered ( $\pm$  SEM) and of rears on the wall ( $\pm$  SEM) indicated in the table are representative of horizontal and vertical exploratory behaviors in the openfield test. For the corrective 3 $\alpha$ -DIOL treatment schedule, PAC or VEH<sub>crem</sub> was injected at D1, D3, D5 and D7; 3 $\alpha$ -DIOL or VEH<sub>hpc</sub> i.p. administrations started 8 days after the end of PAC treatment. For the prophylactic treatment schedule, 3 $\alpha$ -DIOL or VEH<sub>hpc</sub> i.p. was administered every 2 days from D1 while PAC or VEH<sub>crem</sub> treatment, which started from D9, was administered at D9, D11, D13 and D15. Testing was performed at D28, i.e. 1 day after the 7<sup>th</sup> 3 $\alpha$ -DIOL injection in the corrective treatment schedule or 3 days after the 13<sup>th</sup> 3 $\alpha$ -DIOL injection in the prophylactic treatment schedule n=4 per group; ns1, not significant compared to VEH<sub>crem</sub>+VEH<sub>hpc</sub>; ns2, not significant compared to PAC+VEH<sub>hpc</sub>. VEH, vehicle; 3 $\alpha$ -DIOL, 3 $\alpha$ -androstenediol; PAC, paclitaxel.


doi: 10.1371/journal.pone.0080915.t001

Indeed, our behavioral analyses demonstrated that prophylactic or corrective 3 $\alpha$ -DIOL treatment prevented or suppressed PAC-induced neuropathic pain symptoms including cold allodynia, heat-thermal hyperalgesia and mechanical allodynia and hyperalgesia. Interestingly, our histopathological and neurochemical studies demonstrated that 3 $\alpha$ -DIOL reversed to normal the repressed NF200 level observed in PAC-treated rat peripheral nerve axons, suggesting that 3 $\alpha$ -DIOL-based therapy may be effective for the treatment of antineoplastic-evoked axonopathy and/or neuronopathy. Furthermore, our work revealed that 3 $\alpha$ -DIOL prevented or counteracted PAC-induced PGP9.5-like immunoreactivity and IENF density decrease in intraplantar skin. Because epidermal innervation loss constitutes a neuropathological damage frequently occurring in antineoplastic-treated patients [1,52,63], 3 $\alpha$ -DIOL appears as a promising neuroprotective compound against toxic drug-induced alterations of epidermal innervations.

It is well demonstrated that painful neurological symptoms such as allodynia and hyperalgesia evoked by anticancer drugs are generally associated with decreased peripheral nerve

conduction velocity and ectopic discharges [4,13,17,60,64-66]. Consistently, our electrophysiological and behavioral investigations clearly evidenced a significant CV<sub>latency</sub> reduction, NAP peak amplitude decrease and a marked allodynia and hyperalgesia in PAC-treated rats. Interestingly, 3 $\alpha$ -DIOL was also capable of rescuing PAC-treated rats from these electrophysiological and functional alterations evoked by PAC treatment. Moreover, in NF knock-out mice, decreased NAP peak amplitude has been evidenced together with internodal conductance changes that are known to be responsible for nerve conduction block and CV reduction [67]. Low NF expression levels, which are major intrinsic determinants of axon caliber in large fibers, led to axonal diameter reduction and subsequently to CV slowing [67-70]. In perfect accordance with these data, our findings showed a drastic reduction of NF expression level in PAC-treated rat peripheral nerves that was correlated to a significant decrease of large fiber CV and NAP peak amplitude. Therefore, the neuroprotective effect of 3 $\alpha$ -DIOL against PAC-induced painful neuropathological symptoms appears to be dependent on its ability to reverse to normal values decreased NF expression level, reduced CV<sub>latency</sub> and NAP peak amplitude in PAC-treated rat peripheral nerves.

Moreover, our results also revealed that 3 $\alpha$ -DIOL modulate differentially the expression of CNPase, NF200 and PGP9.5 genes. Indeed, in naive or vehicle-treated rats, 3 $\alpha$ -DIOL increased CNPase expression level in peripheral nerves but did not modify the amounts of NF200 and PGP9.5. Interestingly, under neuropathological situations when NF200, PGP9.5 and CNPase levels were repressed by PAC treatment, 3 $\alpha$ -DIOL was able to restore to normal the levels of all of these 3 key proteins controlling peripheral nerve activities and intraepidermal innervation sensory functions. Altogether, these results suggest that CNPase gene expression may be highly sensitive to 3 $\alpha$ -DIOL-evoked intracellular signaling while the sensitivity of NF200 and PGP9.5 genes to 3 $\alpha$ -DIOL may increase only under pathophysiological situations thanks to the activation of various transcription factors evoked by PAC treatment [71-73]. Also, 3 $\alpha$ -DIOL is known to stimulate the central inhibitory system via the GABA<sub>A</sub> receptor, suggesting that 3 $\alpha$ -DIOL may exert an analgesic action which may contribute to suppress PAC-evoked painful symptoms [26-31,46]. Another important point is the fact that 3 $\alpha$ -DIOL may be inter-converted into 5 $\alpha$ -DHT by the enzyme 3 $\alpha$ -hydroxysteroid oxido-reductase abundantly expressed in the nervous system [39,74,75]. Therefore, the treatment with exogenous 3 $\alpha$ -DIOL may generate substantial endogenous concentrations of 3 $\alpha$ -DIOL and 5 $\alpha$ -DHT which can exert both neuroprotective and analgesic effects by activating GABA<sub>A</sub> receptors and androgen nuclear receptors, respectively [27-29,31,32,34,46,76]. The genomic mechanisms activated by 3 $\alpha$ -DIOL treatment (via its metabolite 5 $\alpha$ -DHT) may explain 3 $\alpha$ -DIOL ability to restore to normal values the decreased CNPase, NF200 and PGP9.5 expression levels evoked by PAC treatment in rat peripheral nerves. In support of this hypothesis, it has been shown that androgens, acting via their nuclear receptors, promote NF expression and neuritic process development in peripheral nerves [77]. Furthermore, it has also




**Figure 7. Curative effects of 3 $\alpha$ -DIOL against PAC-induced reduction of sciatic nerve action potential parameters (CV<sub>latency</sub>: A; CV<sub>peak</sub>: B and peak amplitude: C).** Histograms of normalized CV<sub>latency</sub> and CV<sub>peak</sub> show their reduction by PAC and recovery by 3 $\alpha$ -DIOL treatment (A,B). Mean CV values were calculated as % of the mean CV obtained from vehicle-treated rats. (C) Recovery from PAC-induced NAP peak amplitude decrease by 3 $\alpha$ -DIOL treatment. Each value is the mean (+SEM) of NAP peak amplitude obtained from 8 rat sciatic nerves per each group investigated. \*\*  $p < 0.01$ , \*\*\*  $p < 0.005$ . \* vs (VEH<sub>crem</sub> + VEH<sub>hpc</sub>), # vs (PAC + VEH<sub>hpc</sub>). doi: 10.1371/journal.pone.0080915.g007

been demonstrated that, in rodents, androgens promote the expression of myelin proteins including CNPase level [78].

The beneficial effects exerted by 3 $\alpha$ -DIOL against thermal and mechanical allodynic and hyperalgesic symptoms may result from the direct interactions of 3 $\alpha$ -DIOL with GABA<sub>A</sub> receptors inducing therefore and acute analgesia. However,

several indications provided by the present study show that the beneficial action of 3 $\alpha$ -DIOL against PAC-evoked painful neuropathy does not depend on 3 $\alpha$ -DIOL-induced sedation or is not only the consequence of 3 $\alpha$ -DIOL-evoked acute analgesia but results from the combination of 3 $\alpha$ -DIOL neuroprotective ability, nerve tissue repairing capacity and long




**Figure 8. Photomicrographs of sagittal sections of sciatic nerves (A-D,I-L) or intraplantar skins (E-H) dissected from (VEH<sub>crem</sub> + VEH<sub>hpc</sub>)(A,E,I)-, (PAC + VEH<sub>hpc</sub>)(B,F,J)-, (VEH<sub>crem</sub> + 3 $\alpha$ -DIOL)(C,G,K)- or (PAC + 3 $\alpha$ -DIOL)(D,H,L)-treated rats. Nerve sections were labeled with the monoclonal NF200 antibody (A-D) or with the monoclonal anti-CNPase (I-L) revealed with Alexa-488-conjugated donkey anti-mouse. (E-H) Intraplantar skin sections were labeled with the polyclonal anti-PGP9.5 revealed with FITC-conjugated goat anti-rabbit. White arrows indicated intraepidermal nerve fibers. Scale bar, 50  $\mu$ m.**

doi: 10.1371/journal.pone.0080915.g008

term analgesic property. Indeed, we observed that prophylactic 3 $\alpha$ -DIOL treatment prevented neurochemical and functional alterations of peripheral nerves and IENF in PAC-treated rats and these animals did not exhibit painful neuropathological symptoms several days after the end of 3 $\alpha$ -DIOL preventive administration. In addition, one week after withdrawal of 3 $\alpha$ -DIOL corrective treatment, painful neurological symptoms did not reappear in PAC-treated rats. Finally, our openfield investigations revealed that the exploratory behavior and motor activity were similar in control and PAC-treated rats after 3 $\alpha$ -DIOL preventive or corrective treatment, indicating that 3 $\alpha$ -

DIOL beneficial effects against PAC-evoked painful peripheral neuropathy and neurological symptoms did not result from its sedative action.

In conclusion, the present study, which shows that the neuroprotective neurosteroid 3 $\alpha$ -DIOL effectively prevents and suppresses several painful and neuropathological symptoms evoked by PAC, also offers interesting opportunities for the development neurosteroid-based therapies against chemotherapy-induced peripheral neuropathy and painful neurological disorders.



**Figure 9. Neuroprotective effects of 3 $\alpha$ -DIOL corrective treatment against PAC-induced alterations in sciatic nerves and intraplantar skin.** (A-C) Chartbars show NF200 (A) or CNPase (C) expression level in sciatic nerve sections or IENF density in intraplantar skin sections (B) dissected from (VEH<sub>crem</sub> + VEH<sub>hpc</sub>)-, (PAC + VEH<sub>hpc</sub>)-, (VEH<sub>crem</sub> + 3 $\alpha$ -DIOL)- and (PAC + 3 $\alpha$ -DIOL)-treated rats. (A) Each value is expressed as mean (+ SEM) of actual counts for NF200-positive fibers detected in sciatic nerve sections. (B) Each value is expressed as mean (+ SEM) of IENF density (counts for PGP9.5-positive terminals) detected in intraplantar skin sections. (C) Each value is expressed as percent (+ SEM) of CNPase-positive cells bodies detected in sciatic nerve sections of control (VEH<sub>crem</sub> + VEH<sub>hpc</sub>)-treated rats. n=8 per group. \*\*\*  $p < 0.001$  vs (VEH<sub>crem</sub> + VEH<sub>hpc</sub>), ###  $p < 0.001$  vs (PAC + VEH<sub>hpc</sub>).

doi: 10.1371/journal.pone.0080915.g009

## Author Contributions

Conceived and designed the experiments: AMN. Performed the experiments: LM CPM OT. Analyzed the data: LM CPM OT

AMN. Contributed reagents/materials/analysis tools: LM CPM OT AMN. Wrote the manuscript: LM CPM OT AMN.

## References

- Balayssac D, Ferrier J, Descoeur J, Ling B, Pezet D, Eschalier A, Authier N (2011) Chemotherapy-induced peripheral neuropathies: from clinical relevance to preclinical evidence. *Expert Opin Drug Saf* 10: 407-417. doi:10.1517/14740338.2011.543417. PubMed: 21210753.
- Quasthoff S, Hartung HP (2002) Chemotherapy-induced peripheral neuropathy. *J Neurol* 249: 9-17. doi:10.1007/PL00007853. PubMed: 11954874.
- Authier N, Balayssac D, Marchand F, Ling B, Zangarelli A, Descoeur J, Coudore F, Bourinet E, Eschalier A (2009) Animal models of chemotherapy-evoked painful peripheral neuropathies. *Neurotherapeutics* 6: 620-629. doi:10.1016/j.nurt.2009.07.003. PubMed: 19789067.
- Dougherty PM, Cata JP, Cordella JV, Burton A, Weng HR (2004) Taxol-induced sensory disturbance is characterized by preferential impairment of myelinated fiber function in cancer patients. *Pain* 109: 132-142. doi:10.1016/j.pain.2004.01.021. PubMed: 15082135.
- Verstappen CC, Heimans JJ, Hoekman K, Postma TJ (2003) Neurotoxic complications of chemotherapy in patients with cancer: clinical signs and optimal management. *Drugs* 63: 1549-1563. doi:10.2165/00003495-200363150-00003. PubMed: 12887262.
- Cai J, Zheng T, Masood R, Smith DL, Hinton DR, Kim CN, Fang G, Bhalla K, Gill PS (2000) Paclitaxel Induces Apoptosis in AIDS-Related Kaposi's Sarcoma Cells. *Sarcoma* 4: 37-45. doi:10.1155/S1357714X00000074. PubMed: 18521433.
- Croom KF, Dhillon S (2011) Bevacizumab: a review of its use in combination with paclitaxel or capecitabine as first-line therapy for HER2-negative metastatic breast cancer. *Drugs* 71: 2213-2229. doi:10.2165/11207720-000000000-00000. PubMed: 22035518.
- Holmes FA, Walters RS, Theriault RL, Forman AD, Newton LK, Raber MN, Buzdar AU, Frye DK, Hortobagyi GN (1991) Phase II trial of taxol, an active drug in the treatment of metastatic breast cancer. *J Natl Cancer Inst* 83: 1797-1805. doi:10.1093/jnci/83.24.1797. PubMed: 1683908.
- Le XF, Bast RC Jr. (2011) Src family kinases and paclitaxel sensitivity. *Cancer Biol Ther* 12: 260-269. doi:10.4161/cbt.12.4.16430. PubMed: 21646863.
- Singh S, Dash AK (2009) Paclitaxel in cancer treatment: perspectives and prospects of its delivery challenges. *Crit Rev Ther Drug Carrier Syst* 26: 333-372. doi:10.1615/CritRevTherDrugCarrierSyst.v26.i4.10. PubMed: 20001890.
- Young M, Plosker GL (2001) Paclitaxel: a pharmacoeconomic review of its use in the treatment of ovarian cancer. *Pharmacoeconomics* 19: 1227-1259. doi:10.2165/00019053-200119120-00005. PubMed: 11772158.
- Cavaletti G, Bogliun G, Marzorati L, Zincone A, Marzola M, Colombo N, Tredici G (1995) Peripheral neurotoxicity of taxol in patients previously treated with cisplatin. *Cancer* 75: 1141-1150. doi:10.1002/1097-0142(19950301)75:5. PubMed: 7850713.
- Chaudhry V, Rowinsky EK, Sartorius SE, Donehower RC, Cornblath DR (1994) Peripheral neuropathy from taxol and cisplatin combination chemotherapy: clinical and electrophysiological studies. *Ann Neurol* 35: 304-311. doi:10.1002/ana.410350310. PubMed: 7907208.
- Forsyth PA, Balmaceda C, Peterson K, Seidman AD, Brasher P, DeAngelis LM (1997) Prospective study of paclitaxel-induced peripheral neuropathy with quantitative sensory testing. *J Neuro Oncol* 35: 47-53. doi:10.1023/A:1005805907311. PubMed: 9266440.
- Gordon AN, Stringer CA, Matthews CM, Willis DL, Nemunaitis J (1997) Phase I dose escalation of paclitaxel in patients with advanced ovarian cancer receiving cisplatin: rapid development of neurotoxicity is dose-limiting. *J Clin Oncol* 15: 1965-1973. PubMed: 9164208.
- Polomano RC, Mannes AJ, Clark US, Bennett GJ (2001) A painful peripheral neuropathy in the rat produced by the chemotherapeutic drug, paclitaxel. *Pain* 94: 293-304. doi:10.1016/S0304-3959(01)00363-3. PubMed: 11731066.
- Rowinsky EK, Eisenhauer EA, Chaudhry V, Arbus SG, Donehower RC (1993) Clinical toxicities encountered with paclitaxel (Taxol). *Semin Oncol* 20: 1-15. PubMed: 8102012.
- van den Bent MJ, van Raaij-van den Aarsen VJ, Verweij J, Doorn PA, Smitt Sillevis (1997) Progression of paclitaxel-induced neuropathy following discontinuation of treatment. *Muscle Nerve* 20: 750-752. doi:10.1002/(SICI)1097-4598(199706)20:6. PubMed: 9149085.
- Durand JP, Alexandre J, Guillevin L, Goldwasser F (2005) Clinical activity of venlafaxine and topiramate against oxaliplatin-induced disabling permanent neuropathy. *Anti Cancer Drugs* 16: 587-591. doi:10.1097/00001813-200506000-00015. PubMed: 15846125.
- Gamelin L, Boisdron-Celle M, Delva R, Guérin-Meyer V, Ifrah N, Morel A, Gamelin E (2004) Prevention of oxaliplatin-related neurotoxicity by calcium and magnesium infusions: a retrospective study of 161 patients receiving oxaliplatin combined with 5-Fluorouracil and leucovorin for advanced colorectal cancer. *Clin Cancer Res* 10: 4055-4061. doi:10.1158/1078-0432.CCR-03-0666. PubMed: 15217938.
- Lang E, Hord AH, Denson D (1996) Venlafaxine hydrochloride (Effexor) relieves thermal hyperalgesia in rats with an experimental mononeuropathy. *Pain* 68: 151-155. doi:10.1016/S0304-3959(96)03223-X. PubMed: 9252010.
- Ling B, Authier N, Balayssac D, Eschalier A, Coudore F (2007) Behavioral and pharmacological description of oxaliplatin-induced painful neuropathy in rat. *Pain* 128: 225-234. doi:10.1016/j.pain.2006.09.016. PubMed: 17084975.
- Marchand F, Alloui A, Chapuy E, Jourdan D, Pelissier T, Ardid D, Hernandez A, Eschalier A (2003) Evidence for a monoamine mediated, opioid-independent, antihyperalgesic effect of venlafaxine, a non-tricyclic antidepressant, in a neurogenic pain model in rats. *Pain* 103: 229-235. doi:10.1016/S0304-3959(03)00168-4. PubMed: 12791429.
- Meyer L, Patte-Mensah C, Taleb O, Mensah-Nyagan AG (2010) Cellular and functional evidence for a protective action of neurosteroids against vincristine chemotherapy-induced painful neuropathy. *Cell Mol Life Sci* 67: 3017-3034. doi:10.1007/s00018-010-0372-0. PubMed: 20431905.
- Meyer L, Patte-Mensah C, Taleb O, Mensah-Nyagan AG (2011) Allopregnanolone prevents and suppresses oxaliplatin-evoked painful neuropathy: multi-parametric assessment and direct evidence. *Pain* 152: 170-181. doi:10.1016/j.pain.2010.10.015. PubMed: 21071147.
- Belelli D, Lambert JJ (2005) Neurosteroids: endogenous regulators of the GABA(A) receptor. *Nat Rev Neurosci* 6: 565-575. doi:10.1038/nrn1703. PubMed: 15959466.
- Bitran D, Hilvers RJ, Frye CA, Erskine MS (1996) Chronic anabolic-androgenic steroid treatment affects brain GABA(A) receptor-gated chloride ion transport. *Life Sci* 58: 573-583. doi:10.1016/0024-3205(95)02326-7. PubMed: 8632710.
- Frye CA, Duncan JE, Basham M, Erskine MS (1996) Behavioral effects of 3 alpha-androstenediol. II: Hypothalamic and preoptic area actions via a GABAergic mechanism. *Behav Brain Res* 79: 119-130. doi:10.1016/0166-4328(96)00005-8. PubMed: 8883823.
- Frye CA, Park D, Tanaka M, Rosellini R, Svare B (2001) The testosterone metabolite and neurosteroid 3alpha-androstenediol may mediate the effects of testosterone on conditioned place preference. *Psychoneuroendocrinol* 26: 731-750. doi:10.1016/S0306-4530(01)00027-0. PubMed: 1174408411500254.
- Hosie AM, Wilkins ME, da Silva HM, Smart TG (2006) Endogenous neurosteroids regulate GABAA receptors through two discrete transmembrane sites. *Nature* 444: 486-489. doi:10.1038/nature05324. PubMed: 17108970.
- Reddy DS (2008) Mass spectrometric assay and physiological-pharmacological activity of androgenic neurosteroids. *Neurochem Int* 52: 541-553. doi:10.1016/j.neuint.2007.05.019. PubMed: 17624627.
- Frye CA, Babson A, Walf AA (2007) Self-administration of 3alpha-androstenediol increases locomotion and analgesia and decreases aggressive behavior of male hamsters. *Pharmacol Biochem Behav* 86: 415-421. doi:10.1016/j.pbb.2006.05.025. PubMed: 16828856.
- Rhodes ME, Frye CA (2004) Androgens in the hippocampus can alter, and be altered by, ictal activity. *Pharmacol Biochem Behav* 78: 483-493. doi:10.1016/j.pbb.2004.04.020. PubMed: 15251257.
- Edinger KL, Frye CA (2004) Testosterone's analgesic, anxiolytic, and cognitive-enhancing effects may be due in part to actions of its 5alpha-reduced metabolites in the hippocampus. *Behav Neurosci* 118: 1352-1364. doi:10.1037/0735-7044.118.6.1352. PubMed: 15598144.
- Akwa Y, Purdy RH, Koob GF, Britton KT (1999) The amygdala mediates the anxiolytic-like effect of the neurosteroid allopregnanolone



- in rat. *Behav Brain Res* 106: 119-125. doi:10.1016/S0166-4328(99)00101-1. PubMed: 10595427.
36. Ciriza I, Azcoitia I, Garcia-Segura LM (2004) Reduced progesterone metabolites protect rat hippocampal neurons from kainic acid excitotoxicity in vivo. *J Neuroendocrinol* 16: 58-63. doi:10.1111/j.1365-2826.2004.01121.x. PubMed: 14962077.
  37. Ciriza I, Carrero P, Frye CA, Garcia-Segura LM (2006) Reduced metabolites mediate neuroprotective effects of progesterone in the adult rat hippocampus. The synthetic progestin medroxyprogesterone acetate (Provera) is not neuroprotective. *J Neurobiol* 66: 916-928. doi:10.1002/neu.20293. PubMed: 16758493.
  38. Griffin LD, Gong W, Verot L, Mellon SH (2004) Niemann-Pick type C disease involves disrupted neurosteroidogenesis and responds to allopregnanolone. *Nat Med* 10: 704-711. doi:10.1038/nm1073. PubMed: 15208706.
  39. Meyer L, Venard C, Schaeffer V, Patte-Mensah C, Mensah-Nyagan AG (2008) The biological activity of 3 $\alpha$ -hydroxysteroid oxidoreductase in the spinal cord regulates thermal and mechanical pain thresholds after sciatic nerve injury. *Neurobiol Dis* 30: 30-41. doi:10.1016/j.nbd.2007.12.001. PubMed: 18291663.
  40. Patte-Mensah C, Kibaly C, Boudard D, Schaeffer V, Béglié A, Saredi S, Meyer L, Mensah-Nyagan AG (2006) Neurogenic pain and steroid synthesis in the spinal cord. *J Mol Neurosci* 28: 17-31. doi:10.1385/JMN:28:1:17. PubMed: 16632873.
  41. Uzunova V, Sheline Y, Davis JM, Rasmusson A, Uzunov DP, Costa E, Guidotti A (1998) Increase in the cerebrospinal fluid content of neurosteroids in patients with unipolar major depression who are receiving fluoxetine or fluvoxamine. *Proc Natl Acad Sci U S A* 95: 3239-3244. doi:10.1073/pnas.95.6.3239. PubMed: 9501247.
  42. Wang JM, Singh C, Liu L, Irwin RW, Chen S, Chung EJ, Thompson RF, Brinton RD (2010) Allopregnanolone reverses neurogenic and cognitive deficits in mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 107: 6498-6503. doi:10.1073/pnas.1001422107. PubMed: 20231471.
  43. Lappe-Siefke C, Goebbels S, Gravel M, Nicksch E, Lee J, Braun PE, Griffiths IR, Nave KA (2003) Disruption of Cnp1 uncouples oligodendroglial functions in axonal support and myelination. *Nat Genet* 33: 366-374. doi:10.1038/ng1095. PubMed: 12590258.
  44. LeBlanc AC, Pringle J, Lemieux J, Poduslo JF, Mezei C (1992) Regulation of 2',3'-cyclic nucleotide phosphodiesterase gene expression in experimental peripheral neuropathies. *Brain Res. Mol Brain Res* 15: 40-46. doi:10.1016/0169-328X(92)90149-6. PubMed: 1279349.
  45. Baulieu EE, Robel P, Schumacher M (1999) *Contemporary Endocrinology*. Totowa: Humana Press. 378 pp.
  46. Melcangi RC, Garcia-Segura LM, Mensah-Nyagan AG (2008) Neuroactive steroids: state of the art and new perspectives. *Cell Mol Life Sci* 65: 777-797. doi:10.1007/s00018-007-7403-5. PubMed: 18038216.
  47. Mensah-Nyagan AG, Do-Rego JL, Beaujean D, Luu-The V, Pelletier G, Vaudry H (1999) Neurosteroids: expression of steroidogenic enzymes and regulation of steroid biosynthesis in the central nervous system. *Pharmacol Rev* 51: 63-81. PubMed: 10049998.
  48. Zimmermann M (1983) Ethical guidelines for investigations of experimental pain in conscious animals. *Pain* 16: 109-110. doi:10.1016/0304-3959(83)90201-4. PubMed: 6877845.
  49. Hargreaves K, Dubner R, Brown F, Flores C, Joris J (1988) A new and sensitive method for measuring thermal nociception in cutaneous hyperalgesia. *Pain* 32: 77-88. doi:10.1016/0304-3959(88)90026-7. PubMed: 3340425.
  50. Lauria G, Lombardi R, Borgna M, Penza P, Bianchi R, Savino C, Canta A, Nicolini G, Marmiroli P, Cavaletti G (2005) Intraepidermal nerve fiber density in rat foot pad: neuropathologic-neurophysiologic correlation. *J Peripher Nerv Syst* 10: 202-208. doi:10.1111/j.1085-9489.2005.0010210.x. PubMed: 15958131.
  51. Shaw G, Osborn M, Weber K (1986) Reactivity of a panel of neurofilament antibodies on phosphorylated and dephosphorylated neurofilaments. *Eur J Cell Biol* 42: 1-9. PubMed: 3539605.
  52. Siau C, Xiao W, Bennett GJ (2006) Paclitaxel- and vincristine-evoked painful peripheral neuropathies: loss of epidermal innervation and activation of Langerhans cells. *Exp Neurol* 201: 507-514. doi:10.1016/j.expneurol.2006.05.007. PubMed: 16797537.
  53. Sprinkle TJ (1989) 2',3'-cyclic nucleotide 3'-phosphodiesterase, an oligodendrocyte-Schwann cell and myelin-associated enzyme of the nervous system. *Crit Rev Neurobiol* 4: 235-301. PubMed: 2537684.
  54. Yoshino JE, Dinneen MP, Sprinkle TJ, DeVries GH (1985) Localization of 2',3'-cyclic nucleotide 3'-phosphodiesterase on cultured Schwann cells. *Brain Res* 325: 199-203. doi:10.1016/0006-8993(85)90316-6. PubMed: 2983827.
  55. Lee MK, Cleveland DW (1996) Neuronal intermediate filaments. *Annu Rev Neurosci* 19: 167-217. doi:10.1146/annurev.ne.19.030196.001155. PubMed: 8833441.
  56. Frye CA, Walf AA (2009) Depression-like behavior of aged male and female mice is ameliorated with administration of testosterone or its metabolites. *Physiol Behav* 97: 266-269. doi:10.1016/j.physbeh.2009.02.022. PubMed: 19254734.
  57. Frye CA, Edinger KL, Lephart ED, Walf AA (2010) 3 $\alpha$ -androstenediol, but not testosterone, attenuates age-related decrements in cognitive, anxiety, and depressive behavior of male rats. *Front Aging Neurosci* 2: 15. PubMed: 20552051.
  58. Roglio I, Bianchi R, Camozzi F, Carozzi V, Cervellini I, Crippa D, Lauria G, Cavaletti G, Melcangi RC (2009) Docetaxel-induced peripheral neuropathy: protective effects of dihydroprogesterone and progesterone in an experimental model. *J Peripher Nerv Syst* 14: 347-355.
  59. Postma TJ, Vermorken JB, Liefing AJ, Pinedo HM, Heimans JJ (1995) Paclitaxel-induced neuropathy. *Ann Oncol* 6: 489-494. PubMed: 7669713.
  60. Cavaletti G, Tredici G, Braga M, Tazzari S (1995) Experimental peripheral neuropathy induced in adult rats by repeated intraperitoneal administration of taxol. *Exp Neurol* 133: 64-72. doi:10.1006/exnr.1995.1008. PubMed: 7601264.
  61. Wang MS, Davis AA, Culver DG, Glass JD (2002) Wild mice are resistant to paclitaxel (taxol) neuropathy. *Ann Neurol* 52: 442-447. doi:10.1002/ana.10300. PubMed: 12325073.
  62. Fazio R, Quattrini A, Bolognesi A, Bordogna G, Villa E, Previtali S, Canal N, Nemni R (1999) Docetaxel neuropathy: a distal axonopathy. *Acta Neuropathol (Berl)* 98: 651-653. doi:10.1007/s004010051132. PubMed: 10603043.
  63. Boyette-Davis J, Xin W, Zhang H, Dougherty PM (2011) Intraepidermal nerve fiber loss corresponds to the development of taxol-induced hyperalgesia and can be prevented by treatment with minocycline. *Pain* 152: 308-313. doi:10.1016/j.pain.2010.10.030. PubMed: 21145656.
  64. Dougherty PM, Cata JP, Burton AW, Vu K, Weng HR (2007) Dysfunction in multiple primary afferent fiber subtypes revealed by quantitative sensory testing in patients with chronic vincristine-induced pain. *J Pain Symptom Manage* 33: 166-179. doi:10.1016/j.jpainsymman.2006.08.006. PubMed: 17280922.
  65. Tanner KD, Reichling DB, Levine JD (1998) Nociceptor hyper-responsiveness during vincristine-induced painful peripheral neuropathy in the rat. *J Neurosci* 18: 6480-6491. PubMed: 9698336.
  66. Cliffer KD, Siuciak JA, Carson SR, Radley HE, Park JS, Lewis DR, Zlotchenko E, Nguyen T, Garcia K, Tonra JR, Stambler N, Cedarbaum JM, Bodine SC, Lindsay RM, DiStefano PS (1998) Physiological characterization of taxol-induced large-fiber sensory neuropathy in the rat. *Ann Neurol* 43: 46-55. doi:10.1002/ana.410430111. PubMed: 9450768.
  67. Kriz J, Zhu Q, Julien JP, Padjen AL (2000) Electrophysiological properties of axons in mice lacking neurofilament subunit genes: disparity between conduction velocity and axon diameter in absence of NF-H. *Brain Res* 885: 32-44. doi:10.1016/S0006-8993(00)02899-7. PubMed: 11121527.
  68. Hoffman PN, Cleveland DW, Griffin JW, Landes PW, Cowan NJ, Price DL (1987) Neurofilament gene expression: a major determinant of axonal caliber. *Proc Natl Acad Sci U S A* 84: 3472-3476. doi:10.1073/pnas.84.10.3472. PubMed: 3472217.
  69. Hoffman PN, Griffin JW, Gold BG, Price DL (1985) Slowing of neurofilament transport and the radial growth of developing nerve fibers. *J Neurosci* 5: 2920-2929. PubMed: 2414416.
  70. Hoffman PN, Griffin JW, Price DL (1984) Control of axonal caliber by neurofilament transport. *J Cell Biol* 99: 705-714. doi:10.1083/jcb.99.2.705. PubMed: 6204997.
  71. Amato SF, Swart JM, Berg M, Wanebo HJ, Mehta SR, Chiles TC (1998) Transient stimulation of the c-Jun-NH2-terminal kinase/activator protein 1 pathway and inhibition of extracellular signal-regulated kinase are early effects in paclitaxel-mediated apoptosis in human B lymphoblasts. *Cancer Res* 58: 241-247. PubMed: 9443400.
  72. Yang JY, Hung MC (2011) Deciphering the role of forkhead transcription factors in cancer therapy. *Curr Drug Targets* 12: 1284-1290. doi:10.2174/138945011796150299. PubMed: 21443462.
  73. Peters CM, Jimenez-Andrade JM, Kuskowski MA, Ghilardi JR, Mantyh PW (2007) An evolving cellular pathology occurs in dorsal root ganglia, peripheral nerve and spinal cord following intravenous administration of paclitaxel in the rat. *Brain Res* 1168: 46-59. doi:10.1016/j.brainres.2007.06.066. PubMed: 17698044.
  74. Patte-Mensah C, Kibaly C, Mensah-Nyagan AG (2005) Substance P inhibits progesterone conversion to neuroactive metabolites in spinal sensory circuit: a potential component of nociception. *Proc Natl Acad*

- Sci U S A 102: 9044-9049. doi:10.1073/pnas.0502968102. PubMed: 15951421.
75. Patte-Mensah C, Penning TM, Mensah-Nyagan AG (2004) Anatomical and cellular localization of neuroactive 5 alpha/3 alpha-reduced steroid-synthesizing enzymes in the spinal cord. *J Comp Neurol* 477: 286-299. doi:10.1002/cne.20251. PubMed: 15305365.
76. Edinger KL, Frye CA (2005) Testosterone's anti-anxiety and analgesic effects may be due in part to actions of its 5alpha-reduced metabolites in the hippocampus. *Psychoneuroendocrinol* 30: 418-430. doi:10.1016/j.psyneuen.2004.11.001. PubMed: 15721054.
77. Brooks BP, Merry DE, Paulson HL, Lieberman AP, Kolson DL, Fischbeck KH (1998) A cell culture model for androgen effects in motor neurons. *J Neurochem* 70: 1054-1060. PubMed: 9489725.
78. Cerghet M, Skoff RP, Bessert D, Zhang Z, Mullins C, Ghandour MS (2006) Proliferation and death of oligodendrocytes and myelin proteins are differentially regulated in male and female rodents. *J Neurosci* 26: 1439-1447. doi:10.1523/JNEUROSCI.2219-05.2006. PubMed: 16452667.