

Role of tRNA amino acid-accepting end in aminoacylation and its quality control

Xiao-Long Zhou¹, Dao-Hai Du¹, Min Tan¹, Hui-Yan Lei¹, Liang-Liang Ruan¹, Gilbert Eriani² and En-Duo Wang^{1,*}

¹State Key Laboratory of Molecular Biology, Institute of Biochemistry and Cell Biology, Shanghai Institutes for Biological Sciences, The Chinese Academy of Sciences, Shanghai 200031, China and ²Architecture et Réactivité de l'ARN, Institut de Biologie Moléculaire et Cellulaire du CNRS, Université de Strasbourg, 67084 Strasbourg, France

Received May 20, 2011; Revised July 4, 2011; Accepted July 5, 2011

ABSTRACT

Aminoacyl-tRNA synthetases (aaRSs) are remarkable enzymes that are in charge of the accurate recognition and ligation of amino acids and tRNA molecules. The greatest difficulty in accurate aminoacylation appears to be in discriminating between highly similar amino acids. To reduce mischarging of tRNAs by non-cognate amino acids, aaRSs have evolved an editing activity in a second active site to cleave the incorrect aminoacyl-tRNAs. Editing occurs after translocation of the aminoacyl-CCA₇₆ end to the editing site, switching between a hairpin and a helical conformation for aminoacylation and editing. Here, we studied the consequence of nucleotide changes in the CCA₇₆ accepting end of tRNA^{Leu} during the aminoacylation and editing reactions. The analysis showed that the terminal A₇₆ is essential for both reactions, suggesting that critical interactions occur in the two catalytic sites. Substitutions of C₇₄ and C₇₅ selectively decreased aminoacylation keeping nearly unaffected editing. These mutations might favor the regular helical conformation required to reach the editing site. Mutating the editing domain residues that contribute to CCA₇₆ binding reduced the aminoacylation fidelity leading to cell-toxicity in the presence of non-cognate amino acids. Collectively, the data show how protein synthesis quality is controlled by the CCA₇₆ homogeneity of tRNAs.

INTRODUCTION

Protein synthesis is a central process in organisms from all three domains of life, providing the link between the

genetic information encoded in DNA and functional proteins. This process critically relies on the correct formation of aminoacyl-tRNA (aa-tRNA) by aminoacyl-tRNA synthetase (aaRS) to establish the genetic code through rigorous control of the two-step aminoacylation (1,2). The amino acid is first activated with ATP to synthesize the aa-AMP intermediate with the release of pyrophosphate; the amino acid moiety of the intermediate is subsequently transferred to the tRNA bearing the cognate nucleotide triplet (1). Mis-translation arising from disruption in the fidelity of these interactions has profound consequences (3). However, the presence of various types of amino acids and their analogs, and the fact that amino acids only differ in the side chain, greatly challenge the specificity of aaRS. About half of the aaRSs misactivate non-cognate amino acids (4). To solve this problem, the proofreading (editing) mechanism of aaRS has evolved to hydrolyze the mis-products. Editing can occur at the aa-AMP level (pre-transfer editing) and/or mischarged tRNA level (post-transfer editing), depending on the specific aaRS (4).

Leucyl-tRNA synthetase (LeuRS), isoleucyl-tRNA synthetase (IleRS) and valyl-tRNA synthetase (ValRS) belong to the class Ia of aaRSs, characterized by the connective peptide 1 (CP1) and 2 (CP2) in the Rossmann fold nucleotide binding domain where the synthetic active site is located (5). The CP1 domain of LeuRS is located 35 Å away from the Rossmann fold domain and is responsible for post-transfer editing (6,7). Recent studies have revealed that *Escherichia coli*, *Aquifex aeolicus* and human cytoplasmic LeuRSs (*EcLeuRS*, *AaLeuRS* and *hcLeuRS*) all employ three different pathways (tRNA-independent, tRNA-dependent pre-transfer editing and post-transfer editing), but in different proportions relative to the total editing activity, to remove non-cognate amino acids (8,9). Similar results were also found with *E. coli* IleRS and ValRS (10).

X-ray crystal structures of LeuRS and tRNA^{Leu} in aminoacylation and post-transfer editing states clearly

*To whom correspondence should be addressed. Tel: +86 21 5492 1241; Fax: +86 21 5492 1011; Email: edwang@sibs.ac.cn

revealed that the tRNA^{Leu} main body conformation is indistinguishable between the two states (6,11). However, the CCA₇₆-end shifts to the CP1 domain from the aminoacylation active site and is specifically recognized by several conserved residues, including Lys300, Tyr330, Arg344 and Leu327 (6,11). Therefore, these residues in CP1 collectively constitute the tRNA entrance pathway in post-transfer editing (6). *In vitro* studies have shown that the interaction between the Tyr330 and tRNA^{Leu} CCA₇₆-end is critical for both tRNA-dependent pre-transfer editing and post-transfer editing (8). However, the role of other crucial residues in editing and their significance *in vivo* are unclear. In addition, the contribution of the CCA₇₆-end to the aminoacylation and editing still remains elusive.

The CCA₇₆ sequence is conserved at the 3'-end of all mature tRNA molecules to function as the site of amino acid attachment (12). This sequence is gene encoded in *E. coli* and related bacteria or acquired in eukaryotes and maintained by stepwise nucleotide addition by the ubiquitous CCA-adding enzyme (tRNA nucleotidyltransferase, CCCase), which is an unusual RNA polymerase that does not use a nucleic acid template for nucleotide addition (13,14).

The CCA₇₆ sequence is a universal ligand during several critical steps of protein biosynthesis. It is successively recognized by aaRSs (1,12), elongation factor (EF-Tu) (15) and rRNA (16,17). In F. Crick's adaptor hypothesis, the CCA₇₆ sequence is the ultimate adaptor group that carries the amino acid to the decoding center of the ribosome, whereas at the other end of the molecule, the anticodon triplet is needed to fit with the codon triplet of the mRNA. Because all the tRNAs have to fit to the unique ribosome decoding center, evolution has selected a unique CCA₇₆ sequence shared by all tRNAs. Additional constraints for CCA₇₆ conservation result from the interactions with the EF-Tu and the 20 aaRSs. With the later enzymes, the CCA₇₆ acceptor end is the substrate of two successive reactions starting with the aminoacylation reaction then followed by a proofreading reaction catalyzed in a second catalytic site (6,11). A critical conformational change, resulting from the flexibility of the single-stranded CCA₇₆ sequence, is required to allow transition from the synthetic to the editing site (6,11).

The starting point of the present study was to clarify the role of the ubiquitous CCA₇₆-end in the leucine aminoacylation system. Therefore, we generated a series of CCA₇₆-end mutants and tested their impact on the aminoacylation and editing reactions catalyzed by *E. coli* LeuRS (*EcLeuRS*). Furthermore, mutants were generated in the CP1 editing domain of LeuRS in order to disrupt the interaction with the CCA₇₆-end during the editing process. The *in vivo* effect of the mutations was then analyzed in a LeuRS-deficient strain in stress conditions under the pressure of elevated concentrations of non-cognate amino acid. Combining *in vitro* and *in vivo* results of protein and tRNA mutagenesis showed how life can regulate protein synthesis and control the aminoacylation quality by using the universally conserved sequence.

MATERIALS AND METHODS

Materials

L-leucine, dithiothreitol, NTP, 5'-GMP, tetrasodium pyrophosphate, inorganic pyrophosphate, ATP, Tris-HCl, MgCl₂, NaCl, mouse anti-His₆ antibody and activated charcoal were purchased from Sigma (USA). [³H]leucine, [³H]isoleucine and [α -³²P]ATP were obtained from Amersham Biosciences (England). *Pfu* DNA polymerase, the DNA fragment rapid purification kit and a plasmid extraction kit were purchased from Biotech Company (China). KOD-plus mutagenesis kit was obtained from TOYOBO (Japan). T4 ligase and restriction endonucleases were obtained from MBI Fermentas. Phusion high-fidelity DNA polymerase was purchased from New England Biolabs (USA). Ni²⁺-NTA Superflow was purchased from Qiagen, Inc. (Germany). Polyethyleneimine cellulose plates were purchased from Merck (Germany). Pyrophosphatase was obtained from Roche Applied Science (China). Pyrobest DNA polymerase and the dNTP mixture were obtained from Takara (Japan). Oligonucleotide primers were synthesized by Invitrogen (China). *Escherichia coli* BL21 (DE3) cells were purchased from Stratagene (USA). The *E. coli* KL231 strain [*F*⁻, *leuS31(ts)*, *thyA6*, *rpsL120(strR)*, *deoC1*] was purchased from the *E. coli* genetic stock center (CGSC, Yale University, USA) (18). T7 RNA polymerase was purified from an overproduction strain in our laboratory (19).

Gene cloning and mutagenesis

The plasmid pET30a-*EcleuS* encoding *EcLeuRS* with N-terminal His₆-tag previously constructed by our lab (8) was used as the template to construct genes encoding various *EcLeuRS* mutants using KOD-plus mutagenesis kit for the *in vitro* activity measurements. *EcleuS* was amplified, cleaved by NcoI and HindIII and cloned into pTrc99a (pre-cleaved by NcoI and HindIII) to produce pTrc99a-*EcleuS*. The 18 nucleotides encoding His₆-tag at the N-terminus were incorporated into the forward primer during the construction of pTrc99a-*EcleuS*. The pTrc99a-*EcleuS* construct was used as the template to construct genes encoding various *EcLeuRS* mutants using the KOD-plus mutagenesis kit for the *in vivo* complementation assays in the KL231 strain.

In vitro transcription of tRNAs

Six DNA fragments covering the T7 promoter and tRNA gene double strands were synthesized by Invitrogen, phosphorylated and ligated into pUC19 (pre-cleaved by EcoRI and BamHI) to construct the plasmid pUC19-tRNA. To prepare the wild-type (WT) tRNA^{Leu}_{GAG} transcript, the forward primer (5'-ctgcagtaata *cgactactatagccgaggtggtgg*-3', with T7 promoter sequence in italics) and the reverse primer (5'-tggtaccgaggacgg gactgaaccgtaagccctattg-3') were synthesized to amplify the T7 promoter and the gene encoding tRNA^{Leu}_{GAG} using the Phusion high-fidelity DNA polymerase, which does not add an additional adenosine to the PCR product. The PCR product was separated on a 2% agarose gel,

extracted by phenol/chloroform and precipitated in the presence of cold ethanol and 0.3 M NaAc (pH 5.2). The T7 *in vitro* transcription was carried out at 37°C in a 100 μ l reaction mixture containing 40 mM Tris-HCl (pH 8.0), 22 mM MgCl₂, 1 mM spermidine, 5 mM DTT, 0.5% Triton X-100, 60 ng μ l⁻¹ tDNA template, 5 mM ATP, 5 mM CTP, 5 mM GTP, 5 mM UTP, 0.8 U μ l⁻¹ ribonuclease inhibitor, 20 mM GMP and 500 U μ l⁻¹ T7 RNA polymerase. One hour later, 2 U of pyrophosphatase was added to remove the pyrophosphate for 30 min, and then 5 U of DNase I (RNase I free) was added and incubated for 1 h to digest the transcription template. The transcript was then loaded into a 15% PAGE-8M Urea gel of 1-mm thickness and 40 cm length, and the gel was run at constant 25 W for 10 h to carefully remove any non-specific bands. The tRNA was cut from the gel and eluted with 0.5 M NaAc (pH 5.2) at room temperature, ethanol precipitated at -20°C after phenol/chloroform extraction two times and dissolved in 5 mM MgCl₂. The tRNA was denatured at 80°C for 5 min and slowly cooled down to 30°C. The tRNA^{Leu} mutants were prepared by the same procedure, except that the different reverse primer was synthesized and used in preparing the DNA template.

Protein expression and purification

The WT LeuRS (*EcLeuRS*) and its mutants were produced by transformation of *E. coli* BL21 (DE3) cells with the corresponding plasmids. A single colony from each of the transformants was chosen and cultured in 500 ml of 2 \times YT medium at 37°C. When the cells were grown to mid-log phase ($A_{600} = 0.6$), IPTG (isopropyl-1-thio- β -D-galactopyranoside) was added to a final concentration of 0.2 mM, and cultivation continued for 6 h at 22°C. The proteins were purified by Ni²⁺-NTA chromatography, as described previously (8).

Aminoacylation, mis-aminoacylation and deacylation

The kinetic constants of *EcLeuRS* for WT tRNA and its variants were tested in a reaction mixture containing 100 mM Tris-HCl (pH 7.8), 30 mM KCl, 12 mM MgCl₂, 0.5 mM dithiothreitol, 4 mM ATP, 40 μ M [³H]leucine, 0.25–12 μ M *EctRNA*^{Leu} and 5 nM *EcLeuRS* at 37°C. Mis-aminoacylation assays were carried out at 37°C in a reaction mixture containing 100 mM Tris-HCl (pH 7.8), 30 mM KCl, 12 mM MgCl₂, 0.5 mM dithiothreitol, 4 mM ATP, 5 μ M *EctRNA*^{Leu}, 40 μ M [³H]isoleucine and 1 μ M *EcLeuRS*. Preparation of mischarged Ile-tRNA^{Leu} was carried out in a similar system with mis-aminoacylation, except that 20 μ M *EctRNA*^{Leu} and 1 μ M *EcLeuRS*-T252E were used (20). Hydrolytic editing assays of *EcLeuRS* or its mutants were performed at 37°C in 100 mM Tris-HCl (pH 7.5), 30 mM KCl, 12 mM MgCl₂, 0.5 mM dithiothreitol and 1 μ M [³H]Ile-tRNA^{Leu}, and the reactions were initiated with 5 nM enzyme. Nine microliter aliquots of the reaction solution were added to Whatman filter pads and quenched with cold 5% trichloroacetic acid (TCA) at various time intervals. The pads were washed three times for 15 min each with cold 5% TCA and then three times for 10 min each with

100% ethanol. The pads were dried under a heat lamp. The radioactivities of the precipitates were quantified with a scintillation counter (Beckman Coulter).

Charging plateau measurements

Charging plateau measurements of transcripts were performed in 40 μ l reaction mixtures each containing 100 mM Tris-HCl (pH 7.8), 30 mM KCl, 12 mM MgCl₂, 0.5 mM dithiothreitol, 4 mM ATP, 20 μ M [³H]leucine (15 Ci/mmol), 1 μ M tRNA (ΔA_{76} , $\Delta C_{75}A_{76}$, $\Delta C_{74}C_{75}A_{76}$, $\Delta A_{76}/C_{75}A$, $\Delta A_{76}/C_{75}U$ and $\Delta A_{76}/C_{75}G$) and 100 nM *EcLeuRS* at 37°C. Aliquots of 9 μ l of the reaction solution were removed at various time intervals, quenched on Whatman filter pads and processed as mentioned in the procedures above. The blank experiment without tRNA^{Leu} was performed under the same conditions. In another control experiment, RNase I (final 0.25 U/ μ l) was added to the aliquot at various time intervals to digest the aminoacylation product and incubated at 37°C for another 5 min. The plateau value of WT tRNA^{Leu} was >1400 pmol/A₂₆₀.

AMP formation

In the editing Nva reaction of *EcLeuRS*, AMP formation was measured in a reaction mixture containing 100 mM Tris-HCl (pH 7.8), 30 mM KCl, 12 mM MgCl₂, 5 mM dithiothreitol, 5 U ml⁻¹ pyrophosphatase, 3 mM ATP, 20 nM [α -³²P]ATP and 15 mM Norvaline (Nva) in the presence or absence of 5 μ M tRNA^{Leu} or its variants. The reaction was initiated by the addition of 0.2 μ M *EcLeuRS* or its mutants and incubated at 37°C. Aliquots (1.5 μ l) were quenched in 6 μ l of 200 mM NaAc (pH 5.0). The quenched aliquots (1.5 μ l each) were spotted in duplicate on polyethyleneimine cellulose plates pre-washed with water. Separation of Nva-[³²P]AMP, [³²P]AMP and [³²P]ATP was performed by developing thin layer chromatography (TLC) on polyethyleneimine cellulose plates in 0.1 M NH₄Ac and 5% acetic acid. The plates were visualized by phosphorimaging, and the data were analyzed using Multi Gauge Version 3.0 software (FUJIFILM). The gray densities of [³²P]AMP spots were compared with the gray density of known [³²P]ATP concentrations. Rate constants were obtained from graphs of [³²P]AMP formation plotted against time (8).

KL231 complementation assay

Competent KL231 cells were electro-transformed with various plasmids, including pTrc99a (negative control), pTrc99a-*EcleuS*, pTrc99a-*EcleuS*-L327G, -L327R, -K300E, -Y330D, -R344D, -K300E/Y330D, -K300E/R344D, -Y330D/R344D and -K300E/Y330D/R344D. Transformants were grown on solid LB plates supplemented with 100 μ g ml⁻¹ ampicillin and 200 μ g ml⁻¹ thymine at 30°C. A single colony was selected and grown in liquid LB medium supplemented with 100 μ g ml⁻¹ ampicillin, 200 μ g ml⁻¹ thymine and 0.1 mM IPTG at 30°C. The cells harboring these plasmids were diluted and adjusted to the same A_{600} value of 0.2. To test the effect of non-cognate Nva on cell growth, 5 μ l of the diluted mixture were dropped on a solid LB plate

(supplemented with 100 $\mu\text{g ml}^{-1}$ ampicillin, 200 $\mu\text{g ml}^{-1}$ thymine and 0.1 mM IPTG) containing 0, 5, 10, 20, 50, 100 or 200 mM Nva, and the plates were incubated at 42°C to observe the growth of cells.

RESULTS

Mutation of the CCA₇₆-end impacts the tRNA aminoacylation reaction

All tRNA^{Leu} isoacceptors from various species contain the absolutely conserved A₇₃ (12), which functions as the discriminator during recognition by LeuRSs in different species (12,21). However, the effect of the CCA₇₆-end on tRNA aminoacylation has not been reported. Here, each nucleotide of the CCA₇₆-end of tRNA^{Leu}_{GAG} was individually mutated to one of the three other nucleotides (Figure 1A), and the aminoacylation kinetic constants of *Ec*LeuRS for these tRNA^{Leu}s were assayed. In aminoacylation of the tRNA^{Leu} A76U and A76C mutants, the catalytic efficiency of *Ec*LeuRS was obviously decreased (155.8 and 87.2 s⁻¹ mM⁻¹, respectively) as compared with that of the WT tRNA transcript (923.1 s⁻¹ mM⁻¹). The amino acid-accepting activity of

the A76G mutant was too low to accurately measure the kinetics. Similarly, the three C₇₅ mutants reduced the amino acid-accepting activity to 2–46% of the native tRNA^{Leu}, with the C75G mutant showing the greatest decrease (14.6 s⁻¹ mM⁻¹). Additionally, the three tRNA variants from C₇₄ showed obvious decreases in amino acid-accepting activity; their catalytic constants for *Ec*LeuRS ranged from 8% (C74G, 69.4 s⁻¹ mM⁻¹), 41% (C74U, 381.8 s⁻¹ mM⁻¹) to 50% (C74A, 460.3 s⁻¹ mM⁻¹) of that with the WT tRNA^{Leu} (Table 1). All the decreases were mainly due to a reduced k_{cat} value of *Ec*LeuRS and/or changed K_m for these tRNA^{Leu} mutants. These results clearly showed that the CCA₇₆-end mutations negatively impacted the aminoacylation of these mutants to various levels, and the presence of guanosine at each of the 3-nt positions resulted in the lowest level of tRNA aminoacylation.

LeuRS can aminoacylate tRNA^{Leu} lacking the terminal adenosine

Besides point mutations, we also performed progressive deletions from the CCA₇₆-end of tRNA in order to obtain the following three mutants: Δ A₇₆ (ending




Figure 1. Secondary structures of tRNA^{Leu}s and location of the mutations. Cloverleaf structures of tRNA^{Leu}_{GAG} (A) and tRNA^{Leu}_{CAG} (B) used in the present work. The point and deletion mutations on the tRNA^{Leu}_{GAG} are shown.

Table 1. Aminoacylation kinetics of transcripts of tRNA^{Leu}_{GAG} by *Ec*LeuRS

tRNA	K_m (μM)	k_{cat} (s ⁻¹)	k_{cat}/K_m (s ⁻¹ mM ⁻¹)	k_{cat}/K_m (relative)
tRNA ^{Leu} _{GAG}	1.95 ± 0.23	1.80 ± 0.27	923.1	1
A76U	1.99 ± 0.30	0.31 ± 0.04	155.8	0.17
A76G	–	<<0.002	–	–
A76C	3.90 ± 0.48	0.34 ± 0.03	87.2	0.09
C75A	0.38 ± 0.05	0.16 ± 0.03	421.1	0.46
C75U	1.18 ± 0.17	0.10 ± 0.02	84.7	0.09
C75G	1.78 ± 0.22	0.026 ± 0.004	14.6	0.02
C74A	0.63 ± 0.05	0.29 ± 0.04	460.3	0.50
C74U	0.55 ± 0.07	0.21 ± 0.03	381.8	0.41
C74G	0.36 ± 0.05	0.025 ± 0.002	69.4	0.08

with C₇₅), $\Delta C_{75}A_{76}$ (ending with C₇₄) and $\Delta C_{74}C_{75}A_{76}$ (ending with A₇₃) (Figure 1A). In initial charging plateau measurements for accepting activity with a large excess of 1 μ M enzyme, it was surprising to observe that LeuRS could charge ΔA_{76} mutant obviously. As it has been reported that some aaRSs can ligate their cognate amino acids to themselves without tRNA (by self-aminoacylation occurring at high enzyme concentration and excess of adenylate formation) (22–24), we tested the self-aminoacylation of *Ec*LeuRS at different concentrations without tRNA. We found that self-aminoacylation of *Ec*LeuRS was negligible at low concentrations (<300 nM). However, *Ec*LeuRS at higher concentrations >500 nM could be self-aminoacylated and was resistant to RNase I (Supplementary Figure S1). Therefore, we tested the ΔA_{76} charging plateau value by using 100 nM *Ec*LeuRS and found that 70% of ΔA_{76} could be charged with ³H-Leu. In the same conditions, no detectable labeled product could be measured at the absence of tRNA (Figure 2). To further confirm that, the tRNA but not the enzyme itself was ligated with ³H-Leu, we treated the aminoacylation product in the presence of RNase I, and the values dropped down to the same level with that in the absence of tRNA (Figure 2). These results confirmed that LeuRS could aminoacylate tRNA^{Leu} with the terminal A₇₆ deleted. However, reliable aminoacylation kinetics could not be measured because of the very low k_{cat} . However, the other two tRNA mutants, $\Delta C_{75}A_{76}$ and $\Delta C_{74}C_{75}A_{76}$, could not be aminoacylated at 100 nM or higher concentrations of *Ec*LeuRS (data not shown).

To further analyze the ΔA_{76} mutant, we mutated the terminal C₇₅ to the three other nucleotides in order to identify potential interactions that could be more productive in aminoacylation. The resulting mutants, $\Delta A_{76}/C_{75}A$, $\Delta A_{76}/C_{75}U$ and $\Delta A_{76}/C_{75}G$, could be




Figure 2. Aminoacylation of tRNA^{Leu}- ΔA_{76} , $-\Delta A_{76}/C_{75}A$, $-\Delta A_{76}/C_{75}U$ or $-\Delta A_{76}/C_{75}G$ by *Ec*LeuRS. Charging plateaus of *EctRNA*^{Leu}- ΔA_{76} (filled diamond), $-\Delta A_{76}/C_{75}A$ (filled square), $-\Delta A_{76}/C_{75}U$ (filled triangle) or $-\Delta A_{76}/C_{75}G$ (filled inverted triangle) were measured as described in the Materials and methods section. The control (unfilled circle) was performed without tRNA. The charging plateaus were determined using GraphPad Prism software. According to the plateau values (to calculate the amount of tRNA^{Leu} charged leucine) and total tRNA^{Leu} used in the reaction, *EctRNA*^{Leu}- ΔA_{76} , $-\Delta A_{76}/C_{75}A$, $-\Delta A_{76}/C_{75}U$ and $-\Delta A_{76}/C_{75}G$ are aminoacylated to 70, 70, 61 and 35%, respectively. Digestion of aminoacyl-*EctRNA*^{Leu}- ΔA_{76} by RNase I digestion (unfilled triangle) and no product accumulation in control excluded the self-aminoacylation of the enzyme.

aminoacylated by *Ec*LeuRS to plateau levels of 70, 61 and 35%, respectively (Figure 2). Therefore, more productive interactions were not induced by these nucleotides, and here again, the mutant harboring a guanosine mutation was the least active compared to those with the other nucleotides.

Role of CCA₇₆-end in total editing

Usually, more than one ATP molecule is consumed by an aaRS in the presence of a non-cognate amino acid due to repetitive cycles of synthesis–hydrolysis of the non-cognate products. The excess of ATP consumption can be determined by measuring the release of AMP in the TLC assay (25). In the presence of tRNA, the TLC assay measures the global editing activity, including the tRNA-independent and tRNA-dependent pre-transfer editing in addition to the post-transfer editing (8,26). In the absence of tRNA, the assay measures the AMP produced from the sole tRNA-independent pre-transfer editing activity (26).

In the presence of non-cognate Nva, the observed rate constant (k_{obs}) of WT LeuRS for AMP formation without tRNA in the editing reaction was $0.56 \pm 0.07 s^{-1}$, accounting for 13% of that in the presence of tRNA transcript ($4.42 \pm 0.64 s^{-1}$) (Table 2). The tRNA transcript without the modified bases was tested to determine if it was as efficient as the native tRNA in stimulating the editing activity. The k_{obs} for AMP formation in the presence of *EctRNA*^{Leu}_{GAG} purified from the overproducing *E. coli* strain (27) was $5.59 \pm 0.76 s^{-1}$. This value was only slightly larger than that in the presence of transcript ($4.42 \pm 0.64 s^{-1}$), indicating that the modified bases in *EctRNA*^{Leu} did not play a critical role in stimulating editing activity (Table 2).

Different variants of the CCA₇₆-end were also tested for their abilities to stimulate AMP formation in editing reactions with *Ec*LeuRS. The AMP formation rates for A76U, A76G and A76C mutants dropped down to a level similar to that without tRNA, indicating only tRNA-independent pre-transfer editing remained and the tRNA-dependent

Table 2. k_{obs} of AMP formation by *Ec*LeuRS in the presence of Nva and tRNA^{Leu}_{GAG} or its variants

tRNA	k_{obs} (s^{-1})	Relative k_{obs}
No tRNA ^{Leu} _{GAG}	0.56 ± 0.07	0.13
tRNA ^{Leu} _{GAG} (<i>in vivo</i>) ^a	5.59 ± 0.76	1.27
tRNA ^{Leu} _{GAG} (transcript) ^b	4.42 ± 0.64	1
A76U	0.24 ± 0.04	0.05
A76G	0.20 ± 0.04	0.05
A76C	0.48 ± 0.06	0.11
ΔA_{76}	0.71 ± 0.08	0.16
C75A	8.34 ± 1.03	1.89
C75U	6.26 ± 0.81	1.42
C75G	0.64 ± 0.07	0.15
C74A	7.48 ± 0.98	1.69
C74U	8.87 ± 1.34	2.01
C74G	2.65 ± 0.32	0.60

^atRNA^{Leu}_{GAG} obtained from the overproducing *E. coli* strain (27).

^btRNA^{Leu}_{GAG} obtained by T7 *in vitro* transcription.




Figure 3. Misacylation of tRNA-A₇₆ mutants with Ile. (A and B) WT tRNA^{Leu} (filled circle) and mutated derivatives A76U (unfilled circle), A76G (filled inverted triangle) or A76C (unfilled triangle) and ΔA₇₆ (unfilled inverted triangle) were mischarged with non-cognate Ile.

editing was abolished (Table 2). As a consequence of this loss of editing ability, the two A₇₆-mutants of tRNA^{Leu} (A76U and A76C) were obviously mischarged with Ile (Figure 3A). The decreased formation of Ile-tRNA^{Leu} of A76G may be due to its deficiency in accepting the amino acid. Among the other six variants of tRNA^{Leu}, only C75G with a k_{obs} value of $0.64 \pm 0.07 \text{ s}^{-1}$ had tRNA-independent pre-transfer editing and lost the tRNA-dependent editing, indicating that C75G could not play a role in the tRNA-dependent editing; C74G with a k_{obs} value of $2.65 \pm 0.32 \text{ s}^{-1}$ decreased the tRNA-dependent editing to nearly half of that of the WT tRNA^{Leu}; the four mutants of tRNA^{Leu}, C75A, C75U, C74A and C74U stimulated more AMP formation than WT tRNA^{Leu}, with the k_{obs} values of 8.34 ± 1.03 , 6.26 ± 0.81 , 7.48 ± 0.98 and $8.87 \pm 1.34 \text{ s}^{-1}$, respectively (Table 2), indicating that these mutations did not affect the tRNA-dependent editing but rather stimulated it. The six variants of nucleotides 74 and 75 did not form any detectable Ile-tRNA^{Leu} in the misacylation assay (data not shown). Additionally, in the presence of the ΔA₇₆ mutant, *EcLeuRS* had comparable editing activity with that in the absence of tRNA (Table 2). Correspondingly, *EcLeuRS* catalyzed the synthesis of an obvious amount of mischarged ΔA₇₆ mutant (Ile-tRNA^{Leu}-ΔA₇₆) (Figure 3B). These results clearly showed that the terminal A₇₆ is a critical element in the tRNA-dependent editing activity of *EcLeuRS*, and the nucleotides C₇₄ and C₇₅ crucial for aminoacylation do not play an indispensable role to the tRNA-dependent editing of *EcLeuRS*.

Role of CCA₇₆-end in post-transfer editing

It is now universally accepted that after aminoacylation, the CCA₇₆-end of the tRNA is translocated ~35 Å from the aminoacylation active site to the editing active site embedded in the CP1 domain (6). Both tRNA CCA₇₆-end and mischarged amino acid moieties are recognized by the active site of post-transfer editing within the CP1 domain of *LeuRS*. To test the contribution of the CCA₇₆-end to the post-transfer editing, we first mischarged 10 variants of the tRNA with non-cognate ³H-Ile using an editing-deficient *LeuRS*-T252E mutant (20).

The deacylation of the mischarged Ile-tRNA^{Leu} by WT *LeuRS* was then measured. The mutants of tRNA^{Leu} could be mischarged with Ile by the *LeuRS*-T252E mutant. However the three mis-acylated mutants of A₇₆ (A76C, A76G and A76U) were resistant to deacylation catalyzed by *LeuRS* (Figure 4A), indicating the crucial role of this nucleotide in post-transfer editing. This result was also consistent with their inability to stimulate total editing in the TLC assay (Table 2). In addition, Ile-tRNA^{Leu}-ΔA₇₆ could not be hydrolyzed by the post-transfer editing (Figure 4A). This is consistent with the fact that the A₇₆ base is specifically recognized by the main chain carbonyl- and amino-groups of *EcLeuRS* Leu327 in the Nva2AA-containing structure (analog of the post-transfer editing substrate) (28) and *Thermus thermophilus* *LeuRS*-tRNA^{Leu} structure (in post-transfer editing conformation) (6). Obviously, changing the terminal base affects the proper positioning of the CCA₇₆-end into the CP1 domain and thus blocks the hydrolytic editing reaction.

In general, mutations of nucleotides 75 and 74 did not influence the tRNA deacylation properties, except when the guanosine mutations C75G and C74G were introduced (Figure 4B and C). The result suggested that guanine residues at either position of CCA₇₆-end may prevent proper interaction with the enzyme and/or tRNA end translocation.

The tRNA entrance pathway critically contributes to quality control

According to the crystal structure, the CP1 editing domain binds only with CCA₇₆-end residues without any interaction with the main tRNA body. In detail, the equivalent side chains in *EcLeuRS* interacting specifically with the C₇₄ base and the phosphates of C₇₅ and A₇₆ are Lys300, Arg344 and Tyr330, respectively (6). Additionally, the main chain carbonyl- and amino-group of Leu327 interact with the A₇₆ base (Figure 5A). These amino acid residues are absolutely conserved in prokaryotic *LeuRS*s and thus constitute an entrance pathway to orient the aminoacylated CCA₇₆-arm into the editing active site. Previous studies have demonstrated the essential role of Tyr330 of *EcLeuRS* in tRNA-dependent




Figure 4. Effect of the CCA₇₆ mutations on deacylation of mischarged tRNA^{Leu}. Mischarged tRNA mutants from A₇₆, C₇₅ or C₇₄ were prepared by editing-deficient *EcLeuRS*-T252E (20). Mischarged tRNA- Δ A₇₆ mutant was obtained with WT *EcLeuRS*. Deacylation of mischarged WT tRNA^{Leu} was indicated (unfilled circle). The deacylation assay of mischarged Δ A₇₆ (unfilled square), A76U (filled inverted triangle), A76G (unfilled triangle), A76C (filled square) in (A), C75A (filled inverted triangle), C75U (unfilled triangle), C75G (filled square) in (B) or C74A (filled inverted triangle), C74U (unfilled triangle), C74G (filled square) in (C) was performed at 37°C with 1 μ M [³H]Ile-tRNA^{Leu} and 5 nM *EcLeuRS*. The spontaneous hydrolysis of each mischarged tRNA (control) was carried out without enzyme. For clarity, a single representative control is shown (filled circle).




Figure 5. The tRNA entrance pathway contributes critically to the quality control. (A) View of tRNA^{Leu} CCA₇₆-end bound to the CPI domain (gray) (6), showing the binding pattern between C₇₄ (yellow), C₇₅ (cyan), A₇₆ (blue) and the key amino acids (green) in the CPI editing domain (upper). The tRNA main body before nucleotide A₇₃ (orange) and LeuRS main body except for the editing domain were omitted for clarity. The lower panel shows the closer protein-CCA₇₆ interaction network. Distances between the amino-acid residues here studied and the nucleotides are shown. (B) Mischarging of tRNA^{Leu}_{GAG} with non-cognate Ile by WT *EcLeuRS* (filled circle), -Y330D (unfilled circle), -K300E/Y330D (filled inverted triangle), -K300E/R344D (unfilled triangle), -Y330D/R344D (filled square), -K300E/Y330D/R344D (unfilled square) and -R344D (filled diamond) with significant misaminoacylation ability. Mutants that did not accumulate Ile-tRNA^{Leu} are not shown. (C) Deacylation curves of Ile-tRNA^{Leu} by WT *EcLeuRS* (filled circle), -L327G (unfilled circle), -L327R (filled inverted triangle), -K300E/R344D (unfilled triangle), -K300E (filled square) and -R344D (filled diamond). Editing-defective mutants (*EcLeuRS*-Y330D, -K300E/Y330D, -Y330D/R344D and -K300E/Y330D/R344D) are not shown for clarity. Spontaneous hydrolysis without enzyme addition was carried out as a control (unfilled square) as indicated.

editing (8). However, other sites (Leu327, Lys300 and Arg344) and their functions *in vivo* have not been determined. We studied here the roles of residues Leu327, Lys300, Tyr330 and Arg344 by mutating them individually or in combination. The *in vitro* and *in vivo* assays were then performed in order to analyze the effects of these mutations.

The following single, double and triple mutants were constructed in *EcLeuRS* to form *EcLeuRS*-L327G, -L327R, -K300E, -Y330D, -R344D, -K300E/Y330D, -K300E/R344D, -Y330D/R344D and -K300E/Y330D/R344D. All these mutants displayed intact synthetic activities (Leu activation and aminoacylation) as expected from their location in the discrete CP1 editing domain (data not shown). However, *EcLeuRS*-Y330D, -K300E/Y330D, -Y330D/R344D and the triple mutant were completely defective in post-transfer editing (data not shown), and they produced significant amounts of Ile-tRNA^{Leu} (Figure 5B). The other mutants were impaired in post-transfer editing to different extents (Figure 5C), and only *EcLeuRS*-R344D and *EcLeuRS*-K300E/R344D synthesized mischarged tRNA^{Leu}, but obviously at lower levels than that with *EcLeuRS*-Y330D and the derived double and triple mutants (Figure 5B). In the TLC assay, *EcLeuRS*-Y330D, its derived double and triple mutants and *EcLeuRS*-K300E/R344D only retained the tRNA-independent pre-transfer editing activity (Table 3). Altogether, the data showed that the Tyr330 residue plays a major role in tRNA-dependent editing as previously demonstrated (8). The negative effect of Y330D on editing was dominant, and the derived multiple mutants exhibited similar defects. Only the combination mutant *EcLeuRS*-K300E/R344D could reach the same level of tRNA-independent editing of the single *EcLeuRS*-Y330D mutation. Other individual mutants showed more limited decreases in post-transfer editing activity, inducing no measurable tRNA mischarging (data not shown). These *in vitro* data confirmed that mutating the CCA₇₆ entrance pathway of the CP1 domain has a negative effect on the accuracy of tRNA^{Leu} aminoacylation.

To evaluate if these losses of accuracy could affect the fidelity of protein synthesis *in vivo*, complementation

Table 3. k_{obs} of AMP formation by *EcLeuRS* or mutated derivatives in the presence of Nva

<i>EcLeuRS</i>	tRNA ^{Leu} _{GAG} (transcript)	k_{obs} (s ⁻¹)	Relative k_{obs}
WT	+	4.42 ± 0.64	1
WT	-	0.56 ± 0.07	0.13
-L327G	+	1.86 ± 0.26	0.42
-L327R	+	3.60 ± 0.45	0.81
-K300E	+	2.54 ± 0.41	0.57
-Y330D	+	0.59 ± 0.06	0.14
-R344D	+	1.26 ± 0.19	0.29
-K300E/Y330D	+	0.46 ± 0.03	0.10
-K300E/R344D	+	0.82 ± 0.07	0.19
-Y330D/R344D	+	0.49 ± 0.06	0.11
-K300E/Y330D/R344D	+	0.59 ± 0.06	0.14

assays using the *leuS* temperature-sensitive strain KL231 were performed (18). The strain KL231 was transformed with various plasmids encoding *EcLeuRS* and different mutants. At 42°C, all the transformants grew on medium supplemented with 100 µg ml⁻¹ ampicillin and 200 µg ml⁻¹ thymine (Figure 6), and the mutants were well expressed at a level similar to that of WT *EcLeuRS* (Supplementary Figure S2). The growth was then observed at 42°C in the presence of increasing concentrations of Nva in the growth medium. No growth difference could be observed below 20 mM of Nva. However, in the presence of 50 mM Nva, the strains harboring the genes-encoding *EcLeuRS*-Y330D, -K300E/Y330D, -K300E/R344D, -Y330D/R344D and -K300E/Y330D/R344D grew slowly or were inhibited completely, especially those containing the last double and triple mutants (Figure 6). With a 100 mM of Nva no strains containing mutants were able to grow, except for the strain expressing WT enzyme which only grew at a very slow rate. All strains died in the presence of 200 mM Nva.

Altogether, the *in vivo* data are consistent with the *in vitro* results, demonstrating that perturbing the tRNA CCA₇₆-end entry pathway in the CP1 domain has a significantly negative effect on the fidelity of protein biosynthesis.




Figure 6. *In vivo* toxicity resulting from mutations in the CP1 tRNA entrance pathway of *EcLeuRS*. The complementation assay was performed using *E. coli* thermosensitive strain KL231 at 42°C, on solid LB plates supplemented with 100 µg ml⁻¹ ampicillin, 200 µg ml⁻¹ thymine, 0.1 mM IPTG and increasing concentrations of non-cognate Nva. KL231 was transformed with mutated copies of *LeuRS*-encoding genes, and growth was compared with the negative control (empty pTrc99a vector) and positive control (WT *EcLeuRS*). Expression of all *LeuRS*s proteins was controlled by Western blot (Supplementary Figure S2).

Table 4. k_{obs} of AMP formation in the presence of different transcripts of tRNA^{Leu} isoacceptors or mutants

tRNA	k_{obs} (s ⁻¹)	Relative k_{obs}
tRNA _{GAG} ^{Leu}	4.42 ± 0.64	1
tRNA _{GAG} ^{Leu} (G ₁ -C ₇₂)	3.75 ± 0.50	0.85
tRNA _{CAG} ^{Leu}	4.54 ± 0.36	1
tRNA _{CAG} ^{Leu} (G ₁ -U ₇₂)	6.10 ± 0.81	1.34

Different tRNA^{Leu} isoacceptors stimulate LeuRS editing with the same efficiency

In *E. coli*, there are five different tRNA^{Leu} isoacceptors with two types of first base pairs in its accepting stem (Figure 1). tRNA_{GAG}^{Leu} and tRNA_{UAA}^{Leu} harbor the wobble base pair (G₁-U₇₂); while tRNA_{CAG}^{Leu}, tRNA_{CAA}^{Leu} and tRNA_{UAG}^{Leu} contain the standard Watson-Crick one (G₁-C₇₂). It has been shown that *Ec*LeuRS has the same specificity and activity toward different isoacceptors in aminoacylation (29). We transcribed the *Ec*tRNA_{GAG}^{Leu} (with G₁-U₇₂) and tRNA_{CAG}^{Leu} (with G₁-C₇₂) to represent two types of tRNA^{Leu} and compared their capacity to stimulate editing by *Ec*LeuRS. The data showed that they stimulated *Ec*LeuRS editing activity with nearly identical efficiency (4.42 ± 0.64 and 4.54 ± 0.36 s⁻¹, respectively) (Table 4). To explore the possible role of the first base pair in *Ec*LeuRS editing, we mutated the U₇₂ to C₇₂ of tRNA_{GAG}^{Leu} or the C₇₂ to U₇₂ of tRNA_{CAG}^{Leu} to change the first base pair pattern. Measurements of AMP formation showed that the tRNA_{GAG}^{Leu}(G₁-C₇₂) had the editing k_{obs} of 3.75 ± 0.50 s⁻¹, while the tRNA_{CAG}^{Leu}(G₁-U₇₂) mutant stimulated the editing with k_{obs} of 6.10 ± 0.81 s⁻¹ (Table 4). The above results suggested that *Ec*LeuRS editing has no preference among WT isoacceptors with different first base-pairs. However, changing the first base-pair obviously led to distinct editing-stimulating capacities among isoacceptors. Therefore, the first base pair pattern may cooperate with other sequence differences between the two tRNA^{Leu}s to confer the same level of editing-stimulating capacity of the two isoacceptors of WT tRNA^{Leu}.

DISCUSSION

The CCA₇₆-end sequence is critical for both aminoacylation and editing reactions

The CCA₇₆ sequence is conserved at the 3'-end of all mature tRNA molecules. Although it is an indispensable prerequisite for a functional tRNA, very few organisms like *E. coli* encode the CCA₇₆ triplet in their tRNA genes. Most of the time, the CCA₇₆ tail is to be added post-transcriptionally in eukaryotes, archaea and many bacteria, where the tRNA genes do not encode the CCA₇₆ terminus (30). This sequence is acquired and maintained by step-wise nucleotide addition by the ubiquitous tRNA nucleotidyltransferase that synthesizes this specific triplet without a nucleic acid template. The CCA sequence plays key roles during several steps of protein

biosynthesis. During tRNA charging catalyzed by aaRSs, the 2' or 3' hydroxyl of ribose in A₇₆ is the group involved in the esterification reaction with the carboxyl of the amino acid. Although the CCA₇₆-end does not play a critical role in recognition between tRNA and aaRS, it is essential in terms of catalytic efficiency for the aminoacylation reaction (31–33). The universal CCA₇₆ sequence is also involved in EF-Tu binding and during peptide bond formation by interacting with specific nucleotides of the 23S rRNA (17). Little is known about the precise role of the CCA₇₆ nucleotides during the editing reaction catalyzed by aaRSs. It has been shown that only chargeable tRNA^{Val} mutants are able to stimulate the editing reaction of ValRS, suggesting that the enzyme required prior charging of tRNA (31,32,34). Although tRNA is not a strict prerequisite for the editing of both classes I and II aaRSs (26,35–37), its presence strongly stimulates editing (38). Remarkably, both classes of aaRSs use effectively the ability of the CCA₇₆-end of tRNA to switch between a hairpin and a helical conformation during the aminoacylation and editing reactions. Our previous investigations have shown that the interaction between the tRNA^{Leu} CCA₇₆-end and the CPI domain of LeuRS is critical for tRNA-dependent editing (post-transfer and pre-transfer editing) (8). However, the molecular basis of the phenomenon is not yet understood in detail. Here, we extended the investigations to the three CCA₇₆ nucleotides and assayed the effect of mutations on the tRNA-dependent pre-transfer editing and post-transfer editing.

We showed that the universally conserved A₇₆ is essential for tRNA^{Leu} to trigger both aminoacylation and editing reaction of LeuRS. The mutants of tRNA^{Leu} with any base substituted for A₇₆ showed significant decreases of aminoacylation properties and total loss of the ability to stimulate the editing activity and catalyze deacylation of pre-formed Ile-tRNA^{Leu}. As a consequence of this loss of editing activity, two of these mutants (A76U, A76C) were misacylated by Ile. Mutations of the other bases of the CCA₇₆-end had negative effects on the aminoacylation properties, ranging between 2- and 50-fold decreases of the catalytic efficiency of *Ec*LeuRS, with the strongest effect being observed when C₇₅ and C₇₄ were substituted by G₇₅ and G₇₄, respectively. Correspondingly, the mutant A76G was totally inactive, showing that the bulky guanosine substitutions were the least accepted in the synthetic active site. The three guanosine mutants also decreased the deacylation of the mischarged Ile-tRNA^{Leu}, but failed in misacylating tRNA^{Leu} with Ile. Comparison of the global editing activity of *Ec*LeuRS in the presence of the tRNA^{Leu} mutants revealed that four mutations (C75A, C75U, C74A and C74U) led to an increase in the editing up to 2-fold despite the fact that their accepting capacity with Leu were impaired during the aminoacylation reaction. *Ec*LeuRS deacylated these mischarged mutants of tRNA^{Leu} with Ile with unchanged ability, suggesting that these mutants of tRNA^{Leu} preferred binding at the editing active site rather than at the synthetic site. Previous studies have shown that both classes of aaRSs use effectively the ability of the CCA₇₆-end of tRNA to switch

between a hairpin and a helical conformation of aaRSs for aminoacylation and editing. In class I aaRSs, their cognate tRNA CCA₇₆-ends adopt a regular helical conformation into the editing site of aaRSs, and a distorted hairpin conformation into the synthetic site (6). A mirror image is observed in class II aaRSs, as already seen for tRNA binding and amino acid activation. The stimulation of the editing activity of C75A, C75U, C74A and C74U mutants strongly suggests that these mutations may favor the helical conformation, which is more suitable for the editing reaction. Moreover, the fact that these mutated tRNAs are poorly aminoacylated in the synthetic active site may favor stagnation of the non-cognate adenylate into the synthetic active site and its subsequent editing by the tRNA-dependent pre-transfer editing also measured in the global editing assay.

The CCA₇₆-end entrance pathway into editing domain of *Ec*LeuRS critically contributes to protein synthesis fidelity

Despite the importance of the interaction between CCA₇₆-end of tRNA and the CP1 domain of class Ia aaRSs revealed by *in vitro* methods, the corresponding *in vivo* evidence is generally lacking. In the present work, we developed an efficient system with the combination of non-cognate Nva and *leuS* temperature-sensitive *E. coli* strain to assess the significance of LeuRS tRNA-dependent editing *in vivo*. Our evaluation of the tRNA CCA₇₆-end entrance pathway using the *leuS* temperature-sensitive KL231 strain revealed the following aspects. (i) Without or at a low concentration of non-cognate amino acid (e.g. 20 mM Nva here), mutation of pivotal residues controlling protein biosynthesis fidelity had no obvious effect on the bacterial cell viability. The discrimination by EF-Tu and/or the ribosome may have prevented Nva mis-incorporation, or the *E. coli* cells could endure low level of Nva mis-incorporation at Leu codons; (ii) at extremely high concentrations of non-cognate amino acids, even the WT enzyme had no ability to prevent all the mis-activated amino acids from entering the newly synthesized polypeptide on the ribosome, and the mis-translation led to an obvious inhibitory effect on cell growth; (iii) our *in vitro* results were consistent with *in vivo* data, demonstrating the importance of conserved amino acid residues in aaRS and nucleotides in tRNA in protein synthesis quality control, especially under severe environmental stress.

Multiple steps to monitor and control the CCA acceptor sequence

As mentioned above, the trinucleotide CCA₇₆ sequence is present at the 3' terminus of all mature tRNAs. Despite this high conservation, transcripts of *E. coli* tRNA^{Val} with altered 3' termini are readily aminoacylated and can function in polypeptide synthesis (32). Accordingly, the present study performed on LeuRS shows that the aminoacylation reaction admits some sequence flexibility at the CCA₇₆ end. Several CCA-mutated tRNAs can be efficiently aminoacylated, and LeuRS can even aminoacylate a tRNA^{Leu} lacking the terminal adenosine, showing a remarkable plasticity of the synthetic site of

LeuRS and tRNA^{Leu} acceptor end. Several studies suggest that the aminoacylation of similar molecules may occur naturally *in vivo*. Despite the high fidelity of the CCA-adding enzyme, in some conditions, CCCase can add a wrong nucleotide to the 3'-end of the tRNA, leading to CCA-modified tRNAs (39). Moreover, mutations of CCCase may exist *in vivo* with relaxed active sites as those engineered *in vitro* that lead to nucleotide mis-incorporation at the end of tRNA (40). In some bacteria the CCA-adding activity is naturally split into a CC-adding enzyme and an A-adding enzyme (41). Therefore, a tRNA deprived of the terminal A₇₆ might temporarily exist *in vivo* and may be the substrate of LeuRS.

However, if under these exceptional circumstances aminoacylation of CCA-mutated tRNAs can occur, these tRNAs would have to face additional discrimination processes based on EF-Tu recognition (42) and interaction with ribosomal RNA during translation (43). Both processes would monitor and exclude these molecules according to different nucleotide recognition specificities (32).

In summary, the study highlighted the significant role of the conserved nucleotides from the CCA₇₆-end and the strong collaboration with amino acid residues of the CP1 editing domain during the aminoacylation and editing catalysis. The CCA₇₆ nucleotides are critical for both aminoacylation and editing; and in combination with the amino acids located in the entrance path of the editing domain, they critically contribute to the fidelity of protein synthesis. It is a remarkable example of a control mechanism that prevent damaged tRNAs from entering the protein synthesis based on the single discrimination of a triplet sequence that is universally found in all tRNAs.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

ACKNOWLEDGEMENTS

The authors thank Dr Franck Martin for the help during the experiment.

FUNDING

Natural Science Foundation of China (Nos. 30930022 and 31000355); Committee of Science and Technology in Shanghai (No. 09JC1415900); a Visiting Professorship for Senior International Scientists from the Chinese Academy of Sciences (No. 2009S2-19); Programme International de Coopération Scientifique from CNRS (Grant 3606). Funding for open access charge: Natural Science Foundation of China.

Conflict of interest statement. None declared.

REFERENCES

1. Ibba, M. and Söll, D. (2000) Aminoacyl-tRNA synthesis. *Annu. Rev. Biochem.*, **69**, 617–650.
2. Schimmel, P. (1987) Aminoacyl tRNA synthetases: general scheme of structure-function relationships in the polypeptides and recognition of transfer RNAs. *Annu. Rev. Biochem.*, **56**, 125–158.
3. Lee, J.W., Beebe, K., Nangle, L.A., Jang, J., Longo-Guess, C.M., Cook, S.A., Davisson, M.T., Sundberg, J.P., Schimmel, P. and Ackerman, S.L. (2006) Editing-defective tRNA synthetase causes protein misfolding and neurodegeneration. *Nature*, **443**, 50–55.
4. Ling, J., Reynolds, N. and Ibba, M. (2009) Aminoacyl-tRNA synthesis and translational quality control. *Annu. Rev. Microbiol.*, **63**, 61–78.
5. Zhou, X., Zhu, B. and Wang, E. (2008) The CP2 domain of leucyl-tRNA synthetase is crucial for amino acid activation and post-transfer editing. *J. Biol. Chem.*, **283**, 36608–36616.
6. Tukalo, M., Yaremchuk, A., Fukunaga, R., Yokoyama, S. and Cusack, S. (2005) The crystal structure of leucyl-tRNA synthetase complexed with tRNA^{Leu} in the post-transfer-editing conformation. *Nat. Struct. Mol. Biol.*, **12**, 923–930.
7. Chen, J., Guo, N., Li, T., Wang, E. and Wang, Y. (2000) CP1 domain in *Escherichia coli* leucyl-tRNA synthetase is crucial for its editing function. *Biochemistry*, **39**, 6726–6731.
8. Tan, M., Zhu, B., Zhou, X., He, R., Chen, X., Eriani, G. and Wang, E. (2010) tRNA-dependent pre-transfer editing by prokaryotic leucyl-tRNA synthetase. *J. Biol. Chem.*, **285**, 3235–3244.
9. Chen, X., Ma, J., Tan, M., Yao, P., Hu, Q., Eriani, G. and Wang, E. (2011) Modular pathways for editing non-cognate amino acids by human cytoplasmic leucyl-tRNA synthetase. *Nucleic Acids Res.*, **39**, 235–247.
10. Dulic, M., Cvetic, N., Perona, J.J. and Gruic-Sovulj, I. (2010) Partitioning of tRNA-dependent editing between pre- and post-transfer pathways in Class I aminoacyl-tRNA synthetases. *J. Biol. Chem.*, **285**, 23799–23809.
11. Fukunaga, R. and Yokoyama, S. (2005) Aminoacylation complex structures of leucyl-tRNA synthetase and tRNA^{Leu} reveal two modes of discriminator-base recognition. *Nat. Struct. Mol. Biol.*, **12**, 915–922.
12. Giege, R., Sissler, M. and Florentz, C. (1998) Universal rules and idiosyncratic features in tRNA identity. *Nucleic Acids Res.*, **26**, 5017–5035.
13. Pan, B., Xiong, Y. and Steitz, T.A. (2010) How the CCA-adding enzyme selects adenine over cytosine at position 76 of tRNA. *Science*, **330**, 937–940.
14. Hou, Y.M. (2000) Unusual synthesis by the *Escherichia coli* CCA-adding enzyme. *RNA*, **6**, 1031–1043.
15. Nissen, P., Kjeldgaard, M., Thirup, S., Polekhina, G., Reshetnikova, L., Clark, B.F.C. and Nyborg, J. (1995) Crystal structure of the ternary complex of Phe-tRNA^{Phe}, EF-Tu, and a GTP analog. *Science*, **270**, 1464–1472.
16. Selmer, M., Dunham, C.M., Murphy, F.V., Weixlbaumer, A., Petry, S., Kelley, A.C., Weir, J.R. and Ramakrishnan, V. (2006) Structure of the 70S Ribosome Complexed with mRNA and tRNA. *Science*, **313**, 1935–1942.
17. Tamura, K. and Schimmel, P. (2000) Trial for peptide bond formation using model molecules based on the interactions between the CCA sequence of tRNA and 23S rRNA. *Nucleic Acids Symp. Ser.*, **44**, 251–252.
18. Low, B., Gates, F., Goldstein, T. and Söll, D. (1971) Isolation and partial characterization of temperature-sensitive *Escherichia coli* mutants with altered leucyl- and seryl-transfer ribonucleic acid synthetases. *J. Bacteriol.*, **108**, 742–750.
19. Li, Y., Wang, E. and Wang, Y. (1999) A modified procedure for fast purification of T7 RNA polymerase. *Protein Expr. Purif.*, **16**, 355–358.
20. Xu, M., Li, J., Du, X. and Wang, E. (2004) Groups on the side chain of T252 in *Escherichia coli* leucyl-tRNA synthetase are important for discrimination of amino acids and cell viability. *Biochem. Biophys. Res. Commun.*, **318**, 11–16.
21. Yao, P., Zhu, B., Jaeger, S., Eriani, G. and Wang, E. (2008) Recognition of tRNA^{Leu} by *Aquifex aeolicus* leucyl-tRNA synthetase during the aminoacylation and editing steps. *Nucleic Acids Res.*, **36**, 2728–2738.
22. Lorber, B., Kern, D., Gieg, R. and Ebel, J.P. (1982) Covalent attachment of aspartic acid to yeast aspartyl-tRNA synthetase induced by the enzyme. *FEBS Lett.*, **146**, 59–64.
23. Rapaport, E., Yogeewaran, G., Zamecnik, P.C. and Remy, P. (1985) Covalent modification of phenylalanyl-tRNA synthetase with phenylalanine during the amino acid activation reaction catalyzed by the enzyme. *J. Biol. Chem.*, **260**, 9509–9512.
24. Kovaleva, G.K., Moroz, S.G., Favorova, O.O. and Kisselev, L.L. (1978) Tryptophanyl-tRNA synthetase: evidence for an anhydrous bond involved in the tryptophanyl enzyme formation. *FEBS Lett.*, **95**, 81–84.
25. Gruic-Sovulj, I., Uter, N., Bullock, T. and Perona, J.J. (2005) tRNA-dependent aminoacyl-adenylate hydrolysis by a nonediting Class I aminoacyl-tRNA synthetase. *J. Biol. Chem.*, **280**, 23978–23986.
26. Zhu, B., Yao, P., Tan, M., Eriani, G. and Wang, E. (2009) tRNA-independent pretransfer editing by Class I leucyl-tRNA synthetase. *J. Biol. Chem.*, **284**, 3418–3424.
27. Li, Y., Wang, E. and Wang, Y. (1998) Overproduction and purification of *Escherichia coli* tRNA^{Leu}. *Sci. China, C, Life Sci.*, **41**, 225–231.
28. Lincecum, T.L. Jr, Tukalo, M., Yaremchuk, A., Mursinna, R.S., Williams, A.M., Sproat, B.S., Van Den Eynde, W., Link, A., Van Calenbergh, S., Gröli, M. et al. (2003) Structural and mechanistic basis of pre- and posttransfer editing by leucyl-tRNA synthetase. *Mol. Cell*, **11**, 951–963.
29. Li, T., Li, Y., Guo, N., Wang, E. and Wang, Y. (1999) Discrimination of tRNA^{Leu} isoacceptors by the insertion mutant of *Escherichia coli* leucyl-tRNA synthetase. *Biochemistry*, **38**, 9084–9088.
30. Schurer, H., Schiffer, S., Marchfelder, A. and Morl, M. (2001) This is the end: processing, editing and repair at the tRNA 3'-terminus. *Biol. Chem.*, **382**, 1147–1156.
31. Tamura, K., Nameki, N., Hasegawa, T., Shimizu, M. and Himeno, H. (1994) Role of the CCA terminal sequence of tRNA(Val) in aminoacylation with valyl-tRNA synthetase. *J. Biol. Chem.*, **269**, 22173–22177.
32. Liu, M. and Horowitz, J. (1994) Functional transfer RNAs with modifications in the 3'-CCA end: differential effects on aminoacylation and polypeptide synthesis. *Proc. Natl Acad. Sci. USA*, **91**, 10389–10393.
33. Eriani, G. and Gangloff, J. (1999) Yeast aspartyl-tRNA synthetase residues interacting with tRNA(Asp) identity bases connectively contribute to tRNA(Asp) binding in the ground and transition-state complex and discriminate against non-cognate tRNAs. *J. Mol. Biol.*, **291**, 761–773.
34. Tardif, K.D. and Horowitz, J. (2002) Transfer RNA determinants for translational editing by *Escherichia coli* valyl-tRNA synthetase. *Nucleic Acids Res.*, **30**, 2538–2545.
35. Splan, K.E., Ignatov, M.E. and Musier-Forsyth, K. (2008) Transfer RNA modulates the editing mechanism used by class II prolyl-tRNA synthetase. *J. Biol. Chem.*, **283**, 7128–7134.
36. Gruic-Sovulj, I., Rokov-Plavec, J. and Weygand-Durasevic, I. (2007) Hydrolysis of non-cognate aminoacyl-adenylates by a class II aminoacyl-tRNA synthetase lacking an editing domain. *FEBS Lett.*, **581**, 5110–5114.
37. Minajigi, A. and Francklyn, C.S. (2010) Aminoacyl transfer rate dictates choice of editing pathway in threonyl-tRNA synthetase. *J. Biol. Chem.*, **285**, 23810–23817.
38. Dulic, M., Cvetic, N., Perona, J.J. and Gruic-Sovulj, I. (2010) Partitioning of tRNA-dependent editing between pre- and post-transfer pathways in class I aminoacyl-tRNA synthetases. *J. Biol. Chem.*, **285**, 23799–23809.
39. Hou, Y.M. (2000) Unusual synthesis by the *Escherichia coli* CCA-adding enzyme. *RNA*, **6**, 1031–1043.
40. Cho, H.D., Verlinde, C.L. and Weiner, A.M. (2007) Reengineering CCA-adding enzymes to function as (U,G)- or dCdCdA-adding

- enzymes or poly(C,A) and poly(U,G) polymerases.
Proc. Natl Acad. Sci. USA, **104**, 54–59.
41. Tomita, K. and Weiner, A.M. (2001) Collaboration between CC- and A-adding enzymes to build and repair the 3'-terminal CCA of tRNA in *Aquifex aeolicus*. *Science*, **294**, 1334–1336.
42. Nissen, P., Kjeldgaard, M., Thirup, S., Polekhina, G., Reshetnikova, L., Clark, B.F. and Nyborg, J. (1995) Crystal structure of the ternary complex of Phe-tRNA^{Phe}, EF-Tu, and a GTP analog. *Science*, **270**, 1464–1472.
43. Selmer, M., Dunham, C.M., Murphy, F.V.t., Weixlbaumer, A., Petry, S., Kelley, A.C., Weir, J.R. and Ramakrishnan, V. (2006) Structure of the 70S ribosome complexed with mRNA and tRNA. *Science*, **313**, 1935–1942.