

Coadjoint Orbits of Lie Algebras and Cartan Class

Michel GOZE[†] and Elisabeth REMM[‡]

[†] *Ramm Algebra Center, 4 rue de Cluny, F-68800 Rammersmatt, France*

E-mail: goze.rac@gmail.com

[‡] *Université de Haute-Alsace, IRIMAS EA 7499, Département de Mathématiques, F-68100 Mulhouse, France*

E-mail: elisabeth.remm@uha.fr

Received September 13, 2018, in final form December 31, 2018; Published online January 09, 2019

<https://doi.org/10.3842/SIGMA.2019.002>

Abstract. We study the coadjoint orbits of a Lie algebra in terms of Cartan class. In fact, the tangent space to a coadjoint orbit $\mathcal{O}(\alpha)$ at the point α corresponds to the characteristic space associated to the left invariant form α and its dimension is the even part of the Cartan class of α . We apply this remark to determine Lie algebras such that all the nontrivial orbits (nonreduced to a point) have the same dimension, in particular when this dimension is 2 or 4. We determine also the Lie algebras of dimension $2n$ or $2n + 1$ having an orbit of dimension $2n$.

Key words: Lie algebras; coadjoint representation; contact forms; Frobenius Lie algebras; Cartan class

2010 Mathematics Subject Classification: 17B20; 17B30; 53D10; 53D05

1 Introduction

Let G be a connected Lie group, \mathfrak{g} its Lie algebra and \mathfrak{g}^* the dual vector space of \mathfrak{g} . We identify \mathfrak{g} with the Lie algebra of left invariant vector fields on G and \mathfrak{g}^* with the vector space of left invariant Pfaffian forms on G . The Lie group G acts on \mathfrak{g}^* by the coadjoint action. If α belongs to \mathfrak{g}^* , its coadjoint orbit $\mathcal{O}(\alpha)$ associated with this action is reduced to a point if α is closed for the adjoint cohomology of \mathfrak{g} . If not, the coadjoint orbit is an even-dimensional manifold provided with a symplectic structure. From the Kirillov theory, if G is a connected and simply connected nilpotent Lie group, there exists a canonical bijection from the set of coadjoint orbits onto the set of equivalence classes of irreducible unitary representations of this Lie group. In this work, we establish a link between the dimension of the coadjoint orbit of the form α and $\text{cl}(\alpha)$ its class in Elie Cartan's sense. More precisely $\dim \mathcal{O}(\alpha) = 2 \lfloor \frac{\text{cl}(\alpha)}{2} \rfloor$. Recall that the Cartan class of α corresponds to the number of independent Pfaffian forms needed to define α and its differential $d\alpha$ and it is equal to the codimension of the characteristic space [4, 13, 19]. The dimension of $\mathcal{O}(\alpha)$ results in a natural relation between this characteristic space and the tangent space at the point α to the orbit $\mathcal{O}(\alpha)$.

As applications, we describe classes of Lie algebras with additional properties related to its coadjoint orbits. For example, we determine all Lie algebras whose nontrivial orbits are all of dimension 2 or 4 and also the Lie algebras of dimension $2p$ or $2p + 1$ admitting a maximal orbit of dimension $2p$ that is admitting $\alpha \in \mathfrak{g}^*$ such that $\text{cl}(\alpha) \geq 2p$. Notice that the nilpotent Lie algebras classified in Sections 3 and 4 play a role in connection with a so-called inverse problem in representation theory of nilpotent Lie groups (see [6] and [7, Section 5.1]).

2 Dimension of coadjoint orbits and Cartan class

2.1 Cartan class of a Pfaffian form

Let M be a n -dimensional differentiable manifold and α a Pfaffian form on M , that is a differential form of degree 1. The *characteristic space of α at a point $x \in M$* is the linear subspace $\mathcal{C}_x(\alpha)$ of the tangent space $T_x(M)$ of M at the point x defined by

$$\mathcal{C}_x(\alpha) = A(\alpha(x)) \cap A(d\alpha(x)),$$

where

$$A(\alpha(x)) = \{X_x \in T_x(M), \alpha(x)(X_x) = 0\}$$

is the *associated subspace of $\alpha(x)$* ,

$$A(d\alpha(x)) = \{X_x \in T_x(M), i(X_x)d\alpha(x) = 0\}$$

is the *associated subspace of $d\alpha(x)$* and $i(X_x)d\alpha(x)(Y_x) = d\alpha(x)(X_x, Y_x)$.

Definition 2.1. Let α be a Pfaffian form on the differential manifold M . The *Cartan class of α at the point $x \in M$* [9] is the codimension of the characteristic space $\mathcal{C}_x(\alpha)$ in the tangent space $T_x(M)$ to M at the point x . We denote it by $\text{cl}(\alpha)(x)$.

The function $x \rightarrow \text{cl}(\alpha)(x)$ is with positive integer values and is lower semi-continuous, that is, for every $x \in M$ there exists a suitable neighborhood V such that for every $x_1 \in V$ one has $\text{cl}(\alpha)(x_1) \geq \text{cl}(\alpha)(x)$.

The *characteristic system of α at the point x of M* is the subspace $\mathcal{C}_x^*(\alpha)$ of the dual $T_x^*(M)$ of $T_x(M)$ orthogonal to $\mathcal{C}_x(\alpha)$:

$$\mathcal{C}_x^*(\alpha) = \{\omega(x) \in T_x^*(M), \omega(x)(X_x) = 0, \forall X_x \in \mathcal{C}_x(\alpha)\}.$$

Then

$$\text{cl}(\alpha)(x) = \dim \mathcal{C}_x^*(\alpha).$$

Proposition 2.2. *If α is a Pfaffian form on M , then*

- $\text{cl}(\alpha)(x) = 2p + 1$ if $(\alpha \wedge (d\alpha)^p)(x) \neq 0$ and $(d\alpha)^{p+1}(x) = 0$,
- $\text{cl}(\alpha)(x) = 2p$ if $(d\alpha)^p(x) \neq 0$ and $(\alpha \wedge (d\alpha)^p)(x) = 0$.

In the first case, there exists a basis $\{\omega_1(x) = \alpha(x), \omega_2(x), \dots, \omega_n(x)\}$ of $T_x^*(M)$ such that

$$d\alpha(x) = \omega_2(x) \wedge \omega_3(x) + \dots + \omega_{2p}(x) \wedge \omega_{2p+1}(x)$$

and

$$\mathcal{C}_x^*(\alpha) = \mathbb{R}\{\alpha(x)\} + A^*(d\alpha(x)).$$

In the second case, there exists a basis $\{\omega_1(x) = \alpha(x), \omega_2(x), \dots, \omega_n(x)\}$ of $T_x^*(M)$ such that

$$d\alpha(x) = \alpha(x) \wedge \omega_2(x) + \dots + \omega_{2p-1}(x) \wedge \omega_{2p}(x)$$

and

$$\mathcal{C}_x^*(\alpha) = A^*(d\alpha(x)).$$

If the function $\text{cl}(\alpha)(x)$ is constant, that is, $\text{cl}(\alpha)(x) = \text{cl}(\alpha)(y)$ for any $x, y \in M$, we say that the Pfaffian form α is of constant class and we denote by $\text{cl}(\alpha)$ this constant. The distribution

$$x \rightarrow \mathcal{C}_x(\alpha)$$

is then regular and it is an integrable distribution of dimension $n - \text{cl}(\alpha)$, called the characteristic distribution of α . It is equivalent to say that the Pfaffian system

$$x \rightarrow \mathcal{C}_x^*(\alpha)$$

is integrable and of dimension $\text{cl}(\alpha)$.

If $M = G$ is a connected Lie group, we identify its Lie algebra \mathfrak{g} with the space of left invariant vector fields and its dual \mathfrak{g}^* with the space of left invariant Pfaffian forms. Then if $\alpha \in \mathfrak{g}^*$, the differential $d\alpha$ is the 2-differential left invariant form belonging to $\Lambda^2(\mathfrak{g}^*)$ and defined by

$$d\alpha(X, Y) = -\alpha[X, Y]$$

for any $X, Y \in \mathfrak{g}$. It is obvious that any left invariant form $\alpha \in \mathfrak{g}^*$ is of constant class and we will speak on the *Cartan class* $\text{cl}(\alpha)$ of a linear form $\alpha \in \mathfrak{g}^*$. We have

- $\text{cl}(\alpha) = 2p + 1$ if and only if $\alpha \wedge (d\alpha)^p \neq 0$ and $(d\alpha)^{p+1} = 0$,
- $\text{cl}(\alpha) = 2p$ if and only if $(d\alpha)^p \neq 0$ and $\alpha \wedge (d\alpha)^p = 0$.

Definition 2.3. Let \mathfrak{g} be an n -dimensional Lie algebra.

- It is called *contact Lie algebra* if $n = 2p + 1$ and if there exists a contact linear form, that is, a linear form of Cartan class equal to $2p + 1$.
- It is called *Frobenius Lie algebra* if $n = 2p$ and if there exists a Frobenius linear form, that is, a linear form of Cartan class equal to $2p$.

If $\alpha \in \mathfrak{g}^*$ is neither a contact nor a Frobenius form, the characteristic space $\mathcal{C}(\alpha) = \mathcal{C}_e(\alpha)$ at the unit e of G is not trivial and the characteristic distribution on G given by $\mathcal{C}_x(\alpha)$ with $x \in G$ has a constant non-zero dimension. As it is integrable, the subspace $\mathcal{C}(\alpha)$ is a Lie subalgebra of \mathfrak{g} .

Proposition 2.4. Let $\alpha \in \mathfrak{g}^*$ be a linear form of maximal class, that is

$$\forall \beta \in \mathfrak{g}^*, \quad \text{cl}(\alpha) \geq \text{cl}(\beta).$$

Then $\mathcal{C}(\alpha)$ is an abelian subalgebra of \mathfrak{g} .

Proof. Assume that α is a linear form of maximal class. If $\text{cl}(\alpha) = 2p + 1$, there exists a basis $\{\omega_1 = \alpha, \omega_2, \dots, \omega_n\}$ of \mathfrak{g}^* such that $d\alpha = \omega_2 \wedge \omega_3 + \dots + \omega_{2p} \wedge \omega_{2p+1}$ and $\mathcal{C}^*(\alpha)$ is generated by $\{\omega_1, \dots, \omega_{2p+1}\}$. If the subalgebra $\mathcal{C}(\alpha)$ is not abelian, there exists j , $2p + 2 \leq j \leq n$ such that

$$d\omega_j \wedge \omega_1 \wedge \dots \wedge \omega_{2p+1} \neq 0.$$

Then

$$\text{cl}(\alpha + t\omega_j) > \text{cl}(\alpha)$$

for some $t \in \mathbb{R}$. But α is of maximal class. Then $\mathcal{C}(\alpha)$ is abelian. The proof when $\text{cl}(\alpha) = 2p$ is similar. ■

Note that this result can alternatively be derived from the result of M. Duflo and M. Vergne [12].

Recall some properties of the class of a linear form on a Lie algebra. The proofs of these statements are given in [14, 15, 19].

- If \mathfrak{g} is a finite-dimensional nilpotent Lie algebra, then the class of any non-zero $\alpha \in \mathfrak{g}^*$ is always odd.
- A real or complex finite-dimensional nilpotent Lie algebra is never a Frobenius Lie algebra. More generally, an unimodular Lie algebra is non-Frobenius [11, 24].
- Let \mathfrak{g} be a real compact Lie algebra. Any nontrivial $\alpha \in \mathfrak{g}^*$ has an odd Cartan class.
- Let \mathfrak{g} be a complex semisimple Lie algebra of rank r . Then any $\alpha \in \mathfrak{g}^*$ satisfies $\text{cl}(\alpha) \leq n - r + 1$ [15]. In particular, a semisimple Lie algebra is never a Frobenius algebra. A semisimple Lie algebra is a contact Lie algebra if its rank is 1, that is, \mathfrak{g} is isomorphic to $\mathfrak{sl}(2, \mathbb{C})$. In particular, if \mathfrak{g} is a contact real simple Lie algebra, then \mathfrak{g} is isomorphic to $\mathfrak{sl}(2, \mathbb{R})$ or $\mathfrak{so}(3)$.
- The Cartan class of any linear nontrivial form on a simple nonexceptional complex Lie algebra of rank r satisfies $\text{cl}(\alpha) \geq 2r$ [15]. Moreover, if \mathfrak{g} is isomorphic of type A_r , there exists a linear form of class $2r$ which reaches the lower bound.
- Any $(2p + 1)$ -dimensional contact real Lie algebra such that any nontrivial linear form is a contact form is isomorphic to $\mathfrak{so}(3)$.

2.2 Cartan class and the index of a Lie algebra

For any $\alpha \in \mathfrak{g}^*$, we consider the stabilizer $\mathfrak{g}_\alpha = \{X \in \mathfrak{g}, \alpha \circ \text{ad} X = 0\}$ and d the minimal dimension of \mathfrak{g}_α when α lies in \mathfrak{g}^* . It is an invariant of \mathfrak{g} which is called the index of \mathfrak{g} . If α satisfies $\dim \mathfrak{g}_\alpha = d$ then, from Proposition 2.4, \mathfrak{g}_α is an abelian subalgebra of \mathfrak{g} . Considering the Cartan class of α , \mathfrak{g}_α is the associated subspace of $d\alpha$:

$$\mathfrak{g}_\alpha = A(d\alpha),$$

so the minimality is realized by a form of maximal class and we have $d = n - \text{cl}(\alpha) + 1$ if the Cartan class $\text{cl}(\alpha)$ is odd or $d = n - \text{cl}(\alpha)$ if $\text{cl}(\alpha)$ is even. In particular,

- 1) if \mathfrak{g} is a $2p$ -dimensional Frobenius Lie algebra, then the maximal class is $2p$ and $d = 0$,
- 2) if \mathfrak{g} is $(2p + 1)$ -dimensional contact Lie algebra, then $d = n - n + 1 = 1$.

This relation between index and Cartan class is useful to compute sometimes this index. For example we have

Proposition 2.5. *Let L_n or Q_{2p} be the naturally graded filiform Lie algebras. Their index satisfy*

- 1) $d(L_n) = n - 2$,
- 2) $d(Q_{2p}) = 2$.

In fact L_n is defined in a basis $\{e_0, \dots, e_{n-1}\}$ by $[e_0, e_i] = e_{i+1}$ for $i = 1, \dots, n - 2$ and we have $\text{cl}(\alpha) \in \{1, 3\}$ and $d = n - 2$ for any $\alpha \in L_n^*$. The second algebra Q_{2p} is defined in the basis $\{e_0, \dots, e_{2p-1}\}$ by $[e_0, e_i] = e_{i+1}$ for $i = 1, \dots, 2p - 3$, $[e_i, e_{2p-1-i}] = (-1)^{i-1} e_{2p-1}$ for $i = 1, \dots, p - 1$. In this case we have $\text{cl}(\alpha) \in \{1, 3, 2p - 1\}$ for any $\alpha \in Q_{2p}^*$ and $d = 2$. Let us note that a direct computation of these indexes are given in [1].

2.3 The coadjoint representation

Let G be a connected Lie group and \mathfrak{g} its (real) Lie algebra. The adjoint representation of G on \mathfrak{g} is the homomorphism of groups:

$$\text{Ad}: G \rightarrow \text{Aut}(\mathfrak{g})$$

defined as follows. For every $x \in G$, let $A(x)$ be the automorphism of G given by $A(x)(y) = xyx^{-1}$. This map is differentiable and the tangent map to the identity e of G is an automorphism of \mathfrak{g} . We denote it by $\text{Ad}(x)$.

Definition 2.6. The coadjoint representation of G on the dual \mathfrak{g}^* of \mathfrak{g} is the homomorphism of groups:

$$\text{Ad}^*: G \rightarrow \text{Aut}(\mathfrak{g}^*)$$

defined by

$$\langle \text{Ad}^*(x)\alpha, X \rangle = \langle \alpha, \text{Ad}(x^{-1})X \rangle$$

for any $\alpha \in \mathfrak{g}^*$ and $X \in \mathfrak{g}$.

The coadjoint representation is sometimes called the K -representation. For $\alpha \in \mathfrak{g}^*$, we denote by $\mathcal{O}(\alpha)$ its orbit, called the coadjoint orbit, for the coadjoint representation. The following result is classical [23]: any coadjoint orbit is an even-dimensional differentiable manifold endowed with a symplectic form.

Let us compute the tangent space to this manifold $\mathcal{O}(\alpha)$ at the point α . Any $\beta \in \mathcal{O}(\alpha)$ is written $\beta = \alpha \circ \text{Ad}(x^{-1})$ for some $x \in G$. The map $\rho: G \rightarrow \mathfrak{g}^*$ defined by $\rho(x) = \alpha \circ \text{Ad}(x)$ is differentiable and its tangent map at the identity of G is given by $\rho_e^T(X) = \text{i}(X)d\alpha$ for any $X \in \mathfrak{g}$ with $\text{i}(X)d\alpha(Y) = -\alpha[X, Y]$. Then the tangent space to $\mathcal{O}(\alpha)$ at the point α corresponds to $A^*(d\alpha) = \{\omega \in \mathfrak{g}^*, \omega(X) = 0, \forall X \in A(d\alpha)\}$.

Proposition 2.7. Consider a non-zero $\alpha \in \mathfrak{g}^*$. The tangent space to $\mathcal{O}(\alpha)$ at the point α is isomorphic to the dual space $A^*(d\alpha)$ of the associated space $A(d\alpha)$ of $d\alpha$.

Corollary 2.8. Consider a non-zero α in \mathfrak{g}^* . Then $\dim \mathcal{O}(\alpha) = 2p$ if and only if $\text{cl}(\alpha) = 2p$ or $\text{cl}(\alpha) = 2p + 1$.

An immediate application is

Proposition 2.9. Any $(2p+1)$ -dimensional Lie algebra with $\dim \mathcal{O}(\alpha) = 2p$ for all $\alpha \in \mathfrak{g}^* \setminus \{0\}$, is isomorphic to $\mathfrak{so}(3)$ or $\mathfrak{sl}(2, \mathbb{R})$ and then of dimension 3.

Proof. From [15, 19] any contact Lie algebra such that any nontrivial linear form is a contact form is isomorphic to $\mathfrak{so}(3)$. Assume now that any nontrivial form on \mathfrak{g} is of Cartan class equal to $2p$ or $2p + 1$. With similar arguments developed in [15, 19], we prove that such a Lie algebra is semisimple. But in this case, we have seen that \mathfrak{g} is isomorphic to $\mathfrak{sl}(2, \mathbb{R})$ or $\mathfrak{so}(3)$. ■

Remark 2.10. Assume that $\dim \mathfrak{g} = 2p$ and all the nontrivial coadjoint orbits are also of dimension $2p$. Then for any nontrivial $\alpha \in \mathfrak{g}^*$, $\text{cl}(\alpha) = 2p$. If I is a nontrivial abelian ideal of \mathfrak{g} , there exists $\omega \in \mathfrak{g}^*$, $\omega \neq 0$ such that $\omega(X) = 0$ for any $X \in I$. The Cartan class of this form ω is smaller than $2p$. Then \mathfrak{g} is semisimple. But the behavior of the Cartan class on simple Lie algebra leads to a contradiction.

We deduce also from the previous corollary:

Proposition 2.11.

1. If \mathfrak{g} is isomorphic to the $(2p + 1)$ -dimensional Heisenberg algebra, then any nontrivial coadjoint orbit is of dimension $2p$.
2. If \mathfrak{g} is isomorphic to the graded filiform algebra L_n , then any nontrivial coadjoint orbit is of dimension 2.
3. If \mathfrak{g} is isomorphic to the graded filiform algebra Q_n , then any nontrivial coadjoint orbit is of dimension 2 or $n - 2$.
4. If \mathfrak{g} is a $(2p + 1)$ -dimensional 2-step nilpotent Lie algebra with a coadjoint orbit of dimension $2p$, then \mathfrak{g} is isomorphic to the $(2p + 1)$ -dimensional Heisenberg Lie algebra (see [19]).
5. If \mathfrak{g} is a n -dimensional complex classical simple Lie algebra of rank r , then the maximal dimension of the coadjoint orbits is equal to $n - r$ if this number is even, if not to $n - r - 1$ (see [15]).

3 Lie algebras whose coadjoint orbits are of dimension 2 or 0

In this section, we determine all Lie algebras whose coadjoint orbits are of dimension 2 or 0. This problem was initiated in [3, 5]. This is equivalent to say that the Cartan class of any linear form is smaller or equal to 3. If \mathfrak{g} is a Lie algebra having this property, any direct product $\mathfrak{g}_1 = \mathfrak{g} \oplus J$ of \mathfrak{g} by an abelian ideal J satisfies also this property. We shall describe these Lie algebras up to an abelian direct factor, that is indecomposable Lie algebras. It is obvious that for any Lie algebra of dimension 2 or 3, the dimensions of the coadjoint orbits are equal to 2 or 0. We have also seen:

Proposition 3.1. *Let \mathfrak{g} be a simple Lie algebra of rank 1. Then for any $\alpha \neq 0 \in \mathfrak{g}^*$, $\dim \mathcal{O}(\alpha) = 2$. Conversely, if \mathfrak{g} is a Lie algebra such that $\dim \mathcal{O}(\alpha) = 2$ for any $\alpha \neq 0 \in \mathfrak{g}^*$ then \mathfrak{g} is simple of rank 1.*

Recall that the rank of a real semisimple Lie algebra \mathfrak{g} is the dimension of any Cartan subalgebra \mathfrak{h} of \mathfrak{g} . This is well defined since \mathfrak{h} is a Cartan subalgebra if and only if $\mathfrak{h}_{\mathbb{C}}$ is Cartan in the complexified simple Lie algebra $\mathfrak{g}_{\mathbb{C}}$. Now we examine the general case. Assume that \mathfrak{g} is a Lie algebra of dimension greater or equal to 4 such that for any nonzero $\alpha \in \mathfrak{g}^*$ we have $\text{cl}(\alpha) = 3, 2$ or 1.

Assume in a first step that $\text{cl}(\alpha) = 2$ for any nonclosed $\alpha \in \mathfrak{g}^*$. Let α be a non-zero 1-form and $\text{cl}(\alpha) = 2$. Then there exists a basis $\{\alpha = \omega_1, \dots, \omega_n\}$ of \mathfrak{g}^* such that $d\alpha = d\omega_1 = \omega_1 \wedge \omega_2$. This implies $d(d\omega_1) = 0 = -\omega_1 \wedge d\omega_2$. Therefore $d\omega_2 = \omega_1 \wedge \omega$ with $\omega \in \mathfrak{g}^*$. As $\text{cl}(\omega_2) \leq 2$, $\omega_2 \wedge \omega = 0$. If $\{X_1, X_2, \dots, X_n\}$ is the dual basis of $\{\alpha = \omega_1, \dots, \omega_n\}$, then $A(d\omega_1) = \mathbb{K}\{X_3, \dots, X_n\}$ is an abelian subalgebra of \mathfrak{g} . Suppose $[X_1, X_2] = X_1 + aX_2 + U$ where U, X_1, X_2 are linearly independent. The dual form of U would be of class 3 so $U = 0$. Under the assumption of change of basis if $a \neq 0$ we can assume that $[X_1, X_2] = X_1$. So $d\omega_2 = \omega_1 \wedge \omega$ and $\omega \wedge \omega_2 = 0$ imply that $d\omega_2 = 0$. This implies that $A(d\omega_1)$ is an abelian ideal of codimension 2. Let β be in $A(d\omega_1)^*$, with $d\beta \neq 0$. If such a form doesn't exist then $\mathbb{K}X_1 \oplus A(d\omega_1)$ is an abelian ideal of codimension 1. Otherwise $d\beta = \omega_1 \wedge \beta_1 + \omega_2 \wedge \beta_2$ with $\beta_1, \beta_2 \in A(d\omega_1)$. As $\beta \wedge d\beta = d\beta^2 = 0$, $\beta_1 \wedge \beta_2 = 0$ which implies $d\beta = (a\omega_1 + b\omega_2) \wedge \beta$. But $d(d\beta) = 0$ implies $ad\omega_1 \wedge \beta = 0 = a\omega_1 \wedge \omega_2 \wedge \beta$ thus $a = 0$ and $d\beta = b\omega_2 \wedge \beta$. We conclude that $[X_1, A(d\omega_1)] = 0$ and $\mathbb{K}\{X_1\} \oplus A(d\omega_1)$ is an abelian ideal of codimension 1.

Assume now that there exists ω of class 3. There exists a basis $\mathcal{B} = \{\omega_1, \omega_2, \omega_3 = \omega, \dots, \omega_n\}$ of \mathfrak{g}^* such as $d\omega = d\omega_3 = \omega_1 \wedge \omega_2$ and the subalgebra $A(d\omega_3) = \mathbb{K}\{X_3, \dots, X_n\}$ is abelian. If $\{X_1, \dots, X_n\}$ is the dual basis of \mathcal{B} , we can assume that $[X_1, X_2] = X_3$. As $A(d\omega_3)$ is an abelian subalgebra of \mathfrak{g} , for any $\alpha \in \mathfrak{g}^*$ we have $d\alpha = \omega_1 \wedge \alpha_1 + \omega_2 \wedge \alpha_2$ with $\alpha_1, \alpha_2 \in A(d\omega_3)^*$. But

2. Let $U(\mathfrak{g})$ be the universal enveloping algebra of \mathfrak{g} and consider the category $U(\mathfrak{g}) - \mathcal{M}od$ of right $U(\mathfrak{g})$ -module. Then if \mathfrak{g} is a Lie algebra described in this section (that is with coadjoint orbits of dimension 0 or 2) thus any $U(\mathfrak{g})$ -mod satisfy the property that “any injective hulls of simple right $U(\mathfrak{g})$ -module are locally Artinian” (see [21]).
3. The notion of elementary quadratic Lie algebra was introduced by G. Pinczon and R. Ushirobira [25]. They also prove that is \mathfrak{g} is an elementary quadratic Lie algebra, then all coadjoint orbits have dimension at most 2.

4 Lie algebras whose nontrivial coadjoint orbits are of dimension 4

We generalize some results of the previous section, considering here real Lie algebras such that for a fixed $p \in \mathbb{N}$, $\dim \mathcal{O}(\omega) = 2p$ or 0 for any $\omega \in \mathfrak{g}^*$. We are interested, in this section, in the case $p = 2$. The Cartan class of any nonclosed linear form is equal to 5 or 4.

Lemma 4.1. *Let \mathfrak{g} be a Lie algebra whose Cartan class of any nontrivial and nonclosed linear form is 4 or 5. Then \mathfrak{g} is solvable.*

Proof. If \mathfrak{g} is a simple Lie algebra of rank r and dimension n , then the Cartan class of any linear form $\omega \in \mathfrak{g}^*$ satisfies $c \leq n - r + 1$ and this upper bound is reached. Then $n - r + 1 = 4$ or 5 and the only possible case is for $r = 2$ and $\mathfrak{g} = \mathfrak{so}(4)$. Since this algebra is compact, the Cartan class is odd. We can find a basis of $\mathfrak{so}(4)$ whose corresponding Maurer–Cartan equations are

$$\begin{cases} d\omega_1 = -\omega_2 \wedge \omega_4 - \omega_3 \wedge \omega_5, \\ d\omega_2 = \omega_1 \wedge \omega_4 - \omega_3 \wedge \omega_6, \\ d\omega_3 = \omega_1 \wedge \omega_5 + \omega_2 \wedge \omega_6, \\ d\omega_4 = -\omega_1 \wedge \omega_2 - \omega_5 \wedge \omega_6, \\ d\omega_5 = -\omega_1 \wedge \omega_3 + \omega_4 \wedge \omega_6, \\ d\omega_6 = -\omega_2 \wedge \omega_3 - \omega_4 \wedge \omega_5. \end{cases}$$

If each of the linear forms of this basis has a Cartan class equal to 5, it is easy to find a linear form, for example $\omega_1 + \omega_6$, of Cartan class equal to 3. Then \mathfrak{g} is neither simple nor semisimple. This implies also that the Levi part of a nonsolvable Lie algebra is also trivial, then \mathfrak{g} is solvable. ■

4.1 Description of these Lie algebras

A consequence of Lemma 4.1 is that \mathfrak{g} contains a nontrivial abelian ideal. From the result of the previous section, the codimension of this ideal I is greater or equal to 2 and $\mathfrak{g} = \mathfrak{m} \oplus I$ where \mathfrak{m} is a vector subspace of \mathfrak{g} of dimension greater or equal to 2.

Assume in a first time that $\dim \mathfrak{m} = 2$. Since I is a maximal abelian ideal, the Cartan class of any nontrivial linear form is 4 or 5 and the coadjoint nontrivial orbits are of dimension 4. To precise this case it remains to describe the action of \mathfrak{m} on I when we consider the second case. Assume that $\mathfrak{g} = \mathfrak{m} \oplus I$ and $\dim \mathfrak{m} = 2$. Let $\{T_1, T_2\}$ be a basis of \mathfrak{m} . Then $\tilde{\mathfrak{g}} = \mathfrak{g}/\mathbb{K}\{T_2\}$ is a Lie algebra whose any nonclosed linear form is of class 2 or 3. Such Lie algebra is described in Proposition 3.2.

Proposition 4.2. *Let \mathfrak{g} be a Lie algebra, $\mathfrak{g} = \mathfrak{m} \oplus I$ where I is an abelian ideal of codimension 2 and whose coadjoint orbit of any nonclosed linear form is of dimension 4. Let $\{T_1, T_2\}$ be a basis of \mathfrak{m} and $\tilde{\mathfrak{g}} = \mathfrak{g}/\mathbb{K}\{T_2\}$. Then \mathfrak{g} is a one-dimensional extension by a derivation f of $\tilde{\mathfrak{g}}$ such that*

$f(T_1) = 0$, $\text{Im}(f) = \text{Im}(\text{ad } T_1)$ and for any $Y \in \text{Im}(\text{ad } T_1)$, there exist $X_1, X_2 \in I$ linearly independent such that

$$f(X_2) = [T_1, X_1] = Y.$$

Assume now that $\dim \mathfrak{m} = 3$. In this case \mathfrak{g}/I is abelian, that is $[\mathfrak{m}, \mathfrak{m}] \subset I$. Otherwise, there would exist a linear form on \mathfrak{m}^* of Cartan class equal to 2 or 3. This implies that for any $\omega \in \mathfrak{m}^*$ we have $d\omega = 0$ and we obtain, considering the dimension of $[\mathfrak{m}, \mathfrak{m}]$, the following Lie algebras which are nilpotent because the Cartan class is always odd:

1. $\dim[\mathfrak{m}, \mathfrak{m}] = 3$:

$$\begin{cases} d\omega_1 = d\omega_2 = d\omega_3 = d\omega_7 = 0, \\ d\omega_4 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_7, \\ d\omega_5 = \omega_1 \wedge \omega_3 - \omega_2 \wedge \omega_7, \\ d\omega_6 = \omega_2 \wedge \omega_3 + \omega_1 \wedge \omega_7, \end{cases} \quad (4.1)$$

which is of dimension 7 sometimes called the Kaplan Lie algebra or the generalized Heisenberg algebra,

$$\begin{cases} d\omega_1 = d\omega_2 = d\omega_3 = d\omega_7 = d\omega_8 = 0, \\ d\omega_4 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_7, \\ d\omega_5 = \omega_1 \wedge \omega_3 + \omega_2 \wedge \omega_8, \\ d\omega_6 = \omega_2 \wedge \omega_3 + \omega_1 \wedge \omega_7, \end{cases} \quad (4.2)$$

of dimension 8,

$$\begin{cases} d\omega_1 = d\omega_2 = d\omega_3 = d\omega_7 = d\omega_8 = d\omega_9 = 0, \\ d\omega_4 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_7, \\ d\omega_5 = \omega_1 \wedge \omega_3 + \omega_2 \wedge \omega_8, \\ d\omega_6 = \omega_2 \wedge \omega_3 + \omega_1 \wedge \omega_9, \end{cases} \quad (4.3)$$

of dimension 9.

2. $\dim[\mathfrak{m}, \mathfrak{m}] = 2$:

$$\begin{cases} d\omega_1 = d\omega_2 = d\omega_3 = d\omega_7 = d\omega_8 = 0, \\ d\omega_4 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_7, \\ d\omega_5 = \omega_1 \wedge \omega_3 + \omega_2 \wedge \omega_7, \\ d\omega_6 = \omega_1 \wedge \omega_7 + \omega_2 \wedge \omega_8, \end{cases} \quad (4.4)$$

which is of dimension 8,

$$\begin{cases} d\omega_1 = d\omega_2 = d\omega_3 = d\omega_7 = d\omega_8 = d\omega_9 = 0, \\ d\omega_4 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_7, \\ d\omega_5 = \omega_1 \wedge \omega_3 + \omega_2 \wedge \omega_9, \\ d\omega_6 = \omega_1 \wedge \omega_7 + \omega_2 \wedge \omega_8, \end{cases} \quad (4.5)$$

which is of dimension 9.

3. $\dim[\mathfrak{m}, \mathfrak{m}] = 1$

$$\begin{cases} d\omega_1 = d\omega_2 = d\omega_3 = d\omega_7 = d\omega_8 = 0, \\ d\omega_4 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_7, \\ d\omega_5 = \omega_1 \wedge \omega_8 + \omega_2 \wedge \omega_7, \\ d\omega_6 = \omega_1 \wedge \omega_7 + \omega_3 \wedge \omega_8, \end{cases} \quad (4.6)$$

which is of dimension 8.

4. $\dim[\mathfrak{m}, \mathfrak{m}] = 0$

$$\begin{cases} d\omega_1 = d\omega_2 = d\omega_3 = d\omega_7 = d\omega_8 = 0, \\ d\omega_4 = \omega_1 \wedge \omega_7 + \omega_2 \wedge \omega_8, \\ d\omega_5 = \omega_1 \wedge \omega_8 + \omega_3 \wedge \omega_7, \\ d\omega_6 = \omega_2 \wedge \omega_7 + \omega_1 \wedge \omega_8, \end{cases} \quad (4.7)$$

which is of dimension 8.

Assume now that $\dim \mathfrak{m} = 4$. In this case \mathfrak{g}/I is abelian or isomorphic to the solvable Lie algebra whose Maurer–Cartan equations are

$$\begin{cases} d\omega_2 = d\omega_4 = 0, \\ d\omega_1 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_4, \\ d\omega_3 = \omega_3 \wedge \omega_2 + a\omega_1 \wedge \omega_4 \end{cases}$$

with $a \neq 0$. Let us assume that \mathfrak{g}/I is not abelian. Let $\{X_1, \dots, X_n\}$ be a basis of \mathfrak{g} such that $\{X_1, \dots, X_4\}$ is the basis of \mathfrak{m} dual of $\{\omega_1, \dots, \omega_4\}$ and $\{X_5, \dots, X_n\}$ a basis of I . Since I is maximal, then $[X_1, I]$ and $[X_3, I]$ are not trivial. There exists a vector of I , for example, X_5 such that $[X_1, X_5] \neq 0$. Let us put $[X_1, X_5] = Y$ with $Y \in I$ and let be ω its dual form. Then

$$d\omega = \omega_1 \wedge \omega_5 + \theta$$

with $\omega_1 \wedge \omega_5 \wedge \theta \neq 0$ and $\omega_3 \wedge \omega_4 \wedge \theta = \omega_3 \wedge \omega_2 \wedge \theta = 0$ if not there exists a linear form of class greater than 5. This implies that there exists ω_6 independent with ω_5

$$d\omega = \omega_1 \wedge \omega_5 + b\omega_3 \wedge \omega_6$$

with $b \neq 0$. Now the Jacobi conditions which are equivalent to $d(d\omega) = 0$ implies that we cannot have $\omega = \omega_5$ and $\omega = \omega_6$. Then we put $\omega = \omega_7$. This implies

$$d\omega_7 = \omega_1 \wedge \omega_5 + b_2\omega_3 \wedge \omega_6, \quad d\omega_5 = \omega_2 \wedge \omega_5 + b_3\omega_4 \wedge \omega_6, \quad d\omega_6 = \omega_4 \wedge \omega_5 + b_4\omega_2 \wedge \omega_6.$$

We deduce that \mathfrak{g} is isomorphic to the Lie algebra whose Maurer–Cartan equations are

$$\begin{cases} d\omega_2 = d\omega_4 = 0, \\ d\omega_1 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_4, \\ d\omega_3 = \omega_3 \wedge \omega_2 + a_1\omega_1 \wedge \omega_4, \\ d\omega_5 = \omega_2 \wedge \omega_5 + a_2\omega_4 \wedge \omega_6, \\ d\omega_6 = \omega_4 \wedge \omega_5 + a_3\omega_2 \wedge \omega_6, \\ d\omega_7 = \omega_1 \wedge \omega_5 + a_4\omega_3 \wedge \omega_6 \end{cases} \quad (4.8)$$

with $a_1 a_2 a_3 a_4 \neq 0$.

If $\dim \mathfrak{m} \geq 5$, then $\dim A(\omega) = 4$ or 0 and the codimension of I is greater than $n - 4$. Then $\dim \mathfrak{m} \leq 4$.

Proposition 4.3. *Let \mathfrak{g} be a Lie algebra whose coadjoint orbit of any nonclosed linear form is of dimension 4. Then*

- 1) \mathfrak{g} is isomorphic to (4.1), (4.2), (4.3), (4.4), (4.5), (4.6), (4.7), (4.8),
- 2) or \mathfrak{g} admits a decomposition $\mathfrak{g} = \mathfrak{m} \oplus I$ where I is a codimension 2 abelian ideal.

4.2 Classification when $\text{codim } I = 2$

These Lie algebras are described in Proposition 4.2. It remains to classify them up to isomorphism. Let $\{T_1, T_2\}$ be a basis of \mathfrak{m} and $\tilde{\mathfrak{g}} = \mathfrak{g}/\mathbb{K}\{T_2\}$.

• $\dim \mathfrak{g} = 4$. Then $\dim \tilde{\mathfrak{g}} = 3$ and it is isomorphic to one of the two algebras whose Lie brackets are given by

- $[T_1, X_1] = X_2$,
- or $[T_1, X_1] = X_1$, $[T_1, X_2] = aX_2$, $a \neq 0$.

In the first case, it is easy to see that we cannot find derivation of $\tilde{\mathfrak{g}}$ satisfying Proposition 4.2. In the second case the matrix of f in the basis $\{T_1, X_1, X_2\}$ is

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & b & c \\ 0 & d & e \end{pmatrix}.$$

We have no solution if $a \neq 1$. If $a = 1$ then f satisfies

$$(e - b)^2 + 4cd < 0.$$

In particular for $b = e = 0$ and $c = 1$ we obtain

Proposition 4.4. *Any 4-dimensional Lie algebra whose coadjoint orbits of nonclosed linear forms are of dimension 4 is isomorphic to the Lie algebra $\mathfrak{g}_4(\lambda)$ whose Maurer–Cartan equations are*

$$\begin{cases} d\alpha_1 = d\alpha_2 = 0, \\ d\omega_1 = \alpha_1 \wedge \omega_1 + \alpha_2 \wedge \omega_2, \\ d\omega_2 = \alpha_1 \wedge \omega_2 + \lambda\alpha_2 \wedge \omega_1 \end{cases}$$

with $\lambda < 0$.

This Lie algebra corresponds to the Lie algebra M_a^{10} in the classification of W.A. de Graaf (see [10, 14]).

• $\dim \mathfrak{g} = 5$. Let us put $\tilde{\mathfrak{g}} = \mathbb{K}\{T_1\} \oplus I$. Let h_1 be the restriction of $\text{ad } T_1$ to I . It is an endomorphism of I and since $\dim I = 3$, it admits an eigenvalues λ . Assume that $\lambda \neq 0$ and let X_1 be an associated eigenvector. Then, since f is a derivation commuting with $\text{ad } T_1$,

$$[T_1, f(X_1)] = f([T_1, X_1]) = \lambda f(X_1).$$

Then $f(X_1)$ is also an eigenvector associated with λ . By hypothesis X_1 and $f(X_1)$ are independent and λ is a root of order 2. Thus h_1 is semisimple. Let λ_2 be the third eigenvalue. If X_3 is an associated eigenvector, as above $f(X_3)$ is also an eigenvector and λ_2 is a root of order 2 except if $\lambda_2 = \lambda_1$. Then

Lemma 4.5. *If $\dim I$ is odd, then if the restriction h_1 of $\text{ad } T_1$ to I admits a nonzero eigenvalue, we have $h_1 = \lambda \text{Id}$.*

5 Lie algebras with coadjoint orbits of maximal dimension

In this section we are interested by n -dimensional Lie algebra admitting orbits of dimension n if n is even or dimension $n - 1$ if n is odd. This is equivalent to consider Frobenius Lie algebras in the first case and contact Lie algebras in the second case.

5.1 $(2p + 1)$ -dimensional Lie algebras with a $2p$ -dimensional coadjoint orbit

Let \mathfrak{h}_{2p+1} be the $(2p + 1)$ -dimensional Heisenberg algebra. There exists a basis $\{X_1, \dots, X_{2p+1}\}$ such that the Lie brackets of \mathfrak{h}_{2p+1} are $[X_1, X_2] = \dots = [X_{2p-1}, X_{2p}] = X_{2p+1}$, and $[X_i, X_j] = 0$ for $i < j$ and $(i, j) \notin \{(1, 2), \dots, (2p-1, 2p)\}$. If $\{\omega_1, \dots, \omega_{2p+1}\}$ denotes the dual basis of \mathfrak{h}_{2p+1}^* , the Maurer–Cartan equations writes

$$d\omega_{2p+1} = -\omega_1 \wedge \omega_2 - \dots - \omega_{2p-1} \wedge \omega_{2p}, \quad d\omega_i = 0, \quad i = 1, \dots, 2p.$$

Then ω_{2p+1} is a contact form on \mathfrak{h}_{2p+1} and the coadjoint orbit $\mathcal{O}(\omega_{2p+1})$ is of maximal dimension $2p$. Let us note that, in \mathfrak{h}_{2p+1} all the orbits are of dimension $2p$ or are singular. In the following, we will denote by μ_0 the Lie bracket of \mathfrak{h}_{2p+1} .

Definition 5.1. A formal quadratic deformation \mathfrak{g}_t of \mathfrak{h}_{2p+1} is a $(2p+1)$ -dimensional Lie algebra whose Lie bracket μ_t is given by

$$\mu_t(X, Y) = \mu_0(X, Y) + t\varphi_1(X, Y) + t^2\varphi_2(X, Y),$$

where the maps φ_i are bilinear on \mathfrak{h}_{2p+1} with values in \mathfrak{h}_{2p+1} and satisfying

$$\begin{aligned} \delta_{\mu_0}\varphi_1 &= 0, & \varphi_1 \circ \varphi_1 + \delta_{\mu_0}\varphi_2 &= 0, \\ \varphi_2 \circ \varphi_2 &= 0, & \varphi_1 \circ \varphi_2 + \varphi_2 \circ \varphi_1 &= 0. \end{aligned}$$

In this definition δ_μ denotes the coboundary operator of the Chevalley–Eilenberg cohomology of a Lie algebra \mathfrak{g} whose Lie bracket is μ with values in \mathfrak{g} , and if φ and ψ are bilinear maps, then $\varphi \circ \psi$ is the trilinear map given by

$$\varphi \circ \psi(X, Y, Z) = \varphi(\psi(X, Y), Z) + \varphi(\psi(Y, Z), X) + \varphi(\psi(Z, X), Y).$$

In particular $\varphi \circ \varphi = 0$ is equivalent to Jacobi identity and φ , in this case, is a Lie bracket and the coboundary operator writes

$$\delta_\mu\varphi = \mu \circ \varphi + \varphi \circ \mu.$$

Theorem 5.2 ([19]). *Any $(2p+1)$ -dimensional contact Lie algebra \mathfrak{g} is isomorphic to a quadratic formal deformation of \mathfrak{h}_{2p+1} .*

Then its Lie bracket μ writes $\mu = \mu_0 + t\varphi_1 + t^2\varphi_2$ and the bilinear maps φ_1 and φ_2 have the following expressions in the basis $\{X_1, \dots, X_{2p+1}\}$:

$$\begin{aligned} \varphi_1(X_{2k-1}, X_{2k}) &= \sum_{s=1}^{2p} C_{2k-1, 2k}^s X_s, & k &= 1, \dots, p, \\ \varphi_1(X_l, X_r) &= \sum_{s=1}^{2p} C_{l,r}^s X_s, & 1 \leq l < r \leq 2p, & \quad (l, r) \neq (2k-1, 2k), \end{aligned}$$

and

$$\begin{aligned}\varphi_2(X_l, X_r) &= 0, \quad l, r = 1, \dots, 2p, \\ \varphi_2(X_l, X_{2p+1}) &= \sum_{s=1}^{2p} C_{l,2p+1}^s X_s, \quad l = 1, \dots, 2p,\end{aligned}$$

and the nondefined values are equal to 0. Since the center of any deformation of \mathfrak{h}_{2p+1} is of dimension less than or equal to the dimension of \mathfrak{h}_{2p+1} , we deduce:

Corollary 5.3 ([15]). *The center $Z(\mathfrak{g})$ of a contact Lie algebra \mathfrak{g} is of dimension less than or equal to 1.*

5.1.1 Case of nilpotent Lie algebras

If \mathfrak{g} is a contact nilpotent Lie algebra, its center is of dimension 1. In the given basis, this center is $\mathbb{R}\{X_{2p+1}\}$. This implies that $\varphi_2 = 0$.

Proposition 5.4. *Any $(2p+1)$ -dimensional contact nilpotent Lie algebra is isomorphic to a linear deformation $\mu_t = \mu_0 + t\varphi_1$ of the Heisenberg algebra \mathfrak{h}_{2p+1} .*

As consequence, we have

Corollary 5.5. *Any $(2p+1)$ -dimensional contact nilpotent Lie algebra is isomorphic to a central extension of a $2p$ -dimensional symplectic Lie algebra by its symplectic form.*

Proof. Let \mathfrak{t} be the $2p$ -dimensional vector space generated by $\{X_1, \dots, X_{2p}\}$. The restriction to \mathfrak{t} of the 2-cocycle φ_1 is with values in \mathfrak{t} . Since $\varphi_1 \circ \varphi_1 = 0$, it defines on \mathfrak{t} a structure of $2p$ -dimensional Lie algebra. If $\{\omega_1, \dots, \omega_{2p+1}\}$ is the dual basis of the given classical basis of \mathfrak{h}_{2p+1} , then $\theta = \omega_1 \wedge \omega_2 + \dots + \omega_{2p-1} \wedge \omega_{2p}$ is a 2-form on \mathfrak{t} . We denote by d_{φ_1} the differential operator on the Lie algebra $(\mathfrak{t}, \varphi_1)$, that is $d_{\varphi_1}\omega(X, Y) = -\omega(\varphi_1(X, Y))$ for all $X, Y \in \mathfrak{t}$ and $\omega \in \mathfrak{t}^*$. Since μ_0 is the Heisenberg Lie algebra multiplication, the condition $\delta_{\mu_0}\varphi_1 = 0$ is equivalent to $d_{\varphi_1}(\theta) = 0$. It implies that θ is a closed 2-form on \mathfrak{t} and \mathfrak{g} is a central extension of \mathfrak{t} by θ . ■

We deduce:

Theorem 5.6 ([19]). *Let \mathfrak{g} be a $(2p+1)$ -dimensional k -step nilpotent Lie algebra. Then there exists on \mathfrak{g} a coadjoint orbit of dimension $2p$ if and only if \mathfrak{g} is a central extension of a $2p$ -dimensional $(k-1)$ -step nilpotent symplectic Lie algebra \mathfrak{t} , the extension being defined by the 2-cocycle given by the symplectic form.*

Since the classification of nilpotent Lie algebras is known up the dimension 7, the previous result permits to establish the classification of contact nilpotent Lie algebras of dimension 3, 5 and 7 using the classification in dimension 2, 4 and 6 [17]. For example, the classification of 5-dimensional nilpotent Lie algebras with an orbit of dimension 4 is the following:

- \mathfrak{g} is 4-step nilpotent:
 $\mathfrak{n}_5^1: [X_1, X_i] = X_{i+1}, i = 2, 3, 4, [X_2, X_3] = X_5.$
- \mathfrak{g} is 3-step nilpotent:
 $\mathfrak{n}_5^3: [X_1, X_i] = X_{i+1}, i = 2, 3, [X_2, X_5] = X_4.$
- \mathfrak{g} is 2-step nilpotent:
 $\mathfrak{n}_5^6: [X_1, X_2] = X_3, [X_4, X_5] = X_3.$

We shall now study a contact structure in respect of the characteristic sequence of a nilpotent Lie algebra. For any $X \in \mathfrak{g}$, let $c(X)$ be the ordered sequence, for the lexicographic order, of the dimensions of the Jordan blocks of the nilpotent operator $\text{ad } X$. The characteristic sequence of \mathfrak{g} is the invariant, up to isomorphism,

$$c(\mathfrak{g}) = \max \{c(X), X \in \mathfrak{g} - \mathcal{C}^1(\mathfrak{g})\}.$$

Then $c(\mathfrak{g})$ is a sequence of type $(c_1, c_2, \dots, c_k, 1)$ with $c_1 \geq c_2 \geq \dots \geq c_k \geq 1$ and $c_1 + c_2 + \dots + c_k + 1 = n = \dim \mathfrak{g}$. For example,

- 1) $c(\mathfrak{g}) = (1, 1, \dots, 1)$ if and only if \mathfrak{g} is abelian,
- 2) $c(\mathfrak{g}) = (2, 1, \dots, 1)$ if and only if \mathfrak{g} is a direct product of an Heisenberg Lie algebra by an abelian ideal,
- 3) if \mathfrak{g} is 2-step nilpotent then there exists p and q such that $c(\mathfrak{g}) = (2, 2, \dots, 2, 1, \dots, 1)$ with $2p + q = n$, that is p is the occurrence of 2 in the characteristic sequence and q the occurrence of 1,
- 4) \mathfrak{g} is filiform if and only if $c(\mathfrak{g}) = (n - 1, 1)$.

This invariant was introduced in [2] in order to classify 7-dimensional nilpotent Lie algebras. A link between the notions of breath of nilpotent Lie algebra introduced in [22] and characteristic sequence is developed in [26]. If $c(\mathfrak{g}) = (c_1, c_2, \dots, c_k, 1)$ is the characteristic sequence of \mathfrak{g} then \mathfrak{g} is c_1 -step nilpotent.

Assume now that $c_1 = c_2 = \dots = c_l$ and $c_{l+1} < c_l$. Then the dimension of the center of \mathfrak{g} is greater than l because in each Jordan blocks corresponding to c_1, \dots, c_l , the last vector is in $\mathcal{C}^{c_1}(\mathfrak{g})$ which is contained in the center of \mathfrak{g} . We deduce

Proposition 5.7. *Let \mathfrak{g} be a contact nilpotent Lie algebra. Then its characteristic sequence is of type $c(\mathfrak{g}) = (c_1, c_2, \dots, c_k, 1)$ with $c_2 \neq c_1$.*

Example 5.8.

1. If \mathfrak{g} is 2-step nilpotent, then $c(\mathfrak{g}) = (2, \dots, 2, 1, \dots, 1)$ and if \mathfrak{g} is a contact Lie algebra, we have $c(\mathfrak{g}) = (2, 1, \dots, 1)$. We find again the results given in [19] which precise that any 2-step nilpotent $(2p + 1)$ -dimensional contact Lie algebra is isomorphic to the Heisenberg algebra \mathfrak{h}_{2p+1} .
2. If \mathfrak{g} is 3-step nilpotent, then $c(\mathfrak{g}) = (3, 2, \dots, 2, 1, \dots, 1)$ or $c(\mathfrak{g}) = (3, 1, \dots, 1)$. In case of dimension 7, this gives $c(\mathfrak{g}) = (3, 2, 1, 1)$ and $c(\mathfrak{g}) = (3, 1, 1, 1, 1)$. For each one of characteristic sequences there are contact Lie algebras:

- (a) The Lie algebra given by

$$[X_1, X_2] = X_3, \quad [X_1, X_3] = [X_2, X_5] = [X_6, X_7] = X_4,$$

is a 7-dimensional contact Lie algebra of characteristic $(3, 1, 1, 1, 1)$

- (b) The Lie algebras given

$$\begin{aligned} [X_1, X_i] &= X_{i+1}, & i &= 2, 3, 5, & [X_2, X_5] &= X_7, & [X_2, X_7] &= X_4, \\ [X_5, X_6] &= X_4, & [X_5, X_7] &= \alpha X_4 \end{aligned}$$

with $\alpha \neq 0$ are contact Lie algebras of characteristic $(3, 2, 1, 1)$.

3. If \mathfrak{g} is 4-step nilpotent, then $c(\mathfrak{g}) = (4, 3, \dots, 1)$. For example, the 9-dimensional Lie algebra given by

$$\begin{aligned} [X_1, X_i] &= X_{i+1}, & i &= 2, 3, 4, 6, 7, \\ [X_6, X_9] &= X_3, & [X_7, X_9] &= X_4, & [X_8, X_9] &= X_5, \\ [X_2, X_6] &= (1 + \alpha)X_4, & [X_3, X_6] &= X_5, & [X_2, X_7] &= \alpha X_5 \end{aligned}$$

with $\alpha \neq 0$ is a contact Lie algebra with $c(\mathfrak{g}) = (4, 3, 1, 1)$.

Let us note also, that in [27], we construct the contact nilpotent filiform Lie algebras, that is with characteristic sequence equal to $(2p, 1)$.

5.1.2 The non-nilpotent case

It is equivalent to consider Lie algebras with a contact form defined by a quadratic nonlinear deformations of the Heisenberg algebra. We refer to [19] to the description of this class of Lie algebras.

An interesting particular case consists to determine all the $(2p + 1)$ -dimensional Lie algebras ($p \neq 0$), such that all the coadjoint orbits of nontrivial elements are of dimension $2p$.

Lemma 5.9 ([15]). *Let \mathfrak{g} a simple Lie algebra of rank r and dimension n . Then any nontrivial linear form α on \mathfrak{g} satisfies*

$$\text{cl}(\alpha) \leq n - r + 1.$$

Moreover, if \mathfrak{g} is of classical type, we have

$$\text{cl}(\alpha) \geq 2r.$$

In particular, a simple Lie algebra admits a contact form if its rank is equal to 1 and this Lie algebra is isomorphic to $\mathfrak{sl}(2, \mathbb{R})$ or $\mathfrak{so}(3)$. From Proposition 2.9 these algebras are the only Lie algebras of dimension $2p$ or $2p + 1$ whose orbits $\mathcal{O}(\alpha)$ with $\alpha \neq 0$ are of dimension $2p$.

5.2 $2p$ -dimensional Lie algebras with a $2p$ -dimensional coadjoint orbit

Such Lie algebra is Frobenius (see [24]). Since the Cartan class of a linear form on a nilpotent Lie algebra is always odd, this Lie algebra is not nilpotent. In the contact case, we have seen that any contact Lie algebra is a deformation of the Heisenberg algebra. On other words, any contact Lie algebra can be contracted on the Heisenberg algebra. In the Frobenius case, we have a similar but more complicated situation. We have to determine an irreducible family of Frobenius Lie algebras with the property that any Frobenius Lie algebra can be contracted on a Lie algebra of this family.

In a first step, we recall the notion of contraction of Lie algebras. Let \mathfrak{g}_0 be a n -dimensional Lie algebra whose Lie bracket is denoted by μ_0 . We consider $\{f_t\}_{t \in]0,1]}$ a sequence of isomorphisms in \mathbb{K}^n with $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . Any Lie bracket

$$\mu_t = f_t^{-1} \circ \mu_0(f_t \times f_t)$$

corresponds to a Lie algebra \mathfrak{g}_t which is isomorphic to \mathfrak{g}_0 . If the limit $\lim_{t \rightarrow 0} \mu_t$ exists (this limit is computed in the finite-dimensional vector space of bilinear maps in \mathbb{K}^n), it defines a Lie bracket μ of a n -dimensional Lie algebra \mathfrak{g} called a contraction of \mathfrak{g}_0 .

Remark 5.10. Let \mathfrak{L}^n be the variety of Lie algebra laws over \mathbb{C}^n provided with its Zariski topology. The algebraic structure of this variety is defined by the Jacobi polynomial equations on the structure constants. The linear group $\text{GL}(n, \mathbb{C})$ acts on \mathbb{C}^n by changes of basis. A Lie algebra \mathfrak{g} is contracted to the \mathfrak{g}_0 if the Lie bracket of \mathfrak{g} is in the closure of the orbit of the Lie bracket of \mathfrak{g}_0 by the group action (for more details see [8, 14]).

5.2.1 Classification of complex Frobenius Lie algebras up to a contraction

Let \mathfrak{g} be a $2p$ -dimensional Frobenius Lie algebra. There exists a basis $\{X_1, \dots, X_{2p}\}$ of \mathfrak{g} such that the dual basis $\{\omega_1, \dots, \omega_{2p}\}$ is adapted to the Frobenius structure, that is,

$$d\omega_1 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_4 + \dots + \omega_{2p-1} \wedge \omega_{2p}.$$

In the following, we define Lie algebras, not with its brackets, but with its Maurer–Cartan equations. We assume here that $\mathbb{K} = \mathbb{C}$.

Theorem 5.11. *Let $\mathfrak{g}_{a_1, \dots, a_{p-1}}$, $a_i \in \mathbb{C}$ be the Lie algebras defined by*

$$\begin{aligned} d\omega_1 &= \omega_1 \wedge \omega_2 + \sum_{k=1}^{p-1} \omega_{2k+1} \wedge \omega_{2k+2}, & d\omega_2 &= 0, \\ d\omega_{2k+1} &= a_k \omega_2 \wedge \omega_{2k+1}, & 1 \leq k &\leq p-1, \\ d\omega_{2k+2} &= -(1 + a_k) \omega_2 \wedge \omega_{2k+2}, & 1 \leq k &\leq p-1. \end{aligned}$$

Then any $2p$ -dimensional Frobenius Lie algebra can be contracted in an element of the family $\mathcal{F} = \{\mathfrak{g}_{a_1, \dots, a_{p-1}}\}_{a_i \in \mathbb{C}}$. Moreover, any element of \mathcal{F} cannot be contracted in another element of this family.

Proof. See [16] and [19]. ■

Remark 5.12. The notion of principal element of a Frobenius Lie algebra is defined in [24].

In the basis $\{X_1, \dots, X_{2p}\}$ for which $d\omega_1 = \omega_1 \wedge \omega_2 + \sum_{k=1}^{p-1} \omega_{2k+1} \wedge \omega_{2k+2}$, the principal element is X_2 .

Proposition 5.13. *The parameters $\{a_1, \dots, a_{p-1}\}$ which are the invariants of Frobenius Lie algebras up to contraction are the eigenvalues of the principal element of $\mathfrak{g}_{a_1, \dots, a_{p-1}}$.*

5.2.2 Classification of real Frobenius Lie algebras up to contraction

We have seen that, in the complex case, the classification up to contraction of $2p$ -dimensional Lie algebras is in correspondence with the reduced matrix of the principal element. We deduce directly the classification in the real case:

Theorem 5.14. *Let $\mathfrak{g}_{a_1, \dots, a_s, b_1, \dots, b_{2p-s-1}}$, $a_i, b_j \in \mathbb{R}$ be the $2p$ -dimensional Lie algebras given by*

$$\begin{aligned} [X_1, X_2] &= [X_{2k-1}, X_{2k}] = X_1, & k &= 2, \dots, p, \\ [X_2, X_{4k-1}] &= a_k X_{4k-1} + b_k X_{4k+1}, & [X_2, X_{4k}] &= (-1 - a_k) X_{4k} - b_k X_{4k+2}, & k &\leq s, \\ [X_2, X_{4k+1}] &= -b_k X_{4k-1} + a_k X_{4k+1}, \\ [X_2, X_{4k+2}] &= b_k X_{4k} + (-1 - a_k) b_k X_{4k+2}, & k &\leq s, \\ [X_2, X_{4s+2k-1}] &= -\frac{1}{2} X_{4s+2k-1} + b_k X_{4k+2k}, & 2 &\leq k \leq p-2s, \\ [X_2, X_{4s+2k}] &= -b_k X_{4s+2k-1} - \frac{1}{2} X_{4s+2k}, & 2 &\leq k \leq p-2s. \end{aligned}$$

Then any $2p$ -dimensional Frobenius real Lie algebra is contracted in an element of the family $\mathcal{F} = \{\mathfrak{g}_{a_1, \dots, a_s, b_1, \dots, b_{2p-s-1}}\}_{a_i, b_j \in \mathbb{R}}$. Moreover, any element of \mathcal{F} cannot be contracted in other different element of this family.

Corollary 5.15. *Any $2p$ -dimensional Lie algebras described in (5.11) in the complex case and in (5.14) in the real case have an open coadjoint orbit of dimension $2p$. Moreover any $2p$ -dimensional Lie algebras with an open coadjoint orbit of dimension $2p$ is a deformation of a Lie algebra of these families.*

Acknowledgements

The authors would like to thank the referees for their kind advices and useful comments to improve the paper.

References

- [1] Adimi H., Makhlouf A., Index of graded filiform and quasi filiform Lie algebras, *Filomat* **27** (2013), 467–483, [arXiv:1212.1650](#).
- [2] Ancochéa-Bermúdez J.M., Goze M., Classification des algèbres de Lie nilpotentes complexes de dimension 7, *Arch. Math. (Basel)* **52** (1989), 175–185.
- [3] Arnal D., Cahen M., Ludwig J., Lie groups whose coadjoint orbits are of dimension smaller or equal to two, *Lett. Math. Phys.* **33** (1995), 183–186.
- [4] Awane A., Goze M., Pfaffian systems, k -symplectic systems, *Kluwer Academic Publishers*, Dordrecht, 2000.
- [5] Beltiță D., Cahen B., Contractions of Lie algebras with 2-dimensional generic coadjoint orbits, *Linear Algebra Appl.* **466** (2015), 41–63, [arXiv:1401.3272](#).
- [6] Beltiță I., Beltiță D., Coadjoint orbits of stepwise square integrable representations, *Proc. Amer. Math. Soc.* **144** (2016), 1343–1350, [arXiv:1408.1857](#).
- [7] Beltiță I., Beltiță D., On the isomorphism problem for C^* -algebras of nilpotent Lie groups, [arXiv:1804.05562](#).
- [8] Burde D., Degenerations of 7-dimensional nilpotent Lie algebras, *Comm. Algebra* **33** (2005), 1259–1277, [math.RA/0409275](#).
- [9] Cartan E., Les systèmes de Pfaff, à cinq variables et les équations aux dérivées partielles du second ordre, *Ann. Sci. École Norm. Sup. (3)* **27** (1910), 109–192.
- [10] de Graaf W.A., Classification of solvable Lie algebras, *Experiment. Math.* **14** (2005), 15–25, [math.RA/0404071](#).
- [11] Diatta A., Left invariant contact structures on Lie groups, *Differential Geom. Appl.* **26** (2008), 544–552, [math.DG/0403555](#).
- [12] Duflo M., Vergne M., Une propriété de la représentation coadjointe d’une algèbre de Lie, *C. R Acad. Sci. Paris Sér. A-B* **268** (1969), A583–A585.
- [13] Godbillon C., Géométrie différentielle et mécanique analytique, Hermann, Paris, 1969.
- [14] Goze M., Algèbres de Lie de dimension finie, Ramm Algebra Center, available at <http://ramm-algebra-center.monsite-orange.fr>.
- [15] Goze M., Sur la classe des formes et systèmes invariants à gauche sur un groupe de Lie, *C. R. Acad. Sci. Paris Sér. A-B* **283** (1976), A499–A502.
- [16] Goze M., Modèles d’algèbres de Lie frobeniusiennes, *C. R. Acad. Sci. Paris Sér. I Math.* **293** (1981), 425–427.
- [17] Goze M., Bouyakoub A., Sur les algèbres de Lie munies d’une forme symplectique, *Rend. Sem. Fac. Sci. Univ. Cagliari* **57** (1987), 85–97.
- [18] Goze M., Haraguchi Y., Sur les r -systèmes de contact, *C. R. Acad. Sci. Paris Sér. I Math.* **294** (1982), 95–97.
- [19] Goze M., Remm E., Contact and Frobeniusian forms on Lie groups, *Differential Geom. Appl.* **35** (2014), 74–94.
- [20] Goze M., Remm E., k -step nilpotent Lie algebras, *Georgian Math. J.* **22** (2015), 219–234, [arXiv:1502.05016](#).
- [21] Hatipoglu C., Injective hulls of simple modules over nilpotent Lie color algebras, [arXiv:1411.1512](#).
- [22] Khuhirun B., Misra K.C., Stitzinger E., On nilpotent Lie algebras of small breadth, *J. Algebra* **444** (2015), 328–338, [arXiv:1410.2778](#).
- [23] Kirillov A.A., Elements of the theory of representations, *Grundlehren der Mathematischen Wissenschaften*, Vol. 220, Springer-Verlag, Berlin – New York, 1976.
- [24] Ooms A.I., On Frobenius Lie algebras, *Comm. Algebra* **8** (1980), 13–52.
- [25] Pinczon G., Ushirobira R., New applications of graded Lie algebras to Lie algebras, generalized Lie algebras, and cohomology, *J. Lie Theory* **17** (2007), 633–667, [math.RT/0507387](#).
- [26] Remm E., Breadth and characteristic sequence of nilpotent Lie algebras, *Comm. Algebra* **45** (2017), 2956–2966, [arXiv:1605.06583](#).
- [27] Remm E., On filiform Lie algebras. Geometric and algebraic studies, *Rev. Roumaine Math. Pures Appl.* **63** (2018), 179–209, [arXiv:1712.00318](#).