

## Snow season variability in a boreal-Arctic transition area monitored by MODIS data

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 Environ. Res. Lett. 11 125005

(<http://iopscience.iop.org/1748-9326/11/12/125005>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 158.38.175.146

This content was downloaded on 30/11/2016 at 07:38

Please note that [terms and conditions apply](#).

You may also be interested in:

[Impacts of snow season on ground-ice accumulation, soil frost and primary productivity in a grassland of sub-Arctic Norway](#)

Jarle W Bjerke, Hans Tømmervik, Matthias Zielke et al.

[Record-low primary productivity and high plant damage in the Nordic Arctic Region in 2012 caused by multiple weather events and pest outbreaks](#)

Jarle W Bjerke, Stein Rune Karlsen, Kjell Arild Høgda et al.

[Satellite observed changes in the Northern Hemisphere snow cover phenology and the associated radiative forcing and feedback between 1982 and 2013](#)

Xiaona Chen, Shunlin Liang and Yunfeng Cao

[Modeling the influence of snow cover on low Arctic net ecosystem exchange](#)

K A Luus, R E J Kelly, J C Lin et al.

[Changes in growing season duration and productivity of northern vegetation inferred from long-term remote sensing data](#)

Taejin Park, Sangram Ganguly, Hans Tømmervik et al.

[Consequences of changes in vegetation and snow cover for climate feedbacks in Alaska and northwest Canada](#)

E S Euskirchen, A P Bennett, A L Breen et al.

[High Arctic plant phenology is determined by snowmelt patterns but duration of phenological periods is fixed: an example of periodicity](#)

Philipp R Semenchuk, Mark A K Gillespie, Sabine B Rumpf et al.

## Environmental Research Letters



## LETTER

## Snow season variability in a boreal-Arctic transition area monitored by MODIS data

## OPEN ACCESS

## RECEIVED

1 June 2016

## REVISED

25 October 2016

## ACCEPTED FOR PUBLICATION

10 November 2016

## PUBLISHED

29 November 2016

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.


Eirik Malnes<sup>1,3</sup>, Stein Rune Karlsen<sup>1</sup>, Bernt Johansen<sup>1</sup>, Jarle W Bjerke<sup>2</sup> and Hans Tømmervik<sup>2</sup><sup>1</sup> Northern Research Institute (Norut), PO Box 6434 Forskningsparken N-9294 Tromsø, Norway<sup>2</sup> Norwegian institute for nature research (NINA), FRAM—High North Research Centre for Climate and the Environment, PO Box 6606 Langnes, N-9296 Tromsø, Norway<sup>3</sup> Author to whom any correspondence should be addressed.E-mail: [eirik@norut.no](mailto:eirik@norut.no), [stein-rune@norut.no](mailto:stein-rune@norut.no), [bernt.johansen@norut.no](mailto:bernt.johansen@norut.no), [jarle.bjerke@nina.no](mailto:jarle.bjerke@nina.no) and [Hans.Tommervik@nina.no](mailto:Hans.Tommervik@nina.no)**Keywords:** snow, MODIS, first snow free day, last snow free day, climate change, onset of springSupplementary material for this article is available [online](#)**Abstract**

The duration and extent of snow cover is expected to change rapidly with climate change. Therefore, there is a need for improved monitoring of snow for the benefit of forecasting, impact assessments and the population at large. Remotely sensed techniques prove useful for remote areas where there are few field-based monitoring stations. This paper reports on a study of snow season using snow cover area fraction data from the two northernmost counties in Norway, Troms and Finnmark. The data are derived from the daily 500 m standard snow product (MOD10A1) from the NASA Terra MODerate Resolution Imaging Spectroradiometer (MODIS) sensor for the 2000–2010 period. This dataset has been processed with multi-temporal interpolation to eliminate clouds. The resulting cloud-free daily time series of snow cover fraction maps, have subsequently been used to derive the first and last snow-free day for the entire study area. In spring, the correlation between the first snow-free day mapped by MODIS data and snow data from 40 meteorological stations was highly significant ( $p < 0.05$ ) for 36 of the stations, and with a bias of less than 10 days for 34 of the stations. In autumn, 31 of the stations show highly significant ( $p < 0.05$ ) correlation with MODIS data, and the bias was less than 10 days for 27 of the stations. However, in some areas and some years, the start and end of the snow season could not be detected due to long overcast periods. In spring 2002 and 2004 the first snow-free day was early, but arrived late in 2000, 2005 and 2008. In autumn 2009 snowfall arrived more than 7 days earlier in 50% of the study area as compared to the 2000–2010 average. MODIS-based snow season products will be applicable for a wide range of sectors including hydrology, nature-based industries, climate change studies and ecology. Therefore refinement and further testing of this method should be encouraged.

**1. Introduction**

The duration of the snow-free season controls the rhythm of ecosystems and feedback to the climate system. This is particularly the case in boreal and Arctic-alpine areas, characterized by a short and intensive snow-free season. Changes in the timing of snowmelt in spring may influence the timing of the growing season and thereby the growth, seed production and competition between plants, and is therefore

the first indication of changes in plant diversity and vegetation cover (Wipf 2010, Wipf and Rixen 2010, Cooper *et al* 2011). Changes in snow season may influence the migration patterns and survival rate for birds, mammals, and insects at high northern latitudes (Kausrud *et al* 2008) including reindeer husbandry (Riseth *et al* 2011). Moreover, unseasonal snow thaw caused by midwinter warming events often have severe impacts on plants and animals (Bjerke *et al* 2014, Hansen *et al* 2014). Bokhorst *et al* (2016) provide a



recent interdisciplinary review on advances in snow monitoring and modeling, and the impact of snow changes on ecosystems and society in Arctic regions.

Arctic and boreal temperatures are increasing due to positive feedbacks (e.g., albedo-temperature feedback) of increased greenness and shrubification (Xu *et al* 2013) and this is also linked to a shortening of the winter season. Hence, the exact timing of the onset and disappearance of snow is a critical parameter within a range of climatic and ecological studies. A series of methods have been developed to detect the start and end of the snow season. Hall *et al* (2002) used the normalized differential snow index (NDSI) based on Landsat and MODerate Resolution Imaging Spectroradiometer (MODIS) data, while Xu *et al* (2013) used daily records of freeze-thaw data obtained from passive-microwave remote sensing using the scanning multichannel microwave radiometer and special sensor microwave imager sensors with a spatial resolution of 25 km × 25 km in order to monitor the snow- and ice-free season (photosynthetic active period). Liston and Hiemstra (2011) studied changes in the Pan-Arctic snow cover in the period 1979–2009 using a 10 km dataset based mainly on atmospheric reanalysis data, and found that the number of snow days per year decreased on average by 2.5 days per decade. This trend is supported by many similar studies referred to in SWIPA (2011). Dietz *et al* (2012) presented an inventory of snow cover characteristics based on MODIS Terra and Aqua for Europe for the period 2000–2011.

Within the study region (see section 2.1) there exists two operational services that provide regional snow cover data based on satellite and snow services. These are based on hydrological models provided by the hydrological authorities in Norway, Sweden and Finland (see e.g. senorge.no in Saloranta 2012). The

satellite-based services are provided by SYKE (The Finnish Environment Institute) in Finland and Globesar in Norway. The SYKE snow map (Metsämäki *et al* 2012) is based on Terra MODIS data whereas the Globesar service is based on a multi-sensor snow map algorithm using Terra MODIS and Sentinel-1 data (Malnes *et al* 2005, Solberg *et al* 2010). This service was developed by Norut and the Norwegian Computer Center in Norway. Common to both these products is that they only provide data for the melting season (1 April–31 July) and focus on near-real time data for hydrological applications. Due to the near-real time approach, it is much harder to provide cloud gap-filled data sets than when data can be processed afterwards, as is the approach in this study. The main objective of this study is therefore to develop a method for detecting the first and last snow-free day geographically. To assess the accuracy of the method, we compare the outputs with the first and last snow-free days as measured at all meteorological stations within the study area. Finally, we discuss some applications of the dataset.

## 2. Data and methods

### 2.1. Study area

The study area covers the two northernmost counties in Norway, Finnmark and Troms, and nearby areas in Sweden, Finland, and Russia. It extends from approximate 68.1°N, 16.2°E to 70.8°N, 31.8°E (figure 1). A mountain range divides the area into two main climatic regions; (1) a western oceanic part, characterized by high precipitation rates and mild winters and (2) an eastern part characterized by drier and colder winters. The lowland in the western part is dominated by deciduous birch forest whereas in the southeastern

part, lowland areas are covered by coniferous forest. However, the forest line is mostly below 500 m in the south and can reach sea level in the north, leaving large areas with arctic-alpine treeless tundra, within the mountain range and in the north (Moen 1999).

The vegetation in the study area is sparse. The dominating tree species along the coast are sparse birch (*Betula pubescens*) forests which are among the northernmost forests in the world (Wielgolaski and Sonesson 2001). The inner parts of Troms and central northern Finnmark county (Norway) host the world's northernmost Scots pine forests (Wielgolaski and Sonesson 2001). In the north-eastern part, arctic climate conditions (period: 1961–1990), prevail with an average temperature for the warmest summer month July of  $<10^{\circ}\text{C}$  (Tveito *et al* 2000). The lowest monthly winter temperatures are found in the inland part of Finnmark ( $-16^{\circ}\text{C}$ ). Precipitation ranges from around 300 to 500 mm year (Tveito *et al* 2001) in Finnmark and Finland with rather low monthly winter precipitation (15–20 mm) in the continental parts of Finnmark and Finland (total winter precipitation of 87 mm). On the coast in Troms county, however, the monthly winter precipitation can be more than 100 mm. The snow cover is deepest around mid-March, with an average snow depth of 60–90 cm in eastern and northern Finland and at the coast of Troms County. On an annual basis, and especially during winter, inland Finnmark is the coldest and driest area amongst the Nordic countries (Tveito *et al* 2000).

## 2.2. MODIS data processing

This study uses the NASA MODIS/Terra Snow Cover Daily L3 Global 500 m Grid, Version 5 (MOD10A1) for the 2000–2010 period. The product provides snow cover fraction (SCF in the range 0 to 100%) as daily products (Hall *et al* 2002, Riggs *et al* 2006). SCF is calculated from the NDSI, from MODIS bands 4 (545–565 nm) and 6 (1628–1652 nm) using the formula,

$$\begin{aligned} \text{SCF} &= (0.06 + 1.21 \times \text{NDSI}) \\ &\times 100\% \text{ where NDSI} \\ &= (\text{band 4} - \text{band 6}) / (\text{band 4} + \text{band 6}). \end{aligned} \quad (1)$$

The product masks out areas that are covered by clouds, ocean and winter darkness. We converted the product to WGS 1984 UTM Zone 33 N and performed a refinement by applying multi-temporal interpolation techniques to obtain a cloud-free daily time series of snow maps for the study area. In total we have processed daily data sets from 24 February, 2000 (DOY 55) to 1 November, 2010 (DOY 304). With the exceptions of a few days (particularly in 2001, DOY 166–183) we have daily datasets for the whole period. For the period of polar night in the study area from 15–22 November to 20–30 January, we have no information and pixels corresponding to this period are flagged in the MODIS-products. We have, however,

experienced that the products probably underestimate the snow cover fraction during the period before the onset of polar night (1 November–20 November) and during the period after the Sun returns (21 January–21 February). All snow maps have thus been ‘padded’ with 100% snow cover fraction in this period (21 November–20 February) to avoid confusion and create more likely results.

We have used a land mask for the region based on the N50 standard Norwegian map ([www.kartverket.no](http://www.kartverket.no)) and maps of similar quality from Sweden and Finland to better delineate ocean, lakes, rivers and glaciers than can be done using the masks provided by NASA. This dataset has been further processed with multi-temporal interpolation to eliminate clouds (SI: 1.1 *Multitemporal interpolation method*). For each day in the period we made an SCF map by mosaicking, re-projecting and resampling the MOD10A1 data to the fixed grid in UTM WGS z33N projection (see SI: figure S3).

Gap-filling techniques are frequently used in remote sensing data to mitigate lack of data due to clouds or other reasons (darkness, lack of coverage and instrument failure). Several authors (e.g. Høgda *et al* 2013) use the best observations in 7 or 8 day periods to find the most reliable data points. This strategy significantly alleviates the problems due to cloud cover significantly, but at the expense of a coarser temporal resolution. Our strategy maximizes the number of observations, but increases the risk for including noisy samples in the dataset. Such noise can for example occur in the vicinity of clouds. Since our study has a focus on detecting the first and the last snow-free day, we are confident that our approach reduces the sampling bias and increases the temporal accuracy for detection of the first and last snow free days.

Hall *et al* (2010) reviewed several cloud mitigation strategies for the MODIS snow products (see references therein) and also developed their own approach. Several approaches use both MODIS Terra and Aqua to provide an increased number of daily views, in combination with using temporal filtering/interpolations of observations from the days before. Spatial filtering has also been applied.

Foppa and Seiz (2012) and Hüsler *et al* (2014) applied forward/backward gap-filling techniques for MODIS and AVHRR snow maps over the Alps, respectively. They used the average of the closest observation before and after the gap, instead of using a weighted mean (depending on the number of days before and after) as we do in this paper. The advantage of their approach is that it might be more tolerant to noise in the closest observation, in particular during stable snow conditions. The disadvantage is that the temporal information is smeared and the uncertainty in estimates of the transition time between snow-covered and snow-free conditions will increase.



**Table 1.** Rules for detection of first and last snow free days.

Rule	Start, day	End, day	Threshold	Nrule	Age
MODIS, first snow free day	75	230	50%	10	15
Snow depth, first snow free day	75	230	2 cm	10	15
MODIS, last snow free day	264	310	50%	5	15
Snow depth, last snow free day	264	310	2 cm	5	15

### 2.3. Detection of the first and last snow free day

To select the best threshold for detecting the first and last snow-free day we use snow depth data from meteorological stations (see figure 1(b)), downloaded from [www.eklima.no](http://www.eklima.no). Due to changes from manual to automatic measurements at some of the stations, and that the automatic stations often do not provide snow depth data, the number of stations with snow data were reduced from 40 in 2000 to 33 in 2010.

The mean SCF values were calculated for a  $3 \times 3$  pixel area centered on each of the meteorological stations. The day when SCF passes below or above the threshold set at  $SCF = 50\%$  for an extended period (more than 10 days in the spring and more than 5 days in the fall) is set as the first and last snow-free day respectively. We also require that this can only happen in the period from DOY 75 (15 March) to DOY 230 (15 August) for the first snow-free day and in the period from DOY 264 (20 September) to DOY 310 (5 November) for the last snow-free day. These days are chosen to avoid problems due to low solar angles (in March and November) and confusion with random, short-lived snow falls during summer/early autumn. The detection rules are summarized in table 1.

In addition to the rules mentioned above, we also decided not to make a detection of first/last snow-free day for a given pixel if the age of the day for first/last snow free is above the threshold set at 15 days. It was found experimentally in the validation section (section 3.4) that a higher threshold led to increased bias/standard deviation in the first/last snow estimates. Likewise, a lower threshold led to fewer years (in particular autumns), where we were able to estimate the first/last snow.

The rules for detecting the first snow-free day and first snow-covered day in the meteorological dataset is very similar to the rules for satellite data, and was adapted from Vikhamar-Schuler and Hanssen-Bauer (2010). The only exception is that we now set the threshold to 2 cm snow depth. We have found that by systematically varying the threshold that 2 cm results in a minimum for the overall bias in the estimates for first and last snow cover in the two datasets.

These methods are illustrated for the meteorological station Kautokeino in figure 2 and S2. In figure 2, we show a comparison between SCF and snow depth at the Kautokeino meteorological station. Based on a total number of 43 cloud-free SCF detections during the year 2004, we were able to estimate

the first and the last snow-free day with 1–2 days bias in comparison to the snow depth. The middle panel shows age, i.e. the number of days between the estimated SCF and an actual cloud-free measurement of SCF and this is used as an error estimate throughout this paper.

### 2.4. Development of daily snow maps

For each day in the period we made an SCF map by mosaicking, re-projecting and resampling the MOD10A1 data to the fixed grid in UTM WGS z33N projection (see figure S3(a)). In general, the maps have a high cloud fraction (see figure S3(a)). By applying the multi-temporal interpolation method discussed in 2.4, we obtained cloud-free snow maps for the study area. The method described above is now applied to the complete dataset consisting of 4015 ( $11 \times 365$ ) snow maps. For further information about development of daily snow maps see also SI, section 1.3.


### 2.5. Annual maps for first and last day of snow cover and the length of the snow-free season

Based on the methods mentioned above and the multi-temporal MODIS dataset we are now able to provide maps for the first snow-free day in the spring and the last snow-free day in the autumn. In figure 3 we show an example for 2006. Each pixel in the maps is coded with the first/last day of snow. We can also calculate the pixel age, or uncertainty in this estimate of first/last snow cover. Finally, we have also calculated the number of snow-free days per snow season and the number of cloud-free days per year.

## 3. Results

### 3.1. Snow season trends and variability

Snow seasons varied greatly between different years, both with regards to start, end and duration (figure 4). As expected, the alpine areas had much longer seasons than the lowlands. For instance the first snow-free day was delayed for the years 2000, 2005, 2007 and 2008. In contrast, the first snow-free day appeared to arrive very early in the years 2002, 2006 and 2010. The last snow-free day was delayed for the years 2000, 2003 and 2006, while for the years 2001, 2002 and 2009 the snow arrived earlier. The length of the winter season (snow season) was longest for the winters 2004–2005 and 2009–2010 while the longest snow-free seasons (summer season) were observed for the summers 2001,



**Figure 2.** (a) Remotely sensed SCF compared with ground measurements of snow depth for the Kautokeino meteorological station in 2004. Each cloud free detection is also marked with black diamonds, the red lines show detection windows for first and last day of snow.  $\Delta T$  is the time difference between snow transition detected by MODIS and met-station. (b) Pixel age, number of days since last cloud free observation. (c) Snow depth measured at the meteorological station. Triangles show the detected first-and last snow-free day, and vertical dashed lines show the difference between the two estimates.

2002 and 2007. In contrast, the length of the winter season (snow season) was shortest for the winters 2003–2004 and 2008–2009 while the shortest snow-free season (summer season) was observed for the summer 2005. The summer season (snow-free period) is shortest in the mountains and on the Varanger Peninsula (see figure 1(b)) and longest in the forested continental regions (south-east) and along the coast.

### 3.2. Average maps for first and last day of snow cover and the length of the snow free season

After obtaining the first and last day of snow cover for each year, we can also find the average for the first and last day of snow cover. These estimates are shown in figure 5 together with the overall estimate for the length of the snow-free season. As we can observe from figure 5, the coastal parts and the river valleys as well as the more forested areas in Sweden and Finland experience a quite early start of the snow-free period (spring) as well as late start of the snow season in the autumn/winter. Hence, the snow-free period is longest in these parts of the study area.


### 3.3. Annual variability in the snow cover

We have applied the results from sections 3.3 and 3.4 to study the annual variability in the snow cover for the

entire study area and the geographical variability per year. In figure 6, we show the total daily SCF per day for the entire 11 year time series. We also show the mean value for all the years. By comparing the year to year variation we can observe that the start of the snow-free periods in spring 2000 and 2005 (DOY 135) were late while the year 2002 and 2006 had early starts (DOY 105). In the autumn/start of winter; year 2000 and year 2007 had a late start (DOY 290) of the snow season, whereas the first snow fall came early in 2009 (DOY 270).

### 3.4. Assessment of the MODIS based snow cover products


Table 2 shows the relationship between the first and last day of snow cover measured from MODIS data and from the meteorological stations. The number of years with data is less than 11 for most of the stations due to either lack of MODIS data (due to clouds/polar darkness) or lack of snow depth data from the meteorological stations. In spring, the correlation ( $r = 0.79$ ) between the MODIS-based and the meteorological measurements of the last day of snow cover was very high, where 31 of the 40 stations show significant values at the 1% level or better. Only four of the stations did not show significant correlation



( $p > 0.05$ ). The average standard deviation is six days. Only six of stations show a bias of 10 days or more, where the MODIS-based measurements were earlier than the meteorological measurements in these cases. The relationship between MODIS-based and meteorological measurements of the first day with snow in autumn was weaker ( $r = 0.51$ ) compared with the spring period. On average only 6 years of data were available for the stations. However, 27 of the stations show significant values at the 1% level or better, and the average standard deviation is 7 days. Three of the stations have a bias of 10 days or more and for all of these stations the MODIS-based measurements of the first day of snow were earlier than the meteorological measurements. Only 2 of 40 stations have a correlation of less than 0.5 for the first snow-free day. The standard deviations and bias were also higher for the

end of the growing season than for the spring. The reason for this can be found in the number of years applied. On average, we use only 7 of the 11 available years to estimate the last snow-free day. The reason for this is mostly due to late first snowfall in the autumn, which yields a lower probability of detecting the transition. Also, some stations lack snow depth measurements for some of the years. Some of the stations are located in challenging terrain as the stations are often localized close to the lowest point in the  $3 \times 3$  pixel area used for estimating SCF or in coastal areas with high cloud cover probability (e.g. Gamvik, Ytre Holmbukt, Sørkjosen). The Sørkjosen station ( $r = 0.25$ ) also has a high standard deviation and an inspection of this site indicates that the spring of 2006 and 2009 were exceptions, leading to large errors. The snow depth record from 2006 is







questionable since the snow melts from 70 to 0 cm in 1 day on 1 April. The 2009 spring had a normal snowmelt followed by a cold period where the snow

depth was a few centimetres above the threshold. If these two years are excluded, the correlation and SD improves ( $r = 0.97$ ,  $SD = 2.3$ ).





**Figure 5.** Average time of the first (upper left) and last snow-free day (upper right), the length of snow-free season (lower left), and length of the winter season (lower right) for the period 2000–2010.



**Figure 6.** Daily total snow cover fraction for study area per year. The average daily SCF for the period 2000–2010.

Table 2 also shows that the correlation between the MODIS estimate and the snow depth estimate of the last snow-free day is significantly poorer ( $r = 0.65$ ). The average values indicate that the first snow-free day for the entire area is approximately 1 May and that the last snow-free day is around 15 October. Due to the bias in the distribution of meteorological stations, these values are applicable for the coastline.

## 4. Discussion

### 4.1. Critical discussion on methods and results

We are confident that the methods presented here give reasonable results for the estimation of first and last

snow-free day. The time sequences of cloud-free (multi-temporal) snow cover maps as well as the annual maps for first and last snow-free day are reasonable at the scale we are studying (500 m resolution). The quality of the MOD10A1 product is questionable (e.g. Metsämäki *et al* 2015). However, the overall absolute clear-sky accuracy of the well-studied 500 m resolution swath (MOD10\_L2) and daily tile (MOD10A1) products was assessed to be  $>93\%$ , but varies with land-cover type and snow condition (Hall and Riggs 2007). The cloud mask in the MOD10A1 product is probably too conservative most of the time, and we observed frequently that the cloud cover was overestimated. Since the MOD10A1-algorithm was optimized in order to perform on a global scale, this is

**Table 2.** Validation result per station for first and last snow free day. The table show the average first and last day of snow (MOD = MODIS based, MET = based on snow depth) for each station. We also show the correlation  $r$ , the standard deviation SD and the bias BD. The bottom row gives the averages for all stations and years.

Station	Altmasl.	#y	Spring Day			Autumn Day					
			MOD/MET	$r$	SD	BD	#y	MOD/MET	$r$	SD	BD
Bardufoss	76	11	119/113	0.97	2	-6	8	287/290	0.93	4	3
Bones	230	10	126/125	0.98	1	-1	7	292/296	0.86	5	4
Botnhamn	10	11	131/127	0.97	3	-3	8	287/289	0.46	10	2
Børselv	13	9	123/125	0.98	4	2	5	290/294	-0.14	11	4
Čuovddatmohkki	286	11	132/129	0.74	7	-2	9	288/287	0.33	10	-1
Dividalen	228	9	115/122	0.87	4	7	7	295/297	0.57**	5	2
Gamvik	15	10	130/121	0.52**	10	-8	8	286/296	0.35	11	10
Grunnfarnes	3	9	116/110	0.82	8	-6	7	286/292	0.68	7	6
Grunnfjord	7	10	119/114	0.96	4	-4	4	290/291	0.62	2	1
Guolasjohka	50	9	117/103	0.21**	19	-14	2	291/300	0.62	15	9
Hasvik	6	6	119/114	0.89	5	-4	3	287/290	0.99*	5	3
Jergol	230	6	128/127	0.98	4	0	5	293/293	0.62	9	0
Jotkajávri	389	7	146/139	0.84	5	-6	7	283/286	0.42	12	2
Karasjok	129	10	121/124	0.97	2	2	8	292/293	0.40	11	0
Kautokeino	307	11	129/129	0.77	6	0	8	291/290	0.19	12	-1
Kistrand	14	10	128/124	0.89	4	-4	7	291/295	0.70*	6	4
Kunes	22	7	123/122	0.94	3	-1	5	287/293	0.76	6	5
Lebesby	18	11	128/126	0.55*	12	-1	7	292/294	0.95	5	1
Lyngseidet	18	11	115/110	0.78	8	-5	6	286/290	0.92	4	3
Mollesjohka	382	11	131/127	0.88	6	-3	7	289/289	0.91	4	0
Målselv	3	8	139/140	0.81	6	0	7	286/285	0.64*	10	0
Nordstraum	6	10	107/112	0.94	7	5	4	298/291	-0.52	8	-7
Polmak	18	11	128/120	0.73	6	-8	8	291/294	0.92	4	3
Porsa	38	11	133/129	0.66*	12	-4	7	282/292	0.10**	12	9
Port	115	11	130/130	0.97	2	0	8	291/298	0.19**	8	7
Rustefjelbma	10	9	130/124	0.51**	7	-5	7	290/292	0.85	5	1
Setermoen2	114	11	126/123	0.85	5	-3	7	288/288	0.96	2	0
Senja-Laukhella	9	11	120/118	0.86	6	-2	6	289/288	0.89	3	-1
Sihččajávri	382	11	135/137	0.86	6	2	10	287/293	0.78	6	5
Sirbma	51	11	128/118	0.53*	8	-9	8	289/296	0.31**	9	6
Skaidi	62	9	142/131	0.81	6	-11	6	278/289	0.45**	10	11
Skallelv	16	9	125/127	0.83	5	2	3	286/284	0.79	6	-2
Skibotn	45	9	112/100	0.85	6	-11	2	284/291	-1.00	6	7
Skjervøy	5	11	136/127	0.81	7	-9	7	283/289	0.59**	6	5
Sopnesbukt	8	4	125/122	0.86	5	-3	2	279/280	1.00**	1	1
Storsteinnes	27	11	141/126	0.60*	10	-14	4	279/292	-0.47	16	12
Suolovuopmi	377	11	125/122	0.99	1	-3	4	289/291	0.99	1	1
Sørkjosen	6	10	139/134	0.25**	23	-4	8	287/288	0.71*	7	1
Tromsø	100	11	123/126	0.85	6	3	5	288/287	0.55**	13	-1
Valjok	132	6	128/128	0.99	2	0	5	289/294	0.97	5	5
Veidnes	6	10	126/113	0.76	8	-12	7	290/296	0.89	2	5
Vestre-Jakobselv	11	11	123/112	0.79	8	-10	5	292/297	0.79**	4	4
Ytre Holmebukt	25	6	123/113	0.68*	8	-9	4	283/287	0.13	9	4
<b>Mean values</b>	<b>93</b>	<b>9</b>	<b>126/122</b>	<b>0.79</b>	<b>6</b>	<b>-3</b>	<b>6</b>	<b>288/291</b>	<b>0.51</b>	<b>7</b>	<b>3</b>

Unmarked correlation values have significance at the 1% level, \* significance at 5% level, \*\* insignificant result.

expected. Other algorithms have improved the performance for forested areas (Vikhamar and Rune Solberg 2003, Metsämäki *et al* 2012). We also observed occasionally falsely detected snow during summer (related to erroneous cloud classification) and falsely detected areas with low snow cover fraction during the winter period (related to poor light conditions and uncompensated topography).

The method we have developed for detecting first and last day of snow cover could also be applied on improved SCF-time series (e.g. Globesnow2-data, Metsämäki *et al* 2015), but we suspect that the results

will only improve slightly since our method is robust against many of the existing problems in the MOD10A1 products. Problems with conservative cloud estimates are handled by the multi-temporal interpolation scheme. Effects associated with falsely detected snow during summer and too low SCF in mid-winter will seldom be a problem as we have developed detection rules that excludes such detections in the majority of cases.

The linear interpolation approach may also be questioned. It is a well-known fact that the snow depletion curve is seldom linear in studies where mid-

to large-scale hydropower basins with a large altitude gradient are studied (Kolberg and Gottschalk 2010). On the scale we study however, the snow depletion curve will in most cases approach a sigmoid function with a rather steep slope. The steeper the slope, the more important it becomes to have cloud free SCF estimates around the timing of the snow melt. This problem could be alleviated by including SCF estimates from synthetic aperture radars (SAR) in the multi-temporal time series. Malnes *et al* (2005) have developed methods to fuse SAR and optical time series, but the density of SAR time series is seldom as consistent as optical datasets.

We experienced in our study (table 2) that the average correlation between the MODIS estimate and the snow depth estimate of the last snow-free day was significantly poorer ( $r = 0.51$ ) than the correlation for the first snow-free day ( $r = 0.79$ ). We do not attempt to extract trends from the current dataset. An 11 year dataset is too short, and it can easily be seen from e.g. Figure 4 that the individual years with alternating dominating regional patterns of early and late snow cover will dominate any trends at the current stage. Future studies will attempt to use the whole time series of MODIS data (2000–2016) and might provide some more clarity concerning observable trends.

The validation provided in table 2 gives overall trustworthy values for correlation and significance. We may thus state that our methods developed to assess the first and the last dates of snow cover are overall trustworthy. It should however, be emphasized that the distribution of meteorological stations used for the validation are strongly biased towards the coastline and low altitudes.

A weak point in our validation assessment is the method we use for removing outliers. Since MOD10A1 data gives little to no information following the onset of polar darkness, we are forced to remove all winters with late onset of snow. This yields a clear bias towards the years when the onset of snow is early, and as long as we do not have better sensors to capture late snowfalls this will be a clear limitation. However, we believe that the information provided is better than no information, and we thus choose to provide it.

#### 4.2. Ecological interpretation

The averaged map (2000–2011) for the length of the snow-free season can be compared with phenological maps showing the extent of the growing season. The length of the snow-free season is a good proxy for the growing season (Karlsen *et al* 2009) and the photosynthetic active period (Xu *et al* 2013). We find similarities between the map in figure 5 and the maps produced by Karlsen *et al* (2009), Høgda *et al* (2013) and Xu *et al* (2013). However, the start and the end of the growing season cannot be compared exactly since vegetation activity is included in the phenological maps (Høgda

*et al* 2013). In our approach, we detect the start and end of the snow season, hence there can be several days/weeks difference from when the snow-free period starts to when vegetation green-up starts (e.g. bud burst of downy birch). Similarly in autumn, there may be several days/weeks from when the vegetation becomes yellow/brown or leaves have fallen until the start of the snow season. However, it is clear that the map of the duration of the snow-free season (figure 5) has provided valuable information in order to detect the extent of the growing season as well as data for assessing and modeling the reindeer range conditions (Tømmervik 2005, Riseth *et al* 2011). For example the late start of the snow-free period and large snow depth in year 2000 led to starvation of the reindeer in April and May with considerable loss of calves and adults in Finnmark as a result (Tømmervik *et al* 2009, Tveraa *et al* 2013). Tveraa *et al* (2013) analyzed 10 years of MODIS data in order to analyze the effect of vegetation green-up on 19 semi-domesticated reindeer populations in Finnmark County, northern Norway. An early onset and high peak plant productivity had a positive influence on calf autumn body weights and female reproductive success (Tveraa *et al* 2013).

Data on late snowfall in the autumn detected by this method can together with climate data (temperature and precipitation) reveal likelihoods for injury or damage to the vegetation (Bokhorst *et al* 2012, Bjerke *et al* 2014) during the following winter. Mid-winter rain on snow events which result in snow-free areas may lead to damage to the vegetation (Bjerke *et al* 2014). Such events can also be detected by this method if cloud and light conditions are favorable.

In addition to differences in the snow-free season due to land cover (vegetation type), we also notice that height gradient and continentality are major factors affecting the length of the snow-free season and this is consistent with Karlsen *et al* (2009). We find that the longest lasting snow-free areas are located along the coast and in the river valleys as well as in the areas around the Inari lake (Finland), while the shortest snow-free periods are found in the mountains and in the north-eastern parts of Finnmark (Varanger peninsula), and here we also find the sparsest vegetation (Karlsen *et al* 2009).

#### 4.3. Hydrological interpretation

Daily snow maps have been used in assimilation with lumped hydrological models (Engeset *et al* 2003) with limited success, mainly due to the problems with over-calibration against run-off. The authors emphasize the importance of timely snow maps in the melting season when snow reservoirs are uncertain. In particular flood warnings can be issued in such a context. When used in conjunction with available models, snow cover data based on MODIS measurements may improve the runoff forecast significantly when assimilated in



distributed hydrological models for Snake River basin in USA (Andrealis and Lettenmaier 2006).

#### 4.4. Climate interpretation and other application areas

The current study provides detailed maps for the length and variability of the snow season in the study area. Similar studies using MODIS have also provided maps at a coarser scale for the northern hemisphere (Chen *et al* 2015) and Europe (Dietz *et al* 2012). The current dataset is not sufficiently long to derive trends, but the spatial resolution is sufficiently high to show many regional details which can be related to the complex interaction between large scale weather patterns affecting the coastal climate in northern Norway, the topography and the continental sub-arctic climate zones in the interior of the land. The overall picture, where the length of the snow season is one of several factors that determine the living conditions for many plants, show remarkable resemblance to vegetation maps for the same region. Regions of similar snow phenology also have similar vegetation covers. This shows that the maps, when updated on a regular basis, could be used to determine where there are significant and long-lasting deviations from the average that also could indicate ongoing climate changes.

### 5. Conclusions

We have described a method for generating long-term, cloud-free daily time series of the snow cover fraction based on optical medium resolution sensors. The method is applied to the MOD10A1 dataset to provide an 11 year record for our study area in northern Norway. The time series of snow maps shows a convincing development of the snow cover that can be used for generating climatological records as well as in ecological monitoring and assessments. We have calculated annual maps for the first and last snow-free day for each year in the period. The maps give reasonable estimates which are supported by validation against data from 40 meteorological stations. We have also calculated the average first and last snow-free day for the 11 year period, and the average length of the snow free period.

The correlation between the first snow-free day mapped by MODIS data and by snow data from 40 meteorological stations was highly significant ( $p < 0.05$ ) for 36 of the stations, and with a bias of less than 10 days for 34 of the stations. The correlation for the last snow-free day for 31 of the stations was significant ( $p < 0.05$ ), and the bias was less than 10 days for 27 of the stations. The region of interest is located at high latitudes and is affected by low solar angles/polar darkness during the first snowfall in autumn. These conditions are highly noticeable in the dataset, and cannot be totally removed by any algorithm.

When the first snowfall arrives later than onset of polar darkness (1–20 November), it will remain undetected until we acquire new observations in January–February. Our approach places an emphasis on maximizing the number of observations, and provides in our opinion, the best compromise between using only real observations and refraining from guesswork.

The map showing the average length of the snow-free period is consistent with the land cover in the region, and reflects differences due to continentality and height gradients. By inspecting the map manually and comparing it with local observations (meteorology) we find very convincing trends. Some of the regions with earliest snow-free conditions are well known in local municipalities. E.g. the southern part of Tromsø island becomes snow-free earlier than in the northern part. Similarly we also find early snow-free conditions in farmland areas.

### Acknowledgments

Parts of the work were financed by the Norwegian research council funded by the project EWVA (contact number 216434/E10) and the Fram centre project SenSyF-Fram. MODIS Terra snow cover data were retrieved from NSIDC NASA DAAC: National Snow and Ice Data Center. Meteorological data were obtained from the Norwegian Meteorological institute. We thank H Vickers, Norut for skillful help with improving the language in the paper and two anonymous referees.

#### Conflicts of interest

The authors declare no conflict of interest.

### References

- Andreadis K M and Lettenmaier D P 2006 Assimilating remotely sensed snow observations into a macroscale hydrology model *Adv. Water Resour.* **29** 872–86
- Bjerke J W, Karlsen S R, Høgda K A, Malnes E, Jepsen J U, Lovibond S, Vikhamar Schuler D and Tømmervik H 2014 Record-low primary productivity and high plant damage in the nordic arctic region in 2012 caused by multiple weather events and pest outbreaks *Environ. Res. Lett.* **9** 084006
- Bokhorst S, Bjerke J W, Tømmervik H, Preece C and Phoenix G K 2012 Ecosystem Response to climatic change: the importance of the cold season *Ambio*. **41** (Suppl. 3) 246–55
- Bokhorst S *et al* 2016 Changing Arctic snow cover: a review of recent developments and assessment of future needs for observations, modelling and impacts *Ambio* **45** 516–37
- Chen X, Liang S, Cao Y, He T and Wang D 2015 Observed contrast changes in snow cover phenology in northern middle and high latitudes from 2001–2014 *Sci. Rep.* **5** 16820
- Cooper E J, Dullinger S and Semenchuk P R 2011 Late snowmelt delays plant phenology and results in lower reproductive success in the High Arctic *Plant sci.* **180** 157–67
- Dietz A J, Wohner C and Kuenzer C 2012 European snow cover characteristics between 2000 and 2011 derived from improved MODIS daily snow cover products *Remote Sens.* **4** 2432–54
- eKlima—Free access to weather—and climate data from Norwegian Meteorological Institute from historical data to real time observations ([eklima.no](http://eklima.no)) (Accessed: 1 April 2016)

- Engeset R V, Udnaes H C, Guneriussen T, Koren H, Malnes E, Solberg R and Alfnæs E 2003 Improving runoff simulations using satellite-observed time-series of snow covered area *Hydrol. Res.* **34** 281–94
- Foppa N and Seiz G 2012 Inter-annual variations of snow days over Switzerland from 2000–2010 derived from MODIS satellite data *Cryosphere* **6** 331–42
- Hall D K and Riggs G A 2007 Accuracy assessment of the MODIS snow-cover products *Hydrol. Process.* **21** 1534–47
- Hall D K, Riggs G A, Foster J L and Kumar S V 2010 Development and evaluation of a cloud-gap-filled MODIS daily snow-cover product *Remote Sens. Environ.* **114** 496–503
- Hall D K, Riggs G A, Salomonson V V, DiGirolamo N E and Bayr K J 2002 MODIS snow-cover products *Remote Sens. Environ.* **83** 181–94
- Hansen B B, Isaksen K, Benestad R E, Kohler J, Pedersen Å Ø, Loe L E, Coulson S J, Larsen J O and Varpe Ø 2014 Warmer and wetter winters: characteristics and implications of an extreme weather event in the high arctic *Environ. Res. Lett.* **9** 114021
- Hüsler F, Jonas T, Riffler M, Musial J P and Wunderle S 2014 A satellite-based snow cover climatology (1985–2011) for the European alps derived from AVHRR data *Cryosphere* **8** 73–90
- Høgda K A, Tømmervik H and Karlsen S R 2013 Trends in the start of the growing season in Fennoscandia 1982–2011 *Remote Sens.* **5** 4304–18
- Karlsen S R, Høgda K A and Danks F 2009 A satellite based map of onset of birch (*Betula*) flowering in Norway *Aerobiologica* **25** 15–25
- Karlsen S R, Høgda K A, Wielgolaski F E, Tolvanen A, Tømmervik H, Poikolainen J and Kubin E 2009 Growing-season trends in fennoscandia 1982–2006, determined from satellite and phenology data *Clim. Res.* **39** 275–86
- Kausrud K L, Myrsterud A, Steen H, Vik J O, Østbye E, Cazelles B, Framstad E, Eikeset A M, Mysterud I, Solhøy T and Stenseth N C 2008 Linking climate change to lemming cycles *Nature* **456** 93–7
- Kolberg S and Gottschalk L 2010 Interannual stability of grid cell snow depletion curves as estimated from MODIS images *Water Resour. Res.* **46** 1–15
- Liston G E and Hiemstra C A 2011 The changing cryosphere: Pan-Arctic snow trends (1979–2009) *J. Clim.* **24** 5691–712
- Malnes E, Storvold R, Lauknes I, Solbø S, Solberg R, Amlien J and Koren H 2005 Multi-sensor monitoring of snow parameters in Nordic mountainous areas *Proc. 2005 IEEE Int. Geoscience and Remote Sensing Symp., IGARSS'05* vol 3, pp 1951–4
- Metsämäki S, Mattila O-P, Pulliainen J, Niemi K, Luojus K and Böttcher K 2012 An optical reflectance model-based method for fractional snow cover mapping applicable to continental scale *Remote Sens. Environ.* **123** 508–21
- Metsämäki S, Pulliainen J, Salminen M, Luojus K, Wiesmann A, Solberg R and Ripper E 2015 Introduction to globsnow snow extent products with considerations for accuracy assessment *Remote Sens. Environ.* **156** 96–108
- Moen A 1999 *National Atlas of Norway: Vegetation* (Hønefoss, Norway: Norwegian Mapping Authority) pp 141–4
- Arctic Monitoring and Assessment Programme (AMAP) 2011 *Snow, Water, Ice and Permafrost in the Arctic (SWIPA): Climate Change and the Cryosphere* (Oslo, Norway) xii + 538 pp ([www.amap.no/swipa/](http://www.amap.no/swipa/))
- Norwegian Mapping Authority—free access to Norwegian map layers from ([www.kartverket.no](http://www.kartverket.no)) (Accessed: 23 November 2016)
- Riggs G A, Hall D K and Salomonson V V 2006 MODIS Snow Products User Guide to Collection 5 (<http://modis-snow-ice.gsfc.nasa.gov/?c=userguides>)
- Riseth J Å *et al* 2011 Sámi traditional ecological knowledge as a guide to science: snow, ice and reindeer pasture facing climate change *Polar Res.* **47** 202–17
- Saloranta T M 2012 Simulating snow maps for Norway: description and statistical evaluation of the seNorge snow model *Cryosphere* **6** 1323–37
- Solberg R, Koren H, Amlien J, Malnes E, Schuler D V and Orthe N K 2010 The development of new algorithms for remote sensing of snow conditions based on data from the catchment of Øvre Heimdalsvatn and the vicinity *Hydrobiologia* **642** 35–46
- Tveito O E 2000 Nordic temperature maps *Met. Report* no. 9 Meteorological Institute, Oslo, Norway ([http://met.no/filestore/09\\_00.pdf](http://met.no/filestore/09_00.pdf))
- Tveito O E, Førland E J, Alexandersson H, Drebs A, Jónsson T, Tuomenvirta H and Vaarby Laursen E 2001 Nordic climate maps *DNMI Report* 06/01 Oslo, Norway ([http://met.no/filestore/06\\_01.pdf](http://met.no/filestore/06_01.pdf))
- Tømmervik H 2005 Estimation of carrying capacity for wild reindeer in Norway by means of the normalised difference snow index (NDSI) *Proc. SPIE* **5976** 597612
- Tømmervik H, Johansen B, Riseth J Å, Karlsen S R, Solberg B and Høgda K A 2009 Above ground biomass changes in the mountain birch forests and mountain heaths of Finnmarksvidda, Northern Norway, in the period 1957–2006 2009 *Forest Ecol. Manage.* **257** 244–57
- Tveraa T, Stien A, Bårdsen B J and Fauchald P 2013 Population densities, vegetation green-up, and plant productivity: impacts on reproductive success and juvenile body mass in reindeer *PLoS One* **8** e56450
- Vikhamar D and Rune Solberg R 2003 Snow-cover mapping in forests by constrained linear spectral unmixing of MODIS data *Remote Sens. Environ.* **88** 309–23
- Vikhamar-Schuler D and Hanssen-Bauer I 2010 Long-term climate trends of Finnmarksvidda, Northern-Norway *Met. Report* no. 06/2010 ([http://met.no/filestore/Ealat\\_climaterep\\_dvs.pdf](http://met.no/filestore/Ealat_climaterep_dvs.pdf)) (Accessed: 23 November 2016)
- Wielgolaski F E and Sonesson M 2001 *Nordic Mountain Birch Ecosystems: A Conceptual Overview (Man and the Biosphere Series)* ed F E Wielgolaski vol 27 (Paris, France: UNESCO) pp 377–84
- Wipf S 2010 Phenology, growth, and fecundity of eight subarctic tundra species in response to snowmelt manipulations *Plant Ecol.* **207** 53–66
- Wipf S and Rixen C 2010 A review of snow manipulation experiments in Arctic and alpine tundra ecosystems *Polar Res.* **29** 95–109
- Xu L *et al* 2013 Temperature and vegetation seasonality diminishment over Northern Lands *Nat. Clim. Change* **3** 581–6