



REPORT SNO 5205-2006



Biomarkers in monitoring

A review



Photo: Thomas Lang, BFA-Fi, Cuxhaven, DE

Main Office

P.O. Box 173, Kjelsås
 N-0411 Oslo, Norway
 Phone (47) 22 18 51 00
 Telefax (47) 22 18 52 00
 Internet: www.niva.no

**Regional Office,
Sørlandet**

Televeien 3
 N-4879 Grimstad, Norway
 Phone (47) 37 29 50 55
 Telefax (47) 37 04 45 13

**Regional Office,
Østlandet**

Sandvikaveien 41
 N-2312 Ottestad, Norway
 Phone (47) 62 57 64 00
 Telefax (47) 62 57 66 53

**Regional Office,
Vestlandet**

Nordnesboder 5
 N-5008 Bergen, Norway
 Phone (47) 55 30 22 50
 Telefax (47) 55 30 22 51

**Regional Office
Midt-Norge**

P.O. Box 1266
 N-7462, Norway
 Phone (47) 73 54 63 85 / 86
 Telefax (47) 73 54 63 87

Title Biomarkers in monitoring – a review	Serial No. 5205-2006	Date 07/04-2006
	Report No. Sub-No. O-25302	Pages Price 106
Author(s) Ketil Hylland Anders Ruus Jan Fredrik Børseth (IRIS) Renée Bechmann (IRIS) Janina Barsiene (University of Vilnius, Latvia)	Topic group Ecotoxicology	Distribution Open
	Geographical area	Printed NIVA


Client(s) OLF	Client ref. Laurence Pinturier
------------------	-----------------------------------

<p>Abstract</p> <p>Biological effects are increasingly used to monitor impacts of contaminants in marine ecosystems. Biomarkers have been defined as "biochemical, cellular, physiological or behavioural variations in the tissue or body fluids or at the level of whole organism that provide evidence of exposure to chemical pollutants, and may also indicate a toxic effect". The biomarkers reviewed here were PAH bile metabolites, cytochrome P4501A, glutathione S-transferase, markers for DNA damage (adducts, alkaline unwinding, Comet assay), micronucleus formation, peroxisomal proliferation, acetyl cholinesterase inhibition, metallothionein, vitellogenin and delta-aminolevulinic acid dehydratase. Focus for the review was dose-response relationships, confounding factors, links to population effects, baseline values, assessment criteria and quality assurance for the relevant biomarker. Whereas correlative links to population-relevant effects have been found for some biomarkers, e.g. biomarkers for DNA damage, most biomarkers are generally more useful as markers for possible impacts elsewhere in ecosystems.</p>


<p>4 keywords, Norwegian</p> <ol style="list-style-type: none"> 1. biomarker 2. overvåking 3. offshore 4. økologiske effekter 	<p>4 keywords, English</p> <ol style="list-style-type: none"> 1. biomarker 2. monitoring 3. offshore 4. ecological impact
---	---



Ketil Hylland
 Project manager



Kevin V Thomas
 Research manager
 ISBN 82-577-4924-9



Jarle Nygard
 Strategy Director

Biomarkers in monitoring

A review

Preface

The project was initiated by OLF (Unn Orstein) to produce a review of methods relevant to water column monitoring with a focus on interpretation and ecological links. Laurence Pinturier, Total, has been OLF's contact with the project group. The project has been shared between NIVA and IRIS with Ketil Hylland, NIVA, as lead author.

Rogaland Research were responsible for the review of PAH metabolites (Lars-Petter Myhre with inputs from Ketil Hylland) the Comet assay (Renée Bechmann) and micronucleus analyses (Jan Fredrik Børseth and Janina Barsiene). NIVA was responsible for editing, DNA adducts, vitellogenin, metallothionein and ALA-D (Ketil Hylland), cytochrome P4501A activity (Anders Ruus and Ketil Hylland), acetyl cholinesterase inhibition (Anders Ruus and Merete Grung), MXR and peroxisomal proliferation (Merete Grung) and glutathione *S*-transferase (Knut-Erik Tollefsen).

Oslo, 9.3 2006

Ketil Hylland

Contents

Summary	1
1. Introduction	3
1.1 Background – biomarkers	3
1.2 The need for baseline data and assessment criteria	4
1.3 Links between biomarkers and ecologically relevant endpoints	5
1.4 Quality assurance	6
2. PAH metabolites in bile	7
2.1 Introduction	7
2.2 Dose-response (species specific)	8
2.3 Species sensitivity	10
2.4 Relevance of other factors	10
2.5 Links	10
2.6 Background responses	10
2.7 Assessment criteria	10
2.8 Quality assurance	11
3. Cytochrome P4501A activity (EROD)	17
3.1 Introduction	17
3.2 Dose-response	17
3.3 Blue mussel	19
3.4 Species sensitivity	19
3.5 Relevance of other factors	19
3.6 Links	20
3.7 Background responses	20
3.8 Assessment criteria	20
3.9 Quality assurance	21
4. Glutathione <i>S</i>-transferase activity	22
4.1 Introduction	22
4.2 Dose-response	22
4.3 Species sensitivity	25
4.4 Relevance of other factors	25
4.5 Links	26
4.6 Background responses	27
4.7 Assessment criteria	27
4.8 Quality assurance	27
5. DNA damage	29
5.1 Introduction	29
5.2 DNA repair	29
5.3 Alkaline unwinding	29
5.4 DNA adducts	30
5.4.1 Dose-response	30

5.4.2 Species baseline and sensitivity	30
5.4.3 Time-course of adduct development	30
5.4.4 Relevance of other factors	31
5.4.5 Links	32
5.4.6 Background responses	32
5.4.7 Assessment of DNA adduct results	32
5.4.8 Quality assurance	32
6. The Comet assay	40
6.1 Introduction	40
6.2 The Comet assay with invertebrates	44
6.3 The Comet assay with fish red blood cells	45
6.4 Confounding factors	45
6.5 Links	46
6.6 Background responses	46
6.7 Assessment criteria	47
6.8 Quality assurance	47
7. Micronucleus formation	50
7.1 Introduction	50
7.2 Micronucleus analyses <i>in situ</i>	50
7.3 The micronucleus test as laboratory bioassay	51
7.4 Assessment of MN in caged organisms	52
7.5 Target tissues	52
7.5.1 Fish	52
7.5.2 Mussels	53
7.5.3 Other indicator species	54
7.6 Confounding factors	56
7.7 Links	56
7.8 Background responses	56
7.9 Assessment criteria	56
7.10 Quality assurance	56
8. Multixenobiotic response (MXR)	57
8.1 Introduction	57
8.2 Dose-response	57
8.2.1 MXR induction	57
8.2.2 Chemosensitisers	59
8.3 Species sensitivity	59
8.4 Relevance of other factors	59
8.5 Links	60
8.6 Background responses	60
8.7 Assessment criteria	60
8.8 Quality assurance	60
9. Peroxisomal proliferation	61
9.1 General introduction	61
9.2 Dose-response	61
9.3 Species sensitivity	63
9.4 Confounding factors	63
9.5 Links	64
9.6 Background responses	64

9.7 Assessment criteria	64
9.8 Quality assurance	64
10. Acetyl cholinesterase (AChE) inhibition	65
10.1 Introduction	65
10.2 Dose-response	65
10.3 Species sensitivity	67
10.4 Relevance of other factors	67
10.5 Links	68
10.6 Background responses	68
10.7 Assessment criteria	68
10.8 Quality assurance	68
11. Vitellogenin (vtg) induction	69
11.1 Introduction	69
11.2 Dose-response	69
11.3 Relevance of other factors	70
11.4 Links	71
11.5 Background responses	71
11.6 Assessment criteria	72
11.7 Quality assurance	72
12. Metallothionein (MT) concentration	73
12.1 Introduction	73
12.2 Analytical aspects	73
12.3 Metallothionein in mussels	74
12.4 Dose-response in fish	74
12.5 Species sensitivity	75
12.6 Relevance of other factors	75
12.7 Links	76
12.8 Background responses and assessment criteria	76
12.9 Quality assurance	76
13. δ-aminolevulinic acid dehydratase (ALA-D) inhibition	77
13.1 Introduction	77
13.2 Analytical considerations	77
13.3 ALA-D in invertebrates	77
13.4 Dose-response and species sensitivity	77
13.5 Background responses	78
13.6 Links	78
13.7 Other factors	78
13.8 Quality control	79
14. Concluding remarks	80
15. References	82

Summary

Biological effects are increasingly used to monitor the influence of contaminants in marine ecosystems. Biological effects may conveniently be divided into community or population responses, bioassays and biomarkers. While community responses are widely used to address impacts on benthic communities, biomarkers have been used to investigate effects in fish or mussels. This review concerns biomarkers with a focus on methods that have been used in monitoring effluents from offshore activities. The methods included were PAH bile metabolite concentration, cytochrome P4501A activity, glutathione *S*-transferase activity, two methods to address DNA damage (DNA adducts and the Comet method), micronucleus formation, multixenobiotic response, acetyl cholinesterase inhibition, vitellogenin concentration, metallothionein concentration and delta-aminolevulinic acid dehydratase inhibition. Histological assays, including lysosomal stability, were not included in this review.

Biomarkers may be viewed as “biological responses that can be related to an exposure to, or toxic effect of, an environmental chemical or chemicals”. This is a very wide definition, since “biological responses” can refer to responses at any hierarchical level, from molecular processes to the ecosystem. This project used the following definition: “biochemical, cellular, physiological or behavioural variations in the tissue or body fluids or at the level of whole organism that provide evidence of exposure to chemical pollutants, and may also indicate a toxic effect”. In addition, methods were selected for their previous use in monitoring effluents from offshore activities or for their potential in this context.

For each method, the review focused on dose-response relationships, confounding factors, links to effects at population level, baseline levels, assessment criteria and quality assurance.

From previous experience and existing data, all the reviewed biomarkers could conceivably be relevant for use in monitoring offshore effluent environmental impacts. Although it is not possible to establish direct links between responses in any of the biomarkers and population effects, there is support through correlation for a predictive ability for some biomarkers in that regard, e.g. DNA adducts and the Comet assay. Although presumably not critical in adult fish, vitellogenin induction will indicate levels of estrogens that may have population knock-on effects through changes in sex ratio. Similarly, micronucleus formation in mussels indicates the presence of substances that may cause chromosomal damage to other marine organisms and conceivably affect their offspring. Responses in other biomarkers included herein, i.e. cytochrome P4501A activity, metallothionein, AChE and ALA-D, will in most cases indicate contaminant-induced changes, but there are no clear links to population effects. Two of the biomarkers reviewed, peroxisomal proliferation and multixenobiotic resistance, have not been used to a large extent in monitoring and should be further reviewed when more data are available.

Baseline values and assessment criteria could be established for some species for limited range of the methods, i.e. PAH bile metabolites, cytochrome P4501A activity (tentative), vitellogenin, metallothionein and ALA-D.

Quality assurance programmes have until now included PAH bile metabolites, cytochrome P4501A activity, vitellogenin, metallothionein and ALA-D. There have also been some international intercalibration activities for AChE.

1. Introduction

1.1 Background – biomarkers

There is a need to identify whether anthropogenic activities affect natural ecosystems, both because humans have a moral responsibility not to damage the environment and because environmental deterioration may be in conflict with other human uses of the ecosystem. For management purposes there is a need both to be able to quantify the distribution, bioaccumulation and biomagnification of contaminants as well as their effects on marine organisms.

Methods to identify biological effects can be conveniently grouped as shown in Table 1. In addition to the methods shown, biological components are also included in sensors (biosensors) or sensors may be used to monitor biological processes in real-time.

Table 1. A grouping system for biological effects methods.

designation	examples	positive properties	negative properties
ecosystem	none	no generally accepted methods	
population/community (large samples of organisms from defined areas)	benthic community rocky shore community	ecologically relevant	strongly affected by non-contaminant parameters difficult to standardise large natural variability
bioassays (sample or extract from the field is tested in biological system in lab)	Acartia 96-hr test Corophium sediment test Skeletonema 96-hr test fish hepatocyte cytotoxicity test	easy to standardise ecologically relevant endpoints	robust species (not ecologically relevant) not contaminant-specific
biomarkers (health assessment of individual organisms)	EROD DNA adducts metallothionein AChE ALA-D Comet assay	easy to standardise related to individual health contaminant-related species similarities (some methods)	limited ecological relevance species differences (some methods)

There are several definitions for “biomarkers”. According to Peakall (1994) biomarkers are “biological responses that can be related to an exposure to, or toxic effect of, an environmental chemical or chemicals”. This is a very wide definition, since “biological responses” can refer to responses at any hierarchical level, from molecular processes to the ecosystem. Other definitions are more specific, such as that of Mayer et al. (1992) which states that biomarkers are “quantifiable biochemical, physiological, or histological measures that relate in a dose- or time-dependent manner to the degree of dysfunction that contaminants have produced”. The non-governmental organisation English Nature has defined biomarkers as:

“biochemical, cellular, physiological or behavioural variations in the tissue or body fluids or at the level of whole organism that provide evidence of exposure to chemical pollutants, and may also indicate a toxic effect”.

This definition complies well with the selection of methods for this review. In addition, methods were selected for their previous use in monitoring effluents from offshore activities or for their potential in this context.

Any biological effect method should ideally have a number of characteristics, outlined in e.g. Stegeman *et al.* (1992):

- 1 The assay to quantify the biomarker should be sensitive, reliable, and relatively simple;
- 2 baseline data for the concentration/activity of the biomarker should be known in order to be able to distinguish between natural variability (noise) and contaminant induced stress (signal);
- 3 the basic biology/physiology of the test organism should be known so that sources of uncontrolled variation (growth and development, reproduction, food sources) can be minimized;
- 4 all the factors, intrinsic as well as extrinsic, that affect the biomarker should be known;
- 5 it should be established whether changes in biomarker concentration are due to physiological acclimation or to genetic adaptation; and finally,
- 6 changed levels of the biomarker should be correlated with the “health” or “fitness” of the organism.

Very few of the methods used actually comply with all those requirements, but it can be useful to keep such an “ideal” in mind while reviewing different techniques.

1.2 The need for baseline data and assessment criteria

Although it is generally agreed that biological effects results should not be used or assessed singly there are currently no generally agreed frameworks for such assessments. Current activities within OSPAR (WKIMON) may generate a framework; the results from that activity are due in 2006. Until such a framework is available, biomarker methods have to be assessed singly. This means that in any given study, there should be criteria agreed on *a priori* as to whether the result is “an effect” or not. In the scientific literature, and in biomonitoring, this has generally been achieved through hypothesis testing – asking the question “is group B different from the reference or control group”? There could however be biological reasons to state that only biomarker responses at least 5 or 10 times those measured in a reference group would be indicative of an impact.

Depending on the method, there are two strategies for the assessment of single biomarker responses:

- observed biomarker responses can be compared against a global criterion or
- observed biomarker responses must always be compared to responses in a reference group sampled at the same time.

An assessment is obviously simpler if the former strategy is possible; this will however only apply to methods and species for which there is extensive knowledge of all factors that may affect the response (in addition to contaminants). In most cases this will also be methods for which there are limited species differences.

The latter strategy must be chosen for methods for which there is not sufficient information about confounding factors, for which it is not possible to correct for such factors or for which exogenous factors may affect the response in non-predictable ways (but similarly at exposed and reference locations).

1.3 Links between biomarkers and ecologically relevant endpoints

As indicated above there is not a direct link between most biomarker responses and population effects, although it is generally thought that biomarker responses will be predictive of such impacts in one way or another. It is this elusive link that ecotoxicologists have been searching for during the past two decades.

There are however good reasons why this link is so elusive; marine organisms that may be weakened for some reason or other (e.g. contaminants) will not be as apparent as e.g. a weakened fish eagle or polar bear. It will simply disappear, eaten because avoidance responses were too slow or starved to death and then eaten. The implications are that we are always sampling the healthiest part of any population in field campaigns.

It is by definition not possible to measure early and late effects in the same individuals and ecologically relevant parameters (increased mortality, reduced reproductive output) can not be determined under field conditions for most species. One possible way to establish links is through multi-year, -method and -species research studies, but it will still only be causality within a population. There is really only one such situation in the world, the work by NOAA in Puget Sound, recently reviewed by Myers et al. (2003). In their work they have been able to show causality between (petrogenic) PAHs in sediment, PAH-metabolites in bile, cytochrome P4501A induction, DNA adduct formation and development of liver cancer (neoplasms) in flatfish. Since they have been able to include many species they have also been able to see that cancer develops in some species, but not in others. A second possible approach is to perform long-term mesocosm experiments under semi-natural conditions. Such studies will obviously be open to the criticism that fish (or other organisms) were unable to migrate as they normally would. Furthermore, due to space limitations it is never possible to have an appropriate replication in such systems. The most well-known such study was done by RIKZ in the Netherlands. They exposed flounder and plaice to contaminated and reference sediments for two years (Vethaak et al., 1996). From this extensive mesocosm experiment they were able to support a suggested link from PAH in sediment to liver cancer development (Eggens et al., 1996). Some of the steps in between did not fit the model, possibly due to adaptation.

A follow-up issue is the question of whether it is really necessary to find direct links between a response in a fish species and the population of that very same fish, as suggested under item (6) above (the ability to relate a response to health or fitness of the organism under study). A response in the fish indicates that contaminants are present in the ecosystem at sufficient levels to modulate normal metabolism in this species. Other species or life-stages in the ecosystem may be more strongly affected and the biomarker response is simply a marker for this risk. That said, there is obviously a need to qualify such an assessment in terms of which other species could be at risk.

This discussion will be continued for each of the biomarkers in the following chapters.

1.4 Quality assurance

If biomarker results are to be comparable from year to year and between laboratories it is imperative that both internal and external quality assurance procedures are in place. International organisations, e.g. OSPAR, AMAP, ICES, MEDPOL, have focussed on this issue over the past decade.

Over the past 10 years there has been considerable international activity to harmonise protocols and intercalibrate biomarker methods, primarily in the framework of the BEQUALM, first an EU-funded project and now a self-funding venture (<http://www.bequalm.org>).

2. PAH metabolites in bile

2.1 Introduction

Analyses of PAH metabolites in fish bile have been used as a biomarker of exposure to PAH contamination since the early 1980s. The presence of metabolites in bile (and in urine) is the final stage of the biotransformation process whereby lipophilic compounds are transformed to a more soluble form and then passed from the organism in bile or urine. This review will focus on metabolites in bile since this is the method that has been most widely used and is most relevant for offshore monitoring.

As a biomarker of exposure, measuring PAH metabolites in bile has many advantages over other techniques that require sophisticated tissue preparation protocols. The pretreatment of bile samples requires relatively simple dilution steps prior to analysis by direct fluorescence measurement. The bile is diluted in methanol : distilled water (1:1) and fluorescence is measured with a fluorometer. Fixed wavelength fluorescence is a suitable screening method for samples while HPLC/F or GC-MS SIM is utilized for qualitative and quantitative measures.

Bile is generally stored in the gall bladder prior to episodic release into the esophagus where bile salts have a function to perform as part of the digestive process. This period of storage permits a degree of accumulation of metabolites and hence an increase in their concentration. The periodic release of bile does however introduce a variable into the technique, which must be accounted for. The feeding status of fish has been shown to influence both the volume and the density of the bile (Collier and Varanasi, 1991).

The ability of fish to biotransform PAHs into less lipophilic derivatives means that reliance on the detection of parent PAHs alone may lead to an underestimation of the *in vivo* exposure level of PAH in the fish. PAH metabolite detection, on the other hand, represents a quantification of the flux of PAHs streaming through the fish's body. From a toxicological point of view, flux information is more relevant for estimating the actual biotic stress due to PAH exposure, than the body burden data of the unmetabolised parent PAH compounds in tissues (most often liver). Despite this, body burden measurements are still more commonly used within monitoring studies than metabolite determination.




Figure 1. Synchronous fluorescence screening of bile samples from fish at a reference site compared with fish taken at a site polluted with an oil slick. Data courtesy of Endre Aas, IRIS-Akvamiljø.




Figure 2. Fixed fluorescence detection of bile samples from fish at a reference site versus bile from fish at a field site polluted with an oil slick. Data courtesy of Endre Aas, IRIS-Akvamiljø.

2.2 Dose-response (species specific)

The PAH compounds are metabolised rapidly in the organisms and it is the endpoint of this metabolisms that is measured in the bile. The compounds are measured using chemical analysis. A consistent dose-response relationship has been demonstrated in laboratory studies between PAH exposure and the subsequent presence of metabolites in bile (Beyer et al., 1997; Sandvik et al., 1997; Aas et al., 2000). To establish a good dose-response relationship in field studies it is necessary to focus on aspects that influence the excretion of bile.




Figure 3. GC/MS quantification of metabolites of alkylated naphthalene in bile from fish caged downstream from the discharge of produced water at the Tampen field (BECPELAG field study in 2001). Stations G1-G4 are situated in the southern North Sea (Aas et al., in press).

The method requires that bile is available in the gall bladder. Since the fish renew bile as part of normal metabolism and excrete it during digestion, it is important to know about the dietary status of the organism to establish a dose- response relationship. If the fish feed just before sampling, the gall bladder may become more or less empty. After the gall bladder has been emptied it will fill up and metabolites will be concentrated up to a plateau level corresponding to the exposure regime. Consequently the time since last digestion is important for the dose-response relationship. Fish generally have a very efficient metabolic excretion of most PAHs and it has been shown that most of the PAH will excreted after 2 – 8 days following exposure. This means that the PAH metabolites determined in bile will represent exposures on the scale of days and, at most, two weeks.

It has been shown in several field and laboratory studies that there is a good correlation between PAH exposure and bile metabolites (see Table 3). Because of the rapid metabolism and the correlation between bile content and digestive status it is difficult to make a dose-response relationship that can be used to quantify the exposure. Work has been done to try to correlate bile metabolite concentration to digestive status, by correlating it to the amount of protein or biliverdin in the bile. Absorbance at 380 nm is also used (similar to biliverdin) (Hylland, unpublished). This normalisation is not standardised because it has been shown to only explain parts of the variability, but it is recommended to be part of the explaining factors in the interpretation of results. In laboratory studies it is normal to stop the feeding some days before sampling to ensure the bile quality. In field sampling this can be taken into account by letting the fish go some days in tanks before sampling, but this has some logistical challenges.

2.3 Species sensitivity

The responsiveness of bile metabolites towards pollution is shown to be good in several species. In this overview we have focused on cod, haddock, polar cod and sheephead minnows (the latter is a model species from the American continent). The background level differs from species to species so it is important to establish good baseline before using new species. It may be expected that species with fatty livers, i.e. most gadiids, may metabolise PAHs more slowly as more will partition into fat, but this has not been documented experimentally.

2.4 Relevance of other factors

As mentioned above, food availability will affect the concentration of PAH metabolites in bile. In an assessment of data for more than 500 individual cod sampled through five years of national monitoring, variables such as size/age and sex explained some variability in multiple regression models (Ruus et al., 2003). This could be due to different feeding preferences, but also endogenous processes. In addition, the fat-content of the liver (measured as liver-somatic index, LSI) came out as significant, presumably because fat decreases the availability of PAH to the cellular compartments of liver cells.

2.5 Links

PAH metabolites are biomarkers of exposure. Consequently it is only possible to link the PAH metabolites in bile indirectly to effects by correlate it to other expected effects that will appear with exposure to PAHs. But it is important to bear in mind that the occurrence of PAH metabolites in bile indicate a biotransformation process of the corresponding PAHs in the organism. Thus any implications for fitness related to induction of enzymes, such as EROD, are equally appropriate if the final products of this process can be measured in the bile. Establishing links between bile metabolites and fitness parameters will require a similar approach to that taken for the previous biomarkers, chronic, low dose exposures in the laboratory and the examination of endpoints related to growth and reproduction in association with regular analysis of bile profiles. Manipulation of feeding regimes as part of this process may throw additional light on the relationship.

2.6 Background responses

Baseline levels of PAH metabolites have been established for many of the species relevant for monitoring in Norwegian coastal and offshore waters. From Ruus et al. (2003) values for the relevant species are: (all values standardised to absorbance at 380 nm) Atlantic cod: 0.6-4 µg/kg bile, flounder 27-89 µg/kg bile, dab 3.1-34 µg/kg bile, plaice 0.4-3 µg/kg bile (all quantified using HPLC separation and fluorescence detection and quantification). Standardisation at 380 nm is used to remove variability due to bile salts.

2.7 Assessment criteria

It is possible to establish global criteria for individual PAH metabolites. Baseline data for individual species may be used to test against to determine whether fish have been exposed to PAHs. As mentioned above, some variation in PAH metabolites in bile appear to be related to

sex and size/age (Ruus et al., 2003), knowledge of which should be included in the sampling design.

2.8 Quality assurance

A general protocol outlining analytical strategies and their strengths as well as weaknesses has recently become available (Ariese et al., 2005). There have been international intercalibration exercises for the determination of PAH-metabolites in fish bile, arranged in collaboration between an EU-project and QUASIMEME¹. Reference bile samples were generated as part of the aforementioned EU project and are now available at IRMM, JRC, Geel, Belgium (<http://www.irmm.jrc.be/html/homepage.html>).

¹ QUASIMEME – organisation that offers quality assurance for chemical endpoints; <http://www.quasimeme.org>

Table 2. Overview of field and laboratory studies – PAH metabolites measured by fixed fluorescence.

Species	Substance (lab/field)	Test concentrations/area	Exposure time	Metabolite	Baseline	control or reference	exposed /control	reference/comments
Cod (Gadus morhua)	Feral fish	Barents Sea	Baseline					Aas et al., 2003
Cod (Gadus morhua)	Feral fish	Egersund	Baseline non polluted area	Naph type	5.3 ug/ml			Klungsoyr et al. 2003
				Pyren type	0.8 ug/ml			
				BaP type	0.4 ug/ml			
Cod (Gadus morhua)	Feral fish	Steipner	Baseline polluted area?	Naph type	6.1 ug/ml			Klungsoyr et al. 2003
				Pyren type	1.0 ug/ml			
				BaP type	0.5 ug/ml			
Cod (Gadus morhua)	Feral fish	Statfjord	Baseline polluted area?	Naph type	5.9 ug/ml			Klungsoyr et al. 2003
				Pyren type	0.9 ug/ml			
				BaP type	0.3 ug/ml			
Cod (Gadus morhua)	Feral fish	Frøy, ceased installation 10 000 m (ref) 2000 m - 200 m	Baseline polluted area?	Naph type		3.9 ug/ml	1.1 - 1.1	Beyer et al. 2003
				Pyren type		0.6 ug/ml	1.1 - 0.9	
				BaP type		0.3 ug/ml	0.9 - 0.9	
Cod (Gadus morhua)	Feral fish	Barents sea	Baseline	Naph type	2,15 ug/g			Sundt, 2002
				Pyren type	1,63 ug/g			
				BaP type	0,69 ug/g			
Cod (Gadus morhua)	Feral fish	Barents sea	Baseline	Naph type	5,8 ug/g			Aas & Børseth, 2002
				Pyren type	1,7 ug/g			
				BaP type	0,8 ug/g			
Cod (Gadus morhua)	Laboratory	1 ppm crude oil Statfjord B	14 days					Aas et al., 2002
Cod (Gadus morhua)	Laboratory	0.06 - 0.25 - 1 ppm Oil	average 3, 7, 14, 24 days	Naph type		3,9 ug/g	7,5 - 23,7 - 31,4	Skadshheim et al., 2004
				Pyren type		2,6 ug/g	3,6 - 10,6 - 13	
				BaP type		1,0 ug/g	1,7 - 2,4 - 2,2	
Cod (Gadus morhua)	Laboratory	0.06 - 0.25 - 1 ppm Oil	average 3, 17, 31 day	Naph type		53,7 ug/g	0,7 - 2,3 - 2,9	Skadshheim et al., 2004
				Pyren type		7,0 ug/g	1 - 2,9 - 3,3	
				BaP type		1,0 ug/g	1,1 - 1,5 - 1,5	
Cod (Gadus morhua)	Laboratory	Oil 0.06 - 0.25 - 1 ppm	30 days	Naph type		7,1 fi	5,1 - 9,5 - 227,5	Aas et al. 2000
				Pyren type		2 fi	6,4 - 12,7 - 43,3	
				BaP type		0,8 fi	2,3 - 3,6 - 9,6	

Species	Substance (lab/field)	Test concentrations/area	Exposure time	Metabolite	Baseline	control or reference	exposed /control	reference/comments
Cod (Gadus morhua)	Laboratory	PW Oseberg, 1:1000 - 1:200 - 0.2 ppm oil - 0.2 ppm oil + PAH-mix	15 days	Naph type		12.6 ug/ml	1.3 - 2.5 - 3.6 - 5.4	Sundt, 2004
				Pyren type,		4 ug/ml	1.7 - 3.7 - 4.1 - 17.8	
				BaP type,		1.8 ug/ml	1.3 - 1.8 - 1.5 - 2.4	
Cod (Gadus morhua)	Field, Caged	North Sea - Stafjord, 10000 m - 2000m - 500 m German bight G	5.5 weeks	Naph type	7.5 ug/ml	0,7	1.7 - 1.9 - 2.1	Aas et al., in press
				Pyren type	3.1 ug/ml	0,7	1.2 - 1.5 - 1.6	
				BaP type	1.2 ug/ml	0,8	1.2 - 1.1 - 1.2	
Cod (Gadus morhua)	Field, Caged	German bight G4 (Ref) G1 - G2 - G3	5.5 weeks	Naph type	7.5 ug/ml	0,4	0.9 - 0.9 - 1.6	Aas et al., in press
				Pyren type	3.1 ug/ml	0,5	0.8 - 0.9 - 1.7	
				BaP type	1.2 ug/ml	0,7	0.8 - 1 - 1.3	
Cod (Gadus morhua)	Field, Caged	North Sea - Troll, 1000 m - 500m	6 weeks	Naph type	4.6 ug/ml	1,4	1.7 - 2.5	Børseth et al., 2004
				Pyren type	2.4 ug/ml	0,9	1.1 - 1.3	
				BaP type	0.9 ug/ml	1,1	1.1 - 1.3	
Cod (Gadus morhua)	Field, Caged	North Sea - Tampen, 10000 - 2500 - 1000 - 500	6 weeks	Naph type		8.8 ug/ml	1.0 - 1.5 - 1.2 - 1.2	Hyland et al., 2005
				Pyren type				
				BaP type		1.4 ug/ml	0.9 - 0.7 - 0.8 - 0.9	
Haddock (Melanogrammus aeglefinus)	Feral fish	Egersund	Baseline non polluted area	Naph type	5.1 ug/ml			Klungsoyr et al. 2003
				Pyren type	1.4 ug/ml			
				BaP type	0.7 ug/ml			
Haddock (Melanogrammus aeglefinus)	Feral fish	Sleipner	Baseline polluted area?	Naph type	6.8 ug/ml			Klungsoyr et al. 2003
				Pyren type	1.9 ug/ml			
				BaP type	0.8 ug/ml			
Haddock (Melanogrammus aeglefinus)	Feral fish	Stafjord	Baseline polluted area?	Naph type	11.2 ug/ml			Klungsoyr et al. 2003
				Pyren type	2.5 ug/ml			
				BaP type	0.7 ug/ml			
Haddock (Melanogrammus aeglefinus)	Feral fish	Barents sea		Naph type	2.52 ug/g			Sundt, 2004
				Pyren type	1.69 ug/g			
				BaP type	0.77 ug/g			
Haddock (Melanogrammus aeglefinus)	Feral fish	Barents sea		Naph type	2.0 ug/g			Aas & Børseth, 2004
				Pyren type	1.3 ug/g			
				BaP type	0.6 ug/g			
Haddock	Feral fish	Frøy, ceased installation	Baseline polluted	Naph type		5.6 ug/ml	1.3 - 2.2	Beyer et al., 2003

Species	Substance (lab/field)	Test concentrations/area	Exposure time	Metabolite	Baseline	control or reference	exposed /control	reference/comments
(Melanogrammus aeglefinus)		10 000 m (ref) 2000 m - 200 m	area?	Pyren type		1.4 ug/ml	1.4 - 0.7	
				BaP type		0.75 ug/ml	1.8 - 0.6	
Sheepshead minnow	Laboratory	North sea oil A 0.1 - 0.4 - 0.7 ppm	5 weeks	Naph type		6916	2.3 - 6.2 - 9.3	Bechmann et al. 2004
				Pyren type		569	2.5 - 5 - 6.3	
				BaP type		107	4 - 13.1 - 19.2	
Sheepshead minnow	Laboratory	North sea oil B 0.1 - 0.9 - 5.6 ppm	6 weeks	Naph type		18164	1.8 - 4.3 - 12.5	Bechmann et al. 2004
				Pyren type		438	5.6 - 12.6 - 30.8	
				BaP type		110	12.6 - 42.7 - 123.9	
Sheepshead minnow	Laboratory	2 - 14 - 214 ppb	5 weeks	Naph type		267280	0.9 - 2.2 - 18.6	Bechmann et al. 2004
				Pyren type		9926	0.9 - 1.5 - 9.6	
				BaP type		5152.7	3 - 17.4 - 207	
Polar cod (Boreogadus saida)	Laboratory, feral fish 2001, 2002	1.5 ppm StatfjA oil, baseline, control	14 days	Naph type	16.0 ug/g	2	16.9	Sundt & Bechmann, 2004
				Pyren type	0.9 ug/g	5.5	74.4	
				BaP type	0 ug/g	0	1.8	

Table 3. PAH-metabolites in marine fish – measured by GC-MS.

Species	Substance (lab/field)	Test concentrations	Exposure time	Metabolite	Baseline	control or reference	exposed/control	reference
Cod (Gadus morhua)	Feral fish	Barents sea	baseline	Naph sum	150,6 ng/g			Aas & Børseth, 2002
				Phen sum	61,2 ng/g			
				Pyren	4,6 ng/g			
Cod (Gadus morhua)	Feral fish	Barents sea	baseline	Naph sum	1285 ng/g			Sundt, 2004
				Phen sum	220 ng/g			
				Pyren	3,5 ng/g			
Cod (Gadus morhua)	Feral fish	Egersund	Baseline non polluted area	Naph sum	2005,1 ng/g			Klungøy et al. 2003
				Phen sum	230,2 ng/g			
				Pyren	3,9 ng/g			
Cod (Gadus morhua)	Feral fish	Sleipner	Baseline polluted area?	Naph sum	1296,1 ng/g			Klungøy et al. 2003
				Phen sum	197,8 ng/g			
				Pyren	0 ng/g			
Cod (Gadus morhua)	Feral fish	Stafford	Baseline polluted area?	Naph sum	1361,7 ng/g			Klungøy et al. 2003
				Phen sum	351,1 ng/g			
				Pyren	4,0 ng/g			
Cod (Gadus morhua)	Laboratory	0.06 - 0.25 - 1 ppm Oil	average 3, 7, 14, 24 days	Naph sum		2549 ng/g	4,6 - 13,4 - 23,6	Skadsheim et al., 2004
				Phen sum		691 ng/g	7,7 - 22,9 - 34,9	
				Pyren		27 ng/g	7,3 - 16,2 - 25,1	
Cod (Gadus morhua)	Laboratory	0.06 - 0.25 - 1 ppm Oil	average 3, 17, 31 day	Naph sum		5702 ng/g	4 - 13,3 - 12,7	Skadsheim et al., 2004
				Phen sum		377 ng/g	10,5 - 40,3 - 48,7	
				Pyren		5 ng/g	8,6 - 63 - 88,4	
Cod (Gadus morhua)	Field, Caged	North Sea - Stafford, 500 - 2000 - 10000 m		Naph sum		1150 ng/g	3,0 - 2,0 - 1,3	Aas et al., in press
				Phen sum		340 ng/g	3,5 - 2,7 - 2,5	
				Pyren		-	-	
Cod (Gadus morhua)	Field, Caged	North Sea - Troll, 1000 m - 500m	6 weeks	Naph sum	1515,1 ng/g	1,1	1,1 - 1,2	Børseth et al., 2004
				Phen sum	327,2 ng/g	1,6	2,1 - 2,0	
				Pyren	173,2 ng/g	1,2	0,9 - 1,2	
Cod (Gadus morhua)	Field, Caged	North Sea - Tampen, 10000 - 2500 - 1000 - 500	6 weeks	Naph sum		965,3 ng/g	0,9 - 1,7 - 0,9 - 1	Hylland et al., 2005
				Phen sum		934,5 ng/g	1,4 - 3 - 1,8 - 1,5	
				Pyren		3,7 ng/g	0 - 0 - 0,5 - 0,0	
Cod (Gadus morhua)	Field, Caged	North Sea -	5.5 weeks	Naph sum	228 ng/g	0,2	0,9 - 1,1 - 0,9	Aas et al., in press

Species	Substance (lab/field)	Test concentrations	Exposure time	Metabolite	Baseline	control or reference	exposed/control	reference
Cod (<i>Gadus morhua</i>)	Field, Caged	Statfjord, 10000 m – 2000m - 500 m	5.5 weeks	Phen sum	482 ng/g	2.0	3 - 4.5 - 6.7	Aas et al., in press
				Pyren	28 ng/g	10,2	29.5 - 31.1 - 41.5	
				Naph sum	228 ng/g	0,8	1 - 1 - 1.9	
Haddock (<i>Melanogrammus aeglefinus</i>)	Feral fish	Egersund	Baseline non polluted area	Phen sum	482 ng/g	1,0	0.7 - 0.8 - 0.8	Klungsoyr et al. 2003
				Pyren	28 ng/g	0,0	0 - 0 - 0	
				Naph sum	1346.9 ng/g			
Haddock (<i>Melanogrammus aeglefinus</i>)	Feral fish	Sleipner	Baseline polluted area?	Phen sum	526.8 ng/g			Klungsoyr et al. 2003
				Pyren	5.7 ng/g			
				Naph sum	1111.5 ng/g			
Haddock (<i>Melanogrammus aeglefinus</i>)	Feral fish	Statfjord	Baseline polluted area?	Phen sum	331.5 ng/g			Klungsoyr et al. 2003
				Pyren	10.4 ng/g			
				Naph sum	1279.7 ng/g			
Haddock (<i>Melanogrammus aeglefinus</i>)	Feral fish	Barents sea		Phen sum	331.9 ng/g			Sundt, 2004
				Pyren	3.1 ng/g			
				Naph sum	1474 ng/g			
Polar cod (<i>Boreogadus saida</i>)	Laboratory, feral fish 2001, 2002	1.5 ppm Statfjord oil, baseline, control	14 days	Phen sum	165 ng/g			Sundt & Bechmann, 2004
				Pyren	0			
				Naph sum	1330 ng/g	1,3	114	
				Phen sum	538 ng/g	0,9	90	
				Pyren	52 ng/g	14,6	60	

3. Cytochrome P4501A activity (EROD)

3.1 Introduction

EROD (7-Ethoxyresorufin-*O*-deethylase) is a specific cytochrome P450 reaction where ethoxyresorufin is the substrate (Burke and Mayer 1974). Cytochrome P450 1A catalyse the deethylation of 7-ethoxyresorufin to resorufin. CYP activity in microsome fractions can be quantified from the amount of resorufin produced.

The cytochrome P450 system is a large superfamily of enzymes with several hundred forms comprising more than 250 different families, further divided into subfamilies. The CYP system is a highly diversified set of proteins and is found in bacteria, plants, lower eukaryotes and in animals. Members of the P450 subfamily CYP1A are particularly important in the metabolism of many pollutants. In the case of planar molecules, such as polycyclic aromatic hydrocarbons (PAHs) isoenzymes of CYP1A are responsible for the insertion of oxygen into the molecule, which is the first oxidative step in the biotransformation process (termed 'phase I'; Williams, 1974). The introduced oxygen creates a functional group for attachment of larger polar molecules of endogenous origin in a 'phase II' reaction (which includes e.g. glutathione *S*-transferases).

In addition to being substrates for biotransformation, planar compounds, such as PAHs, can also interact with cytochrome P450 1A as inducers, by binding to the cytosolic Ah (aryl hydrocarbon)-receptor. EROD is a tool used to quantify this induction. The induction of cytochrome P450 enzymes in fish liver was first suggested as an indicator of environmental contamination in the 1970s by Payne (1976). It has later gained widespread use, and the results of the most relevant studies for this review will be addressed in the following.

3.2 Dose-response

Whyte *et al.* (2000) rank chemicals according to the level of EROD activity they induce in treated or exposed fish when compared with untreated or control fish. Contaminants that induce EROD less than 10-fold above control levels are considered "weak" inducers, 10- to 100-fold are "moderate" inducers, and chemicals that elicit > 100-fold induction are considered "strong" inducers. Dioxins, planar PCBs and PAHs (benzo[a]pyrene) are categorised as "strong" inducers.

Over 25 studies have observed induction of hepatic EROD by benzo[a]pyrene in 15 species of fish (Whyte *et al.* 2000). However, in the present overview, results for Gadoid species are focussed on, due to their relevance in offshore monitoring studies in the Norwegian sector.

Table 4. Dose-response, background response and sensitivity in gadoid fish.

species	substance(s)	lowest-highest concs	exposure time	baseline/control (level/activity)	induction (fold)	reference
Experimental studies						
Polar cod <i>Boreogadus saida</i> juvenile	Crude oil (Oseberg C)	~200 mg/kg (i.p. inj.)	10 and 21 d post inj.	~30 pmol/min/mg	~8 and ~2.5 (245 and 80 pmol/min/mg)	(George et al. 1995)
Polar cod <i>Boreogadus saida</i> male	Crude oil (Oseberg C)	~200 mg/kg (oral)	21 d post exposure	28 pmol/min/mg ± 6 (n=12)	~5 (132 ± 14 pmol/min/mg)	(George et al. 1995)
Polar cod <i>Boreogadus saida</i> female	Crude oil (Oseberg C)	~200 mg/kg (oral)	21 d post exposure	8 pmol/min/mg ± 2 (n=14)	~5 (42 ± 6 pmol/min/mg)	(George et al. 1995)
Polar cod <i>Boreogadus saida</i> juvenile	β-naphthoflavone	50 mg/kg (i.p. inj.)	21 d post inj.	~30 pmol/min/mg	~12.5 (380 pmol/min/mg)	(George et al. 1995)
Cod, <i>Gadus morhua</i> juvenile	2,3,7,8-TCDD	0.008 mg/kg oral dose twice, d 0 and d 4	9 and 17 d post exposure	55.4 (d 9) and 91.4 (d 17) pmol/min/mg	~4 and ~3 (230 and 277 pmol/min/mg)	(Hektoen et al. 1994)
Cod, <i>Gadus morhua</i> juvenile	PCB-105	10 mg/kg oral dose twice, d 0 and d 4	measure at d 9 and d 17	55.4 (d 9) and 91.4 (d 17) pmol/min/mg	1.5 and 1.2	(Bernhoft et al. 1994)
Cod, <i>Gadus morhua</i> juvenile	β-naphthoflavone	100 mg/kg (i.p. inj. at d 0 and d 4)	measure at d 7	84 pmol/min/mg ± 8 (n=5)	~13 (1074 ± 340 pmol/min/mg)	(Goksoyr et al. 1987)
Cod, <i>Gadus morhua</i>	β-naphthoflavone	100 mg/kg (2 i.p. inj.)	measure 3-4 d after last injection	40 pmol/min/mg	~72 (2870 pmol/min/mg)	(Goksoyr et al. 1991)
Cod, <i>Gadus morhua</i> juvenile	Crude oil (North Sea)	0.06 – 1 ppm	30 days	~2 pmol/min/mg	~ 2- 5.5 (~ 4 – 11 pmol/min/mg)	(Aas et al. 2000)
Field studies						
Rockling, <i>Ciliata mustella</i>	Crude oil (Gulfaks; M.V. Braer spill, Shetland)	85000 tons spill 129 ± 38 ng /g dry wt. of PAHs (selected 2- and 3-ring) detected in muscle.	3 months after spill	~160 pmol/min/mg ± 50	~9 (1480 pmol/min/mg)	(George et al. 1995)
Roundnose grenadier, <i>Coryphaenoides rupestris</i>	i.a. PAHs and PCBs			260 ± 20 (Male) ~170 (Female) pmol/min/mg	~2 (530 ± 70 (male) and ~350 (female) pmol/min/mg)	(Lindesjoo et al. 1996)
Burbot, <i>Lota lota</i>	Bleached-kraft mill			20 pmol/min/mg	1.7, not sign.	(Kloppersams and

species	substance(s)	lowest-highest concs	exposure time	baseline/control (level/activity)	induction (fold)	reference
(Freshwater)	effluents (i.a. dioxins)			± 10	(34 ± 23 pmol/min/mg)	Benton 1994)
Hake, <i>Urophycis</i> spp.	Pollution (PAH) from oil platforms (Gulf of Mexico) <100m from platforms			10.9 ± 6.4 and 11.7 ± 10.5 pmol/min/mg (>3000 m from platforms)	<1 (10.6 ± 3.8 and 10.5 ± 7.1 pmol/min/mg)	(McDonald et al. 1996)

Table 4 continued.

<i>species</i>	<i>substance(s)</i>	<i>exposure time</i>	<i>baseline/control (level/activity)</i>	<i>induction (fold)</i>	<i>reference</i>
Caging studies					
Cod, <i>Gadus morhua</i> juvenile	i.a. PAH and PCB from industry (smelters)	4 weeks	~55 pmol/min/mg (farthest from source)	~3 (closest to source)	(Goksoyr <i>et al.</i> 1994)
Cod, <i>Gadus morhua</i> juvenile	i.a. PAH and PCB from industry (smelters)	3 months	16 ± 3 pmol/min/mg (farthest from source)	~2 (closest to source)	(Beyer <i>et al.</i> 1996)

3.3 Blue mussel

Cytochrome P450-related activity is less well described in mussels or other invertebrates compared to fish. The assay most commonly used is benzo[a]pyrene hydroxylase (BaPH), for which there is some experience from both research and monitoring (e.g. Michel *et al.*, 2001).

3.4 Species sensitivity

As will be apparent from the previous section, PAH-related induction of metabolising enzymes (e.g. cytochrome P4501A) is expressed to a much larger extent in fish compared to most invertebrates. It is not entirely clear whether this is because of the assay (e.g. substrate) or whether it is the biochemistry of the organisms. High bioaccumulation of PAHs in, for example, mussels compared to fish support the latter view.

3.5 Relevance of other factors

Several factors have been shown to affect hepatic EROD, both endogenous and exogenous. The most important endogenous factors for most fish species are developmental stage (juvenile-mature), gender, reproductive status and age, all of which can be controlled through sampling design. In addition, environmental temperature has been shown to affect EROD (Sleiderink *et al.*, 1995; Lange *et al.*, 1999). Seasonal cycles in EROD induction have been observed for e.g. rainbow trout (Förlin & Haux 1990), flounder (Nissen-Lie 1997; Hylland *et al.*, 1998), salmon (Larsen *et al.*, 1992), most likely due to both to changes in water temperature and reproductive cycles (which it is not really possible to separate in the field).

Several species have baseline EROD activities within the same order of magnitude among different studies/measurements and also show greater than 10-fold EROD induction after contaminant exposure (Whyte *et al.* 2000). These are, however, mostly freshwater species. Developmental stage of the fish is very important. The main age-related factors are time of exposure/accumulation, food selection and reproductive stage.

The mechanism for CYP1A suppression in spawning females is related to 17 β -estradiol (E₂) (or xenoestrogen) levels. The hormone controls the induction of vitellogenin (VTG; egg yolk protein) production during gonadal recrudescence. Some of the inter-gender differences during spawning can be attributed to increased levels of CYP isoenzymes in males rather than suppression of levels in females.

Dietary factors can be important for the induction of CYP1A. Firstly, of course, AhR ligands can be presented to the organism through the food. Secondly, proper nutrition is a prerequisite for enzyme systems to function properly. Hylland *et al.* (1995) reported an elimination of EROD response (i.e. to control levels) in BaP-treated flounder deprived of food for one month.

3.6 Links

Cytochrome P4501A (CYP1A) activity is one of the parameters which have been included in a causality model of how PAHs in sediments may cause effects on fish that lead to the development of liver cancer (neoplasia) (Myers *et al.*, 2003). There is therefore evidence for a correlation between CYP1A activity, DNA adduct formation and development of neoplasia. Changes in EROD may also affect normal steroid metabolism, but evidence for such effects is tenuous.

3.7 Background responses

Baseline levels of EROD in four marine species have been estimated from results derived from the Norwegian monitoring programme (Ruus *et al.*, 2003). The baseline value for Atlantic cod has been set to 9-95 pmol/min/mg protein including fish from the Norwegian west coast and if only fish from the Barents sea had been included, the values would have been 9-25 pmol/min/mg protein. For flounder, baseline values are in the range 10-43 pmol/min/mg protein, for dab 123-529 pmo/min/mg protein and for plaice 33-146 pmol/min/mg protein. The fish were sampled from reference locations (i.e. no known local sources of contamination) in the autumn, the data includes males and females and the water temperature at the sampling locations was 9-11°C.

3.8 Assessment criteria

As many factors are known to influence EROD and it is not possible to correct for all in the design, it is advisable always to include an appropriate reference group in studies that include EROD as an endpoint. Experience suggests that an EROD value in most marine species above twice the upper limit of baseline values indicate an ecosystem influenced by planar organic contaminants.

3.9 Quality assurance

Cytochrome P4501A is possibly the most widely used biomarker. There have been two international intercalibrations for the method, both within BEQUALM. The intercalibrations have pinpointed variability relating to most steps in the analytical process, excepting possibly the enzyme kinetic analysis itself. It is imperative that laboratories have internal quality assurance procedures, e.g. use internal reference samples with all batches of analyses.

4. Glutathione *S*-transferase activity

4.1 Introduction

Glutathione *S*-transferase enzymes (GSTs) are a group of cytosolic enzymes that are involved in phase-II biotransformation of a large number of pollutants (Lindstrom-Seppä *et al.*, 1986). These enzymes catalyse the conjugation of glutathione (GSH) to lipophilic compounds with electrophilic centres in order to neutralise and excrete chemicals capable of causing toxic effects. To date, several groups of enzymes (isoenzymes) have been reported and number of isoenzymes and their characteristics vary between different animals and tissues (Fitzpatrick and Sheehan, 1993; Gallagher *et al.*, 1996; Martinez-Lara *et al.*, 1997; Angelucci *et al.*, 2000; Novoa-Valinas *et al.*, 2002). Glutathione *S*-transferase is one the most efficient phase-II biotransformation pathways for potentially toxic chemicals in vertebrates and invertebrates and has been characterized in various species including polychaetes, clams, gastropods, mussels, crustaceans and fish (see Table 5 for details).

Many of the GST isoenzymes can be induced by pollutants and increased hepatic GST activity has been reported in several studies after exposure to chemicals (Perez-Lopez *et al.*, 2002). The induction of GST has thus been suggested to be adopted for use as a biomarker for exposure to chemicals such as such as PAHs, PCBs and Dioxins (Van der Oost *et al.*, 2003).

Total GST level is commonly measured by means of a catalytic assay involving the conversion of the substrate 1-chloro-2,4-dinitrobenzene (CDNB). The activity of individual isoenzymes, which is expressed as specific catalytic activity, may in a similar fashion be measured with the use of isoenzyme specific substrates. Although GST activity has been documented in various tissues, determination of GST is predominantly performed on tissues that display high biotransformation activity. This applies in particular to use of liver tissues in vertebrates and the hepatopancreas/digestive gland in invertebrates, although gill GST is of importance as the first line defence towards waterborne pollutants. Measurement of GST is most commonly performed on the cytosolic fraction (CF) obtained after high speed centrifugation.

4.2 Dose-response

Dose-response relationships are commonly obtained for the induction of total GST activity following exposure to phase-II substrates under controlled laboratory conditions (experimental studies), sampling of wild fish (field studies) and caging of fish (caging deployment studies). Although many pollutants are documented to induce GST activity, proper dose-response relationships are difficult to derive from the reported data due to lack of measured concentrations. In general, GST is only weakly inducible with an induction factor of up three times in laboratory studies and up to five times in field studies. GST activity is inhibited compared to control groups in several studies with pure chemicals and mixtures as well as under field experiments. Table 5 shows the dose-response relationships for selected fish and invertebrates after exposure to pure chemicals and environmental mixtures. This biomarker has been measured in both fish and invertebrates. It may be both up- and down regulated as a result of contaminant exposure.

Table 5. Dose-response, background response and induction of GST in fish.

species	Tissue	substance(s) Field/caging locality	concentration	exposure time (days)	baseline/control (nmol/min/mg protein)	induction (%)	reference
Experimental studies							
Olive Flounder, <i>Paralichthys olivaceus</i>	Liver CF	Phenanthrene	0.5, 1, 2 uM (nominal)	14, 28	650±70 (14d) 640±30 (28d)	0 (14d, L-CF) 0 (28d, L-CF)	(Jee and Kang, 2005)
	Gill CF	Phenanthrene	0.5, 1, 2 uM (nominal)	14, 28	290±20 (14d) 360±50 (24d)	+300 (14d) +258 (28d)	(Jee and Kang, 2005)
	Kidney CF	Phenanthrene	0.5, 1, 2 uM (nominal)	14, 28	230±20 (14d) 250±30 (28 d)	0 (14d) 0 (28d)	(Jee and Kang, 2005)
Atlantic cod <i>Gadus morhua</i>	Liver S9	4-t-butylphenol 4-n-pentylphenol 4-n-hexylphenol 4-n-heptylphenol	0,02-80 µg/kg (nominal, oral)	28	952 ± 153	-20% (only lowest concentration)	(Hasselberg <i>et al.</i> , 2004)
Pacific herring, <i>Clupea pallasii</i>	Liver CF	WSF crude oil	∑PAH 9.7-99.3 µg/L. (measured, unstable)	2.4, 4	101.1± 8.5 (2.4d) 103.8±3.6 (4d)	+69 (2.4d) 0 (4d)	(Kennedy and Farrell, in press)
Rainbow trout <i>Onchorynchus mykiss</i>	Liver CF	Nonylphenol	66 and 220, µg/l (nominal)	7,14,21	68.8 ± 9.7 (7d) 83.7 ± 11.5 (14d)	+200 (7d, 220 µg/l) +127 (7d, 66 µg/l) -24 (21d, 66µg/l)	(Uguz <i>et al.</i> , 2003)
Rainbow trout <i>Onchorynchus mykiss</i>	Liver CF	BNF CB-77 CB-126 (i.p. injection)	50mg/kg (BNF) 0.1, 1, 5 mg/kg (CB-77, CB-126) (nominal)	6	310±96	-18 (BNF) +75 (CB-77, 0.1) +55 (CB-77, 1) +23 (CB-77, 5) +47 (CB-126, 5) +74 (CB-126, 1) +23 (CB-126, 5)	(Huuskonen <i>et al.</i> , 1996)
Rainbow trout <i>Onchorynchus mykiss</i>	Liver CF	Arochlor-1254 (i.p. injection)	5 mg/kg (nominal)	63	493±147	-23 (BNF) +29 (CB-77, 0.1) +8 (CB-77, 1) -25 (CB-77, 5) -31 (CB-126, 5) -18 (CB-126, 1) +2 (CB-126, 5)	(Huuskonen <i>et al.</i> , 1996)
Field studies							
Flounder <i>Platichthys flesus</i>	Liver S9	Southern Baltic Sea	n.a.	Feral	86.6±39.8	+48%	(Napierska and Podolska, 2005)
Cage deployment studies							
Atlantic cod <i>Gadus morhua</i>	Liver CF	Troll Oil field, North Sea	n.a.	42	~750	Approx. +13% (1000 m station)	(Børseth and Tollefsen, 2004)

species	Tissue	substance(s) Field/caging locality	concentration	exposure time (days)	baseline/control (nmol/min/mg protein)	induction (%)	reference
Atlantic cod <i>Gadus morhua</i>	Liver CF	Statfjord B, North Sea	n.a.	42	~950	Approx. -29% (males, 1000 m station)	(Hylland <i>et al.</i> , 2005)

n.a. – not analysed, i.p. -intrapertitoneal, WSF - water soluble fraction of crude oil, CF – cytosolic fraction, S9 – S9 fraction obtained after high speed centrifugation.

Table 6. Dose-response, background response and induction of GST in invertebrate species.

species	Tissue	substance(s) Field/caging locality	concentration	exposure time (days)	baseline/control (nmol/min/mg protein)	induction (%)	reference
Experimental studies							
Mediterranean mussels <i>Mytilus galloprovincialis</i>	Gill S9	BaP (feed)	50 mg B[a]P:kg d.w. (theoretical)	28	~170	Approx. -9	(Akcha <i>et al.</i> , 2000)
<i>Lymnaea palustris</i>	Whole Body CF	Atrazine	5, 25, 125 µg/L (nominal)	21	~1040-1125	-16,4 (5 µg/l) -14,9 (25 µg/l) -14,3 (125 µg/l)	(Baturó and Lagadic, 1996)
Shore crab <i>Carcinus maenas</i>	Hepatopan creas S9	Cypermethrin	50, 200, 500ng/l (nominal) Variable measured levels	7	~38	+46 (50 ng/l) +27 (200 ng/l) +53 (500 ng/l)	(Gowland <i>et al.</i> , 2002)
Field studies							
<i>Nereis diversicolor</i>	Digestive Gland S9	Cádiz bay (Spain)	n.a.	Feral	25.4±3.6	+200	(Perez <i>et al.</i> , 2004)
<i>Scrobicularia plana</i>	Digestive Gland S9	Cádiz bay (Spain)	n.a.	Feral	392±69	0	(Perez <i>et al.</i> , 2004)
Cage deployment studies							
Blue mussels <i>Mytilus edulis</i>	Hepatopan creas S9	Loch Leven (Scotland)	Mussel concentration ΣPAH 15-927 ng/g w.w.	Variable (study design)	10.2	+456	(Gowland <i>et al.</i> , 2002)
Green lipped mussels <i>Perna viridis</i>	Digestive Gland S9	Various harbours, (China)	Mussel concentration 0-300 ng PAH/g d.w. 0-90 ngBaP/g d.w.	30	61-95	Approx. +40	(Cheung <i>et al.</i> , 2001)
	Gill S9	Various harbours, (China)	Mussel concentration 0-300 ng PAH/g d.w. 0-90 ngBaP/g d.w.	30	90-120	Approx. +45	(Cheung <i>et al.</i> , 2001)

n.a. – not analysed, i.p. -intrapertitoneal, WSF - water soluble fraction of crude oil, CF – cytosolic fraction, S9 – Sp fraction obtained after high speed centrifugation, BNF – beta-naphthoflavone; CB - chlorobifenyyl.

4.3 Species sensitivity

No studies are available that address sensitivity of GST to exposure to pollutants directly, although it seems that GST are induced to a higher degree in invertebrates than in vertebrates. This difference in sensitivity may be caused by the lower constitutive levels of GST in invertebrates compared to vertebrates. Up to 8 times differences in basal levels of GST has been documented in fish species (Förlin *et al.*, 1995), thus suggesting that interspecies differences in GST induction sensitivity may also be relevant for fish. Laboratory studies and field studies with English sole (*Parophrys vetulus*) and starry flounder (*Platichthys stellatus*) confirm that susceptibility towards contaminant-associated hepatic lesions is partly related to differences in constitutive levels of GST in these fish (Collier *et al.*, 1992).

4.4 Relevance of other factors

GST activity is constitutively expressed and basic activity may be affected by various factors related to the physiological state of the organisms such as food availability, diet gender and maturation status. Several reports document that GST activity may be affected by co-exposure to other pollutants, algal toxins, and but also to infections by pathogen organisms such as *Listeria monocytogene*. All these factors may potentially influence the ability to determine effects in the field either as a result of inhibition of GST, leading to false negatives, or to induction of GST, leading to false positives. Table 7 depicts factors that are known to influence GST activity in fish and vertebrates.

Table 7. Relevance of other factors.

Species	Tissue	Factor	Response	Comment	Reference
Rainbow trout <i>Oncorhynchus mykiss</i>	Liver CF	Food deprivation	Decreased GST activity	Reduced to 50% control activity	(Blom <i>et al.</i> , 2000)
Various fish species	Liver CF	Diet composition	No change	GST activity is not affected by diet composition.	(DeKoven <i>et al.</i> , 1992; Guderley <i>et al.</i> , 2003; Morrow <i>et al.</i> , 2004)
Blue mussels, <i>Mytilus edulis</i>	Digestive Gland & Gill S9	Seasonal variation	Variable response	~200% difference between high and low activity throughout year	(Manduzio <i>et al.</i> , 2004)
Sockeye salmon (<i>Oncorhynchus nerka</i>)	Liver S9	Maturation status	Decrease	GST decrease at spawning	(Kennish <i>et al.</i> , 1992)
Flounder <i>Platichthys flesus</i> Bullrout <i>Myoxocephalus scorpius</i>	Liver S9	Gender	GST levels dependent on gender	Males display higher GST activity than females	(Ruus <i>et al.</i> , 2002; Napierska and Podolska, 2005)
Arctic Charr, <i>Salvelinus alpinus</i> .	Liver CF	Chemical co-exposure	Inhibition of GST	TBT inhibited GST activity 37-60 %	(Padros <i>et al.</i> , 2003)
Mediterranean mussels <i>Mytilus galloprovincialis</i> .	Gill CF	Chemical co-exposure	Inhibition of GST	Retene inhibited GST activity	(Gravato <i>et al.</i> , 2004)
Atlantic salmon, <i>Salmo salar</i>	Liver CF	Exposure to toxic algae	Induction of GST	Paralytic shellfish poisoning toxins induce GST (~250%)	(Gubbins <i>et al.</i> , 2000)
Carp <i>Cyprinus carpio</i>	Liver Kidney Gill CF	Bacterial infections	Inhibition of GST	Listeria monocytogenes infection reduce GST activity	(Chambras <i>et al.</i> , 1999; Dautrempuits <i>et al.</i> , 2002)

4.5 Links

Since GST activity is a biomarker for exposure rather than effect, direct linkage between alteration in GST levels and effects on health, population parameters and community composition are limited. Alterations in GST activity may affect the ability to detoxify potentially toxic chemicals and consequently affect these parameters indirectly. Several studies support this assumption:

Loss or inhibition of hepatic GST activity may increase susceptibility to liver neoplasms and to DNA damage by environmental or endogenous chemicals that are normally detoxified by GST (Stalker *et al.*, 1991; Kirby *et al.*, 1995; Gravato and Santos, 2003). Laboratory and field studies by Collier *et al.* (1992) suggest that the prevalence in contaminant-associated liver lesions is related to a combination of induction of Phase-I biotransformation pathways causing bioactivation of certain PAHs and lower constitutive levels of detoxification activity by GST. In European flounder (*Platichthys flesus L.*) that has developed liver carcinomas, expression of multi-xenobiotic resistance (MXR), phase-I and Phase-II biotransformation activities (including GST) were induced as clonal adaptation to pollution exposure (Köhler *et al.*, 1998; Koehler *et al.*, 2004).

GST levels were found to increase in parallel with DNA adducts in livers of gold-lined sea bream (*Rhabdosargus sarba*) after intraperitoneal exposure to benzo[*a*]pyrene (Xu *et al.*, 2001). In vitro studies with hepatocytes from rainbow trout show that GST activity is unregulated and DNA integrity, DNA damage and prevalence of apoptotic cells were increased after exposure to wastewater (Faverney *et al.*, 2001; Risso-de Faverney *et al.*, 2001). GST induction was found to increase in parallel with lipid peroxidation and DNA damage in the digestive glands of the mangrove mussel (*Mytella guyanensis*) after exposure to polluted waters (Torres *et al.*, 2002). In other studies, GST levels were found to be correlated to exposure to PAHs such as benzo[*a*]pyrene, but DNA adducts were not (Akcha *et al.*, 2000; Bocquene *et al.*, 2004).

4.6 Background responses

Background responses vary within animal groups and individual species. Table 8 indicates ranges and average/median levels of GST activity for selected vertebrates and invertebrates.

4.7 Assessment criteria

There is insufficient information about factors that affect GST to develop assessment criteria at this time.

4.8 Quality assurance

The analysis of GST activity using CDNB is straightforward and most laboratories use the procedures outlined by Habig *et al.* (1974). Laboratories that analyse for the protein must establish in-house quality assurance procedures.

Table 8. Background levels of GST in different tissues of vertebrates and invertebrates.

Animal group	Species	Tissue	Background response (nmol/min/mg protein)
Vertebrates	Olive Flounder, <i>Paralichthys olivaceus</i>	Liver CF	640-650 (average 645)
		Gill CF	290-360 (average 325)
		Kidney CF	230-250 (average 240)
	Rainbow trout, <i>Onchorynchus mykiss</i>	Liver CF	68.8-493 (median 310)
	Atlantic cod <i>Gadus morhua</i>	Liver CF	750-952 (median 950)
	Flounder <i>Platichthys flesus</i>	Liver S9	86.6
	Pacific herring, <i>Clupea pallasii</i>	Liver CF	101.1-103.8 (average 102)
Invertebrates	Mediterranean mussels <i>Mytilus galloprovincialis</i>	Gill S9	170
	Blue mussels <i>Mytilus edulis</i>	Hepatopancreas S9	10.2
	Green lipped mussels <i>Perna viridis</i>	Digestive Gland S9	61-95 (average 78)
		Gill S9	90-120 (average 105)
	Shore crab <i>Carcinus maenas</i>	Hepatopancreas S9	~38
	<i>Nereis diversicolor</i>	Digestive Gland S9	25.4
	<i>Scrobicularia plana</i>	Digestive Gland S9	392
<i>Lymnaea palustris</i>	Whole body CF	1040-1125 (average 1083)	

5. DNA damage

5.1 Introduction

It is of vital importance to the survival of the cell and of the organism that DNA is kept intact and unchanged. All cells therefore have an extensive system of quality control and repair involved in the replication and transcription of DNA. Since normal cellular processes will generate damage to DNA continuously, any impairment of repair systems will obviously also potentially result in DNA damage. In addition to intracellular processes, there are immunological mechanisms (in vertebrates) that will identify and kill cells with DNA modifications that may be deleterious to the organism (i.e. cancer). DNA damage may simply cause cell death, but may also lead to carcinogenic changes in the cell, structural damage to chromosomes and mutagenic effects.

There are three main direct or indirect mechanisms that may lead DNA damage – breaks, adducts and impaired repair. There are a range of methods in ecotoxicology that address breaks and adduct formation, but few have been established to assess repair mechanisms. A distinction is commonly made between methods that identify chromosomal breaks and those that are used to quantify all breaks in DNA. It is also possible to distinguish between one- and two-strand breaks in the latter category. Micronucleus formation (see section 7) is the most widely used method to address chromosomal breaks that result in the formation of satellite DNA. The two most widely used assays for breaks are alkaline unwinding and the Comet assay. The former will be discussed briefly in this section, the Comet assay will be treated in more detail in section 6.

5.2 DNA repair

Although presumably critical in the maintenance of DNA integrity, DNA repair has not been extensively studied in fish. The results found by Willett et al. (2001) may indicate part of the problem – they found no detectable nucleotide excision repair (NER) in hepatocytes from two catfish species following fairly harsh UV-treatment. There are however other studies in which such activity has been detected, e.g. Walter et al. (2001), David et al. (2004). This is obviously a field which needs further research.

5.3 Alkaline unwinding

As mentioned above, different methods are being used to address DNA damage in aquatic organisms. Different methods may yield results which differ by many orders of magnitude (Collins 2005) and it is obviously critical that comprehensive documentation is available for each study. Alkaline unwinding assays can be set up to distinguish between single and double breaks and have been used in fish (Shugart 1988, Everaarts 1995, Liepelt et al. 1995, Feng et al. 2003) and in mussels (Accomando et al. 1991, Nacci et al. 1992, Ching et al. 2001). The principle of the technique is the same as that used in the Comet assay, but the latter has the advantage of measuring effects on single cells. There is not really sufficient information to review alkaline unwinding in terms of sensitivity or dose-dependency for species relevant to North Sea monitoring. Results from alkaline unwinding have failed to relate to contaminant exposure in e.g. mussels (Siu et al. 2003).

5.4 DNA adducts

There is a considerable number of ecotoxicological studies that have used DNA adducts as a measure of genotoxic effects. There is also a range of different methods and units to present results, despite the existence of an internationally acknowledged manual (Reichert et al., 1999). In the tables below, only studies that use the Nuclease P1 enrichment technique have been considered.

5.4.1 Dose-response

There is a limited number of toxicokinetic studies with DNA adducts or alkaline unwinding as endpoints. It is also difficult to use results from studies earlier than the mid-1990s due to obvious developments in both experimental and analytical protocols. The majority of studies use fish or mussels sampled in the field. In general there is only limited knowledge of exposure history in field studies.

There is not always a straightforward dose-response relationship. In some studies, e.g. Ericson et al. (1999), decreased concentrations of DNA adducts were found in intestine at dietary exposures higher than 150 μmol PAH-mixture/kg, presumably due to cytotoxic effects. Adduct concentrations in the liver of that study remained high even at higher levels (but did not increase further). An overview of field and laboratory studies with fish can be found in Table 9 and Table 10, studies with mussels can be found in Table 11 and Table 12.

5.4.2 Species baseline and sensitivity

Species differences in DNA adduct concentrations may derive from different capacities of phase-1 or phase-2 metabolism, as suggested for two flatfish species by Anulacion et al. (1998) or by differences in the rate of which adducts are cleared from tissues, as suggested for two species of catfish by Ploch et al. (1998).

5.4.3 Time-course of adduct development

The time-course of adduct development has been studied both following dietary exposure and exposure through water.

In a study by Ericson et al. (1999), DNA adducts were found to be significantly elevated in PAH-fed pike (*Esox lucius*) after 3 weeks (the first sampling point) and then increased beyond the last feeding (day 50) until day 83 for adducts in intestine, liver and gills. There were still significantly (> 100 -fold) increased concentrations of DNA adducts in the three tissues after 128 days (2.5 months after last exposure). A second study with pike explored shorter time scales and dose-dependency. Following a single i.p. injection of a PAH-mixture, adducts increased in liver until day 9, but then dropped to day 12. In the same experiment, adducts increased in intestine as well and levelled off on day 9. In an accompanying feeding experiment, a single bolus with a PAH-mixture resulted in maximally increased adduct levels in intestine already after three days, decreasing on day 12. The largest mean increase of hepatic adducts was seen on day 6.

In a study reported by Aas et al. (2000), DNA adducts in the liver of Atlantic cod were apparent after only 3 days at the highest exposure level, dosed through water (0.94 mg/L dispersed oil; 0.6 ppb PAH). The adduct levels increased until 30 days, at the time of which there was also a significant increase in adducts in a second group exposed to 0.04 mg/L dispersed oil.

In the American freshwater species mummichog (*Fundulus heteroclitus*), DNA adducts increased in all tissues (liver, anterior kidney and spleen) until 32 days after a single injection of benzo[*a*]pyrene (12 mg/kg) (Rose et al., 2001). Levels of adducts decreased significantly by day 96 (Rose et al., 2001).

5.4.4 Relevance of other factors

Fish

Carpenter et al. (1995) found weak temperature effects *in vitro* (higher adduct levels at higher temperatures), but generally similar adduct formation when assays were run at acclimation temperature.

The results found by Akcha et al. (2003) show that there may indeed be detectable levels of DNA adducts in all dab populations in an area, but also indicated that there could possibly be a sex difference in the sensitivity to adduct formation.

External agents other than PAHs may also cause adduct formation. Armstrong et al. (2002) found that UV-radiation would cause DNA damage (although not measured as adducts) in medaka (*Oryzias latipes*). Other contaminants, e.g. CBs, may also cause adducts as shown by e.g. Dubois et al. (1995).

Some studies have also shown that DNA adducts will increase at different rates in different tissues (e.g. Ericson et al., 1999). The liver is the main tissue used in most cases and there is not sufficient data to suggest that other tissues be used at the moment. A study is ongoing to investigate this possibility with Atlantic cod (Balk, personal communication).

There is no clear evidence that spawning affect DNA adducts in other tissues or for seasonality in the formation or concentration of DNA adducts.

Mussels

Not surprisingly, DNA adducts in blue mussels have been shown to have a seasonal variability (Skarphedinsdottir et al. 2005), presumably due to spawning activity and variations in food availability (i.e. fat content) through the year. In the same study evidence was presented that intertidal mussels have higher concentrations of DNA adducts than subtidal mussels, not correlated to carcinogenic PAHs in the tissues, but to a larger extent to total PAH (Skarphedinsdottir et al. 2005).

There may also be differences between organs in the response to contaminants, as found by Pisoni et al. (2004), in which there was good agreement between results for gills and hepatopancreas as concerned the reference location, but results for contaminated locations varied between sampling campaigns.

5.4.5 Links

DNA adducts have been used as an intermediate between chemical or biochemical endpoints, i.e. PAH-metabolites and cytochrome P4501A activity, and histological changes in tissues, i.e. neoplasia or cancers (Myers et al., 2003). There is thus evidence that increased DNA adducts are linked to cancer in fish, but still unclear whether this will actually decrease growth or reproductive output. Recent data has indicated that adduct formation is found at concentrations similar to those that will cause reproductive effects in marine organisms (Bechmann, Sanni & Smit, personal communication).

5.4.6 Background responses

Aas et al. (2003) investigated background levels in a range of marine species from the Norwegian Sea, i.e. polar cod (*Boreogadus saida*), daubed shanny (*Leptoclinus maculatus*), sea tadpole (*Careproctus reinhardti*), Atlantic spiny lumpsucker (*Eumicrotremus spinosus*), black seasnail (*Paraliparis bathybius*), Arctic rockling (*Gaidropsarus argentatus*), doubleline eelpout (*Lycodes eudipleurostictus*), polar sculpin (*Cottunculus microps*), Greenland halibut (*Reinhardtius hippoglossoides*), Atlantic cod (*Gadus morhua*) and capelin (*Mallotus villosus*). Levels were below detection limit (from 0.4 to 1.3 nmol adducts/mol DNA) for most of individuals and species. Polar sculpin was the only species for which all sampled individuals had detectable adduct levels (mean \pm SD: 1.74 \pm 1.04 nmol adducts/mol DNA). Balk (unpublished) has indicated that background level of DNA adducts in the liver of haddock (*Melanogrammus aeglefinus*) is in a range similar to that found for other fish species, but with a distinct zone on the sheet found in all individuals. This zone is then subtracted for all to quantify contaminant-related adducts (Balk, unpublished).

In blue mussels the baseline concentration of adducts appear to be below detection limit for the normally used ^{32}P post labelling method (Skarphedinsdottir et al., 2005). This means that any adducts present were found at concentrations so low as to be inseparable from the analytical control.

5.4.7 Assessment of DNA adduct results

DNA adducts is one of the methods for which it is possible to set global reference values (using baseline values). Data as those generated by Aas et al. (2003) could be used to establish such limits, which lie in the range 0.5-1.5 nmol adduct/mol DNA for most, but not all, species. Although DNA adducts is the one biomarker methods closest to manifest disease (liver cancer) it is still not feasible to set absolute levels since other factors also affect the levels of liver cancers in fish (if that would be an acceptable "ecologically relevant" endpoint). Presumably the most useful approach is to indicate to which extent DNA adduct values in a given study exceed baseline values. A lower trigger level would be twice the baseline value.

5.4.8 Quality assurance

The measurement of DNA adducts through ^{32}P -postlabelling is a complex analysis with many steps, many of which are crucial to the final result (Dolcetti et al., 2002). As the results above indicate, results from different laboratories are not necessarily directly comparable, most commonly because they use different protocols or because they express results differently.

DNA adducts was included as one component in the EU-funded BEQUALM project and an intercalibration between different laboratories was performed by ITM, Stockholm University (led by Lennart Balk). There are plans to include DNA adducts in the ongoing BEQUALM activity, led by CEFAS (UK; project office) and NIVA (responsible for biomarkers).

Table 9. Overview of field studies in which DNA adducts were analysed using nuclease P1 enrichment (fish).

species	ecosystem	tissue	substance(s)	lowest-highest concs	baseline/control (level/activity)	induction (fold)	reference
brown bullhead (<i>Ameiurus nebulosus</i>)	freshwater	liver	field		42 nmol/mol	25-30	(Pinkney et al. 2004)
mummichog (<i>Fundulus heteroclitus</i>)	freshwater	liver	field (creosote)		<2 nmol/mol DNA	>100	(Rose et al. 2000)
mummichog (<i>Fundulus heteroclitus</i>)	freshwater	anterior kidney	field (creosote)		<2 nmol/mol DNA	>100	(Rose et al. 2000)
mummichog (<i>Fundulus heteroclitus</i>)	freshwater	spleen	field (creosote)		<2 nmol/mol DNA	>100	(Rose et al. 2000)
mummichog (<i>Fundulus heteroclitus</i>)	freshwater	pooled blood	field (creosote)		<2 nmol/mol DNA	>50 (only one sample)	(Rose et al. 2000)
perch (<i>Perca fluviatilis</i>)	freshwater	liver	field (pulp mill)	2 km and 8 km from source	<0.5 nmol/mol DNA	>30	(Ericson & Larsson 2000)
perch (<i>Perca fluviatilis</i>)	freshwater	intestine	field (pulp mill)	2 km and 8 km from source	<0.5 nmol/mol DNA	>50	(Ericson & Larsson 2000)
perch (<i>Perca fluviatilis</i>)	freshwater	liver	field (creosote)	various distances from source	0.2 nmol/mol DNA	>100	(Ericson et al. 1999)
perch (<i>Perca fluviatilis</i>)	freshwater	liver	field (smelter)	various distances from source	5 nmol/mol DNA	3	(Ericson et al. 1998)
perch (<i>Perca fluviatilis</i>)	freshwater	gill	field (smelter)	various distances from source	0.3 nmol/mol DNA	5	(Ericson et al. 1998)
perch (<i>Perca fluviatilis</i>)	freshwater	head kidney	field (smelter)	various distances from source	1 nmol/mol DNA	5	(Ericson et al. 1998)
perch (<i>Perca fluviatilis</i>)	freshwater	trunk kidney	field (smelter)	various distances from source	0.8 nmol/mol DNA	10	(Ericson et al. 1998)
Atlantic cod (<i>Gadus morhua</i>)	marine	liver	field (smelter)	various distances from source	<0.5 nmol/mol DNA	none	(Næs et al. 1999)
Atlantic cod (<i>Gadus morhua</i>)	marine	liver	field (smelter)	various distances from source	<0.5 nmol/mol DNA	>100	(Aas et al. 2003)
Atlantic cod (<i>Gadus morhua</i>)	marine	liver	caging-offshore, 5 wk	gradient from Staffjord	0.3 nmol/mol DNA	1.5-2 α	Balk et al., in press
corkwing wrasse (<i>Symphodus melops</i>)	marine	liver	field (smelter)	various distances from source	27 nmol/mol DNA	>100	(Aas et al. 2003)
dab (<i>Limanda limanda</i>)	marine	liver	field	no assoc sed pah	0.1 adducts/10-8 nucleotides α	>10	(Akcha et al. 2003)

species	ecosystem	tissue	substance(s)	lowest-highest concs	baseline/control (level/activity)	induction (fold)	reference
dab (<i>Limanda limanda</i>)	marine	liver	field	no assoc sed pah	4 adducts/10-8 nucleotides	6	(Lyons et al. 2000)
dab (<i>Limanda limanda</i>)	marine	liver	field	no difference between sites	0.2-0.4 adducts/10-8 DNA nucleotides	none	(Akcha et al. 2004)
flounder (<i>Platichthys flesus</i>)	marine	liver	field		4 adducts/10-8 nucleotides	20	(Lyons et al. 2004)
flounder (<i>Platichthys flesus</i>)	marine	liver	field		10 adducts/10-8 nucleotides	3	(Lyons et al. 1999)
grass goby (<i>Zosterisessor ophiocephalus</i>)	marine	liver	field	field	0.055 adducts/10-8 DNA nucleotides	2*	(Venier & Zampieron 2005)
grass goby (<i>Zosterisessor ophiocephalus</i>)	marine	intestine	field	field	0.371 adducts/10-8 DNA nucleotides	7-8*	(Venier & Zampieron 2005)
grass goby (<i>Zosterisessor ophiocephalus</i>)	marine	gill	field	field	0.068 adducts/10-8 DNA nucleotides	approx 30	(Venier & Zampieron 2005)
haddock (<i>Melanogrammus aeglefinus</i>)	marine	liver	field-offshore production	three areas	4 nmol/mol DNA (Egersund bank)	4.5	Hylland et al., 2006
<i>Lipophrys pholis</i>	marine	ovaries/testes	field-oil spill		n/a	3	(Lyons et al. 1999)
<i>Lipophrys pholis</i>	marine	liver	field-oil spill		n/a	1,2	(Lyons et al. 1997)
<i>Lipophrys pholis</i>	marine	gill	field-oil spill		n/a	2*	(Lyons et al. 1997)
sculpin (<i>Myoxocephalus scorpius</i>)	marine	liver	field-harbours	field	1 nmol/mol DNA	15-30	Stephensen et al., 2000

Table 10. Overview of laboratory studies in which DNA adducts were analysed using nuclease P1 enrichment (fish).

species	ecosystem	tissue	substance(s)	lowest-highest concs	exposure time	baseline/control (level/activity)	induction (fold)	reference
Atlantic cod (<i>Gadus morhua</i>)	marine	liver	dispersed oil	1 mg/L dispersed oil	2 weeks	1.6 nmol/mol DNA	10	(Aas et al. 2003)
Atlantic cod (<i>Gadus morhua</i>)	marine	liver	dispersed oil	0.94 mg/L dispersed oil	3, 16, 30	not provided	110 nmol/mol DNA	(Aas et al. 2000)
Atlantic cod (<i>Gadus morhua</i>)	marine	liver	dispersed oil	0.04 mg/L dispersed oil	day 3, 16, 30	not provided	10 nmol/mol DNA	(Aas et al. 2000)
brown bullhead (<i>Ameiurus nebulosus</i>)	freshwater	liver	injection	20 mg BaP/kg	1, 3, 7, 14, 45 d	0.6 fmol adduct/mg DNA	6x	(Ploch et al. 1998)
channel catfish (<i>Ictalurus nebulosus</i>)	freshwater	liver	injection	20 mg BaP/kg	1, 3, 7, 14, 45 d	0.5 fmol adduct/mg DNA	1.8x	(Ploch et al. 1998)
English sole (<i>Pleuronectes vetulus</i>)	marine	liver	manipulated sediments	PAH gradient in sediments	2 wk, 5 wk	5 nmol/mol DNA	>100 (5wk), >30 (2 wk)	(French et al. 1996)
flounder (<i>Platichthys flesus</i>)	brackish water	leukocytes	injection	10, 50 mg BaP/kg	10 d	n/a	>80 (pooled)	Malmström et al., 2000
flounder (<i>Platichthys flesus</i>)	brackish water	liver	injection	10, 50 mg BaP/kg	10 d	n/a	>40	Malmström et al., 2000
flounder (<i>Platichthys flesus</i>)	marine	liver	PAH-mix/diet	1, 5, 50 mg/kg	1 mo, 6 mo	n/a	2 add/10-8 (5 mg/kg), 6 add/10-8 (50 mg/kg)	Reynolds et al., 2003
killifish (<i>Fundulus grandis</i>)	freshwater	liver	injection	5 mg BaP/kg	2, 4, 8, 14 d	<0.1 adducts/10-8 nucleotides	>15 (all time-points)	(Willett et al. 1995)
killifish (<i>Fundulus grandis</i>)	freshwater	liver	injection	1, 5, 15, 50 mg BaP/kg	4 d	<0.1 adducts/10-8 nucleotides	>50 (15, 15 mg/kg)	(Willett et al. 1995)
mummichog (<i>Fundulus heteroclitus</i>)	freshwater	pooled blood	BaP/injection	15 mg/kg BaP	8, 16, 32, 96 d	<2 nmol/mol DNA	>20 (32 d)	(Rose et al. 2001)
mummichog (<i>Fundulus heteroclitus</i>)	freshwater	anterior kidney	BaP/injection	14 mg/kg BaP	8, 16, 32, 96 d	6 nmol/mol DNA	>10 (32 d)	(Rose et al. 2001)
mummichog (<i>Fundulus heteroclitus</i>)	freshwater	spleen	BaP/injection	13 mg/kg BaP	8, 16, 32, 96 d	10 nmol/mol DNA	4 (32 d)	(Rose et al. 2001)
mummichog (<i>Fundulus heteroclitus</i>)	freshwater	liver	BaP/injection	12 mg/kg BaP	8, 16, 32, 96 d	11 nmol/mol DNA	10 (32 d)	(Rose et al. 2001)
pike (<i>Esox lucius</i>)	freshwater	liver	BaP, BkF, DMC/diet	0.180 mmol/kg PAH-mix, single	1, 3, 6, 9 d	<0.5-2.8 nmol/mol DNA	>150 (6 d)	(Ericson & Balk 2000)

species	ecosystem	tissue	substance(s)	lowest-highest concs	exposure time	baseline/control (level/activity)	induction (fold)	reference
pike (Esox lucius)	freshwater	intestine	BaP, BkF, DMC/diet	0.1 mmol/kg PAH-mix x 5 feed	21, 59, 66, 83, 128 d	<0.5 nmol/mol DNA	>100 after 59 d	(Ericson & Balk 2000)
pike (Esox lucius)	freshwater	intestine	BaP, BkF, DMC/diet	0.180 mmol/kg PAH-mix, single feed	1, 3, 6, 9 d	<0.5-1.5 nmol/mol DNA	>150 (6 d)	(Ericson & Balk 2000)
pike (Esox lucius)	freshwater	liver	BaP, BkF, DMC, injection	0.120 mmol/kg PAH-mix, single injection	1, 3, 6, 9, 12 d	<0.5-2.8 nmol/mol DNA	>15 (9 d)	(Ericson & Balk 2000)
pike (Esox lucius)	freshwater	liver	BaP, BkF, DMC/diet	37.5, 150, 300, 600 µmol/kg PAH-mix	9 d	<0.5-2.8 nmol/mol DNA	>100 (100, 200 µmol/kg)	(Ericson & Balk 2000)
pike (Esox lucius)	freshwater	intestine	BaP, BkF, DMC/diet	37.5, 150, 300, 600 µmol/kg PAH-mix	9 d	<0.5-1.5 nmol/mol DNA	>100 (100 µmol/kg)	(Ericson & Balk 2000)
pike (Esox lucius)	freshwater	brain	BaP, BkF, DMC/diet	0.1 mmol/kg PAH-mix x 5	21, 59, 66, 83, 128 d	<0.5 nmol/mol DNA	>100 after 59 d	(Ericson & Balk 2000)
pike (Esox lucius)	freshwater	liver	BaP, BkF, DMC/diet	0.1 mmol/kg PAH-mix x 5	21, 59, 66, 83, 128 d	<0.5 nmol/mol DNA	>100 after 59 d	(Ericson & Balk 2000)
pike (Esox lucius)	freshwater	intestine	BaP, BkF, DMC, injection	0.120 mmol/kg PAH-mix, single injection	1, 3, 6, 9, 12 d	<0.5-1.5 nmol/mol DNA	>5 (9, 12 d)	(Ericson & Balk 2000)
pike (Esox lucius)	freshwater	gill	BaP, BkF, DMC/diet	0.1 mmol/kg PAH-mix x 5	21, 59, 66, 83, 128 d	<0.5 nmol/mol DNA	>100 after 59 d	(Ericson & Balk 2000)
polar cod (Boreogadus saida)	marine	liver	dispersed oil	1 mg/L dispersed oil	2 weeks	1.2 nmol/mol DNA	10	(Aas et al. 2003)
rainbow trout (Oncorhynchus mykiss)	freshwater	liver	effluent pulp mill	2% dilution	50 d	<0.5 nmol/mol DNA	>10	(Lindesjoo et al. 2002)
zebrafish (Danio rerio)	freshwater	liver	BaP/water	0.2 mg/kg - 3 d + 1 mg/kg - 4 d	7 d	n/a	4 adducts/10-8 nucleotides	(Hsu & Deng 1996)
zebrafish (Danio rerio)	freshwater	testis	BaP/water	0.2 mg/kg - 3 d + 1 mg/kg - 4 d	7 d	n/a	0.2 adducts/10-8 nucleotides	(Hsu & Deng 1996)
zebrafish (Danio rerio)	freshwater	brain	BaP/water	0.2 mg/kg - 3 d + 1 mg/kg - 4 d	7 d	n/a	4 adducts/10-8 nucleotides	(Hsu & Deng 1996)

<i>species</i>	<i>ecosystem</i>	<i>tissue</i>	<i>substance(s)</i>	<i>lowest-highest concs</i>	<i>exposure time</i>	<i>baseline/control (level/activity)</i>	<i>induction (fold)</i>	<i>reference</i>
zebrafish (<i>Danio rerio</i>)	freshwater	intestine	BaP/water	0.2 mg/kg - 3 d + 1 mg/kg - 4 d	7 d	n/a	13 adducts/10 ⁸ nucleotides	(Hsu & Deng 1996)

Table 11. Overview of field studies in which DNA adducts were analysed using nuclease P1 enrichment (mussels).

species	tissue	substance(s)	lowest-highest concs	exposure time	baseline/control (level/activity)	induction (fold)	reference
<i>Mytilus galloprovincialis</i>	gill	field	Venice	field	detection limit	(0.400 adducts/10-8 nucleotides)	(Venier & Zampieron 2005)
<i>Mytilus galloprovincialis</i>	gill	field	harbour	field	0.5-1	5.0-10	(Pisoni et al. 2004)
<i>Mytilus galloprovincialis</i>	hepatopancreas	field	harbour	field	<0.1	>50	(Pisoni et al. 2004)
<i>Mytilus edulis</i>	gill	field	harbours	field	0.1-0.5 nmol/mol DNA	2-30 (depending on season)	(Skarphedinsdottir et al. 2005)
<i>Mytilus edulis</i>	gill	field	harbours	field		>20 (seasonally higher)	(Ericson et al. 2002)
<i>Mytilus edulis</i>	hepatopancreas	field	harbours	field		2-3 (less in August)	(Ericson et al. 2002)
<i>Mytilus edulis</i>	gill	caged	harbours	6 wk		2 (March, but not August)	(Ericson et al. 2002)
<i>Mytilus edulis</i>	hepatopancreas	caged	harbours	6 wk		2 (March, but not August)	(Ericson et al. 2002)

Table 12. Overview of laboratory studies in which DNA adducts were analysed using nuclease P1 enrichment (mussels).

species	tissue	substance(s)	lowest-highest concs	exposure time	baseline/control (level/activity)	induction (fold)	LOEC	reference
<i>Mytilus edulis</i>	gill	BaP/diet	50, 100 mg/kg DW	3, 10, 21 + 10 d depuration	n/a	>30 (both concentrations; 21 d)		(Akcha et al. 2000)
<i>Mytilus edulis</i>	gill	BaP/water	5, 50, 100 µg/L	4 d	1.6 nmol/mol DNA	approx 3	5 µg/L	(Skarphedinsdottir et al. 2003)
<i>Mytilus edulis</i>	hepatopancreas	BaP/water	5, 50, 100 µg/L	4 d	1.4 nmol/mol DNA	none	none	(Skarphedinsdottir et al. 2003)
<i>Mytilus edulis</i>	gill	BaP/water	17 µg/L	6 d + 28 d depuration	0.5 nmol/mol DNA	approx 7 (6 d), none (34 d)		(Skarphedinsdottir et al. 2003)
<i>Perna viridis</i>	hepatopancreas	BaP/water	0.3, 3, 30 µg/L	1, 3, 6, 12, 18, 24 d	n/a	approx 6, 5, 2 (30, 3, 0.3 µg/L) 3 d, none (18, 24 d)		(Ching et al. 2001)
<i>Mytilus galloprovincialis</i>	gill	BaP/water	0.5-1000 µg/L	2, 3 d	0.07-0.12 adducts/10-8 nucleotides	approx 5 (2, 3 d)	50 µg/L (2 d), 5 µg/L (3 d)	(Canova et al. 1998)
<i>Mytilus galloprovincialis</i>	hepatopancreas	BaP/water	0.5-1000 µg/L	2, 3 d	0.05-0.06 adducts/10-8 nucleotides	approx 5 (2 d), approx 10 (3 d)	50 µg/L (2 d), 5 µg/L (3 d)	(Canova et al. 1998)

6. The Comet assay

6.1 Introduction

The alkaline single cell gel electrophoresis or the Comet assay is a technique for measuring DNA strand breaks and thereby DNA damage. The assay involves detection, under alkaline conditions, of cell DNA fragments which, on electrophoresis, migrate from the nuclear core, resulting in a “Comet with tail” formation Singh *et al.* (1988). The Comet assay was developed by Singh *et al.* for use on human cells but has later been applied on many species of fish and aquatic invertebrates. In contrast to other similar alkaline treatment assays the Comet assay incorporates the microscopic examination of damage to individual cell nuclei. The Comet assay works because DNA strand breaks create fragments or supercoiled DNA-loops that, when embedded in an agarose gel, migrate in an electric field. The strand breaks damage the higher order, tightly-packed structure of DNA, which also allows migration outside the region of the cell nucleus. Alkaline treatment facilitates the unwinding and denaturation of the DNA molecules, allowing for the sensitive detection of single-strand damage. In human DNA, it has been estimated that the sensitivity of the Comet to detect strand breaks is in the area of (one break in 10^{10} daltons). The Comet assay principles are illustrated in Figure 4.

The Comet assay is a sensitive, rapid and cost-efficient technique for the detection of DNA strand breaks, which is ideally suited as a non-specific biomarker of genotoxicity in fish and other aquatic species (Mitchelmore & Chipman, 1998). The Comet assay offers advantage over cytogenetic assays since there is no requirement for the detailed knowledge of the karyotype, thus enabling it to be easily transferred between species. The Comet assay also has the added advantage of being applicable to non-dividing cells (Dixon & Wilson, 2000). The results of Lee *et al.* (2004) indicate that there is an association between reproduction abnormalities and increased DNA strand breaks as a result of grass shrimp exposure to estuarine sediments receiving highway runoff (PAH). In our experiments we have observed that at the level where significant increase in the level of strand breaks was detected in adult mussels, reduced fitness was observed in the larvae (Baussant, 2004).




Figure 4. The principles of the Comet assay as a biomarker tool. Comets with no damage, moderate damage and severe damage are illustrated.

The level of DNA strand breaks is usually presented as the percentage of DNA migrated out of the nucleus and into the Comet tail region (% DNA in tail), but tail moments and tail lengths are also used. Figure 5 shows an example of results from thanalysis of Comet assay in haemocytes of mussels exposed to styrene (Mamaca *et al.*, 2005). The results show that significant differences between control and exposed animals can be detected both at the individual and the cell level. Only 5% of cells from control mussels had more than 12% DNA in the Comet tail, compared to 50% of the cells from styrene exposed mussels.

When mussels are exposed to low concentrations of oil and other compounds there is more overlap in response between control and exposed animals. Generally the level of DNA strand breaks in exposed cells is less than twice as high as in the corresponding control (Table 14 and Table 15). For details on methods see Aas *et al.* (2002), Taban *et al.* (2004), and Bechmann (2004) and Baussant (2004).




Figure 5. Example of results from the Comet assay analysis. Mussels (*Mytilus edulis*) exposed to styrene (data from Mamaca *et al.*, 2005). **A:** Frequency distribution of % DNA in the Comet tail of cells from control mussels and mussels exposed to styrene (9 mussels x 50 cells from each treatment). Cells from all the replicate mussels from each treatment are combined. **B:** Comparison of mean % DNA in the Comet tail of cells from each of the 9 mussels analysed from each treatment.

Table 13 gives an overview of studies where the Comet assay has been used for biomonitoring in the field. Mussels and different fish species have been used, and in several studies significant differences between contaminated and clean sites were detected. It seems like the Comet assay works best if caged animals are used, or if the objective is to study the effects of a spill. In some cases it was not possible to detect any increase in the level of DNA strand breaks in animals living in an environment chronically exposed to contaminants. Since laboratory results also indicate that long time exposure may reduce the response detected in the Comet assay, caging may be the best choice for monitoring off shore discharges from the oil industry (and for mussels that is also the only possibility).

Table 13. Overview of results from biomonitoring studies using the Comet on different species of aquatic invertebrates and fish collected in the field. In the “result” column symbols are used to indicate whether differences in DNA damage between contaminated and clean sites were observed: + : Significant differences in DNA damage between clean and contaminated sites, ÷ : No difference.

Species	Type of contamination	Result	Reference
Mussel (<i>Mytilus edulis</i>)	Genotoxic waste water	÷	Rank (1999)
Mussel (<i>Mytilus edulis</i>)	PAH polluted site (Loch Leven, Scotland)	÷	Large <i>et al.</i> (2002)
Mussel (<i>Mytilus edulis</i>)	Urban /industrial contaminated site (PCB, PAH)	Caged mussels: + Indigenous mussels : ÷	Shaw <i>et al.</i> (2002)
Mussel (<i>Mytilus edulis</i>)	Naval Station San Diego, PAH, Caging experiment	+	Steinert <i>et al.</i> (1998b)
Mussel (<i>Mytilus edulis</i>)	Caging experiment San Diego Bay, PAH	+	Steinert <i>et al.</i> (1998a)
Mussel (<i>Mytilus edulis</i>)	Reykjavik harbour PAH	+	Halldorsson <i>et al.</i> (2004)
<i>Mytilus sp.</i>	The French coast	÷	Akcha <i>et al.</i> (2004)
Zebra mussel (<i>Dreissena polymorpha</i>)	Caging experiment Municipal wastewater	+	Pavlica <i>et al.</i> (2001)
<i>Mytilus galloprovincialis</i>	Highly eutrophicated coastal lagoon	+	(Frenzilli <i>et al.</i> , 2001)
The intertidal butterfish (<i>Pholis gunnellus</i>)	Industrial contamination	÷	Bombail <i>et al.</i> (2001)
Chub (<i>Leuciscus cephalus</i>)	River Rhone	+	Devaux <i>et al.</i> (1998)
Bullheads (<i>Ameiurus nebulosus</i>) Carp (<i>Cyprinus carpio</i>)	PAH, PCB	+	Pandurangi <i>et al.</i> (1995)
Smallmouth bass (<i>Micropterus dolomieu</i>)	Androscoggin River	+	(Chamberland <i>et al.</i> , 2002)
Chub (<i>Leuciscus cephalus</i>)	Moselle River (France)	÷	(Flammarion <i>et al.</i> , 2002)
California halibut (<i>Paralichthys californicus</i>)	Southern California bays and harbours	+	Brown <i>et al.</i> (2004)
Sacramento sucker (<i>Catostomus occidentalis</i>)	Agricultural runoff Field-caging experiments + lab	+	Whitehead <i>et al.</i> (2004)
Eelpout (<i>Zoarces viviparus</i>)	River Gota estuary Bunker oil spill	+	Frenzilli <i>et al.</i> (2004)

A large number of laboratory *in vitro* and *in vivo* studies have been done using different types of cells from fish and invertebrates. There are, however, few studies that are directly relevant for the oil industry. The two most relevant references are Frenzilli *et al.* (2004) and (Hamoutene *et al.*, 2002). A recent study by Frenzilli *et al.* (2004) showed that the Comet assay can be used to study effects of oil spills on fish. Following a bunker oil (10-100 tonnes) spill Frenzilli *et al.* (2004) analysed the level of DNA strand breaks in nucleated erythrocytes of the eelpout (*Zoarces viviparus*) from the Gothenburg harbour area and a clean reference site. The results showed a high level of damaged DNA, paralleled by a peak in bile PAH metabolites, in fish from the most impacted site, 3 weeks after the oil spill. A significant recovery was observed in specimens from the spill site, 5 months later, but not in fish caught in the middle part of Gothenburg harbour, which is chronically subjected to harbour related pollution Frenzilli *et al.* (2004).

Hamoutene *et al.* exposed mussels (*Mytilus edulis*) and clams (*Mya arenaria*) to dispersions and water soluble fractions of an Arabian crude oil. Exposure to hydrocarbon fractions had no significant impact on clams. However, an increase in DNA damage was observed at $p < 0.1$ with digestive gland cells of mussels exposed to

aqueous fractions of a light crude oil (but no effect on haemocytes) (Hamoutene *et al.*, 2002).

6.2 The Comet assay with invertebrates

The Comet assay has proved to be a sensitive biomarker for detecting DNA strand breaks in bivalves and sea urchins exposed to oil in the laboratory. Increased levels of DNA strand breaks have been observed with increasing exposure concentration of oil (3 – 1000 $\mu\text{g/L}$ oil). The significant results obtained with mussels, sea urchins and Iceland scallops exposed to relatively low concentrations of oil in the lab indicate that the comet assay may be used in biomonitoring of produced water discharges and oil spills.

The mean level of DNA strand breaks in haemocytes of control mussels and coelomocytes of control sea urchins (measured as percent DNA in comet tail) has been compared to the level of DNA strand breaks in mussels and sea urchins exposed to oil (Figure 6). All groups of oil exposed mussels and sea urchins have been grouped together to get an indication of whether comet assay has potential to be used for monitoring produced water discharges/oil spills in the field. The tested oil concentrations were in the range 3 – 1000 $\mu\text{g/L}$ oil (see table 2 and 3 for details). There was a significantly higher level of DNA strand breaks in haemocytes of oil exposed mussels (14.3, $n = 8$) and coelomocytes of oil exposed sea urchins (16.8, $n = 10$) than in control cells from both species (mussels; 8.7, $n = 17$, sea urchins; 11.3, $n = 11$) (Figure 6).

It would be considerably more practical to be able to analyse frozen samples if the Comet assay should be used in off shore monitoring surveys. Scott Steinert (CSC Biomarker Laboratory, San Diego, USA) and Farida Akcha (IFREMER, France) have procedures for freezing cells in a way that apparently does not affect the results from the Comet assay (*personal communication*). This procedure needs to be established and evaluated in oil exposure experiments to determine whether freezing will reduce the sensitivity of the Comet assay in such exposures.




Figure 6. Laboratory controls compared to all responses in all groups of oil exposed mussels – oil concentrations in the range 3 – 1000 $\mu\text{g/L}$, including recovery after high exposure – see table x). Field values are not included. The level of DNA strand breaks in control and oil exposed animals was compared with each pair Student t-test. Groups with different letters are significantly different ($p < 0.05$).

6.3 The Comet assay with fish red blood cells

The study by Frenzilli *et al.* (2004) showed that the Comet assay can be used to study effects of oil spills on fish. The Comet assay has been used on many species of fish, but we have not been able to find any papers where comet assay has been used on blood cells from cod. Our experience from doing comet assay on blood cells from cod is that the controls are more variable than blood cells from corkwing wrasse and invertebrates (bivalves and sea urchins). There was, however, significantly increased level of DNA strand breaks in blood cells from cod exposed to 1000 $\mu\text{g/l}$ oil (Taban *et al.* 2000). The procedure for doing comet assay on blood cells from cod needs further optimization. However, since two other genotoxicity tests (DNA adducts and micronuclei assay) are able to detect DNA damage in oil exposed cod, it is possible that these tests are better biomarkers for biomonitoring of cod than comet assay. It may be more important to invest further effort in establishing the procedure for testing frozen cells from invertebrates using the comet assay. For the invertebrates comet assay seem to be a sensitive tool to detect exposure to genotoxic compounds also in the field.

6.4 Confounding factors

Akcha *et al.* (2004b) have shown that biotic (age, sex) and abiotic (sampling site and period) factors affect the extent of DNA damage measured by the Comet assay in dab (*Limanda limanda*) collected in the field. These results should also be taken into consideration when using the Comet assay on invertebrates.

After long exposures the responses measured by the Comet assay may be reduced. The results from Shaw *et al.* (2002) and Large *et al.* (2002) (and our results; Baussant, 2004) indicate that mussels may be able to adapt to the exposure. Possibly due to adaptation/induced repair mechanisms. Hence if no response is detected in wild organisms this does not necessarily mean that they have not been exposed.

6.5 Links

The Comet assay indicates damage to DNA. There are some data that indicate effects on reproduction at similar concentrations as those that cause increased DNA fragmentation as measured using the Comet assay (Bechmann & Sanni, unpublished), but no direct evidence as yet.

6.6 Background responses

The mean level of DNA strand breaks in cells from control animals has been compared (Figure 7). The mean control level of DNA strand breaks (measured as percent DNA in Comet tail) for mussel haemocytes was 8.7 (n = 17), for scallop haemocytes 10.5 (n = 5), for sea urchin coelomocytes 11.3 (n = 11) and for mussel sperm 13.8 (n = 4). The results show that the variability in the control level of DNA strand breaks is low. To detect the small changes between control and exposed animals at low exposure concentrations, controls are run together with each exposed groups to ensure that all samples have been treated in exactly the same way.




Figure 7. Mean percent DNA in comet tail of the control groups in laboratory exposures performed at RF-Akvamiljø 2001-2005. The control level for the different cell types was compared with Each pair Student t-test. Groups with different letters are significantly different ($p < 0.05$).

6.7 Assessment criteria

There are no established assessment criteria for Comet used for fish or invertebrates.

6.8 Quality assurance

There has been no international intercalibration for the Comet assay as used in ecotoxicology.

Table 14. Selected results from comet assay analysis performed on mussel (*Mytilus edulis*) haemocytes.

Substance (lab/field)	Test concentrations	Exposure time	Response (+/-)	Baseline/control (% DNA in Comet tail)	Exposed/ Control	Reference
North Sea Oil (lab)	1000 µg/L + recovery	2 + 4 weeks	+	8.9 – 8.5	1.6 – 1.3	Aas et al. (2002) Baussant (2004)
	3, 15, 63 µg/L	1 month	+	7.7 – 7.3 – 7.9	1.4 – 1.9 – 1.7	
		7 months	+	12.6 – 10.3 – 6.5	1.3 – 2.6 – 1.3	
Styrene (lab)	200 µg/L	10 days	+	9.5	1.8	Mameca et al. (2005)
Bisphenol A (lab)	59 µg/L	3 weeks	+	9.2	1.3	Taban et al. (2003)
Diallyl phthalate (lab)	38 µg/L	3 weeks	-	8.8	0.9	
PBDE-47 (lab)	0.23 µg/L	3 weeks	-	9.3	1.0	Børseth et al. (2004)
Oil spill (field)	Rocknes (Ramsvik, Songathella)	1 + 7 months	(+?)	15.3/15.6 – 6.3/6.5	1)	
Used water based drilling mud (lab)	0.5, 2, 20 mg/L	3 weeks	+ (0.5, 2 mg/L)	7.6 – 7.2 – 7.8	1.2 – 1.2 – 1.1	Unpublished results from the PROOF drilling mud project (159183)
Barite particles (lab)	23 mg/L	3 weeks	+	10.5	1.1	
Metal mix (Pb+Cd+Cu+Zn) (lab)	Sum metals: 62 µg/L	3 weeks	-	8.0	1.0	Berland & Samni (2005)
Drilling mud (field; Sleipner)	Reference 20 m, exposed 20 m + 40 m	5 weeks	+ (40 m)	7.7 – 13.5	1.1 – 1.4	

¹⁾ 1 month after the accident both stations had higher DNA damage than average for laboratory controls, but after 7 months the mean level of DNA damage was lower than in average laboratory controls. This may indicate that mussels at both stations were affected after 1 month, and that repair was going on after 7 months. The lack of a good field reference stations makes it difficult to conclude.

Table 15. Overview of results from comet assay analysis performed on mussel (*Mytilus edulis*) embryo, mussel sperm, scallop (*Pecten maximus*, *Chlamys islandica*) haemocytes, sea urchin (*Strongylocentrotus droebachiensis*) coelomocytes and sperm and corkwing wrasse blood cells.

Species Tissue	Substance (lab/field)	Test concentrations	Exposure time	Response (+/-) p<0.05: +	Baseline/control (% DNA in Comet tail)	Exposed/ Control	Reference
<i>Mytilus edulis</i> Mussel embryo	North Sea Oil (lab)	63 µg/L (parental + embryo exposure)	7 months (parents) + 1 day (embryo)	+	8.4	1.8	Baussant (2004)
	North Sea Oil (lab)	1000 µg/L	2 weeks	+	14.3	2.5	Aas et al. (2002)
<i>M. edulis</i> Sperm	Bisphenol A (lab)	59 µg/L	4 weeks	+	12.6	1.3	Taban et al. (2003)
	Diallyl phthalate (lab)	38 µg/L	4 weeks	+	10.8	1.5	
	PBDE-47 (lab)	0.23 µg/L	4 weeks	+	17.4	1.4	
Scallop (<i>Pecten maximus</i>) Haemocytes	Used water based drilling mud (lab)	0.5, 2, 20 mg/L		+(2, 20 mg/L)	11.3 – 10.2 – 9.1	0.9 – 1.1 – 1.2	Unpublished results from the PROOF drilling mud project (159183)
	Barite particles (lab)	23 mg/L		+	9.7	1.3	
	Metal mix (Pb+Cd+Cu+Zn) (lab)	Sum metals: 62 µg/L	3 weeks	-	12.2	1.0	
Iceland scallop (<i>Chlamys islandica</i>) Haemocytes	Goliat oil (lab)	2, 14, 64 µg/L	4 weeks	+(14, 64 µg/L)	(15.8), 9.3, 10.0	(0.9), 1.1, 1.9	Baussant (2004)
	Sea urchin (<i>Strongylocentrotus droebachiensis</i>) Coelomocytes	1000 µg/L		2 weeks	+	12.9	1.8
Recovery after 1000 µg/L			3 weeks recovery	+	11.5	1.5	
29, 85 µg/L			1 months	+	11.7 – 9.3	1.4 – 2.0	
4, 29, 85 µg/L			4 months	+	11.8	1.4 – 1.2 – 1.2	
4, 29, 85 µg/L			7 months	+	10.9 – 12.7 – 13.4	1.3 - 1.2 - 1.3	
<i>S. droebachiensis</i> Sperm	North Sea Oil (lab)	1000 µg/L	2 weeks	+	9.2	1.9	Aas et al. (2002)
Corkwing wrasse (<i>Symphodus melops</i>) Blood cells	Styrene (lab)	200 µg/L	10 days	+	7.0	1.6	Mamaca et al. (2005)
	Metals (field)	Visnes	-	+	13.5	1.6	Bjømstad et al. (2002)

7. Micronucleus formation

7.1 Introduction

Markers of genotoxic effects are at high priority due to reflection of damage to genetic material of organisms (Moore et al. 2004). Different methods has been developed for the detection of both double- and single-strand breaks of DNA, DNA-adducts, micronuclei formation, chromosome aberrations. The assessment of chemical induced genetic damage has been widely utilized for prediction of the genotoxic, mutagenic and carcinogenic potencies of a range of substances, however these investigations have mainly been restricted to human or mammals (Siu et al. 2004).

With a growing concern over the presence of genotoxins in the aquatic media, there is a rising need to elaborate sensitive methods for the assessment of genetic damage in indigenous marine or freshwater organisms. The application of cytogenetic assays on ecologically relevant species offers the chance to perform early tests on health in relation to exposure to contaminants. During the last decade an increasing attention was paid to the micronucleus test, which is simple to perform, sensitive enough and a fast test to detect genomic alterations due to clastogenic effects and impairments of mitotic spindle caused by aneuploidogenic poisons (Heddle et al. 1983; 1991; MacGregor 1991; Seelbach et al. 1993; Kramer 1998; Zoll-Moreux and Ferrier 1999).

The marine environment becomes a sink for various mutagenic/genotoxic compounds in complex environmental mixtures that can have adverse health effects on indigenous biota (Hose and Brown 1998). Three methodologically different approaches has been used for the genotoxicity assessment in marine environment: analysis of micronuclei (MN) in indigenous organisms *in situ*, assessment of MN in caged organisms in the gradient of contamination and validation of MN assay in laboratory experimental exposure.

7.2 Micronucleus analyses *in situ*

Some studies have shown that mutagenic PAHs can be responsible for between 10-20%, or sometimes up to 70-80% of sediment mutagenic activity in marine media (LaRocca et al. 1996; White et al. 1998). There are studies that describe increased environmental genotoxicity in zones affected by an oil spill (Parry et al. 1997; Harvey et al. 1999; Pietrapiana et al. 2002; Barsiene et al. 2004). Significant elevation of micronuclei level in mussels 30 days post-oil spill and persistence of the cytogenetic damage up to 100 days later has been described (Lyons et al. 1997). Elevated levels of the micronuclei were detected in flounder and blue mussels 8 months after oil spill from Būtingė oil terminal in the Baltic Sea (Barsiene et al. 2004; 2005b; 2005c). More frequent cytogenetic damage has been described in molluscs inhabiting the marine port and oil terminal areas in the Baltic Sea (Barsiene and Lovejoy 2000; Barsiene 2002).

In marine and estuarine ecosystems, most often environmental genotoxicity investigations have been performed using bivalves from genus *Mytilus* (Scarpato et al. 1990; Brunetti et al. 1992; Bolognesi et al. 1996; Burgeot et al. 1996; Barsiene 2002; Dailianis et al. 2003;

Izquierdo et al. 2003; 2004; Bolognesi et al. 2004; Kalpaxis et al. 2004; Hagger et al. 2005; Baršienė et al. 2005c; 2005d). In most of the studies the increased levels of MN have been shown in polluted areas. Considering that MN incidences in molluscs reflects the action of clastogenic and aneuploidogenic substances, this parameter was suggested to use as marker of cytogenetic damage in marine monitoring programs (Bolognesi et al. 1996; Izquierdo et al. 2003; Barsiene et al. 2004; Bolognesi et al. 2004; Kalpaxis et al. 2004; Baršienė et al. 2005c; 2005d).

The multiple application of micronuclei analysis in various fish and mollusk species inhabiting different sites of the North Sea, Baltic Sea and areas of the North-Atlantic showed that the technique is highly sensitive tool to monitor the cumulative effects caused by genotoxic compounds distributed in the area of exposure *in situ* (Barsiene and Lovejoy 2000; Barsiene 2002; 2004; 2005a; 2005b)(refer also to Baršienė et al., unpublished data; Schiedek et al., in press; Kopecka et al., in press).

7.3 The micronucleus test as laboratory bioassay

The approach to measure genotoxic effects in the aquatic environment was mainly elaborated in freshwater fish and marine mollusk species. Nevertheless in some studies it served as a promising method in marine fish treated with crude oil, produced water from oil platforms, different alkylphenols, PAHs, endocrine disruptors (Gravato and Santos 2002; Maria et al. 2002b; 2003; Baršienė et al. 2005b; 2006c). The treatment of Atlantic cod to different concentrations (0.06, 0.25, 0.5 and 1 ppm) of crude oil from Statfjord B platform showed an elevated level of MN in liver after 3-day treatment and the highest micronuclei induction after 14-day exposure (Baršienė et al. unpublished data - *Biosea programme*). In terms of methodology of crude oil genotoxicity assessment, very important finding of our experimental work was that the efficient recovery occurs after 2 weeks. In overall it means, that genotoxic effects of hazardous crude oil compounds arise early after agent appears in the fish environment and comparatively fast elimination of cytogenetic damage when genotoxic agents are absent in the habitat of the fish. Stepwise Regression analysis of influence of fish weight, liver weight, oil concentration and exposure time on MN formation showed, that the concentration of oil is the most important factor in treatment and recovery groups of whole cod cohort studied (Baršienė et al. unpublished data - *Biosea programme*). Furthermore, crude oil from different platforms, as well as temperature of exposure, has potency to modify the pattern of MN induction in fish (Baršienė et al., unpublished data - *Biosea programme*).

In mussels, cells with micronuclei were found to increase in the gills after treatment with zinc chloride (Majone et al. 1988), benzo[*a*]pyrene and vincristine (Burgeot et al. 1995), mitomycin, colchicine (Majone et al. 1987; 1990), and mercury chloride (Bolognesi et al. 1999). Elevated levels of micronucleated cells have been described in gills or hemolymph of marine molluscs treated with benzo[*a*]pyrene (Burgeot et al. 1995; Venier et al. 1997; Siu et al. 2004), dimethylbenzol[*a*]anthracene (Bolognesi et al. 1996), bisphenol A, diallyl phthalate and tetrabromodifenyl ether-47 (Baršienė et al. 2005b). The highest induction of micronuclei and nuclear buds was observed in mussels and perch after 10-day treatment with 0.5 ppm of crude oil (Baršienė et al. 2006a). Moreover the results of the MN assays showed clear dose- and time-dependent responses to benzo[*a*]pyrene exposure in the bivalve *Perna viridis*. It was

pointed, that the micronuclei test provides a convenient, highly sensitive, non-invasive monitoring tool of genotoxic assessment in the aquatic environment (Siu et al. 2004).

7.4 Assessment of MN in caged organisms

Some of the latest biomonitoring programmes involve environmental genotoxicity assessment in caged marine organisms. The micronucleus test was used in mussels *Mytilus galloprovincialis* caged in selected sites along Gulf of Patras (Kalpaxis et al. 2004) and along the Ligurian coast of the Mediterranean Sea (Bolognesi et al. 2004). The approach was approved as reliable tool for coastal biomonitoring. Micronuclei analysis was performed in cod and mussels caged in different zones around oil drilling platforms in the North Sea (Baršienė et al. unpublished data). There was a gradient of micronuclei induction in hemolymph of mussels caged in different distance from the Statfjord B oil platform. A 2-3-fold increase of MN was detected in mussels immersed at 500 and 1000 meters downstream the platform, when compared with mussel group before the caging experiment. Similar pattern of MN induction was found in cod liver cells, whilst the response was at a lower level than in mussels. Thus, the response to genotoxins showed a good discriminatory power in identification of pollution gradient and the approach was suspected as sensitive tool in biomonitoring for the evaluation of potential ecological risk with regard to marine pollution by crude oil or produced water effluents (Hylland et al. 2005). Comparing responses to genotoxic action in caged and indigenous mussels suggests that wild populations are able to accumulate higher content of contaminants in their tissues and express higher level of time-integrated response to genotoxic agents (Bolognesi et al. 2004). This finding confirms our previous studies on environmental mutagenicity using indigenous and caged molluscs (Baršienė & Bučinskienė, 2001).

Overall, the cytogenetic damage serves as a promising method to monitor environmental contamination with oil genotoxic substances because aquatic organisms activate oil and respond to genotoxic metabolites at low, environmentally realistic concentrations. Hazardous effects of different PAHs arise mainly as a result of oxidative biotransformation producing highly DNA-reactive metabolites. These metabolites are recognized as carcinogenic and mutagenic compounds (Torres-Bugarin et al. 1998; Woodhead et al. 1999). Mechanisms of PAH metabolic transformation have been studied and genotoxic potency of metabolites was confirmed in fish (Pacheco and Santos 1997; Pacheco and Santos 2001; Maria et al. 2002a; Maria et al. 2002b; Teles et al. 2003).

7.5 Target tissues

7.5.1 Fish

Tissue and species-specific response to crude oil was shown in Atlantic cod and turbot. Induction of micronuclei was significantly higher in turbot compared to Atlantic cod. The frequencies of micronuclei in cephalic kidney from all treated groups of turbot and cod were about 2-fold higher than in mature erythrocytes from peripheral blood (Baršienė et al. 2006c). In teleost fish, the main hemopoietic tissue is a cephalic kidney, but fish peripheral blood has been traditionally used as an endpoint in genotoxicity assessment. Since MN can arise after cell division, one disadvantage of the MN test in peripheral blood is the comparatively low response to genotoxic agents. It is not yet known, what proportion of injured erythroblasts

could appear in the peripheral blood circulating system, what extent of DNA repair there is and how long a complete red blood cell turnover in fish would be (Buschini et al. 2004). Treatment with Cytocholasin B, cytokinesis blocker, demonstrated, that in *Tilapia rendalli*, *Oreochromis niloticus* and *Cyprinus carpio* immature kidney erythrocyte cells start to circulate in peripheral blood after 2-4 days. Nevertheless, the rate of cell proliferation varied between fish species (Grisolia and Cordeiro 2000).

Liver was used as a tissue directly affected by oil metabolic components, which have been described as mutagenic agents in fish (Stegeman et al. 1993). Hepatic micronuclei approach has been considered a sensitive tool for the evaluation of genotoxic effects in contaminated sites (Williams and Metcalfe 1992; Rao et al. 1997; Arcand-Hoy and Metcalfe 2000; Pietrapiana et al. 2002). In Atlantic cod, hepatic micronucleus assay was shown being a more sensitive system for the assessment of oil genotoxicity than in peripheral blood erythrocytes (Baršienė et al. 2005a). Cristaldi et al., (2004) demonstrated selective removal of micronucleated erythrocytes from peripheral blood in spleen. Thus, micronuclei counting in peripheral blood can give the wrong information about fish exposure to genotoxic compounds. On the other hand, the comparison of genotoxicity levels in fish liver and peripheral blood showed that cytogenetic changes in peripheral blood reflect recent exposure as a result of the more rapid turnover of blood cells (Tell-Krakoc et al., 2001).

In 5 of the 9 studies where (freshwater and marine) fish were sampled at clean and contaminated sites, the authors conclude that there were differences in MN (Table 16; Tripathy and Das 1995; Minissi et al. 1996; Rao et al. 1997; Arcand-Hoy and Metcalfe 2000; Pietrapiana et al. 2002). A relevant caging experiment with fish has also been done. Tuvikene et al. did not observe a higher frequency of MN in rainbow trout caged at a PAH contaminated site, although the PAH concentration in the fish was high (Tuvikene et al. 1999).

Exposure of Atlantic cod to crude oil resulted in significantly increased frequency of micronucleated erythrocytes and other nuclear abnormalities in head kidney (Aas et al. 2002).

7.5.2 Mussels

In the same specimens of mussels, a much higher MN induction (up to 8.85 MN/1000 cells) has been observed in gills compared to haemocytes (Dolcetti and Venier 2002). Lower MN response in mussel haemocytes than in gill cells was also described by Dailianis et al. (2003). Dose-response relationships between contaminants and MN incidences was shown in gill cells of mussels experimentally treated to certain contaminants as well as pollution gradient-response pattern that was found in mussels from Wismar Bay and Lithuanian coast of the Baltic Sea and from southern Norwegian fjords (Barsiene et al. 2004; 2005b; 2005c; 2006b; Schiedek et al, in press).

Three field studies with invertebrates (mussels and snails) have been found where the MN test has been successfully used to separate contaminated and clean sites (Bolognesi et al. 1996; Barsiene and Lovejoy 2000; 2002). Positive results have also been observed in laboratory experiments (Wrisberg and Rhemrev 1992; Bolognesi et al. 1996), and in a mesocosm experiment (Wrisberg and Rhemrev 1992), but the results of laboratory and field studies by

Burgeot et al. indicate that the MN test are not yet applicable for biomonitoring of genotoxic exposure of molluscs (Burgeot et al. 1995; 1996).

7.5.3 Other indicator species

The nereid marine polychaete *Platynereis dumerilii*, marine tubeworm *Pomatoceros lamarckii* and sea urchins has been used for the evaluation of environmental genotoxicity. Marine worm *Platynereis dumerilii* served as indicator species in assessment of genotoxins in municipal sewage effluents (Jha et al. 1997), genotoxic, cytotoxic properties of TBT (Hagger et al. 2002), benzo[*a*]pyrene and other genotoxins (Jha et al. 1996). Colchicine-induced effects were analyzed in *Pomatoceros lamarckii* (Wilson et al. 2002). Hose studied both anaphase aberrations and MN in sea urchin (*Strongylocentrotus purpuratus*) embryos exposed to benzo[*a*]pyrene in the laboratory (Hose 1985). MN were present at 1 µg/L benzo[*a*]pyrene and higher (anaphase aberration was more sensitive) (Hose 1985). Saotome et al. have developed a micronucleus assay for use in sea urchin embryos (Saotome et al. 1999) (but Hose used the MN test on sea urchins 14 years earlier). The test chemicals (mitomycin C, vinblastine and 1-beta-D-arabinofuranosylcytosine) induced clear micronuclei dose-dependently in *Clypeaster japonicus* and *Hemicentrotus pulcherrimus* (Saotome et al. 1999). The MN test in sea urchins has been improved to consider osmolarity and salinity in laboratory test conditions (Saotome and Hayashi 2003).

Analysis of interspecies variation in fish, mussel and crab DNA damage caused by genotoxic compounds *in situ* showed species specific susceptibility to environmental genotoxins (Bihari and Fafandel 2004). Thus, the multi-species approach in impact and risk assessment should be linked to harmful effects of pollution in sentinel organisms to their ecological consequences. In the context of ecological genotoxicity or ecosystem health, genotoxicity biomarkers are also being used to link cytogenetic damage through to higher levels, i.e. pathology with reduced reproductive success of marine organisms inhabiting zones affected by incidental or operational oil spills from all activities related to transport or processing of petroleum products.

We have not been able to find any field studies where the MN test has been used on sea urchins, but the results from the *BioSea programme* on micronuclei in oil exposed cod and mussel indicate that the micronucleus assay can be useful for detecting genetic damage caused by oil (at least in the laboratory). The study performed by Pietrapiana et al. (2002) indicates that the micronucleus assay also can be used to detect chromosome damage in fish exposed to oil in the field.

Table 16. Overview of results from biomonitoring studies using cytogenetic assays (tests measuring effects at the chromosome level) on different species of freshwater and marine fish collected in the field; + : significant difference between exposed and control; ÷ : no significant difference between exposed and control; ? : results not clear.

Test	Species	Type of contamination	Result	Reference
Chromosome aberration test	Pacific herring <i>Clupea pallasii</i>)	Exxon Valdez oil spill	+	Hose and Brown 1998
			+	Hose et al. 1996
			+	Norcross et al. 1996
			but recovery + caging exp.	Kocan et al. 1996
	Winter flounder <i>Pseudopleuronectes americanus</i>)	The North Cape oil spill	+	Rovero et al. 1999
Micronucleus	<i>Barbus plebejus</i>) freshwater fish	Polluted river Tiber),	+	Minissi et al. 1996
	Freshwater fish	Paper mill effluents	+	Tripathy and Das 1995
	Bullheads <i>Ameiurus nebulosus</i>)	Hamilton Harbour	+	Rao et al. 1997
	Various fish	The Noril'sk mining and smelting complex	?	Zakhidov et al. 1996.
	Freshwater: <i>Carassius sp.</i> , <i>Zacco platypus</i> Marine: <i>Leiognathus nuchalis</i> , <i>Ditrema temmincki</i>	River, harbour	?	Hayashi et al. 1998
	Caged rainbow trout <i>Oncorhynchus mykiss</i>) and roach <i>Rutilus rutilus</i>)	The oil shale industry, PAH	÷	Tuvikene et al. 1999.
	Brown bullheads <i>Ameiurus nebulosus</i>) freshwater)	Genotoxic contaminants in the field	+/- seasonal differences	Arcand-Hoy and Metcalfe 2000
	<i>Tilapia rendalli</i> , <i>Oreochromis niloticus</i> and <i>Cyprinus carpio</i>	Sewage treatment plant discharges	÷	Grisolia and Starling 2001
	<i>Lepidorhombus boscii</i> , <i>Merluccius merluccius</i> and <i>Mullus barbatus</i>	The "Haven" oil spill	+ species differences	Pietrapiana et al. 2002
	Eel <i>Anguilla anguilla</i>)	Heavy metal pollution (river)	÷	Sanchez-Galan et al. 2001

Table 17. Overview of results from biomonitoring studies using micronucleus assessment on different species of aquatic invertebrates collected in the field; + : significant difference between exposed and control; ÷ : no significant difference between exposed and control; ? : results not clear.

Species	Type of contamination	Result	Reference
<i>Mytilus edulis</i>	produced water, Statfjord	+	Hylland et al., 2005 (WCM)
Mussel <i>M. galloprovincialis</i>	French Mediterranean coast	÷	Burgeot et al. 1996.
Oyster <i>C. gigas</i>	Cadmium and copper	÷	Burgeot et al. 1995
Mussel <i>M. edulis</i>	Contaminated sediment mesocosm	?	Wrisberg and Rhemrev 1992
5 species of freshwater bivalves and snails	Contaminated port area	+	Barsiene and Lovejoy 2000
Mussel <i>M. edulis</i>	Baltic Sea at Butinge oil terminal	+	Barsiene 2002
Mussel <i>M. galloprovincialis</i>	The La Spezia gulf, Ligurian sea	+	Bolognesi et al. 1996

7.6 Confounding factors

There is limited knowledge about the non-contaminant factors that may affect micronucleus formation.

7.7 Links

Micronucleus formation indicates chromosomal breaks, known to result in teratogenesis (effects on offspring) in mammals. There is however limited knowledge of relationships between micronucleus formation and effects on offspring in aquatic organisms.

7.8 Background responses

There is not sufficient data to establish the background response for micronucleus formation, but baseline levels appear to be low in most studies.

7.9 Assessment criteria

There are no established assessment criteria for micronucleus assessment. Results from the WCM 2003 and 2004 indicate that it should be feasible to establish such criteria, but this requires some more data.

7.10 Quality assurance

There has been no international intercalibration for micronucleus assessment as used in ecotoxicology.

8. Multixenobiotic response (MXR)

8.1 Introduction

For many organisms it is observed that tissue levels of certain contaminants are lower than levels observed in the environment. It has been suggested that this is mediated by the MXR phenomenon, which is similar to the multi drug resistance MDR phenomenon that was first observed in tumour cell lines resistant to anti-cancer drugs. The MXR mechanism acts as an energy dependent pump that translocates a wide variety of structurally and functionally diverse compounds. These tend to be moderately hydrophobic and planar natural products which are often substrates for or metabolites of detoxification enzymes such as cytochromes P450 (CYPs). The protein responsible for this transport function is the transmembrane P-glycoprotein (Pgp), which is found in specialized epithelial tissues such as gut, liver, kidney and blood vessels at the blood brain barrier. Although Pgps have been intensively studied in relation to cancer biology, less is known regarding multixenobiotic resistance in wild populations of organisms, and there is similarly little information on environmentally relevant Pgp substrates.

Immunoblot and immunohistochemical studies of Pgp expression can be performed in a variety of organisms. The most successful and widely used antibody is C219, which identifies all known Pgps. A list of aquatic organisms and the tissues in which C219 reactive proteins have been detected is provided in Bard (2000).

Two methods can be used to measure MXR transport activity: bioaccumulation assays and efflux assays. In the first case, the accumulation of a marked substrate (radioactive or fluorescent e.g. rhodamine dye), is measured with or without the presence of a competitive inhibitor in translucent living cells or in tissue homogenates. In efflux assays, the rate of efflux of a previously accumulated (marked) substrate into the medium is measured in the presence or absence of a competitive inhibitor. The efflux rate is directly proportional to Pgp transport activity. This assay is simple and inexpensive and animals are not injured by the procedure.

8.2 Dose-response

8.2.1 MXR induction

Most studies have reported induction of multixenobiotic resistance transport activity and elevated Pgp protein levels in

- field populations of pollutant exposed aquatic organisms
- transplantation of organisms to different polluted sites
- laboratory exposure experiments

Reviews of MXR activity and relevant studies has been provided by several authors recently (Bard 2000; Daughton and Ternes 1999; Epel 1998; Galgani 2000; Jha 2004; Livingstone *et al.* 2000; Minier *et al.* 1999; Moore *et al.* 2004; Pain and Parant 2003; van der Oost *et al.* 2003).

Induction of the MXR system has been reported after exposure to several pollutants, among them diesel oil, metal pollution, PAH, alkylphenols, pesticides PCB and polluted sediments (Table 18 and Table 19).

Table 18. Induction of MXR in bivalves.

Species and tissue	Substance	Concentration	Exposure time (days)	Response	Reference
Method: Measurement of protein expression					
<i>Mytilus edulis</i> Blood cells	Vincristine		19-29	1,5-2,5	Minier and Moore 1996
<i>Mytilus californianus</i> Gill tissue	Chlorthal/PCB	5 ppm	3	+	Eufemia and Epel 1998
<i>Mytilus californianus</i> Gill tissue	Chlorthal/PCP/DDE	1-5 ppm	1-3	+	Eufemia and Epel 2000
<i>Mytilus californianus</i> Gill tissue	CdCl ₂ NaAsO ₃	0,5 1-5 ppm	1-3	+	Eufemia and Epel 2000
<i>Crassostrea gigas</i> Gill tissue	Contaminated sites (12 sites)		30	1 site	Ringwood <i>et al.</i> 1999
Method: Reduction of accumulation					
<i>Mytilus edulis</i> Blood cells	Vincristine	5-10 µg/ml	19-29	35-48%	Minier and Moore 1996
<i>Mytilus galloprovincialis</i> Gill tissue	Contaminated site		3	62%	Kurelec 1995
<i>Mytilus galloprovincialis</i> Gill tissue	Rhodamine 123 Diesel 2 oil		4	45-50%	Smital and Kurelec 1998
<i>Mytilus californianus</i> Gill tissue	Chlorthal/PCB	5 ppm	3	>75%/ 50%	Eufemia and Epel 1998
<i>Mytilus californianus</i> Gill tissue	Chlorthal/PCP/DDE	1-5 ppm	1-3	>50%	Eufemia and Epel 2000
<i>Mytilus californianus</i> Gill tissue	CdCl ₂ NaAsO ₃	0,5 1-5 ppm	1-3	50% 40-80%	Eufemia and Epel 2000
<i>Corbicula fluminea</i> Gill tissue	Diesel-2 oil		3	43%	Kurelec <i>et al.</i> 1996
<i>Corbicula fluminea</i> Gill tissue	Sediments from the Rhine river		3	28%	Kurelec <i>et al.</i> 1996
Method: increase of efflux					
<i>Dreissena polymorpha</i> Whole organism	Contaminated site		21	×4,5	Parant and Pain 2001

Part of the table is based on Pain & Parant (2003).

Table 19. Induction of MXR in fish.

Species and tissue	Substance	Concentration	Exposure time (days)	Response	Reference
Method: Measurement of increased protein expression					
<i>Oncorhynchus mykiss</i> Liver	Dieldrine	5, 15 ppm	70	-	Curtis <i>et al.</i> 2000
<i>Oncorhynchus mykiss</i> Liver	Prochloraz, Nonylphenol diethoxylate	10, 100µg/l 100µg/l +33 µg/l	7-14	-	Sturm <i>et al.</i> 2001
<i>Fundulus heteroclitus</i> Liver	Creosot contaminated site vs. unpolluted site			× 2,65-3,25	Cooper <i>et al.</i> 1999
<i>Fundulus heteroclitus</i> Liver	3-methylcholanthrene	100 ml/kg	<i>ip</i> -injections	-	Cooper <i>et al.</i> 1999
<i>Ictalurus punctatus</i> Intestine and liver	VIN BNF BaP TCB	20 µg/kg diet 10-50 mg/kg diet 0.5- mg/kg diet 1-10 mg/kg diet	10	-	Doi <i>et al.</i> 2001
Method: Reduction of accumulation					
<i>Oncorhynchus mykiss</i> Hepatocytes	Prochloraz, Nonylphenol diethoxylate	10, 100µg/l 100µg/l +33 µg/l	7-14	50-76%	Sturm <i>et al.</i> 2001

8.2.2 Chemosensitisers

The protective role of the MXR defence system appears to be fragile. As opposed to MXR induction, it has also been demonstrated that many classes of chemicals are capable of competitively inhibit the MXR function Kurelec et al. (1997). These compounds are referred to as chemosensitisers and can be both naturally occurring compounds and pollutants. Chemosensitisers can saturate the MXR system, hence promoting the accumulation of potentially toxic xenobiotics. The relevance of MXR chemosensitisers has been investigated in several studies (see table below) and the results showed that their presence led to the apparition of toxic effects in organisms exposed to toxic compounds below the non-observed effect concentration.

The inhibitors of MXR activity has been described by Smital *et al.* (2004). In this article, it is described that pesticides, fragrances, microbial degradation products and natural inhibitors possess the ability to inhibit the important function the MXR defensive system. Other relevant work is presented in Table 20.

Table 20. Effect of chemosensitisers (inhibition of MXR activity).

Species and tissue	Factor	Inhibitory effect	Reference
<i>Dreissena polymorpha</i> Gill tissue	Sava river (high conc. chemosensitisers) compared to water with low conc.	×2 number of mutagens produced	(Britvic and Kurelec 1999)
<i>Mytilus edulis</i> <i>Lytechinus pictus</i> Embryo	Extracts of <i>Egrecia</i>	+	(Eufemia <i>et al.</i> 2002)
Fibroblasts	Extracts from landfill (household and industrial)	+	(Kurelec <i>et al.</i> 1998)
<i>Mytilus californianus</i> Gill tissue	Musk ketone Musk xylene	+	(Luckenbach <i>et al.</i> 2004)
<i>Suberites domuncula</i> Single cells <i>Corbicula fluminea</i> Gill tissue	Extracts of "marine snow" (amorphous aggregates)	+	(Muller <i>et al.</i> 1998)
<i>Corbicula fluminea</i> Gill tissue	Verapamil Staurosporine	+	(Waldmann <i>et al.</i> 1995)

8.3 Species sensitivity

In bivalves, results from in the field studies showed that it was possible to detect evidence of MXR activity in quite every organism investigated so far (for review see Pain & Parant, 2003).. The literature indicate that MXR expression in aquatic organisms such as bivalves, worms and sponges can reliably be used as and indicator of contaminant exposure and /or resistance (van der Oost et al., 2003). For fish, examples of interactions have been reported, but further characterization of wild fish populations is required.

8.4 Relevance of other factors

The main difficulty of using biomarkers in a monitoring programme is the interference of natural environmental factors with the biological responses. *In situ* biomarker responses can be altered or increased by physical or chemical parameters, hence, making the interpretation rather difficult. It is, therefore, essential to characterise the physical and chemical state of the sampling sites and to study natural variations in biomarker responses in order to differentiate

a pollution-induced stress from the background response and to correct for natural variations (Minier et al., 2002). Seasonal variations of MXR activity in polluted and unpolluted sites have been studied, as well as the effect of salinity, temperature and oxygen supply. The hypothesis that the defence system could be affected by environmental parameters such as temperature or algal proliferation has been proposed and supported by results of laboratory studies (Table 21). In addition, measured MXR activity is variable at the species level and even at the population level (Pain & Parant, 2003).

Table 21. The relevance of non-contaminant factors on MXR in bivalves.

Species and tissue	Factor	Response	Reference
Method: Measurement of increased protein expression			
<i>Mytilus edulis</i> Digestive gland and gill tissue	temperature salinity oxygen supply	- + +	Luedeking and Koehler 2004
<i>Mytilus galloprovincialis</i> Digestive gland	Seasonal variations	×10	Minier <i>et al.</i> 2000
<i>Crassostrea virginica</i> Gill or mantle tissue	Temperature, seasonal variation	×3	Keppler and Ringwood 2001
Method: increase of efflux			
<i>Dreissena polymorpha</i> Whole organism	Fuel oil 2		Pain & Parant, 2003

8.5 Links

MXR activity could conceivably increase the efflux of components required by the cell and thereby cause cellular damage, but this has not been documented experimentally. Further research is required to clarify the relative importance of induction as compared to chemosensitisation and both in relation to cell and individual health.

8.6 Background responses

Baseline responses have not been established for MXR activity for any species yet.

8.7 Assessment criteria

The assay still needs to be clearly defined; as yet there is no authoritative protocol for the determination of the various components of the system(s). Although shown to be affected by oil components in earlier work, it is premature to include MXR in offshore monitoring programmes at the moment.

8.8 Quality assurance

There has been no quality assurance activity in relation to MXR measurements as yet.

9. Peroxisomal proliferation

9.1 General introduction

Peroxisomes are ubiquitous single-membrane limited versatile organelles essential for β -oxidation of fatty acids, especially very long-chain fatty acids. Peroxisomes are highly heterogeneous organelles that vary in size, shape and protein composition depending on species, tissue, cell, metabolic state and developmental stage. In general, peroxisomes are single membrane-bounded organelles with a finely granular electron-dense matrix. They have a diameter ranging from 0.1 to 1.5 μm , and are more abundant in tissues active in lipid metabolism in general. Reviews of cell biology of peroxisomes in aquatic organisms and use as biomarker are given by Cancio and Cajaraville (2000) and Cajaraville *et al.* (2003).

Peroxisomes acquired relevance with the discovery that clofibrate, a pharmaceutical drug to lower the serum levels of lipids in patients with hypercholesterolemia, caused a massive proliferation of peroxisomes in rodents (Hess *et al.* 1965). It has later been shown that it is a link between the proliferation process and carcinogenesis in rodents (Reddy *et al.* 1980). Peroxisomal proliferation is usually accompanied by induction of all three enzymes of the peroxisomal β -oxidation pathway, namely acylCoA oxidase (AOX), peroxisomal hydratase-dehydrogenase or multifunctional enzyme (PH), and thiolase (Cajaraville *et al.* 2003). High-fat diets, cold adaptation, vitamin E deficiency, riboflavin deficiency, genetic obesity, diabetes and starvation can also lead to peroxisomal proliferation in rodents (Bentley *et al.* 1993). Disappearance of the peroxisomes proliferators results in a quick return of peroxisomes to basal numbers and enzyme activity levels. The most abundant peroxisomal enzyme is catalase, which is often used as a marker enzyme.

In addition to fibrates and other hypolipidemic drugs, peroxisomal proliferators comprise a heterogeneous group of compounds including phthalates, certain classes of pesticides, steroids, solvents, industrial chemicals, food flavours, therapeutic drugs, polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs) and natural products (Bentley *et al.* 1993; Cancio and Cajaraville 2000). One common feature of these compounds or their metabolic derivatives is a hydrophobic-lipophilic backbone with an acidic function, generally a carboxylic group. As peroxisomal proliferators are non-genotoxic and non-mutagenic compounds, their carcinogenicity has been ascribed to a number of different mechanisms including increased oxidative stress and DNA damage, increased DNA synthesis and mitogenesis, promotion of initiated liver foci and suppression of apoptosis (Cajaraville *et al.* 2003). Although there is a strong correlation between peroxisomal proliferation and hepatocarcinogenesis in rodent, a causal link remains to be established.

9.2 Dose-response

Aquatic organisms are exposed to a variety of contaminants in their environment. Some of these are known for their ability to cause peroxisomal proliferation, and increasing evidence suggests that aquatic organisms such as bivalves and fish are responsive to these environmental pollutants (Table 22 and Table 23).

Table 22. Effects on peroxisomal proliferation parameters in fish.

Species and tissue	Substance	Conc	Exposure time (days)	Response Peroxisomal proliferation	Response Catalase	Reference
<i>Anguilla anguilla</i> Liver	Dinitro-o-cresol	5, 50 250 µl/l	28	+	×1.4	(Braunbeck and Volkl 1991)
<i>Ictalurus punctatus</i> Liver	BKME	10, 20, 40%	14		×2-3	(Mather-Mihaich and Di Giulio 1991)
<i>Oncorhynchus mykiss</i> Liver	Endosulfan + disulfoton	50 mg/l 1 or 5 µg/l	18-34	×2.7 (transient)	None	(Arnold <i>et al.</i> 1995)
<i>Oncorhynchus mykiss</i>	Ciprofibrate injection	25, 25, 35 mg/kg	21	×2.3	×3.5	(Yang <i>et al.</i> 1990)
<i>Oncorhynchus mykiss</i> Liver	Atrazine	10, 20, 40, 80, 160 µg/l		+		(Oulmi <i>et al.</i> 1995)
<i>Oncorhynchus mykiss</i> Liver	Linuron	30 µg/l		+		(Oulmi <i>et al.</i> 1995)
<i>Sparus aurata</i>	Dieldrine (injection)	0.15-1 mg/kg	2 and 7		× 2.7 (7 days)	(Pedrajas <i>et al.</i> 1996)
<i>Fundulus heteroclitus</i> liver	2,4-dichlorophenoxyacetic acid	0.01, 0.1 and 1 ppm	7, 14 or 21	+		(Ackers <i>et al.</i> 2000)
<i>Brachydanio rerio</i> Liver	Lindane	40, 80, 110, 130, 150 µg/l			Decrease	(Braunbeck <i>et al.</i> 1990)
<i>Anguilla anguilla</i> Hepatocytes	Dinitro-o-cresol	5, 50, 250 µg/l	28	+	+	(Braunbeck and Volkl 1991)
<i>Cyprinus carpio</i> liver	PFOS (injection)	16-864 ng/g	1 and 5	-		(Hoff <i>et al.</i> 2003)
<i>Oncorhynchus mykiss</i> liver Transcriptome responses (microarray)	Pyrene Carbon tetrachloride	25, 100 µg/l 5,10,15 µl/l	4	 down regulation	suppression	 (Krasnov <i>et al.</i> 2005)
<i>Danio rerio</i> liver	Estradiol Dibutylphthalate Methoxychlor 4-t-octylphenol ethynylestradiol	10 µg/l 500 µg/l 100 µg/l 500 µg/l 10 µg/l	7 and 15	+ + + + +	Up. (15d) Down (7d)	(Ortiz-Zarragoitia and Cajaraville 2005)
<i>Solea ovata</i>	B[a]P (injection)	0.1, 0.5, 1, 10 mg/kg	7	+ (1mg/kg and higher)		(Au <i>et al.</i> 1999)

Part of the table is based on Cancio *et al.* (2001).

Table 23. Effects on peroxisomal proliferation parameters in bivalves.

Species and tissue	Substance	Concentration	Exposure time (days)	Response Peroxisomal proliferation	Response Catalase	Reference
<i>Mytilus edulis</i> Digestive gland	PAHs or PCBs		30	+	×2 (PCBs)	(Krishnakumar <i>et al.</i> 1997)
<i>Mytilus edulis</i> Digestive gland	Benzo[a]pyrene Menadione	50 ppb 1 ppm	2, 6, 19		×1.2 (transient) ×1.4	(Livingstone <i>et al.</i> 1990)
<i>Mytilus galloprovincialis</i> Digestive gland	WAF lubricant oil WAF ural crude oil WAF maya crude oil	0.6, 6, 40%	21, 49, 91	3.8-5 2-3 1.4-1.5	×2,25 ×1.9 ×1.8	(Cajaraville <i>et al.</i> 1997)
<i>Mytilus galloprovincialis</i> Digestive gland	WAF lubricant oil WAF (injected) Clofibrate Clofibrate (injected)	5% 100 µl 5% 2.5 mg/l 1,5 mg/100µl	1, 7, 21	Decrease ×1.45-1.6 Decrease ×1.12-2.7	None Decrease None Decrease	(Cancio <i>et al.</i> 1998)

	DEHP DEHP (injected) B[a]P B[a]P (injected)	60 µg/l 100µg/100µl 300µg/l 500µg/100µl		Decrease None None None	None None None None	
<i>Mytilus galloprovincialis</i> Digestive gland	B[a]P DEHP Cd Mix B[a]P and Cd		21	+ Reduction -	+ +	(Orbea <i>et al.</i> 2002)
<i>Mytilus galloprovincialis</i> Digestive gland	Different sites (PAH burden) – transplant study		29	×4 (polluted vs. nonpolluted)		(Cajaraville <i>et al.</i> 2003)

Part of the table is based on Cancio *et al.* (2000).

In addition to exposure studies with bivalves, field studies have demonstrated differences in the number and size of peroxisomes of *Mytilus* sp. collected in sites with different concentrations of organic pollutants (Cancio *et al.*, 2000). In a reciprocal transplant experiment with mussels, it was demonstrated that the peroxisome proliferation response may be reversible.

In field studies with fish from sites with different pollutant concentrations have been compared, contradictory results have been obtained with activities of peroxisomal antioxidant enzymes.

9.3 Species sensitivity

The first studies of peroxisomes in the aquatic environment was in fish, and in fish there are data indicating that fibrates hypolipidemic drugs, various pesticides, bleached kraft pulp and paper mill effluents (BKMEs), PAHs, phthalates, alkylphenols and estrogens can cause peroxisome proliferation. An overview of these studies has been provided by Cajaraville *et al.* (2003). Later research has focussed on mussels (*Mytilus* sp) because these filter-feeding bivalves are used worldwide as a bioindicator and sentinel species of environmental pollution in marine and estuarine areas. Peroxisomes in the digestive gland of mussels share several characteristics with vertebrate liver peroxisomes, but they also show some special features.

9.4 Confounding factors

The detailed knowledge of seasonal variations in peroxisomal parameters is essential to be able to detect changes induced by contaminants. In addition, peroxisomes may respond differently to pollutants depending on the time of the year. Several studies have investigated this, and some of them are presented in the table below. In addition to seasonal fluctuations and population-specific responses, gender of animals could be another possible confounding factor (see table for details).

Table 24. Overview of factors that may affect peroxisomal proliferation in various organisms.

Species and tissue	Factor	Response	Reference
<i>Mytilus galloprovincialis</i> Digestive gland	Seasonal variation	× 8 (spring vs. winter)	(Cancio <i>et al.</i> 1999)
<i>Leuciscus idus melanotus</i> hepatocytes	Low and high temperatures (14 and 28 °C)	+	(Braunbeck <i>et al.</i> 1987)
<i>Mytilus galloprovincialis</i> Digestive gland	Seasonal variation PAH burden (different sites)	+ (uncontamin. site)	(Orbea <i>et al.</i> 1999)
<i>M. galloprovincialis</i> <i>Crassostrea</i> sp <i>Carcinus maenas</i> <i>Mugil cephalus</i>	Seasonal variation PAH/PCB body burden (different sites)	- (season var. mussel) + (season var. oyster) + (season var. crab) + (season var. fish)	(Orbea <i>et al.</i> 2002)
<i>Salmo trutta</i>	Seasonal variation	+ (more pronounced in females)	(Rocha <i>et al.</i> 1999)
<i>Fundulus heteroclitus</i>	Starvation (96 h) gender	Decrease	(Haasch 2002)
<i>Mugil cephalus</i> hepatocytes	Seasonal variation	Level higher in summer	(Orbea <i>et al.</i> 1999)
<i>Danio rerio</i>	Gender Developmental stage	No significant difference observed	(Ibabe <i>et al.</i> 2005)

9.5 Links

The potential deleterious effects on reproduction, development and growth are unknown, providing a challenge for future investigations (Cajaraville *et al.*, 2003). In rodents, peroxisome proliferators are known to adversely affect reproduction and development. This opens a new field of research aiming to establish possible connections between peroxisome proliferation and endocrine disruption (Ortiz-Zarragoitia *et al.*, 2005).

9.6 Background responses

There are still only a limited number of laboratories that have sufficient competence to perform the required analyses. Baseline responses have been defined by the University of the Basque Country for mussel, but it is not clear whether the baseline will be similar in the North Sea.

9.7 Assessment criteria

Criteria have been established for Spain, but as mentioned above it remains to be clarified whether they can be applied to mussels from other geographical areas.

9.8 Quality assurance

Internal quality assurance guidelines exist, e.g. SOP for procedures. No international intercalibration has been performed for the relevant methods.

10. Acetyl cholinesterase (AChE) inhibition

10.1 Introduction

In vertebrates, acetylcholine (ACh) acts as an excitatory transmitter for voluntary muscle in the somatic nervous system. ACh also serves as both a preganglionic and a postganglionic transmitter in the parasympathetic nervous system. Cholinesterase enzymes (ChE) are responsible for the removal of ACh from the synaptic cleft by hydroxylation. Cholinesterases are usually divided into the acetylcholinesterases (AChEs) and the butylcholinesterases (BChEs). In fish, the tissues containing most AChE are brain and muscle.

Among the AChE inhibitors, the most common are a) organophosphate insecticides (OPs) such as terbufos, trichlorfon and chlorpyrifos b) carbamate insecticides such as carbofuran and carbaryl and c) synthetic pyrethroid compounds. OPs have become one of the most widely used classes of insecticides worldwide because of their relatively non-persistent characteristics in the environment. Although rapidly degraded, the substances lack target specificity and have high acute toxicity toward many nontarget vertebrate and invertebrate species. OPs and carbamates are applied in agricultural and residential uses, and can also be accidentally introduced into the environment through overspray, aerial drift or runoff. Thus, many terrestrial as well as aquatic organisms may be at risk for intoxication caused by exposure to these compounds.

This biomarker could be affected by e.g. biocides used in oil and gas production (and hence released with produced water). It is a sensitive marker for a range of substances that may have neurotoxic effects.

Tissue residues of AChE inhibitors associated with reduced survival appear to be highly variable, ranging from 0.00004 mmol/kg to 29 mmol/kg (Barron *et al.* 2002). It has been suggested that brain AChE inhibition levels of >70% are associated with mortality in most species (Fulton and Key 2001).

10.2 Dose-response

Most of the studies involving AChE inhibitors have been performed with insecticides, but some studies have also focussed on the effect of classical pollutants on the activity of AChE (Table 25 and Table 26).

Table 25. Dose-response, background response and sensitivity in fish species (brain tissue if nothing else mentioned).

species	substance(s)	concentration	exposure time (days)	baseline/control (level/activity)	inhibition %	reference
Experimental studies						
Salmon, <i>Salmo salar</i>	Fenitrothion	0.005-0.213 mg/L	7		12 – 58	(Morgan <i>et al.</i> 1990)
Coho Salmon, <i>Oncorhynchus kisutch</i>	Chlorpyrifos	0.6-2.5 µg/L (nominal) 0.3-1.7 µg/L initial measured)	4	Brain: 23.2 ± 0.5 (SE) Muscle: 38.9 ± 1.4 (SE) µmol/min/g	lowest conc: Brain: 23% Muscle: 12% highest conc: Brain: 64% Muscle: 67%	(Sandahl <i>et al.</i> 2005)
larval rainbow trout (<i>Oncorhynchus mykiss</i>)	Diazinon	250-1000 µg/L	1 and 4	~13 and ~11.5 µmol/min/g	lowest conc: ~30% ~21% Highest conc: ~38% ~30% lowest conc: ~64% ~72% Highest conc: ~64% ~41%	(Beauvais <i>et al.</i> 2000)
	Malathion	20-40 µg/L		~12.5 and ~13 µmol/min/g		
<i>Oncorhynchus mykiss</i>	Polluted sediments, mainly PAH	5.8 ng/g d.w.	14		No significant decrease	(Sanchez-Hernandez <i>et al.</i> 1998)
Field studies						
Menhaden, <i>Brevoortia tyrannus</i>	OPs				~50% comp. to reference site	(Williams and Sova 1966)
Flounder muscle	Sites in the Baltic, organic pollutants			94-315 nmol/min/mg		(Kopecka <i>et al.</i> 2004)
<i>Anguilla anguilla</i> muscle	PAH and organochlorine content in fish			Relationship between volatile PAHs and activity of AChE		(Roche <i>et al.</i> 2002)
Cage deployment studies						
<i>Fundulus heteroclitus</i>	Azinphosmethyl	0.00-7.00 µg/L	4		98% at highest conc. (mortality 17%)	Fulton & Key, 2001

Table 26. Dose-response, background response and sensitivity in bivalves.

species	substance(s)	concentration	exposure time (days)	baseline/control (level/activity)	inhibition %	reference
Experimental studies						
<i>Crassostrea gigas</i> gills	Dichlorpyros	0.1-1.0 mg/L	0-48 h		83% (6h) 87% (4h)	(LeBris <i>et al.</i> 1995)
<i>Mytilus galloprovincialis</i> gills	iso-OMPA	10 ⁴ M	15 min	24.2 ± 3.5 SEM nmol/min/mg	~15% ~58%	(Escartin and Porte 1997)
	Eserine	10 ⁶ M, 10 ⁵ M and 10 ⁴ M		24.2 ± 3.5 SEM nmol/min/mg	~70% and ~80%	
<i>Mytilus galloprovincialis</i>	B[a]P in food	50 mg/kg dry weight mussel	28		Significantly depressed	(Akcha <i>et al.</i> 2000)

species	substance(s)	concentration	exposure time (days)	baseline/control (level/activity)	inhibition %	reference
gills						
Field studies						
<i>Mytilus edulis</i> gills	Sites in the Baltic, organic pollutants			15-38 nmol/min/mg		(Kopecka <i>et al.</i> 2004)
<i>Mytilus edulis</i>	Oil spill after "Erika" accident		1 year after accident		Significant reduction	(Bocquene <i>et al.</i> 2004)
<i>Mytilus galloprovincialis</i> hemolymph	Portuguese coast			30-70 nmol/min/mg	Decreased in polluted site vs. unpolluted	(Moreira and Guilhermino 2005)
<i>Mytilus galloprovincialis</i> Digestive gland and Gills	West coast of France	PAH, PCB and metals in mussels determined			Inhibition by unidentified compound	(Romeo <i>et al.</i> 2003)

10.3 Species sensitivity

Both fish and mussels have been successfully employed in studies of interactions of OPs with AChE response. However, it seems that much of the work with classical pollutants (PAH, PCB, metals) are done with bivalves in field studies. In a comparative study, Galgani & Bocquené (1990) found that AChE from whole mussel (*Mytilus edulis*) was less sensitive to five OPs than the enzyme from the muscle tissues of the shrimp *Palaemon serratus* and two other fish species.

10.4 Relevance of other factors

Seasonal variations of AChE activity in polluted and unpolluted sites have been studied (Table 27). In addition, the effect of fish length and the location of the enzymes in the cells have been studied.

Table 27. Factors that have been found to affect AChE in fish and mussels.

Species and tissue	Factor	Response	Comment	Reference
<i>Leuciscus cephalus</i>	Fish length	+	The larger fish had the lower ChE-activities	Flammarion <i>et al.</i> 2002
<i>Crassostrea gigas</i>	Responsiveness of different ChE to OPs	+	Soluble enzymes insensitive, membrane bound highly sensitive	Bocquene <i>et al.</i> 1997
<i>Sciaenops acellatus</i> <i>Fundulus heteroclitus</i>	Relationship between azinophosmethyl-induced AChE inhibition and mortality		Inhibition of AChE in the peripheral nervous system may be a better predictor of OP-induced mortality than brain inhibition	VanDolah <i>et al.</i> 1997
<i>Mytilus galloprovincialis</i>	Seasonal variation (2 year study)		Variation of AChE seemed less variable with season than other biomarkers studied	Bodin <i>et al.</i> 2004
<i>Mytilus galloprovincialis</i> hemolymph	Seasonal variation (1 year study)	+	AChE activity showed an apparent seasonal trend in 3 of 4 sites	Moreira and Guilhermino 2005
<i>Mytilus galloprovincialis</i> gills and digestive gland	Seasonal variation	+	Strong seasonal variation from June to October of all parameters was noted.	Romeo <i>et al.</i> 2003

10.5 Links

It has been suggested that brain AChE inhibition levels of >70% are associated with mortality in most species (Fulton & Key, 2001). However, some species may tolerate much higher levels (>90%) of brain AChE inhibition. Sublethal effects on stamina have been reported for some estuarine marine fish in association with brain AChE inhibition levels as low as 50% (Fulton & Key, 2001).

Reduced swimming stamina has been shown for e.g. sheepshead minnow (Cripe et al., 1984) Swimming stamina was affected in fish exposed to the organophosphorus insecticide EPN at 4.1 and 2.2 µg/l (reduced by 54% and 43%, respectively).

Brain AChE inhibition and reductions in spontaneous swimming and feeding activity were significantly correlated in Coho Salmon (*Oncorhynchus kisutch*) exposed to chlorpyrifos (Sandahl et al., 2005).

Changes in swimming speed and distance of larval rainbow trout (*Oncorhynchus mykiss*) were significantly correlated with changes in ChE activity after exposure to diazinon and malathion (Beauvais et al., 2000).

10.6 Background responses

There is not sufficient data to set baseline responses for AChE for species relevant to the North Sea.

10.7 Assessment criteria

AChE is one of the methods with an obvious link to higher levels effects. A fish that swims slowly or is disoriented will have a much lower probability of survival than a healthy fish. Such values could be modelled without too much effort with the right input data. Unfortunately such data are not available for North Sea species.

It is less clear whether AChE inhibition in mussels will affect e.g. clearance rate, which would have direct consequences for fitness.

As indicated above, a percentage inhibition of AChE could be set as a limit, but effects on behaviour or swimming stamina is highly species dependent. There is not sufficient data to set assessment criteria for North Sea species.

10.8 Quality assurance

An internationally agreed protocol has been developed for AChE determination (Bocquené & Galgani, 1998). International intercalibrations have been performed within specific projects.

11. Vitellogenin (vtg) induction

11.1 Introduction

In all egg-laying vertebrates, including fish, most of the material that will eventually constitute the egg is synthesised in the liver of the female. This includes both the precursor-protein for egg yolk, vitellogenin, and precursor proteins for the eggshell (Oppen-Berntsen et al. 1992). Since the proteins are synthesised in the liver they need to be transported with blood to the gonads, which means that at the time of ovarian development the concentration of vitellogenin in the blood plasma of female fish can be phenomenal; up to 70% of the total protein in the plasma can be vitellogenin. Synthesis of vitellogenin is regulated by the steroid estrogenic hormone estradiol. High levels of estradiol mean high output of vitellogenin from the liver. For some reason, the gene for vitellogenin is also available for transcription in male fish, but it is rarely turned on because estradiol-levels are generally low in male fish. There are some fish species in which the male has naturally somewhat higher estradiol concentrations in the blood, which means that they also synthesise vitellogenin (to no use). This has been found for plaice (Scott et al., 2000) and for some wrasse species (Hylland, unpublished).

Early in the 1990s, fish physiologists that studied reproduction in freshwater fish in UK rivers observed male fish with high levels of vitellogenin in the blood and even some individuals with oocytes mixed into their testes. Further studies showed that substances in sewage effluents were responsible, which included natural estrogens (from the female half of the human population), synthetic estrogens (from pharmaceuticals) and industrial or household chemicals.

In the past decade it has been established that most sewage effluents are estrogenic and that many chemicals have estrogenic or anti-estrogenic activity, including substances in produced water effluents. Vitellogenin is close to a perfect biomarker in that the response is well-characterised and limited to substances with estrogenic (or anti-estrogenic) activity. Following exposure to estrogens, the response measured in plasma may increase million-fold compared to controls, although 10-fold or 100-fold increases are generally more commonly seen under environmental conditions (Scott & Hylland, 2002).

11.2 Dose-response

As for most of the biomarkers there is most data on freshwater fish. For the purpose of this review the main focus will be on marine fish. All measurements of vitellogenin referred to in this review have been done on plasma. It is possible to analyse for vitellogenin in liver as well, but concentrations are generally much lower than in plasma. Studies on gene expression naturally use liver tissue. Relevant studies are indicated below (Table 28 and Table 29).

Table 28. North Sea relevant field studies on vitellogenin responses (all studies with male fish).

<i>species</i>	<i>areas</i>	<i>exposure time</i>	<i>concentration at reference location</i>	<i>induction (fold)</i>	<i>reference</i>
Atlantic cod (<i>Gadus morhua</i>)	caging - Staffjord gradient	5 wks	30-50 ng/mL	5 (trend, ns)	Scott et al., 2006a
Atlantic cod (<i>Gadus morhua</i>)	North, Norwegian, Barents sea	field	100-1000 ng/mL	5-10	Scott et al. 2006b
Atlantic cod (<i>Gadus morhua</i>)	Skagerrak coastal sites	field	n/a	4	Hylland et al., 1996
flounder (<i>Platichthys flesus</i>)	UK estuaries	field	200 ng/mL	5-1000 (different estuaries)	Kirby et al. 2004
flounder (<i>Platichthys flesus</i>)	Dutch coastal areas	field	500 ng/mL	2-3	Vethaak et al. 2005
flounder (<i>Platichthys flesus</i>)	North Sea, UK estuaries	field	100 ng/mL	10-4 - 10-6	Allen et al. 1999

Table 29. Selected studies on dose-dependency of vitellogenin following exposure to contaminants; all studies with male fish except Hylland (unpublished), which used juvenile fish.

<i>species</i>	<i>ecosystem</i>	<i>substance</i>	<i>lowest-highest concs</i>	<i>exposure time</i>	<i>baseline/control (level/activity)</i>	<i>induction (fold)</i>	<i>LOEC</i>	<i>reference</i>
Atlantic cod (<i>Gadus morhua</i>)	marine	water, 4-NP	5, 20, 100 µg/L	1 wk	100 ng/mL	15	20 mg/kg	Hylland, unpublished
Atlantic salmon (<i>Salmo salar</i>)	freshwater	inject, BPA	1, 5, 25, 125 mg/kg	2 wk	n/a	>5	25 mg/kg	Arukwe et al. 2000
eelpout (<i>Zoarces viviparus</i>)	marine	inject, 4-NP	10, 100 mg/kg	25 d	1-5 mg/mL	1000	10 mg/kg	Christiansen et al. 1998
flounder (<i>Platichthys flesus</i>)	marine	oral, 4-OP	0.5, 1, 2.5, 5, 7.5, 10, 25 mg/kg	11 d	40-80 ng/mL	>500	5 mg/kg	Madsen et al. 2003
flounder (<i>Platichthys flesus</i>)	marine	oral, 4-OP	10, 50, 100 mg/kg	11 d	40-80 ng/mL	>10-6	n/r	Madsen et al. 2003
flounder (<i>Platichthys flesus</i>)	marine	inject, 4-NP	10, 50, 100, 150, 200 mg/kg	2 wk	n/a	>5	10 mg/kg	Christensen et al. 1999

11.3 Relevance of other factors

As mentioned above, the vitellogenin response is thought to be limited to estrogenic or anti-estrogenic substances. In contrast to other biomarkers other endogenous or exogenous factors do not appear to have a strong influence. In a series of studies, juvenile cod were tested at NIVAs research facility at Solbergstrand for its suitability as an indicator species for vitellogenin research and monitoring. One of the issues to be resolved was whether there would be a seasonal variation in the sensitivity of juvenile cod to estrogens. There was some seasonal variation in the baseline level of the protein, but no variation in the response following treatment with estrogens (Figure 8). Other studies showed that starvation did not affect the induction and neither did differences in acclimation temperature, salinity or changes in temperature (Hylland & Aspholm, unpublished).




Figure 8. Vitellogenin in juvenile cod exposed for one week to 5 ng/L estradiol (top) or 50 $\mu\text{g/L}$ 4-nonylphenol (bottom) through water; the figure shows median, quartiles and 10/90 percentiles. Note logarithmic y-axis.

11.4 Links

Elevated concentrations of plasma vitellogenin in male or juvenile fish (which would be low) indicates the presence of estrogens in the environment. Although exceedingly high concentrations of vitellogenin in the plasma may clog the kidneys of fish and cause mortality due to kidney malfunction and that there is some metabolic cost in producing the protein, it is unlikely that the presence of vitellogenin in itself will have strong negative effects. The presence of estrogens in sufficient quantities to cause induction of vitellogenin may affect larval stages of fish in the environment. Studies on model species such as zebrafish indicate that exposure to estrogens may skew the sex ratio in favour of female fish even at very low concentrations (Andersen et al. 2003, Örn et al. 2003). In fact, the concentrations of model estrogens needed to change the sex ratio was lower than that needed to induce vitellogenin in those studies.

11.5 Background responses

The background response of vitellogenin in North Sea species has until now been defined by the quality of the analytical procedures used for analyses of the protein, i.e. RIA and ELISA, since vitellogenin levels in male and juvenile fish are very low for most species (on the order

of 10-100 ng/mL). As indicated above there is a seasonal variability in plasma vitellogenin, even in juveniles (and presumably males); 1 µg vitellogenin/mL plasma appears to be the upper limit for plasma vitellogenin in Atlantic cod from unpolluted areas (Scott et al., in press; Hylland, unpublished observations). The concentration would have to be determined using a quantitative immunoassay (ELISA, RIA).

11.6 Assessment criteria

As mentioned above, changed sex ratios have been seen in experimental studies at concentrations below those for which there was an increase in vitellogenin. Sex ratios have only been determined in field studies for species with suitable reproductive physiology, such as eelpout (*Zoarces viviparus*). The high sensitivity of some species to estrogens indicate that some caution should be exercised and that a three-fold increase over the upper limit of the baseline for cod (i.e. 3 µg/mL) would indicate the presence of estrogens in the environment.

11.7 Quality assurance

A widely available protocol for the analysis of plasma vitellogenin in fish is available (Scott & Hylland, 2002) and there are commercial kits available for relevant species, e.g. Atlantic cod (<http://www.biosense.com>). In 2004-2005 an international intercalibration was performed within BEQUALM, the results of which indicated that laboratories that used similar methods (quantitative ELISA) had comparable results.

12. Metallothionein (MT) concentration

12.1 Introduction

The low-molecular-weight protein metallothionein (MT) is present in most cells in vertebrates and some invertebrates. It is a peculiar protein in that around a third of the protein is made up by the sulphhydryl-containing amino acid cysteine, which contributes directly to the binding of 7-12 metal atoms in each MT molecule (Klaassen et al. 1999). Early studies indicated that concentrations of MT would increase following metal exposure (especially to cadmium) and it was then suggested that metal detoxification was the major function of the protein. Later work has shown that MT is involved in the regulation of the intracellular availability of zinc (Zn) and/or copper (Cu), which to some implied a possible role in total gene-regulation (through regulating Zn availability). Metallothionein is induced in cells exposed to free radicals and do indeed appear to protect against damaging effects from e.g. ionising radiation or radical-generating agents (Kagi & Schaffer 1988). The production of MT knock-out mice definitely established that the protein is not essential to survival, at least not in mammals, but will provide protection from toxic metals such as cadmium (Masters et al. 1994).

In addition to Cu and Zn, MT binds non-essential metals such as cadmium (Cd), mercury (Hg) and silver (Ag). The synthesis of the protein is induced by elevated intracellular concentrations of the metals mentioned above. Metallothionein induction is a response to elevated intracellular metal concentrations and the protein has for that reason been suggested as a biomarker for environmental metal contamination (Engel & Roesijadi 1987, Haux & Förlin 1989, George & Olsson 1994).

12.2 Analytical aspects

The methods most commonly used to quantify MT in tissues exploit some of the characteristics of this protein: its metal content, the number of sulphhydryl groups, its size and/or its heat-stability. In addition, various immunochemical assays using MT-specific antisera have been established in the past 10-15 years (Roesijadi et al. 1988, Hogstrand et al. 1989, Hylland 1999). More recently, metallothionein gene expression (mRNA) has been quantified in fish species using rtPCR (Cheung et al. 2004, Tom et al. 2004). Similar methods have also been developed for mussels (Dondero et al. 2005). Quantification of the protein, and not mRNA, should theoretically be most useful for monitoring purposes for two reasons. Firstly, the physiological and toxicological significance of a response will depend upon the concentration of MT, and not MT mRNA, present in the cell. Secondly, the half-life of MT mRNA is much shorter than the half-life of MT and the time-span of a measurable response thus shorter. On the other hand, recent studies do suggest that expression studies (rtPCR) may also be relevant for monitoring (see George et al. 2004).

The existing methods for quantification of MT in both fish and invertebrates (mussels) may conveniently be divided into four categories: (1) quantification by metal-content, (2) metal-substitution, (3) quantification of sulphhydryl groups and (4) immunoassays. Quantification of sulphhydryl groups is either done using a spectrophotometric method (viarengo2000) or electrochemically using a polarograph (Olafson & Olsson 1991).

12.3 Metallothionein in mussels

As for many other biomarkers, mussels have been favourite invertebrate species for metallothionein studies (Noël-Lambot 1976, Bebianno & Langston 1991, Cajaraville et al. 2000, Mourgaud et al. 2002). Metallothionein in mussels have been suggested for monitoring and there have been international intercalibration exercises (Viarengo et al. 2000). Recently, rtPCR techniques were developed to quantify the two isoforms in mussels, MT10 and MT20 (Dondero et al. 2005). As for fish, different quantification methods have been used and a recent study has showed that they give entirely different results (Zorita et al. 2005). Metallothionein was determined in mussels during the BECPELAG workshop with the most widely used method (polarography), but there were no differences between locations (Erk & Raspor, in press). The general impression is that metal-derived increases in mussel MT is of a much smaller magnitude than increases that can be seen in fish, although (Viarengo et al. 1999) contends that mussel MT is more suited for monitoring than fish MT due to its closer link to external metal stress. The ICES working group on biological effects of contaminants (WGBEC) has not recommended MT in mussels for monitoring, but has included it with “promising methods” (WGBEC, 2005).

12.4 Dose-response in fish

Although MT is found in detectable amounts in most tissues, there is limited data on organs other than the liver. As for most other biomarkers, there is a wealth of studies in freshwater fish, but less for marine species. As mentioned above, it is not really feasible to compare studies with different methods.

Table 30. Selected studies of metallothionein in marine fish species; MT measured with pulse polarography in all studies except Sulaiman et al., who used the Cd-saturation assay.

<i>species</i>	<i>tissue</i>	<i>lowest-highest concs</i>	<i>exposure time</i>	<i>baseline/control (level/activity)</i>	<i>induction (fold)</i>	<i>reference</i>
Atlantic cod (<i>Gadus morhua</i>)	liver	field	field	4-7 µg/mg protein	none	Holth, 2004
Atlantic cod (<i>Gadus morhua</i>)	liver	caging – 4 locations	3 mo	1 µg/mg protein	none	Beyer et al., 1997
Atlantic cod (<i>Gadus morhua</i>)	liver	reference locations	field	6.5-16 µg/mg protein	none	Ruus et al., 2003
dab (<i>Limanda limanda</i>)	liver	reference locations	field	7.2-13 µg/mg protein	none	Ruus et al., 2003
flounder (<i>Platichthys flesus</i>)	gill	caging – 4 locations	3 mo	0.8 µg/mg protein	none	Beyer et al., 1996
flounder (<i>Platichthys flesus</i>)	kidney	caging – 4 locations	3 mo	0.4 µg/mg protein	none	Beyer et al., 1996
flounder (<i>Platichthys flesus</i>)	liver	field	field season	7-16 (female), 5-12 (male) µg/mg protein	none	Nissen-Lie, 1997
flounder (<i>Platichthys flesus</i>)	liver	field	field	6-11 µg/mg protein	none	Holth, 2004
flounder (<i>Platichthys flesus</i>)	liver	field	field gradient	20 µg/g ww	19	Sulaiman et al., 1991
flounder (<i>Platichthys flesus</i>)	liver	caging – 4 locations	3 mo	1 µg/mg protein	none	Beyer et al., 1996
flounder (<i>Platichthys flesus</i>)	liver	reference locations	field	8.4-15 µg/mg protein	none	Ruus et al., 2003
plaice (<i>Pleuronectes platessa</i>)	liver	reference locations	field	6.5-14 µg/mg protein	none	Ruus et al., 2003

12.5 Species sensitivity

There is limited knowledge of how metallothionein in tissues of different fish or invertebrate species respond to metal stress. For the marine species relevant in the North Sea area there appear to be surprisingly little difference in baseline hepatic MT levels. Both design and analytical protocols differ between studies and it is rarely feasible to compare species directly.

12.6 Relevance of other factors

There are species differences in how endogenous and exogenous factors other than metal stress affect MT in fish tissues, as indicated in Ruus et al. (2003). In a comprehensive multiple regression analysis for Atlantic cod (including more than 600 individual fish sampled at 8 sites over 5 years), both organic contaminants and essential metals came up as highly significant, in addition to size and a site-related factor that was not reflected in tissue-levels of metals or organochlorines. Similar regression models for flatfish included essential metals, but also tissue concentrations of organic contaminants and size. All samples were collected at the same time of the year (September-October) and water temperatures were surprisingly similar (within 2-3°C) at the different locations sampled through a five-year period.

The relevance of various factors that affect hepatic MT in flounder was assessed by Nissen-Lie (1997) in a seasonal study over 1 year. As for cod, hepatic concentrations of the essential metals copper and zinc appeared to drive (or be driven by?) hepatic concentrations of MT. Metallothionein in the liver varied through the year in both male and female fish, apparently related to the demand for Zn in gonad development and maturation. As for cod, results from Lacorn et al. (2001) for dab similar identify season as an important factor for hepatic MT, but also suggest that water temperature is relevant. The latter can not easily be distinguished from season, so no clear conclusions can be drawn. The main conclusions in Rotchell et al. (2001) were that age and season are important for variations in hepatic MT in flounder.

The results in Hylland et al. (1996) indicated that starvation had less influence on MT than on other biomarkers in flounder fed cadmium-contaminated polychaetes. There was also surprisingly little effect of co-exposure of cadmium with BaP or PCB in that study. Corresponding results were found in the study by Sandvik et al. (1997), but fish treated with BaP prior to Cd appeared to have a lower capacity to synthesise MT than fish who only received Cd or PCB prior to Cd. There are unfortunately no similar studies with Atlantic cod or any other gadiid.

12.7 Links

As mentioned in the introduction to this section, metallothionein appears to be a household protein in many organisms (at least in mammals) with a range of functions, but none really essential. It is not clear whether this is the case for fish. Fish have higher levels of metals in the liver than most mammals and some fish species have phenomenally high concentrations of metals under normal conditions (Hogstrand et al., 1996; Hogstrand & Haux, 1996; Thompson et al., 2003). Data for flounder and cod from unpolluted to moderately polluted areas support a role for metallothionein in trace metal (zinc and possibly copper) metabolism in marine fish, but non-essential metals such as cadmium do not appear to have a strong effect (in contrast to the situation in some freshwater ecosystems). Metallothionein is however one of the genes most clearly expressed in rtPCR studies and the protein appears to have a function in a general stress response, as much or more related to organic contaminants as to metals. There are however no clear links from metallothionein induction to individual health or reproduction. Some freshwater studies have shown reduced growth at metal-levels which also induce MT (e.g. Roch et al., 1984), but such levels are rarely if ever encountered in marine ecosystems.

12.8 Background responses and assessment criteria

As long as an appropriate species is used, sampling is performed during the appropriate time period and individuals selected on the basis of sex and size, absolute assessment criteria may be used for hepatic metallothionein. The main dataset used in this context has been derived from the JAMP monitoring programme (Ruus et al., 2003), but supplemented with other studies, e.g. Nissen-Lie (1997) for flounder.

12.9 Quality assurance

Metallothionein was included in BEQUALM and intercalibration between 14 laboratories run on liver samples from cod. The results clearly showed some of the inherent problems of the analytical procedures. Metal-saturation assays and polarographic analyses produced remarkably similar results, whereas immunochemical assays (ELISA) and the colorimetric assay produced results that deviated from the overall median

13. δ -aminolevulinic acid dehydratase (ALA-D) inhibition

13.1 Introduction

Many metals are in fact essential to all organisms, e.g. Cu, Zn, Fe, Mn and Mo, but there are also metals for which no biological function is known such as Hg, Cd, Au, Ag and Pb. Non-essential metals generally have much higher toxicity than the essential metals. Uptake, storage and excretion are also less well controlled for the non-essential than for essential metals.

One of the most important toxic mechanisms of non-essential metals is interaction with and inhibition of enzymes, especially enzymes with metal co-factors. Δ -aminolevulinic acid dehydratase (ALA-D, E.C. 4.2.1.24) is such an enzyme, which has zinc as a co-factor (Granick et al., 1972). This enzyme catalyses a step in the synthesis pathway for heme and is found in bacteria, plants and invertebrates as well as in vertebrates. Heme is incorporated in macromolecules such as hemoglobin and cytochromes. In mammals and birds, inhibition of ALA-D may lead to anaemia since it is one of the rate-limiting enzymes in heme (and hence hemoglobin) synthesis. This does not appear to be the case for fish (Larsson et al. 1985). The reason for ecotoxicological interest in ALA-D is its inhibition by Pb, even at very low exposure levels (Hodson et al. 1984, Haux & Förlin 1988).

ALA-D has been used to investigate effects of metals (mainly lead) in rainbow trout (Hodson 1976, Hodson et al. 1977, Hodson et al. 1978, Addison et al. 1990, Sordyl & Osterland 1990), longear sunfish (Dwyer et al. 1988), perch (Haux et al. 1985), salmon (Johansson-Sjöbeck & Larsson 1979), flounder (Johansson-Sjöbeck & Larsson 1978), carp (Nakagawa et al. 1995, Nakagawa et al. 1998), various catostomid species (Schmitt et al. 1984), Atlantic cod (Ruus et al. 2003) and grey mullet (Krajnovic-Ozretic & Ozretic 1980). As will be apparent from this list, the method has predominantly been used with freshwater species.

13.2 Analytical considerations

There has been a tradition to express results for ALA-D in terms of blood cell volume (e.g. hematocrit), but more recent recommendations suggest to use protein (Hylland, 1999). As for other methods, it is not possible to compare different studies because values have not been reported in the same format.

13.3 ALA-D in invertebrates

As mentioned above, ALA-D is also present in invertebrates and it is possible to adapt the existing protocol for fish. There is limited knowledge of effects in invertebrates, although ALA-D would be assumed to be sensitive to metals in invertebrates as well.

13.4 Dose-response and species sensitivity

ALA-D activity can be determined in liver, kidney, spleen and red blood cells in teleosts (see (Hodson et al. 1984). The sensitivity, timing and specificity of ALA-D inhibition in response to lead exposure have been reviewed (Hodson et al. 1984). For rainbow trout, red blood cell

ALA-D activity may be inhibited following exposure to less than 5 µg/L of lead through water (Haux et al. 1986) and blood lead concentration of 300 µg/L (Hodson et al. 1982), although other studies indicate a lower sensitivity (Burden et al. 1998). In rainbow trout, ALA-D inhibition is evident 1-2 days following exposure to lead and there is a clear dose-response relationship (Hodson et al. 1982). The half-life of the response will depend on the amount of lead accumulated in other tissues (in equilibrium with blood), but will generally be of the order of months. No dose-response studies have been done with marine species. It is however likely that responses will be similar to those observed for rainbow trout taking into account that lead is generally less bioavailable in marine systems than in freshwater.

ALA-D activity in red blood cells of Atlantic cod (*Gadus morhua*), flounder (*Platichthys flesus*) and dab (*Limanda limanda*) has been used in the Norwegian national monitoring programme for 5 years (1997-2001; Ruus et al. 2003). In that programme, ALA-D was generally found to be inhibited in blood of fish from areas with known urban impact and in one area with a known metal impact (Ruus et al. 2003). In fact, results for ALA-D in that study indicated that there could be lead toxicity in individual cod with hepatic lead levels below the detection limit for routine analyses. Despite its specificity, ALA-D has not been widely used in monitoring. It has been included in OSPAR's list of recommended methods to monitor biological effects of metals (JAMP, 1998). In addition to the above, Holth (2004) investigated ALA-D in Atlantic cod and flounder in southern Norway. He found clear inhibition of the enzyme in both species sampled in the inner Oslofjord compared to individuals collected from the outer Oslofjord. Individuals from both species collected in the outer Oslofjord had values in the range of baseline values (see below).

13.5 Background responses

From the analyses done under Norwegian JAMP, typical values from reference areas are: Atlantic cod 15-21 ng PBG/min/mg protein, plaice 13-21 ng PBG/min/mg protein, flounder 13-21 ng PBG/min/mg protein and dab 10-20 ng PBG/min/mg protein.

13.6 Links

As mentioned above, inhibition of ALA-D will lead to anaemia in birds and humans. Initial studies with very high inhibition of the enzyme in fish did not find such effect in freshwater fish and it is thought that the enzyme is not rate-limiting for heme synthesis in fish (and hence inhibition will not necessarily lead to less heme produced). There is thus no clear link to higher level effects in fish.

13.7 Other factors

Lead is generally found in the environment as the inorganic form, but the alkylated form may be present. Alkyllead does not affect ALA-D strongly and there may thus be a situation where there is little inhibition, but high blood lead levels (Hodson et al. 1984). Data on the influence of metals other than Pb are contradictory (Jackim 1973, Dwyer et al. 1988, Sordyl & Osterland 1990). There is some evidence that very high levels of Cd may inhibit the enzyme in flounder (Johansson-Sjöbeck & Larsson 1978). Zinc may ameliorate inhibition by lead, and appears to affect fish red blood cell ALA-D *in vivo* (Schmitt et al. 1984, Schmitt et al. 1993, Schmitt et al. 2005). The influence has been found to be weak and variable in rainbow trout (Hodson et al. 1984).

13.8 Quality control

There exists no commercially available standard reference material for fish ALA-D. Bovine ALA-D is available commercially and may be used to test the assay. The pH of the assay needs to be established for each fish species (and will differ from the pH optimum of bovine ALA-D). Suitable internal reference materials (blood supernatants from relevant fish species) should be prepared by each laboratory undertaking ALA-D analyses. This material should be aliquotted and stored at -80°C for use with each batch of analyses. Control charts should be prepared during the analyses.

ALA-D was one of the methods included in the EU-funded project BEQUALM (Biological effects quality assurance in marine monitoring programmes; project PL97-3587). The low number of laboratories that participated in the exercise (5) precludes general conclusions on the use of the method. The exercise did show that homogenous test samples for intercalibration purposes can be prepared from red blood cells diluted in buffer.

14. Concluding remarks

A range of biomarkers relevant for monitoring offshore Norway have been reviewed. Histological biomarkers, including lysosomal stability, were not included.

The review shows that a range of methods should be considered for inclusion in the water column monitoring programme offshore (PAH bile metabolites, GST, DNA adducts, the Comet assay, micronucleus formation, AChE, vtg, ALA-D), some need more research (peroxisomal proliferation, MXR) and two have either been found to be not very useful (CYP1A activity) or not very relevant (metallothionein) (Table 31).

Table 31. Overview of methods included and their characteristics; relevant species are gadiids and/or blue mussels; all factors shown to affect the biomarker in question are indicated under “confounding factors”, the most important are shown in bold.

biomarker	datasets available for relevant species	link to population effects	confounding factors	assessment criteria	quality assurance	consider for WCM
PAH bile metabolites	extensive	indicates exposure	recent feeding sex, size/age	yes	QUASIMEME	yes
cytochrome P4501A (CYP1A)	extensive	contaminant-related, but no clear link	sex env temperature season, age	yes	BEQUALM	as protein, not activity
glutathione S-transferase (GST)	limited	contaminant-related, but no clear link	sex, season starvation	no	no	yes
DNA adducts	good	correlative link to cancer and population effects	sex season (mussel)	yes	BEQUALM	yes
Comet assay	limited	possibly correlative link to cancer and population effects	sex? sample handling age, starvation, adaptation	no	no	yes
micronucleus formation (MN)	limited	expected from mammalian studies, but not shown	unknown	no	no	yes
acetyl cholinesterase (AChE)	very limited	contaminant-related, but no clear link	other contaminants?	no	other	yes
peroxisomal proliferation	limited	uncertain	reproductive status sex	no	no	research
multixenobiotic resistance (MXR)	limited	uncertain	unknown	no	no	research
metallothionein (MT)	good	presumably contaminant-related	other contaminants sex season	yes	BEQUALM	no
vitellogenin (vtg)	good	indicates level of xenoestrogens that may have population effects	(only use male or juvenile fish) season	yes	BEQUALM	yes
delta-aminolevulinic acid dehydratase (ALA-D)	good	contaminant-related, but no clear link	other metals?	yes	BEQUALM	if metals

From the range of biomarkers included in this review there are some for which links to population effects is largely unknown or not established (peroxisomal proliferation, MXR,

MT), one that indicates exposure to a specific class of contaminants (PAH metabolites), three for which there is some evidence in the literature for links with putative population effects (CYP1A, GST, ALA-D), three for which links could be expected from studies with other species or life stages (micronucleus formation, Comet assay, vitellogenin) and one for which there is correlative evidence of such links (DNA adducts).

15. References

Chapter 1 (Introduction)

- Eggens M.L., Vethaak A.D., Leaver M.J., Horbach G.J.M.J., Boon J.P., Seinen W. 1996. Differences in CYP1A response between flounder (*Platichthys flesus*) and plaice (*Pleuronectes platessa*) after long-term exposure to harbor dredged spoil in a mesocosm study. *Chemosphere*. 32:1357–1380
- Mayer, F. L., Versteeg, D. J., McKee, M. J., Folmar, L. C., Graney, R. L., McCume, D. C., Rattner, B. A. 1992. Physiological and nonspecific biomarkers. In *Biomarkers. Biochemical, physiological, and histological markers of anthropogenic stress*. Eds: Huggett, R. J., Kimerle, R. A., Mehrle Jr., P. M. & Bergman, H. L., pp. 5-85, Lewis Publishers.
- Myers MS, Johnson LL, Collier TK (2003) Establishing the causal relationship between polycyclic aromatic hydrocarbon (PAH) exposure and hepatic neoplasms and neoplasia-related liver lesions in English sole (*Pleuronectes vetulus*). *Hum Ecol Risk Assess* 9:67-94
- Peakall, D. B. 1994. The role of biomarkers in environmental assessment. 1. Introduction. *Ecotoxicology*. 3:157-160.
- Stegeman, J. J., Brouwer, M., Di Giulio, R. T., Förlin, L., Fowler, B. A., Sanders, B. M., Van Veld, P. A. 1992. Molecular responses to Environmental contamination: Enzyme and protein systems as indicators of chemical exposure and effect. In: *Biomarkers. Biochemical, physiological, and histological markers of anthropogenic stress*. Eds: Huggett, R. J., Kimerle, R. A., Mehrle Jr., P. M. & Bergman H. L., pp. 235-335, Lewis Publishers.
- Vethaak AD, Jol JG, Meijboom A, Eggens ML, Rheinallt T, Wester PW, van de Zande T, Bergman A, Dankers N, Ariese F, Baan RA, Everts JM, Opperhuizen A, Marquenie JM. 1996. Skin and liver diseases induced in flounder (*Platichthys flesus*) after long-term exposure to contaminated sediments in large-scale mesocosms. *Environ Health Perspect*. 104(11):1218-29.

Chapter 2 (PAH-metabolites)

- Ariese, F., Beyer, J., Jonsson, G., Visa, C.P., Krahn, M.M. 2005. Review of analytical methods for determining metabolites of polycyclic aromatic compounds (PACs) in fish bile. *ICES Techniques in Marine Environmental Sciences*, 39: 1-41.
- Ariese, F., Kok, S.J., Verkaik, M., Gooijer, C., Velthorst, N.H., Hofstraat, J.W., 1993. Synchronous fluorescence spectrometry of fish bile: A rapid screening method for the biomonitoring of PAH exposure. *Aquatic Toxicology* 26, 273-286.
- Bechmann, R. K. et al. 2004. SPE 86676: Bioaccumulation, Biomarker Responses, and Effects on Reproduction in Fish Exposed to a Mixture of PAHs (Polycyclic Aromatic Hydrocarbons) and to Dispersed Oil. 7th SPE Int. Conf. on HSE in Oil and Gas Exploration and Production, Calgary, Alberta, Canada (29-31 March 2004), SPE.
- Børseth et al. Rocknes RF - 2004/165.
- Beyer et al. 2003. Biomarkers in Fish and Invertebrates at a Decommissioned Oil and Gas Production site in the North Sea. RF-AM report 2003/018
- Børseth et al. 2004. Analysis report - Norwegian Water Column Survey 2003.
- Collier, T.K. and Varanasi, U. 1991. Hepatic activities of xenobiotic metabolizing enzymes and biliary levels of xenobiotics in English sole (*Parophrys vetulus*) exposed to environmental contaminants. *Archives of Environmental Contamination and Toxicology* 20: 463-473
- Hylland, K. et al. 2004. Water Column Monitoring 2004 - Summary Report, NIVA - Norwegian Institute for Water Research: 50 + appendixes.
- Jonsson, G., 1999. Determination of PAH metabolites in bile by HPLC/Fluorescence. Akvamiljø, Stavanger, Norway, report no. pp. 13.

- Jonsson, G., Beyer, J., Wells, D., Ariese, F. 2003. The application of HPLC-F and GC-MS to the analysis of selected hydroxy polycyclic hydrocarbons in two certified fish bile reference materials. *Journal of Environmental Monitoring*, 2003. 5: p. 513-520.
- Klungstøyr et al. 2003. Contamination of fish in the North Sea by the offshore oil and gas industry, Summary report to NFR, NFR project nr. 152231/720.
- Lin, E.L.C., Cormier, S.M., Torsella, J.A., 1996. Fish biliary polycyclic aromatic hydrocarbon metabolites estimated by fixed-wavelength fluorescence: Comparison with HPLC-fluorescent detection. *Ecotoxicology and environmental safety* 35, 16-23.
- Ruus, A., Hylland, K., Green, N. 2003. Joint Assessment and Monitoring Programme (JAMP). Biological Effects Methods, Norwegian Monitoring 1997-2001. SFT Report no. 869/03. TA-1948/2003, 139 p.
- Skadsheim, A, et al. 2004. Cod Exposed to crude oils: Uptake, metabolism and biomarker effects. Report AM-2004/005
- Sundt, R.S. 2004. Comparative cod exposure 2004, experiment report, Report AM-2004/032.
- Sundt, R. C. 2004. BIOSEA Biomarker field levels and Screening test with Arctic species - Version 1. Stavanger, Norway, Akvamiljø.
- Sundt, R.C & Bechmann, R.K. 2004. Biomarker background levels in marine species- Part 1. Report AM 2004/004.
- Aas, E., T. Baussant, et al. (2000). "PAH metabolites in bile, cytochrome P4501A and DNA adducts as environmental risk parameters for chronic oil exposure: a laboratory experiment with Atlantic cod." *Aquatic Toxicology* 51(2): 241-258.
- Aas, E., Beyer, J., Goksøyr, A. 2000. Fixed wavelength fluorescence (FF) of bile as a monitoring tool for polyaromatic hydrocarbon exposure in fish: An evaluation of compound specificity, inner filter effect and signal interpretation. *Biomarkers* 5, 9-23.
- Aas E. & Børseth, J. F., 2002. Baseline levels of biomarker responses in Atlantic cod and capelin in the Barents Sea, 2001 - 2002. Report AM 2002/014.
- Aas, E et al. 2002. Biomarkers for genotoxic effects in marine organisms project report 2001-2002. AM 2002/13.
- Aas et al. PAH metabolites and metals in bile from caged Atlantic cod (*Gadus morhua*) and feral fish in the North Sea used in environmental monitoring, SETAC special publication (in press).

Chapter 3 (Cytochrome P4501A activity – EROD)

- Bernhoft, A., Hektoen, H., Skaare, J. U., and Ingebrigtsen, K. (1994). Tissue Distribution And Effects On Hepatic Xenobiotic-Metabolizing Enzymes Of 2,3,3',4,4'-Pentachlorobiphenyl (Pcb-105) In Cod (*Gadus-Morhua*) And Rainbow-Trout (*Oncorhynchus-Mykiss*). *Environmental Pollution* 85, 351-359.
- Beyer, J., Sandvik, M., Hylland, K., Fjeld, E., Egaas, E., Aas, E., Skare, J. U., and Goksoyr, A. (1996). Contaminant accumulation and biomarker responses in flounder (*Platichthys flesus* L) and Atlantic cod (*Gadus morhua* L) exposed by caging to polluted sediments in Sorfjorden, Norway. *Aquatic Toxicology* 36, 75-98.
- Burke, M. D., and Mayer, R. T. (1974). Ethoxyresorufin: Direct fluorimetric assay of a microsomal O-dealkylation which is preferentially inducible by 3-methylchoalanthrene. *Drug metabolism and Disposition* 2, 583-588.
- Förlin L, Haux C (1990) Sex differences in hepatic cytochrome P-450 monooxygenase activities in rainbow trout during an annual reproductive cycle. *J. Endocrinol.* 124:207-213
- Forlin, L., Lemaire, P., and Livingstone, D. R. (1995). Comparative-Studies of Hepatic Xenobiotic-Metabolizing and Antioxidant Enzymes in Different Fish Species. *Marine Environmental Research* 39, 201-204.

- George, S. G., Christiansen, J. S., Killie, B., and Wright, J. (1995). Dietary Crude-Oil Exposure During Sexual-Maturation Induces Hepatic Mixed-Function Oxygenase (Cyp1a) Activity At Very-Low Environmental Temperatures In Polar Cod *Boreogadus-Saida*. *Marine Ecology-Progress Series* 122, 307-312.
- George, S. G., Wright, J., and Conroy, J. (1995). Temporal Studies Of The Impact Of The Braer Oilspill On Inshore Feral Fish From Shetland, Scotland. *Archives Of Environmental Contamination And Toxicology* 29, 530-534.
- Goksoyr, A., Andersson, T., Buhler, D. R., Stegeman, J. J., Williams, D. E., and Forlin, L. (1991). Immunochemical Cross-Reactivity Of Beta-Naphthoflavone-Inducible Cytochrome P450 (P450ia) In Liver-Microsomes From Different Fish Species And Rat. *Fish Physiology And Biochemistry* 9, 1-13.
- Goksoyr, A., Andersson, T., Hansson, T., Klungsoyr, J., Zhang, Y., and Forlin, L. (1987). Species Characteristics Of The Hepatic Xenobiotic And Steroid Biotransformation Systems Of 2 Teleost Fish, Atlantic Cod (*Gadus-Morhua*) And Rainbow-Trout (*Salmo-Gairdneri*). *Toxicology And Applied Pharmacology* 89, 347-360.
- Goksoyr, A., Beyer, J., Husoy, A. M., Larsen, H. E., Westrheim, K., Wilhelmsen, S., and Klungsoyr, J. (1994). Accumulation And Effects Of Aromatic And Chlorinated Hydrocarbons In Juvenile Atlantic Cod (*Gadus-Morhua*) Caged In A Polluted Fjord (Sorfjorden, Norway). *Aquatic Toxicology* 29, 21-35.
- Hasselberg, L., Meier, S., Svardal, A., Hegelund, T., and Celander, M. C. (2004). Effects of alkylphenols on CYP1A and CYP3A expression in first spawning Atlantic cod (*Gadus morhua*). *Aquatic Toxicology* 67, 303-313.
- Hektoen, H., Bernhoft, A., Ingebrigtsen, K., Skaare, J. U., and Goksoyr, A. (1994). Response Of Hepatic Xenobiotic-Metabolizing Enzymes In Rainbow-Trout (*Oncorhynchus-Mykiss*) And Cod (*Gadus-Morhua*) To 2,3,7,8-Tetrachlorodibenzo-P-Dioxin (2,3,7,8-Tcdd). *Aquatic Toxicology* 28, 97-106.
- Hylland K, Sandvik M, Skåre JU, Beyer J, Egaas E, Goksøy A (1996) Biomarkers in flounder (*Platichthys flesus*): an evaluation of their use in pollution monitoring. *Mar. Environ. Res.* 42:223-227.
- Kloppersams, P. J., and Benton, E. (1994). Exposure Of Fish To Biologically Treated Bleached-Kraft Effluent.2. Induction Of Hepatic Cytochrome-P4501a In Mountain Whitefish (*Prosopium-Williamsoni*) And Other Species. *Environmental Toxicology And Chemistry* 13, 1483-1496.
- Lange U, Goksoyr A, Siebers D, Karbe L (1999) Cytochrome P450 1A-dependent enzyme activities in the liver of dab (*Limanda limanda*): kinetics, seasonal changes and detection limits. *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology* 123:361-371
- Larsen, H.E., Celander, M. and Goksøy, A. (1992) The cytochrome P450 system of Atlantic salmon (*Salmo salar*): II. Variations in hepatic catalytic activities and isozyme patterns during an annual reproductive cycle. *Fish Physiol. Biochem.* 10, 291-301.
- Lindesjoo, E., Husoy, A. M., Petterson, I., and Forlin, L. (1996). Histopathological and immunohistochemical studies in roundnose grenadier (*Coryphaenoides rupestris*) in the Skagerrak, North Sea. *Marine Environmental Research* 42, 229-233.
- Livingstone, D. R., Lemaire, P., Matthews, A., Peters, L. D., Porte, C., Fitzpatrick, P. J., Forlin, L., Nasci, C., Fossato, V., Wootton, N., and Goldfarb, P. (1995). Assessment Of The Impact Of Organic Pollutants On Goby (*Zosterisessor Ophiocephalus*) And Mussel (*Mytilus-Galloprovincialis*) From The Venice Lagoon, Italy - Biochemical-Studies. *Marine Environmental Research* 39, 235-240.
- McDonald, S. J., Willett, K. L., Thomsen, J., Beatty, K. B., Connor, K., Narasimhan, T. R., Erickson, C. M., and Safe, S. H. (1996). Sublethal detoxification responses to contaminant exposure associated with offshore production platforms. *Canadian Journal of Fisheries and Aquatic Sciences* 53, 2606-2617.
- Michel, X., Mora, P., Garrigues, P., Budzinski, H., Raoux, C., and Narbonne, J. F. (2001). Cytochrome P450 dependent activities in mussel and fish from coastal marine environment: Field studies on the French coast of the Mediterranean sea. *Polycyclic Aromatic Compounds* 18, 307-324.
- Myers MS, Johnson LL, Collier TK (2003) Establishing the causal relationship between polycyclic aromatic hydrocarbon (PAH) exposure and hepatic neoplasms and neoplasia-related liver lesions in English sole (*Pleuronectes vetulus*). *Hum Ecol Risk Assess* 9:67-94

- Nissen-Lie TR (1997) Årstidsvariasjon i spormetallmetabolismen til skrubbe, *Platichthys flesus* L., fra Hvalerområdet. MSc Thesis, University of Oslo. [in Norwegian]
- Payne, J. F. (1976). Field evaluation of benzopyrene hydroxylase induction as a monitor for marine petroleum pollution. *Science* 191, 945-946.
- Peters, L. D., Nasci, C., and Livingstone, D. R. (1998). Variation in levels of cytochrome P4501A, 2B, 2E, 3A and 4A-immunopositive proteins in digestive gland of indigenous and transplanted mussel *Mytilus galloprovincialis* in Venice Lagoon, Italy. *Marine Environmental Research* 46, 295-299.
- Ruus, A., Hylland, K., W., G. N., and (2003). Joint Assessment and Monitoring Programme (JAMP). Biological Effects Methods, Norwegian Monitoring 1997-2001. Norwegian State Pollution Monitoring Programme Report no. 869/03. Ta-no. 1948/2003.
- Ruus, A., Sandvik, M., Ugland, K. I., and Skaare, J. U. (2002). Factors influencing activities of biotransformation enzymes, concentrations and compositional patterns of organochlorine contaminants in members of a marine food web. *Aquatic Toxicology* 61, 73-87.
- Shaw, J. P., Large, A. T., Chipman, J. K., Livingstone, D. R., and Peters, L. D. (2000). Seasonal Variation in mussel *Mytilus edulis* digestive gland cytochrome P4501A- and 2E-immunoidentified protein levels and DNA strand breaks (Comet assay). *Marine Environmental Research* 50, 405-409.
- Sleiderink H, Beyer J, Everaarts JM, Boon JP (1995) Influence of temperature on cytochrome P450 1A in dab (*Limanda limanda*) from the southern North Sea: Results from field surveys and a laboratory study. *Mar. Environ. Res.* 39:67-71.
- Sole, M., Peters, L. D., Magnusson, K., Sjolín, A., Granmo, A., and Livingstone, D. R. (1998). Responses of the cytochrome P450-dependent monooxygenase and other protective enzyme systems in digestive gland of transplanted common mussel (*Mytilus edulis* L.) to organic contaminants in the Skagerrak and Kattegat (North Sea). *Biomarkers* 3, 49-62.
- Stagg, R., and McIntosh, A. 1998. Biological effects of contaminants: Determination of CYP1A-dependent mono-oxygenase activity in dab by fluorimetric measurement of EROD activity. *ICES Techniques in Marine Environmental Science*, 23, 16 pp.
- Whyte, J. J., Jung, R. E., Schmitt, C. J., and Tillitt, D. E. (2000). Ethoxyresorufin-O-deethylase (EROD) activity in fish as a biomarker of chemical exposure. *Critical Reviews in Toxicology* 30, 347-570.
- Williams, R. T. (1974). Inter-species variations in the metabolism of xenobiotics. *Biochem. Soc. T.* 2, 359-377.
- Williams, T. G., Lockhart, W. L., Metner, D. A., and Harbicht, S. (1997). Baseline studies in the Slave River, NWT, 1990-1994.3. MFO enzyme activity in fish. *Science of the Total Environment* 197, 87-109.
- Aas, E., Baussant, T., Balk, L., Liewenborg, B., and Andersen, O. K. (2000). PAH metabolites in bile, cytochrome P4501A and DNA adducts as environmental risk parameters for chronic oil exposure: a laboratory experiment with Atlantic cod. *Aquatic Toxicology* 51, 241-258.

Chapter 4 (DNA damage)

- Accomando R, Viarengo A, Bordone R, Taningher M, Canesi L, Orunesu M (1991) A Rapid Method for Detecting DNA Strand Breaks in *Mytilus-Galloprovincialis* Lam Induced by Genotoxic Xenobiotic Chemicals. *Int J Biochem* 23:227-229
- Akcha F, Hubert FV, Pfohl-Leszkwicz A (2003) Potential value of the comet assay and DNA adduct measurement in dab (*Limanda limanda*) for assessment of in situ exposure to genotoxic compounds. *Mutat Res* 534:21-32
- Akcha F, Izuel C, Venier P, Budzinski H, Burgeot T, Narbonne JF (2000) Enzymatic biomarker measurement and study of DNA adduct formation in benzo[a]pyrene-contaminated mussels, *Mytilus galloprovincialis*. *Aquat Toxicol* 49:269-287

- Akcha F, Leday G, Pfohl-Leszkowicz A (2004) Measurement of DNA adducts and strand breaks in dab (*Limanda limanda*) collected in the field: effects of biotic (age, sex) and abiotic (sampling site and period) factors on the extent of DNA damage. *Mutat Res* 552:197-207
- Anulacion BF, Myers MS, Willis ML, Collier TK (1998) Quantitation of CYP1A expression in two flatfish species showing different prevalences of contaminant-induced hepatic disease. *Mar Environ Res* 46:7-11
- Armstrong TN, Reimschuessel R, Bradley BP (2002) DNA damage, histological changes and DNA repair in larval Japanese medaka (*Oryzias latipes*) exposed to ultraviolet-B radiation. *Aquat Toxicol* 58:1-14
- Balk, L, Liewenborg, B, Larsen, BK, Aas, E, Hylland, K, Sanni, S. Large hydrophobic adducts and strand breaks analyzed in hepatic DNA from Atlantic cod (*Gadus morhua*) caged at the Statfjord oil field. In: Biological effects of contaminants in pelagic ecosystems. Hylland, K, Vethaak, AD, Lang, T (Eds), in press.
- Canova S, Degan P, Peters LD, Livingstone DR, Voltan R, Venier P (1998) Tissue dose, DNA adducts, oxidative DNA damage and CYP1A-immunopositive proteins in mussels exposed to waterborne benzo[a]pyrene. *Mutat Res* 399:17-30
- Carpenter HM, Zhang Q, Elzahr C, Selivonchick DP, Brock DE, Curtis LR (1995) In-Vitro and in-Vivo Temperature Modulation of Hepatic-Metabolism and DNA Adduction of Aflatoxin-B1 in Rainbow-Trout. *J Biochem Toxicol* 10:1-10
- Ching EWK, Siu WHL, Lam PKS, Xu LH, Zhang YY, Richardson BJ, Wu RSS (2001) DNA adduct formation and DNA strand breaks in green-lipped mussels (*Perna viridis*) exposed to benzo[a]pyrene: Dose- and time-dependent relationships. *Mar Pollut Bull* 42:603-610
- Collins AR (2005) Assays for oxidative stress and antioxidant status: applications to research into the biological effectiveness of polyphenols. *Am J Clin Nutr* 81:261S-267S
- David WM, Mitchell DL, Walter RB (2004) DNA repair in hybrid fish of the genus *Xiphophorus*. *Comparative Biochemistry and Physiology* 138C:301-309
- Dolcetti L, Dalla Zuanna L, Venier P (2002) DNA adducts in mussels and fish exposed to bulky genotoxic compounds. *Mar Environ Res* 54:481-486
- Dubois M, Pfohl-Leszkowicz A, Grosse Y, Kremers P (1995) DNA adducts and P450 induction in human, rat and avian liver cells after exposure to polychlorobiphenyls. *Mutat Res* 345:181-190
- Ericson G, Balk L (2000) DNA adduct formation in northern pike (*Esox lucius*) exposed to a mixture of benzo[a]pyrene, benzo[k]fluoranthene and 7H-dibenzo[c,g]carbazole: time-course and dose-response studies. *Mutat Res* 454:11-20
- Ericson G, Larsson A (2000) DNA adducts in perch (*Perca fluviatilis*) living in coastal water polluted with bleached pulp mill effluents. *Ecotox Environ Safe* 46:167-173
- Ericson G, Liewenborg B, Lindesjoo E, Naf C, Balk L (1999) DNA adducts in perch (*Perca fluviatilis*) from a creosote contaminated site in the River Angermanalven, Sweden. *Aquat Toxicol* 45:181-193
- Ericson G, Lindesjoo E, Balk L (1998) DNA adducts and histopathological lesions in perch (*Perca fluviatilis*) and northern pike (*Esox lucius*) along a polycyclic aromatic hydrocarbon gradient on the Swedish coastline of the Baltic Sea. *Can J Fish Aquat Sci* 55:815-824
- Ericson G, Noaksson E, Balk L (1999) DNA adduct formation and persistence in liver and extrahepatic tissues of northern pike (*Esox lucius*) following oral exposure to benzo[a]pyrene, benzo[k]fluoranthene and 7H-dibenzo[c,g]carbazole. *Mutat Res* 427:135-145
- Ericson G, Skarphedinsdottir H, Dalla Zuanna L, Svavarsson J (2002) DNA adducts as indicators of genotoxic exposure in indigenous and transplanted mussels, *Mytilus edulis* L. from Icelandic coastal sites. *Mutat Res* 516:91-99
- Everaarts JM (1995) DNA integrity as a biomarker of marine pollution: Strand breaks in seastar (*Asterias rubens*) and Dab (*Limanda limanda*). *Mar Pollut Bull* 31:431-438
- Feng T, Guo XQ, Zheng WY, Yuan DX (2003) Detection of DNA strand breaks in the liver of *Boleophthalmus pectinirostris* treated with benzo(a)pyrene. *B Environ Contam Tox* 71:263-269

- French BL, Reichert WL, Hom T, Nishimoto M, Sanborn HR, Stein JE (1996) Accumulation and dose-response of hepatic DNA adducts in English sole (*Pleuronectes vetulus*) exposed to a gradient of contaminated sediments. *Aquat Toxicol* 36:1-16
- Hsu T, Deng FY (1996) Studies on the susceptibility of various organs of zebrafish (*Brachydanio rerio*) to benzo(a)pyrene-induced DNA adduct formation. *Chemosphere* 33:1975-1980
- Hylland, K., Beyer, J., Berntssen, M., Klungsoyr, J., Lang, T., Balk, L. 2006. May persistent organic pollutants affect fish populations in the North Sea? *J. Toxicol. Environ. Hlth. Part A*, 69:125-138.
- Liepelt A, Karbe L, Westendorf J (1995) Induction of DNA Strand Breaks in Rainbow-Trout *Oncorhynchus-Mykiss* under Hypoxic and Hyperoxic Conditions. *Aquat Toxicol* 33:177-181
- Lindesjoo E, Adolfsson-Erici M, Ericson G, Forlin L (2002) Biomarker responses and resin acids in fish chronically exposed to effluents from a total chlorine-free pulp mill during regular production. *Ecotox Environ Safe* 53:238-247
- Lyons BP, Harvey JS, Parry JM (1997) An initial assessment of the genotoxic impact of the Sea Empress oil spill by the measurement of DNA adduct levels in the intertidal teleost *Lipophrys pholis*. *Mutat Res* 390:263-268
- Lyons BP, Stentiford GD, Green M, Bignell J, Bateman K, Feist SW, Goodsir F, Reynolds WJ, Thain JE (2004) DNA adduct analysis and histopathological biomarkers in European flounder (*Platichthys flesus*) sampled from UK estuaries. *Mutat Res* 552:177-186
- Lyons BP, Stewart C, Kirby MF (1999) The detection of biomarkers of genotoxin exposure in the European flounder (*Platichthys flesus*) collected from the River Tyne Estuary. *Mutat Res* 446:111-119
- Lyons BP, Stewart C, Kirby MF (2000) 32P-postlabelling analysis of DNA adducts and EROD induction as biomarkers of genotoxin exposure in dab (*Limanda limanda*) from British coastal waters. *Mar Environ Res* 50:575-579
- Malmström, C.M., Miettinen, S., Bylund, G. 2000. DNA adducts in liver and leukocytes of flounder (*Platichthys flesus*) experimentally exposed to benzo[a]pyrene. *Aquatic Toxicology* 48: 177–184
- Myers MS, Johnson LL, Collier TK (2003) Establishing the causal relationship between polycyclic aromatic hydrocarbon (PAH) exposure and hepatic neoplasms and neoplasia-related liver lesions in English sole (*Pleuronectes vetulus*). *Hum Ecol Risk Assess* 9:67-94
- Nacci D, Nelson S, Nelson W, Jackim E (1992) Application of the DNA Alkaline Unwinding Assay to Detect DNA Strand Breaks in Marine Bivalves. *Mar Environ Res* 33:83-100
- Næs K, Hylland K, Oug E, Förlin L, Ericson G (1999) Accumulation and effects of aluminum smelter-generated polycyclic aromatic hydrocarbons on soft-bottom invertebrates and fish. *Environ Toxicol Chem* 18:2205-2216
- Pinkney AE, Harshbarger JC, May EB, Reichert WL (2004) Tumor prevalence and biomarkers of exposure and response in brown bullhead (*Ameiurus nebulosus*) from the Anacostia River, Washington, DC and Tuckahoe River, Maryland, USA. *Environ Toxicol Chem* 23:638-647
- Pisoni M, Cogotzi L, Frigeri A, Corsi I, Bonacci S, Iacocca A, Lancini L, Mastrototaro F, Focardi S, Svelto M (2004) DNA adducts, benzo(a)pyrene monooxygenase activity, and lysosomal membrane stability in *Mytilus galloprovincialis* from different areas in Taranto coastal waters (Italy). *Environ Res* 96:163-175
- Ploch SA, King LC, Di Giulio RT (1998) Comparative time-course of benzo[a]pyrene-DNA adduct formation, and its relationship to Cyp1A activity in two species of catfish. *Mar Environ Res* 46:345-349
- Ploch SA, King LC, Kohan MJ, Di Giulio RT (1998) Comparative in vitro and in vivo benzo[a]pyrene-DNA adduct formation and its relationship to CYP1A activity in two species of ictalurid catfish. *Toxicol Appl Pharm* 149:90-98
- Reynolds W.J.¹; Feist S.W.; Jones G.J.; Lyons B.P.; Sheahan D.A.; Stentiford G.D. 2003. Comparison of biomarker and pathological responses in flounder (*Platichthys flesus* L.) induced by ingested polycyclic aromatic hydrocarbon (PAH) contamination. *Chemosphere*, 52: 1135-1145.

- Rose WL, French BL, Reichert WL, Faisal M (2000) DNA adducts in hematopoietic tissues and blood of the mummichog (*Fundulus heteroclitus*) from a creosote-contaminated site in the Elizabeth River, Virginia. *Mar Environ Res* 50:581-589
- Rose WL, French BL, Reichert WL, Faisal M (2001) Persistence of benzo[a]pyrene--DNA adducts in hematopoietic tissues and blood of the mummichog, *Fundulus heteroclitus*. *Aquat Toxicol* 52:319-328
- Shugart LR (1988) Quantitation of Chemically-Induced Damage to DNA of Aquatic Organisms by Alkaline Unwinding Assay. *Aquat Toxicol* 13:43-52
- Siu WHL, Hung CLH, Wong HL, Richardson BJ, Lam PKS (2003) Exposure and time dependent DNA strand breakage in hepatopancreas of green-lipped mussels (*Perna viridis*) exposed to Aroclor 1254, and mixtures of B[a]P and Aroclor 1254. *Mar Pollut Bull* 46:1285-1293
- Skarphedinsdottir H, Ericson G, Dalla Zuanna L, Gilek M (2003) Tissue differences, dose-response relationship and persistence of DNA adducts in blue mussels (*Mytilus edulis* L.) exposed to benzo[a]pyrene. *Aquat Toxicol* 62:165-177
- Skarphedinsdottir H, Ericson G, Halldorsson HP, Svavarsson J (2005) Seasonal and intertidal impact on DNA adduct levels in gills of blue mussels (*Mytilus edulis* L.). *Environ Pollut* 136:1-9
- Stephensen E. L., Svavarsson J.; Sturve J.; Ericson G.; Adolfsson-Erici M.; Forlin L. 2000. Biochemical indicators of pollution exposure in shorthorn sculpin (*Myoxocephalus scorpius*), caught in four harbours on the southwest coast of Iceland. *Aquatic Toxicology*, 48: 431-442
- Venier P, Zampieron C (2005) Evidence of genetic damage in grass gobies and mussels from the Venice lagoon. *Environ Int* 31:1053-1064
- Walter RB, Sung HM, Obermoeller RD, Mitchell DL, Intano GW, Walter CA (2001) Relative base excision repair in Xiphophorus fish tissue extracts. *Mar Biotechnol* 3:S50-S60
- Willett KL, Lienesch LA, Di Giulio RT (2001) No detectable DNA excision repair in UV-exposed hepatocytes from two catfish species. *Comparative Biochemistry and Physiology* 128C:349-358
- Willett K, Steinberg M, Thomsen J, Narasimhan TR, Safe S, McDonald S, Beatty K, Kennicutt MC (1995) Exposure of Killifish to Benzo[a]Pyrene - Comparative Metabolism, DNA Adduct Formation and Aryl-Hydrocarbon (Ah) Receptor Agonist Activities. *Comparative Biochemistry and Physiology* 112B:93-103
- Aas E, Baussant T, Balk L, Liewenborg B, Andersen OK (2000) PAH metabolites in bile, cytochrome P4501A and DNA adducts as environmental risk parameters for chronic oil exposure: a laboratory experiment with Atlantic cod. *Aquat Toxicol* 51:241-258
- Aas E, Liewenborg B, Grosvik BE, Camus L, Jonsson G, Borseth JF, Balk L (2003) DNA adduct levels in fish from pristine areas are not detectable or low when analysed using the nuclease P1 version of the P-32-postlabelling technique. *Biomarkers* 8:445-460

Chapter 5 (Glutathione S-transferase)

- Akcha, F., Izuel, C., Venier, P., Budzinski, H., Burgeot, T. and Narbonne, J. (2000). Enzymatic biomarker measurement and study of DNA adduct formation in benzo. *Aquat Toxicol* 49, 269-287.
- Angelucci, S., Sacchetta, P., Moio, P., Melino, S., Petruzzelli, R., Gervasi, P. and Di Ilio, C. (2000). Purification and characterization of glutathione transferases from the sea bass (*Dicentrarchus labrax*) liver. *Arch Biochem Biophys* 373, 435-441.
- Baturo, W. and Lagadic, L. (1996). Benzo[a]pyrene hydroxylase and glutathione S-transferase activities as biomarkers in *Lymnaea palustris* (Mollusca, Gastropoda) exposed to atrazine and hexachlorobenzene in freshwater mesocosms. *Environ Toxicol Chem* 15, 771-781.
- Blom, S., Andersson, T. B. and Forlin, L. (2000). Effects of food deprivation and handling stress on head kidney 17 α -hydroxyprogesterone 21-hydroxylase activity, plasma cortisol and the activities of liver detoxification enzymes in rainbow trout. *Aquatic Toxicol* 48, 265-274.

- Bocquene, G., Chantreau, S., Clerendeau, C., Beausir, E., Menard, D., Raffin, B., Minier, C., Burgeot, T., Leszkowicz, A. P. and Narbonne, J. F. (2004). Biological effects of the "Erika" oil spill on the common mussel (*Mytilus edulis*). *Aquat Living Res* 17, 309-316.
- Børseth, J. F. and Tollefsen, K. E. (2004). Water Column Monitoring 2003 - Final Report. Report RF no. 2004/039, Rogaland Research, Stavanger.
- Chambras, C., Marionnet, D., Taysse, L., Deschaux, P., Moreau, J. and Bosgiraud, C. (1999). Xenobiotic-metabolizing enzymes in carp (*Cyprinus carpio*) liver, spleen, and head kidney following experimental *Listeria monocytogenes* infection. *J Toxicol Environ Health A* 56, 205-219.
- Cheung, C. C., Zheng, G. J., Li, A. M., Richardson, B. J. and Lam, P. K. (2001). Relationships between tissue concentrations of polycyclic aromatic hydrocarbons and antioxidative responses of marine mussels, *Perna viridis*. *Aquat Toxicol* 52, 189-203.
- Collier, T. K., Singh, S. V., Awasthi, Y. C. and Varanasi, U. (1992). Hepatic xenobiotic metabolizing enzymes in two species of benthic fish showing different prevalences of contaminant-associated liver neoplasms. *Toxicol Appl Pharmacol* 113, 319-324.
- Dautremepuits, C., Betoulle, S. and Vernet, G. (2002). Antioxidant response modulated by copper in healthy or parasitized carp (*Cyprinus carpio* L.) by *Ptychobothrium* sp. (Cestoda). *Biochim Biophys Acta* 1573, 4-8.
- DeKoven, D. L., Nunez, J. M., Lester, S. M., Conklin, D. E., Marty, G. D., Parker, L. M. and Hinton, D. E. (1992). A purified diet for medaka (*Oryzias latipes*): refining a fish model for toxicological research. *Lab Anim Sci* 42, 180-189.
- Faverney, C. R., Devaux, A., Lafaurie, M., Girard, J. P. and Rahmani, R. (2001). Toxic effects of wastewaters collected at upstream and downstream sites of a purification station in cultures of rainbow trout hepatocytes. *Arch Environ Con Tox* 41, 129-141.
- Fitzpatrick, P. J. and Sheehan, D. (1993). Separation of multiple forms of glutathione S-transferase from the blue mussel, *Mytilus edulis*. *Xenobiotica* 23, 851-861.
- Förlin, L., Lemaire, P. and Livingstone, D. R. (1995). Comparative-studies of hepatic xenobiotic-metabolizing and antioxidant enzymes in different fish species. *Mar Environ Res* 39, 201-204.
- Gallagher, E. P., Stapleton, P. L., Slone, D. H., Schlenk, D. and Eaton, D. L. (1996). Channel catfish glutathione S-transferase isoenzyme activity toward (+/-)-anti-benzo[a]pyrene-trans-7,8-dihydrodiol-9,10-epoxide. *Aquat Toxicol* 34, 135-150.
- Gowland, B. T., Moffat, C. F., Stagg, R. M., Houlihan, D. F. and Davies, I. M. (2002). Cypermethrin induces glutathione S-transferase activity in the shore crab, *Carcinus maenas*. *Mar Environ Res* 54, 169-177.
- Gowland, B. T. G., McIntosh, A. D., Davies, I. M., Moffat, C. F. and Webster, L. (2002). Implications from a field study regarding the relationship between polycyclic aromatic hydrocarbons and glutathione S-transferase activity in mussels. *Mar Environ Res* 54, 231-235.
- Gravato, C., Oliveira, M. and Santos, M. A. (2004). Genotoxic effects and oxidative stress responses induced by retene in marine mussels (*Mytilus galloprovincialis*). *Fres Environ Bull* 13, 795-800.
- Gravato, C. and Santos, M. A. (2003). Genotoxicity biomarkers' association with B(a)P biotransformation in *Dicentrarchus labrax* L. *Ecotoxicol Environ Saf* 55, 352-358.
- Gubbins, M. J., Eddy, F. B., Gallacher, S. and Stagg, R. M. (2000). Paralytic shellfish poisoning toxins induce xenobiotic metabolising enzymes in Atlantic salmon (*Salmo salar*). *Mar Environ Res* 50, 479-483.
- Guderley, H., Lapointe, D., Bedard, M. and Dutil, J. D. (2003). Metabolic priorities during starvation: enzyme sparing in liver and white muscle of Atlantic cod, *Gadus morhua* L. *Comp Biochem Physiol A* 135, 347-356.
- Habig WH, Pabst MJ, Jakoby WB (1974) Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *JBiolChem* 249:7130-7139.

- Hasselberg, L., Meier, S. and Svardal, A. (2004). Effects of alkylphenols on redox status in first spawning Atlantic cod (*Gadus morhua*). *Aquat Toxicol* 69, 95-105.
- Huuskonen, S., LindstromSeppa, P., Koponen, K. and Roy, S. (1996). Effects of non-ortho-substituted polychlorinated biphenyls (Congeners 77 and 126) on cytochrome P4501A and conjugation activities in rainbow trout (*Oncorhynchus mykiss*). *Comp Biochem Physiol C* 113, 205-213.
- Hylland, K., Marigomez, I., Ruus, A., Balk, L., Sundt, R. C., Abrahamsson, A., Feist, S. and Baršienė, J. (2005). Water Column Monitoring 2004 - Summary Report. REPORT SNO 4993-2005, Norwegian Institute for Water Research (NIVA).
- Jee, J. H. and Kang, J. C. (2005). Biochemical changes as enzymatic defense system of phenanthrene exposure in olive flounder, *Paralichthys olivaceus*. *Physiol Res* 54, 585-592.
- Kennedy, C. J. and Farrell, P. F. (In press). Ion homeostasis and interrenal stress responses in juvenile Pacific herring, *Clupea pallasii*, exposed to the water-soluble fraction of crude oil. *J Exp Mar Biol Ecol*.
- Kennish, J. M., Bolinger, R. A., Chambers, K. A. and Russell, M. L. (1992). Xenobiotic Metabolizing Enzyme-Activity in Sockeye-Salmon (*Oncorhynchus nerka*) during Spawning Migration. *Mar Environ Res* 34, 293-298.
- Kirby, G. M., Stalker, M. J., Gordon, S., Quinn, B. A., van Schooten, F. J. and Hayes, M. A. (1995). Influences of chronic cholangiohepatitis and cholestasis on hepatic metabolism of benzo[a]pyrene in white suckers (*Catostomus commersoni*) from industrially polluted areas of Lake Ontario. *Carcinogenesis* 16, 2923-2929.
- Koehler, A., Alpermann, T., Lauritzen, B. and Van Noorden, C. J. F. (2004). Clonal xenobiotic resistance during pollution-induced toxic injury and hepatocellular carcinogenesis in liver of female flounder (*Platichthys flesus* L.). *Acta Histochemica* 106, 155-170.
- Köhler, A., Lauritzen, B., Bahns, S., George, S. G., Forlin, L. and Van Noorden, C. J. F. (1998). Clonal adaptation of cancer cells in flatfish liver to environmental contamination by changes in expression of P-gp related MXR, CYP450, GST-A and G6PDH activity. *Mar Environ Res* 46, 191-195.
- Lindstrom-Seppa, P. and Pesonen, M. (1986). Biotransformation enzymes in fish as tools for biomonitoring aquatic environment. *Acta Biol Hung* 37, 85-92.
- Manduzio, H., Monsinjon, T., Galap, C., Leboulenger, F. and Rocher, B. (2004). Seasonal variations in antioxidant defences in blue mussels *Mytilus edulis* collected from a polluted area: major contributions in gills of an inducible isoform of Cu/Zn-superoxide dismutase and of glutathione S-transferase. *Aquatic Toxicol* 70, 83-93.
- Martinez-Lara, E., George, S. G., Lopez-Barea, J. and Barcena, J. A. (1997). Purification and characterization of multiple glutathione transferase isoenzymes from grey mullet liver. *Cell Mol Life Sci* 53, 759-768.
- Morrow, M. D., Higgs, D. and Kennedy, C. J. (2004). The effects of diet composition and ration on biotransformation enzymes and stress parameters in rainbow trout, *Oncorhynchus mykiss*. *Comp Biochem Physiol C* 137, 143-154.
- Napierska, D. and Podolska, M. (2005). Biomarkers of contaminant exposure: results of a field study with flounder (*Platichthys flesus*) from the southern Baltic Sea. *Mar Pollut Bull* 50, 758-767.
- Novoa-Valinas, M. C., Perez-Lopez, M. and Melgar, M. J. (2002). Comparative study of the purification and characterization of the cytosolic glutathione S-transferases from two salmonid species: Atlantic salmon (*Salmo salar*) and brown trout (*Salmo trutta*). *Comp Biochem Physiol C* 131, 207-213.
- Padros, J., Pelletier, E. and Ribeiro, C. O. (2003). Metabolic interactions between low doses of benzo[a]pyrene and tributyltin in arctic charr (*Salvelinus alpinus*): a long-term in vivo study. *Toxicol Appl Pharmacol* 192, 45-55.
- Perez-Lopez, M., Novoa-Valinas, M. C. and Melgar-Riol, M. J. (2002). Glutathione S-transferase cytosolic isoforms as biomarkers of polychlorinated biphenyl (Arochlor-1254) experimental contamination in rainbow trout. *Toxicol Lett* 136, 97-106.

- Perez, E., Blasco, J. and Sole, M. (2004). Biomarker responses to pollution in two invertebrate species: *Scrobicularia plana* and *Nereis diversicolor* from the Cadiz bay (SW Spain). *Mar Environ Res* 58, 275-279.
- Risso-de Faverney, C., Devaux, A., Lafaurie, M., Girard, J. P. and Rahmani, R. (2001). Toxic effects of wastewaters collected at upstream and downstream sites of a purification station in cultures of rainbow trout hepatocytes. *Arch Environ Contam Toxicol* 41, 129-141.
- Ruus, A., Sandvik, M., Ugland, K. I. and Skaare, J. U. (2002). Factors influencing activities of biotransformation enzymes, concentrations and compositional patterns of organochlorine contaminants in members of a marine food web. *Aquat Toxicol* 61, 73-87.
- Stalker, M. J., Kirby, G. M., Kocal, T. E., Smith, I. R. and Hayes, M. A. (1991). Loss of glutathione S-transferases in pollution-associated liver neoplasms in white suckers (*Catostomus commersoni*) from Lake Ontario. *Carcinogenesis* 12, 2221-2226.
- Torres, M. A., Testa, C. P., Gaspari, C., Masutti, M. B., Panitz, C. M., Curi-Pedrosa, R., de Almeida, E. A., Di Mascio, P. and Filho, D. W. (2002). Oxidative stress in the mussel *Mytella guyanensis* from polluted mangroves on Santa Catarina Island, Brazil. *Mar Pollut Bull* 44, 923-932.
- Uguz, C., Iscan, M., Erguven, A., Isgor, B. and Togan, I. (2003). The bioaccumulation of nonylphenol and its adverse effect on the liver of rainbow trout (*Onchorynchus mykiss*). *Environ Res* 92, 262-270.
- Van der Oost, R., Beyer, J. and Vermeulen, N. P. E. (2003). Fish bioaccumulation and biomarkers in environmental risk assessment: a review. *Environ Toxicol Pharmacol* 13, 57-149.
- Xu, L., Chen, J., Zhang, Y., Cheung, C. C. and Lam, P. K. (2001). Biomarker studies on gold-lined sea bream (*Rhabdosargus sarba*) exposed to benzo[a]pyrene. *Water Sci Technol* 43, 155-160.

Chapter 6 (Comet assay)

- Aas, E., R. K. Bechmann, B. K. Larsen, J. Barsiene, J. Lazutka and S. Sanni (2002). Biomarkers for genotoxic effects in marine organisms. Project report 2001-2002. Stavanger, Norway, Akvamiljø.
- Akcha, F., G. Leday, et al. 2004b. Measurement of DNA adducts and strand breaks in dab (*Limanda limanda*) collected in the field: effects of biotic (age, sex) and abiotic (sampling site and period) factors on the extent of DNA damage. *Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis* 552 (1-2): 197-207.
- Akcha, F., A. Tanguy, et al. 2004a. Measurement of DNA single-strand breaks in gill and hemolymph cells of mussels, *Mytilus* sp., collected on the French Atlantic Coast. *Marine Environmental Research* 58(2-5): 753-756.
- Baussant, T. 2004. Biological effects of chronic levels of dispersed oil in North Sea and Arctic conditions: Biomarker responses and early life stage tests in two Mollusc species. Version 2. Report AM-2004/017.
- Bechmann, R.K. 2002. Biomonitoring of genetic damage. A literature review. Report AM-02/010.
- Bechmann, R.K., T. Baussant, B.K. Larsen, A. Skadsheim, S. Bamber, S. Sanni. 2004. BioSea JIP summary report of results from laboratory experiments. Report AM 2004/031.
- Bechmann, R.K., Taban, I.C., Aas, E., Sundt, R., Torgrimsen, S., Sanni, S. DNA damage in mussels and sea urchins exposed to crude oil. Manuscript in prep.
- Bechmann, R.K. Taban, I.C. 2004. Effects of Oil on Sea Urchins. Biomarker Responses and Early Life Stage Tests. Version 2. Report AM-2004/019.
- Bechmann, R. K., S. Torgrimsen, E. Aas, R. Sundt, I. C. Taban, S. Sanni. DNA damage in mussels and sea urchins exposed to crude oil. Measurement of DNA strand breaks by the Comet assay. Platform presentation at the SETAC 23rd annual meeting in North America, Salt Lake City, USA, 16 – 20 November 2002.
- Berland, H. and Sanni, S. 2005. Experimental validation of drilling effects in the field. Report from the NFR PROOF ERA validation project. In prep.

- Bjørnstad, A., Larsen, B.K., Taban, I.C., Bechmann, R.K., Torggrimsen, S., Andersen, O.K. 2002. Assessment of DNA damage in Mussels, Crabs and Fish. Poster and platform presentation at the BEEP Workshop, Athens 6th – 9th November 2002. RF-2002/449.
- Bombail, V., Aw, D., Gordon, E., and Batty, J. 2001. Application of the Comet and micronucleus assays to butterfish (*Pholis gunnellus*) erythrocytes from the Firth of Forth, Scotland. *Chemosphere* 44: 383-392.
- Brown, J. S. and S. A. Steinert. 2004. DNA damage and biliary PAH metabolites in flatfish from Southern California bays and harbors, and the Channel Islands. *Ecological Indicators* 3(4): 263-274.
- Børseth, J.F., Eriksen, V. and Tvedten, Ø.F. 2004. Oljesøl Rocknes-forliset. Biologiske effekter. Rapport RF – 2004/165.
- Chamberland, K., Lindroth, B.A., and Whitaker, B. (2002) Genotoxicity in Androscoggin River smallmouth bass. *Northeastern Naturalist* 9: 203-212.
- Devaux, A., Flammarion, P., Bernardon, V., Garric, J., and Monod, G. (1998) Monitoring of the chemical pollution of the river Rhone through measurement of DNA damage and cytochrome P4501A induction in chub (*Leuciscus cephalus*). *Marine Environmental Research* 46: 257-262.
- Dixon, D.R., and Wilson, J.T. 2000. Genetics and marine pollution. *Hydrobiologia* 420: 29-43.
- Flammarion, P., Devaux, A., Nehls, S., Migeon, B., Noury, P., and Garric, J. (2002) Multibiomarker responses in fish from the Moselle River (France). *Ecotoxicology and Environmental Safety* 51: 145-153.
- Frenzilli, G., Nigro, M., Scarcelli, V., Gorbi, S., and Regoli, F. (2001) DNA integrity and total oxyradical scavenging capacity in the Mediterranean mussel, *Mytilus galloprovincialis*: a field study in a highly eutrophicated coastal lagoon. *Aquatic Toxicology* 53: 19-32.
- Frenzilli, G., V. Scarcelli, et al. 2004. DNA damage in eelpout (*Zoarces viviparus*) from Goteborg harbour. *Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis* 552(1-2): 187-195.
- Halldorsson, H. P., G. Ericson, et al. (2004). DNA strand breakage in mussels (*Mytilus edulis* L.) deployed in intertidal and subtidal zone in Reykjavik harbour. *Marine Environmental Research* 58(2-5): 763-767.
- Large, A.T., Shaw, J.P., Peters, L.D., McIntosh, A.D., Webster, L., Mally, A., and Chipman, J.K. (2002) Different levels of mussel (*Mytilus edulis*) DNA strand breaks following chronic field and acute laboratory exposure to polycyclic aromatic hydrocarbons. *Marine Environmental Research* 54: 493-497.
- Lee, R. F., K. A. Maruya, et al. 2004. Exposure of grass shrimp to sediments receiving highway runoff: Effects on reproduction and DNA. *Marine Environmental Research* 58(2-5): 713-717.
- Lemiere, S., C. Cossu-Leguille, et al. 2004. Genotoxicity related to transfer of oil spill pollutants from mussels to mammals via food. *Environmental Toxicology* 19(4): 387-395.
- Lemiere, S., C. Cossu-Leguille, et al. 2005. DNA damage measured by the single-cell gel electrophoresis (comet) assay in mammals fed with mussels contaminated by the 'Erika' oil-spill. *Mutation Research-Genetic Toxicology and Environmental Mutagenesis* 581(1-2): 11-21.
- Mamaca, E., Bechmann, R.K., Torggrimsen, S., Aas, E., Bjørnstad, A., Baussant, T. and Le Floch, S. 2005. The neutral red lysosomal retention assay and Comet assay on haemolymph cells from mussels (*Mytilus edulis*) and fish (*Symphodus melops*) exposed to styrene. *Aquatic Toxicology*. 10:191-201.
- Mitchelmore CL, and Chipman JK. 1998. DNA strand breakage in aquatic organisms and the potential value of the comet assay in environmental monitoring. *Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis* 399: 135-147.
- Pandrangi, R., Petras, M., Ralph, S., and Vrzoc, M. (1995) Alkaline single cell gel (Comet) assay and genotoxicity monitoring using bullheads and carp. *Environmental and Molecular Mutagenesis* 26: 345-356.
- Pavlica, M., Klobucar, G.I.V., Mojas, N., Erben, R., and Papes, D. (2001) Detection of DNA damage in haemocytes of zebra mussel using Comet assay. *Mutation Research-Genetic Toxicology and Environmental Mutagenesis* 490: 209-214.
- Rank, J. 1999. Use of Comet assay on the blue mussel, *Mytilus edulis*, from coastal waters in Denmark. *Neoplasma* 46: 9-10.

- Shaw, J.P., Large, A.T., Livingstone, D.R., Doyotte, A., Renger, J., Chipman, J.K., and Peters, L.D. (2002) Elevation of cytochrome P450-immunopositive protein and DNA damage in mussels (*Mytilus edulis*) transplanted to a contaminated site. *Marine Environmental Research* 54: 505-509.
- Singh, N. P., McCoy, M. T., Tice, R. R. and Schneider, E. L. 1988. A simple technique for quantitation of low-levels of DNA damage in individual cells. *Experimental Cell Research* 175, 184-191.
- Steinert, S.A., Montee, R.S., and Sastre, M.P. 1998. Influence of sunlight on DNA damage in mussels exposed to polycyclic aromatic hydrocarbons. *Marine Environmental Research* 46: 355-358.
- Steinert, S.A., StreibMontee, R., Leather, J.M., and Chadwick, D.B. 1998. DNA damage in mussels at sites in San Diego Bay. *Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis* 399: 65-85.
- Taban, I. C., E. Aas, T. Baussant, R. K. Bechmann, B. K. Larsen, R. C. Sundt, A. Nævdal, S. Torgrimsen, G. Ericson, and J. F. Børseth. 2002. Positive control experiment - Biomarker responses in a range of marine organisms following exposure to dispersed crude oil AM-2002/008. Akvamiljø as.
- Taban, I. C., R. K. Bechmann, S. Torgrimsen, T. Baussant, and S. Sanni. 2004. Detection of DNA damage in mussels and sea urchins exposed to crude oil using comet assay. *Marine Environmental Research* 58:701-705.
- Taban I.C., Bechmann R.K., Torgrimsen S., Sundt R.C., Bjørnstad A., Larsen B.K. and Andersen O.K. 2003. DNA damage in mussels following exposure to three potential endocrine disruptors using the comet assay and the alkaline unwinding assay. Poster and platform presentation at BEEP Workshop, Barcelona, 4.-6. December 2003.
- Whitehead, A., K. M. Kuivila, et al. 2004. Genotoxicity in native fish associated with agricultural runoff events. *Environmental Toxicology and Chemistry* 23(12): 2868-2877.

Chapter 7 (Micronucleus assay)

- Arcand-Hoy, L. D. and C. D. Metcalfe (2000). Hepatic micronuclei in brown bullheads (*Ameiurus nebulosus*) as a biomarker for exposure to genotoxic chemicals. *Journal of Great Lakes Research* 26: 408-415.
- Barsiene, J. (2002). Genotoxic impacts in Klaipeda Marine port and Butinge oil terminal areas (Baltic Sea). *Marine Environmental Research* 54: 475-479.
- Baršienė, J., L. Andreikėnaitė and A. Rybakovas (2006a). Cytogenetic damage in perch (*Perca fluviatilis* L.) and Duck mussel (*Anodonta anatina* L.) exposed to crude oil. *Ekologija* (in press).
- Baršienė, J., V. Dedonytė, A. Rybakovas, L. Andreikėnaitė and O. Andersen (2005b). Induction of micronuclei in Atlantic cod (*Gadus morhua*) and turbot (*Scophthalmus maximus*) after treatment with bisphenol A, diallyl phthalate and tetrabromodiphenyl ether-47. *Ekologija*, No 4.
- Baršienė, J., V. Dedonytė, A. Rybakovas, L. Andreikėnaitė and O. K. Andersen (2006c). Investigation of micronuclei and other nuclear abnormalities in peripheral blood and kidney of marine fish treated with crude oil. *Aquatic Toxicology*. (in press).
- Baršienė, J., V. Dedonytė, A. Rybakovas, K. Broeg, L. Forlin, J. Gercken, J. Kopecka and L. Balk (2005a). Environmental mutagenesis in different zones of the Baltic Sea. *Acta Zoologica Lituanica*. 15: 90-95.
- Barsiene, J., J. Lazutka, J. Syvokiene, V. Dedonyte, A. Rybakovas, E. Bagdonas, A. Bjornstad and O. K. Andersen (2004). Analysis of micronuclei in blue mussels and fish from the Baltic and North Seas. *Environmental Toxicology*. 19: 365-371.
- Baršienė, J., K. K. Lehtonen, A. Koehler, K. Broeg, P. J. Vuorinen, T. Lang, J. Pempkowiak, L. Balk, J. Šyvokiene, V. Dedonyte, A. Rybakovas, R. Repečka, H. Vuontisjärvi and J. Kopecka (2005d). Biomarker responses in flounder (*Platichthys flesus*) and mussel (*Mytilus edulis*) in the Klaipėda-Būtingė Area (Baltic Sea). *Marine Pollution Bulletin*. (in press).
- Barsiene, J. and D. B. Lovejoy (2000). Environmental genotoxicity in Klaipeda port area. *International Review of Hydrobiology*. 85: 663-672.

- Baršienė, J., D. Schiedek, A. Rybakovas, J. Šyvokienė, J. Kopecka and L. Forlin (2005c). Cytogenetic and cytotoxic effects in gill cells of the blue mussel *Mytilus* spp. from different zones of the Baltic Sea. Marine Pollution Bulletin. (in press).
- Baršienė, J., J. Šyvokienė and A. Bjornstad (2006b). Induction of micronuclei and other nuclear abnormalities in mussels exposed to bisphenol A, diallyl phthalate and tetrabromodiphenyl ether-47. Aquatic Toxicology. (in press).
- Bihari, N. and M. Fafandel (2004). Interspecies differences in DNA single strand breaks caused by benzo(a)pyrene and marine environment. Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis. 552: 209-217.
- Bolognesi, C., G. Frenzilli, C. Lasagna, E. Perrone and P. Roggieri (2004). Genotoxicity biomarkers in *Mytilus galloprovincialis*: wild versus caged mussels. Mutation Research. 552: 153-162.
- Bolognesi, C., E. Landini, P. Roggieri, R. Fabbri and A. Viarengo (1999). Genotoxicity biomarkers in the assessment of heavy metal effects in mussels: Experimental studies. Environmental and Molecular Mutagenesis. 33: 287-292.
- Bolognesi, C., R. Rabboni and P. Roggieri (1996). Genotoxicity biomarkers in *M. galloprovincialis* as indicators of marine pollutants. Comp Biochem Physiol C-Pharmacol toxicol endocrin. 113: 319-323.
- Brunetti, R., M. Gabriele, P. Valerio and O. Fumagalli (1992). The Micronucleus Test - Temporal Pattern of Base-Line Frequency in *Mytilus-Galloprovincialis*. Marine Ecology-Progress Series. 83: 75-78.
- Burgeot, T., E. His and F. Galgani (1995). The micronucleus assay in *Crassostrea gigas* for the detection of seawater genotoxicity. Mutation Research-Genetic Toxicology. 342: 125-140.
- Burgeot, T., S. Woll and F. Galgani (1996). Evaluation of the micronucleus test on *Mytilus galloprovincialis* for monitoring applications along French coasts. Marine Pollution Bulletin. 32: 39-46.
- Buschini, A., A. Martino, B. Gustavino, M. Monfrinotti, P. Poli, C. Rossi, A. Santoro, A. M. Dorr and M. Rizzoni (2004). Comet assay and micronucleus test in circulating erythrocytes of *Cyprinus carpio* specimens exposed in situ to lake waters treated with disinfectants for potabilization. Mutation Research-Genetic Toxicology and Environmental Mutagenesis. 557: 119-129.
- Cristaldi, M., L. A. Ieradi, I. Udriou and R. Zilli (2004). Comparative evaluation of background micronucleus frequencies in domestic mammals. Mutation Research-Genetic Toxicology and Environmental Mutagenesis. 559: 1-9.
- Dailianis, S., G. P. Domouhtsidou, E. Raftopoulou, M. Kaloyianni and V. K. Dimitriadis (2003). Evaluation of neutral red retention assay, micronucleus test, acetylcholinesterase activity and a signal transduction molecule (cAMP) in tissues of *Mytilus galloprovincialis* (L.), in pollution monitoring. Marine Environmental Research. 56: 443-470.
- Dolcetti, L. and P. Venier (2002). Susceptibility to genetic damage and cell types in Mediterranean mussels. Marine Environmental Research. 54: 487-491.
- Gravato, C. and M. A. Santos (2002). Juvenile sea bass liver P450, EROD induction, and erythrocytic genotoxic responses to PAH and PAH-like compounds. Ecotoxicology and Environmental Safety. 51: 115-127.
- Gravato, C. and M. A. Santos (2003). Genotoxicity biomarkers' association with B(a)P biotransformation in *Dicentrarchus labrax* L. Ecotoxicology and Environmental Safety. 55: 352-358.
- Grisolia, C. K. and C. M. T. Cordeiro (2000). Variability in micronucleus induction with different mutagens applied to several species of fish. Genetics and Molecular Biology. 23: 235-239.
- Grisolia, C. K. and F. Starling (2001). Micronuclei monitoring of fishes from Lake Paranoa, under influence of sewage treatment plant discharges. Mutation Research-Genetic Toxicology and Environmental Mutagenesis. 491: 39-44.
- Hagger, J. A., F. A. Atienzar and A. N. Jha (2005). Genotoxic, cytotoxic, developmental and survival effects of tritiated water in the early life stages of the marine mollusc, *Mytilus edulis*. Aquatic Toxicology. 74: 205-217.

- Hagger, J. A., A. S. Fisher, S. J. Hill, M. H. Depledge and A. N. Jha (2002). Genotoxic, cytotoxic and ontogenetic effects of tri-n-butyltin on the marine worm, *Platynereis dumerilii* (Polychaeta : Nereidae). *Aquatic Toxicology*. 57: 243-255.
- Harvey, J. S., B. P. Lyons, T. S. Page, C. Stewart and J. M. Parry (1999). An assessment of the genotoxic impact of the Sea Empress oil spill by the measurement of DNA adduct levels in selected invertebrate and vertebrate species. *Mutation Research-Genetic Toxicology and Environmental Mutagenesis*. 441: 103-114.
- Hayashi, M., T. Ueda, K. Uyeno, K. Wada, N. Kinoshita and K. Saotome (1998). Development of genotoxicity assay systems that use aquatic organisms. *Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis*. 399: 125-133.
- Heddle, J. A., M. C. Cimino, M. Hayashi, F. Romagna, M. D. Shelby, J. D. Tucker, P. Vanparrys and J. T. Macgregor (1991). Micronuclei as an Index of Cytogenetic Damage - Past, Present, and Future. *Environmental and Molecular Mutagenesis*. 18: 277-291.
- Heddle, J. A., M. Hite, B. Kirkhart, K. Mavournin, J. T. Macgregor, G. W. Newell and M. F. Salamone (1983). The Induction of Micronuclei as a Measure of Genotoxicity - a Report of the United-States Environmental-Protection-Agency Gene-Tox Program. *Mutation Research*. 123: 61-118.
- Hose, J. E.-b. (1985). Potential Uses of Sea Urchin Embryos for Identifying Toxic Chemicals: Description of a Bioassay Incorporating Cytologic, Cytogenetic and Embryologic Endpoints. *Journal of Applied Toxicology*. 5: 245-254.
- Hose, J. E. and E. D. Brown (1998). Field applications of the piscine anaphase aberration test: lessons from the Exxon Valdez oil spill. *Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis*. 399: 167-178.
- Hose, J. E., D. McGurk, G. D. Marty, D. E. Hinton, E. D. Brown and T. T. Baker (1996). Sublethal effects of the Exxon Valdez oil spill on herring embryos and larvae: Morphological, cytogenetic, and histopathological assessments, 1989-1991. *Canadian Journal of Fisheries and Aquatic Sciences*. 53: 2355-2365.
- Izquierdo, J. I., G. Machado, F. Ayllon, V. L. d'Amico, L. O. Bala, E. Vallarino, R. Elias and E. Garcia-Vazquez (2003). Assessing pollution in coastal ecosystems: a preliminary survey using the micronucleus test in the mussel *Mytilus edulis*. *Ecotoxicology and Environmental Safety*. 55: 24-29.
- Jha, A. N., T. H. Hutchinson, J. M. Mackay, B. M. Elliott and D. R. Dixon (1996). Development of an in vivo genotoxicity assay using the marine worm *Platynereis dumerilii* (Polychaeta: Nereidae). *Mutation Research-Environmental Mutagenesis and Related Subjects*. 359: 141-150.
- Jha, A. N., T. H. Hutchinson, J. M. Mackay, B. M. Elliott and D. R. Dixons (1997). Evaluation of the genotoxicity of municipal sewage effluent using the marine worm *Platynereis dumerilii* (Polychaeta: Nereidae). *Mutation Research-Genetic Toxicology and Environmental Mutagenesis*. 391: 179-188.
- Kalpaxis, D. L., C. Theos, M. A. Xaplanteri, G. P. Dinos, A. V. Catsiki and M. Leotsinidis (2004). Biomonitoring of Gulf of Patras, N. Peloponnesus, Greece. Application of a biomarker suite including evaluation of translation efficiency in *Mytilus galloprovincialis* cells. *Environmental Research*. 94: 211-220.
- Kocan, R. M., J. E. Hose, E. D. Brown and T. T. Baker (1996). Pacific herring (*Clupea pallasii*) embryo sensitivity to Prudhoe Bay petroleum hydrocarbons: Laboratory evaluation and in situ exposure at oiled and unoled sites in Prince William Sound. *Canadian Journal of Fisheries and Aquatic Sciences*. 53: 2366-2375.
- Kramer, P. J. (1998). Genetic toxicology. *Journal of Pharmacy and Pharmacology*. 50: 395-405.
- LaRocca, C., L. Conti, R. Crebelli, B. Crochi, N. Iacovella, F. Rodriguez, L. TurrioBaldassarri and A. diDomenico (1996). PAH content and mutagenicity of marine sediments from the Venice lagoon. *Ecotoxicology and Environmental Safety*. 33: 236-245.
- Lyons, B. P., J. S. Harvey and J. M. Parry (1997). An initial assessment of the genotoxic impact of the Sea Empress oil spill by the measurement of DNA adduct levels in the intertidal teleost *Lipophrys pholis*. *Mutation Research-Genetic Toxicology and Environmental Mutagenesis*. 390: 263-268.

- MacGregor, J. T. (1991). Micronucleus Assay Protocols. Mutation Research. 259: 123-125.
- Majone, F., C. Beltrame and R. Brunetti (1988). Frequencies of Micronuclei Detected on *Mytilus-Galloprovincialis* by Different Staining Techniques after Treatment with Zinc-Chloride. Mutation Research. 209: 131-134.
- Majone, F., R. Brunetti, O. Fumagalli, M. Gabriele and A. G. Levis (1990). Induction of Micronuclei by Mitomycin-C and Colchicine in the Marine Mussel *Mytilus-Galloprovincialis*. Mutation Research. 244: 147-151.
- Majone, F., R. Brunetti, I. Gola and A. G. Levis (1987). Persistence of Micronuclei in the Marine Mussel, *Mytilus-Galloprovincialis*, after Treatment with Mitomycin-C. Mutation Research. 191: 157-161.
- Maria, V. L., A. C. Correia and M. A. Santos (2002a). *Anguilla anguilla* L. biochemical and genotoxic responses to benzo a pyrene. Ecotoxicology and Environmental Safety. 53: 86-92.
- Maria, V. L., C. Gravato, A. C. Correia and M. A. Santos (2002b). Biotransformation and genotoxicity responses to PAHs in two teleost species. Fresenius Environmental Bulletin. 11: 609-615.
- Minissi, S., E. Ciccotti and M. Rizzoni (1996). Micronucleus test in erythrocytes of *Barbus plebejus* (Teleostei, Pisces) from two natural environments: A bioassay for the in situ detection of mutagens in freshwater. Mutation Research - Genetic Toxicology. 367: 245-251.
- Moore, M. N., M. H. Depledge, J. W. Readman and D. R. P. Leonard (2004). An integrated biomarker-based strategy for ecotoxicological evaluation of risk in environmental management. Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis. 552: 247-268.
- Norcross, B. L., J. E. Hose, M. Frandsen and E. D. Brown (1996). Distribution, abundance, morphological condition, and cytogenetic abnormalities of larval herring in Prince William Sound, Alaska, following the Exxon Valdez oil spill. Canadian Journal of Fisheries and Aquatic Sciences. 53: 2376-2387.
- Pacheco, M. and M. A. Santos (1997). Induction of EROD activity and genotoxic effects by polycyclic aromatic hydrocarbons and resin acids on the juvenile eel (*Anguilla anguilla* L.). Ecotoxicology and Environmental Safety. 38: 252-259.
- Pacheco, M. and M. A. Santos (2001). Tissue distribution and temperature-dependence of *Anguilla anguilla* L. EROD activity following exposure to model inducers and relationship with plasma cortisol, lactate and glucose levels. Environment International. 26: 149-155.
- Parry, J. M., J. S. Harvey and B. P. Lyons (1997). The application of genetic toxicology in the analysis of the consequences of a major marine pollution incident. Mutation Res. Supplement 1: S91.
- Pietrapiana, D., M. Modena, P. Guidetti, C. Falugi and M. Vacchi (2002). Evaluating the genotoxic damage and hepatic tissue alterations in demersal fish species: a case study in the Ligurian Sea (NW-Mediterranean). Marine Pollution Bulletin. 44: 238-243.
- Rao, S. S., T. Neheli, J. H. Carey and V. W. Cairns (1997). Fish hepatic micronuclei as an indication of exposure to genotoxic environmental contaminants. Environmental Toxicology and Water Quality. 12: 217-222.
- Rovero, F., R. N. Hughes and C. Chelazzi (1999). Cardiac and behavioural responses of mussels to risk of predation by dogwhelks. Animal Behaviour. 58: 707-714.
- Sanchez-Galan, S., A. R. Linde, F. Ayllon and E. Garcia-Vazquez (2001). Induction of micronuclei in eel (*Anguilla anguilla* L.) by heavy metals. Ecotoxicology and Environmental Safety. 49: 139-143.
- Saotome, K. and M. Hayashi (2003). Application of a sea urchin micronucleus assay to monitoring aquatic pollution: influence of sample osmolality. Mutagenesis. 18: 73-76.
- Saotome, K., T. Sofuni and M. Hayashi (1999). A micronucleus assay in sea urchin embryos. Mutation Research - Genetic Toxicology and Environmental Mutagenesis. 446: 121-127.
- Scarpato, R., L. Migliore, G. Alfinitocognetti and R. Barale (1990). Induction of Micronuclei in Gill Tissue of *Mytilus-Galloprovincialis* Exposed to Polluted Marine Waters. Marine Pollution Bulletin. 21: 74-80.
- Seelbach, A., B. Fissler, A. Strohmusch and S. Madle (1993). Development of a Modified Micronucleus Assay In vitro for Detection of Aneugenic Effects. Toxicology in Vitro. 7: 185-193.

- Siu, W. H. L., J. Cao, R. W. Jack, R. S. S. Wu, B. J. Richardson, L. Xu and P. K. S. Lam (2004). Application of the comet and micronucleus assays to the detection of B a P genotoxicity in haemocytes of the green-lipped mussel (*Perna viridis*). *Aquatic Toxicology*. 66: 381-392.
- Stegeman, J. J., M. N. Moore and M. E. Hahn (1993). Special Issue - Responses of Marine Organisms to Pollutants .2. *Marine Environmental Research*. 35: R3-R4.
- Teles, M., M. Pacheco and M. A. Santos (2003). *Anguilla anguilla* L. liver ethoxyresorufin O-deethylation, glutathione S-transferase, erythrocytic nuclear abnormalities, and endocrine responses to naphthalene and beta-naphthoflavone. *Ecotoxicology and Environmental Safety*. 55: 98-107.
- Torres-Bugarin, O., A. D. Anda-Casillas, M. P. Ramirez-Munoz, J. Sanchez-Corona, J. M. Cantu and G. Zuniga (1998). Determination of diesel genotoxicity in firebreathers by micronuclei and nuclear abnormalities in buccal mucosa. *Mutation Research-Genetic Toxicology and Environmental Mutagenesis*. 413: 277-281.
- Tripathy, A. P. and R. K. Das (1995). River water contaminated with paper-mill effluent induces micronuclei in peripheral erythrocytes of fish. *Current Science*. 68: 766-767.
- Tuvikene, A., S. Huuskonen, K. Koponen, O. Ritola, U. Mauer and P. Lindstrom-Seppa (1999). Oil shale processing as a source of aquatic pollution: Monitoring of the biologic effects in caged and feral freshwater fish. *Environmental Health Perspectives*. 107: 745-752.
- Venier, P., S. Maron and S. Canova (1997). Detection of micronuclei in gill cells and haemocytes of mussels exposed to benzo[a]pyrene. *Mutation Research-Genetic Toxicology and Environmental Mutagenesis*. 390: 33-44.
- White, P. A., J. B. Rasmussen and C. Blaise (1998). Genotoxic substances in the St. Lawrence system I: Industrial genotoxins sorbed to particulate matter in the St. Lawrence, St. Maurice, and Saguenay Rivers, Canada. *Environmental Toxicology and Chemistry*. 17: 286-303.
- Williams, R. C. and C. D. Metcalfe (1992). Development of an In vivo Hepatic Micronucleus Assay with Rainbow-Trout. *Aquatic Toxicology*. 23: 193-202.
- Wilson, J. T., D. R. Dixon and L. R. J. Dixon (2002). Numerical chromosomal aberrations in the early life-history stages of a marine tubeworm, *Pomatoceros lamarckii* (Polychaeta : Serpulidae). *Aquatic Toxicology*. 59: 163-175.
- Woodhead, R. J., R. J. Law and P. Matthiessen (1999). Polycyclic aromatic hydrocarbons in surface sediments around England and Wales, and their possible biological significance. *Marine Pollution Bulletin*. 38: 773-790.
- Wrisberg, M. N. and R. Rhemrev (1992). Detection of genotoxins in the aquatic environment with the mussel *Mytilus edulis*. *Water Science and Technology*. 25: 317-324.
- Zakhidov, S. T., Y. V. Chebotareva, K. A. Savvaitova and V. A. Maksimov (1996). A cytogenetic study of hemopoietic cells in fish from water bodies of the Norilo-Pyasinsk water system (Taimyr). *Izvestiya Akademii Nauk Seriya Biologicheskaya*: 10-15.
- Zoll-Moreux, C. and V. Ferrier (1999). The Jaylet test (newt micronucleus test) and the micronucleus test in xenopus: Two in vivo tests on amphibia evaluation of the genotoxicity of five environmental pollutants and of five effluents. *Water Research*. 33: 2301-2314.
- Aas, E., J. Barsiene, J. Lazutka and R. Sanger (2002). Micronuclei analyses of blue mussels and cod following exposure to dispersed crude oil. *AM 2002/011*: 33.

Chapter 8 (Multixenobiotic response)

- Bard, S. M. (2000). Multixenobiotic resistance as a cellular defense mechanism in aquatic organisms. *Aquatic Toxicol* 48, 357-389.

- Britvic, S., and Kurelec, B. (1999). The effect of inhibitors of multixenobiotic resistance mechanism on the production of mutagens by *Dreissena polymorpha* in waters spiked with premutagens. *Aquatic Toxicology* 47, 107-116.
- Cooper, P. S., Vogelbein, W. K., and Van Veld, P. A. (1999). Altered expression of the xenobiotic transporter P-glycoprotein in liver and liver tumours of mummichog (*Fundulus heteroclitus*) from a creosote-contaminated environment. *Biomarkers* 4, 48-58.
- Curtis, L. R., Hemmer, M. J., and Courtney, L. A. (2000). Dieldrin induces cytosolic [H-3]7,12-dimethylbenz[a]anthracene binding but not multidrug resistance proteins in rainbow trout liver. *Journal Of Toxicology And Environmental Health-Part A* 60, 275-289.
- Daughton, C. G., and Ternes, T. A. (1999). Pharmaceuticals and personal care products in the environment: Agents of subtle change? *Environmental Health Perspectives* 107, 907-938.
- Doi, A. M., Holmes, E., and Kleinow, K. M. (2001). P-glycoprotein in the catfish intestine: inducibility by xenobiotics and functional properties. *Aquat Toxicol* 55, 157-70.
- Epel, D. (1998). Use of multidrug transporters as first lines of defense against toxins in aquatic organisms. *COMP BIOCHEM PHYS A* 120, 23-28.
- Eufemia, N., Clerte, S., Girshick, S., and Epel, D. (2002). Algal products as naturally occurring substrates for p-glycoprotein in *Mytilus californianus*. *Marine Biology* 140, 343-353.
- Eufemia, N. A., and Epel, D. (1998). The multixenobiotic defense mechanism in mussels is induced by substrates and non-substrates: Implications for a general stress response. *Marine Environmental Research* 46, 401-405.
- Eufemia, N. A., and Epel, D. (2000). Induction of the multixenobiotic defense mechanism (MXR), P-glycoprotein, in the mussel *Mytilus californianus* as a general cellular response to environmental stresses. *Aquatic Toxicol* 49, 89-100.
- Galgani, F. (2000). Biological effects of contaminants on marine organisms: the contribution of biotechnology. *International Journal Of Environment And Pollution* 13, 34-44.
- Jha, A. N. (2004). Genotoxicological studies in aquatic organisms: an overview. *Mutation Research-Fundamental And Molecular Mechanisms Of Mutagenesis* 552, 1-17.
- Keppler, C. J., and Ringwood, A. H. (2001). Expression of P-glycoprotein in southeastern oysters, *Crassostrea virginica*. *Marine Environmental Research* 52, 81-96.
- Kurelec, B. (1995). Reversion Of The Multixenobiotic Resistance Mechanism In Gills Of A Marine Mussel *Mytilus-Galloprovincialis* By A Model Inhibitor And Environmental Modulators Of P170-Glycoprotein. *Aquatic Toxicology* 33, 93-103.
- Kurelec, B. (1997). A new type of hazardous chemical: the chemosensitizers of multixenobiotic resistance. *Environ Health Perspect* 105 Suppl 4, 855-60.
- Kurelec, B., Britvic, S., Pivcevic, B., and Smital, T. (1998). Fragility of multixenobiotic resistance in aquatic organisms enhances the complexity of risk assessment. *Marine Environmental Research* 46, 415-419.
- Kurelec, B., Lucic, D., Pivcevic, B., and Krca, S. (1995). Induction And Reversion Of Multixenobiotic Resistance In The Marine Snail *Monodonta-Turbinata*. *Marine Biology* 123, 305-312.
- Kurelec, B., Waldmann, P., and Zahn, R. K. (1996). The modulation of protective effects of the multixenobiotic resistance mechanism in a clam *Corbicula fluminea*. *Marine Environmental Research* 42, 383-387.
- Livingstone, D. R., Chipman, J. K., Lowe, D. M., Minier, C., Mitchelmore, C. L., Moore, M. N., Peters, L. D., and Pipe, R. K. (2000). Development of biomarkers to detect the effects of organic pollution on aquatic invertebrates: recent molecular, genotoxic, cellular and immunological studies on the common mussel (*Mytilus edulis* L.) and other mytilids. *International Journal of Environment and Pollution* 13, 56-91.
- Luckenbach, T., Corsi, I., and Epel, D. (2004). Fatal attraction: synthetic musk fragrances compromise multixenobiotic defense systems in mussels. *Mar Environ Res* 58, 215-9.
- Luedeking, A., and Koehler, A. (2004). Regulation of expression of multixenobiotic resistance (MXR) genes by environmental factors in the blue mussel *Mytilus edulis*. *Aquat Toxicol* 69, 1-10.

- Minier, C., Borghi, V., Moore, M. N., and Porte, C. (2000). Seasonal variation of MXR and stress proteins in the common mussel, *Mytilus galloprovincialis*. *Aquatic Toxicology* 50, 167-176.
- Minier, C., Eufemia, N., and Epel, D. (1999). The multi-xenobiotic resistance phenotype as a tool to biomonitor the environment. *Biomarkers* 4, 442-454.
- Minier, C., and Moore, M. N. (1996). Induction of multixenobiotic resistance in mussel blood cells. *Marine Environmental Research* 42, 389-392.
- Moore, M. N., Depledge, M. H., Readman, J. W., and Leonard, D. R. P. (2004). An integrated biomarker-based strategy for ecotoxicological evaluation of risk in environmental management. *Mutation Research-Fundamental And Molecular Mechanisms Of Mutagenesis* 552, 247-268.
- Muller, W. E. G., Riemer, S., Kurelec, B., Smolaka, N., Puskaric, S., Jagic, B., Muller-Niklas, G., and Queric, N. V. (1998). Chemosensitizers of the multixenobiotic resistance in amorphous aggregates (marine snow): etiology of mass killing on the benthos in the Northern Adriatic? *Environmental Toxicology and Pharmacology* 6, 229-238.
- Pain, S., and Parant, M. (2003). Multixenobiotic defence mechanism (MXDM) in bivalves. *Comptes Rendus Biologies* 326, 659-672.
- Pain, S., and Parant, M. (2003). Response of multixenobiotic defence mechanism in *Dreissena polymorpha* exposed to environmental stress. *Chemosphere* 52, 1105-1113.
- Parant, M., and Pain, S. (2001). Potential use of multixenobiotic defense mechanism (MXDM) in *Dreissena polymorpha* as a biomarker for the monitoring of freshwater pollution. *Water Research* 35, 3743-3748.
- Ringwood, A. H., Conners, D. E., Keppler, C. J., and DiNovo, A. A. (1999). Biomarker studies with juvenile oysters (*Crassostrea virginica*) deployed in-situ. *Biomarkers* 4, 400-414.
- Smital, T., and Kurelec, B. (1998). The activity of multixenobiotic resistance mechanism determined by rhodamine B-efflux method as a biomarker of exposure. *Marine Environmental Research* 46, 443-447.
- Smital, T., Luckenbach, T., Sauerborn, R., Hamdoun, A. A., Vega, R. L., and Epel, D. (2004). Emerging contaminants - pesticides, PPCPs, microbial degradation products and natural substances as inhibitors of multixenobiotic defense in aquatic organisms. *Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis* 552, 101-117.
- Sturm, A., Cravedi, J. P., and Segner, H. (2001). Prochloraz and nonylphenol diethoxylate inhibit an *mdr1*-like activity in vitro, but do not alter hepatic levels of P-glycoprotein in trout exposed in vivo. *Aquat Toxicol* 53, 215-28.
- van der Oost, R., Beyer, J., and Vermeulen, N. P. E. (2003). Fish bioaccumulation and biomarkers in environmental risk assessment: a review. *Environmental Toxicology and Pharmacology* 13, 57-149.
- Waldmann, P., Pivcevic, B., Muller, W. E., Zahn, R. K., and Kurelec, B. (1995). Increased genotoxicity of acetylaminofluorene by modulators of multixenobiotic resistance mechanism: studies with the fresh water clam *Corbicula fluminea*. *Mutat Res* 342, 113-23.

Chapter 9 (Peroxisomal proliferation)

- Ackers, J. T., Johnston, M. F., and Haasch, M. L. (2000). Immunodetection of hepatic peroxisomal PMP70 as an indicator of peroxisomal proliferation in the mummichog, *Fundulus heteroclitus*. *Mar Environ Res* 50, 361-5.
- Arnold, H., Pluta, H., and Braunbeck, T. (1995). Simultaneous exposure of fish to endosulfan and disulfoton in vivo - ultrastructural, stereological and biochemical reactions in hepatocytes of male rainbow-trout (*Oncorhynchus-mykiss*). *AQUAT TOXICOL* 33, 17-43.
- Au, D., Wu, R., Zhou, B., and Lam, P. (1999). Relationship between ultrastructural changes and EROD activities in liver of fish exposed to Benzo[a]pyrene. *ENVIRON POLLUT* 104, 235-247.
- Bentley, P., Calder, I., Elcombe, C., Grasso, P., Stringer, D., and Wiegand, H. J. (1993). Hepatic peroxisome proliferation in rodents and its significance for humans. *Food Chem Toxicol* 31, 857-907.

- Braunbeck, T., Gorgas, K., Storch, V., and Volkl, A. (1987). Ultrastructure of hepatocytes in golden ide (*Leuciscus idus melanotus* L.; Cyprinidae: Teleostei) during thermal adaptation. *Anat Embryol (Berl)* 175, 303-13.
- Braunbeck, T., Gorge, G., Storch, V., and Nagel, R. (1990). Hepatic steatosis in zebra fish (*Brachydanio rerio*) induced by long-term exposure to gamma-hexachlorocyclohexane. *Ecotoxicol Environ Saf* 19, 355-74.
- Braunbeck, T., and Volkl, A. (1991). Induction of biotransformation in the liver of Eel (*Anguilla anguilla* L.) by sublethal exposure to dinitro-o-cresol: an ultrastructural and biochemical study. *Ecotoxicol Environ Saf* 21, 109-27.
- Cajaraville, M., Orbea, A., Marigomez, I., and Cancio, I. (1997). Peroxisome proliferation in the digestive epithelium of mussels exposed to the water accommodated fraction of three oils. *COMP BIOCHEM PHYS C* 117, 233-242.
- Cajaraville, M. P., Cancio, M., Ibabe, A., and Orbea, A. (2003). Peroxisome proliferation as a biomarker in environmental pollution assessment. *Microscopy Research and Technique* 61, 191-202.
- Cancio, I., and Cajaraville, M. P. (2000). Cell biology of peroxisomes and their characteristics in aquatic organisms. *Int Rev Cytol* 199, 201-93.
- Cancio, I., Ibabe, A., and Cajaraville, M. P. (1999). Seasonal variation of peroxisomal enzyme activities and peroxisomal structure in mussels *Mytilus galloprovincialis* and its relationship with the lipid content. *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 123, 135-144.
- Cancio, I., Orbea, A., Volkl, A., Fahimi, H. D., and Cajaraville, M. P. (1998). Induction of peroxisomal oxidases in mussels: Comparison of effects of lubricant oil and benzo(a)pyrene with two typical peroxisome proliferators on peroxisome structure and function in *Mytilus galloprovincialis*. *Toxicology and Applied Pharmacology* 149, 64-72.
- Hess, R., Staubli, W., and Riess, W. (1965). Nature of the hepatomegalic effect produced by ethylchlorophenoxy-isobutyrate in the rat. *Nature* 208, 856-8.
- Hoff, P. T., Van Dongen, W., Esmans, E. L., Blust, R., and De Coen, W. M. (2003). Evaluation of the toxicological effects of perfluorooctane sulfonic acid in the common carp (*Cyprinus carpio*). *AQUAT TOXICOL* 62, 349-359.
- Haasch, M. L. (2002). Effects of vehicle, diet and gender on the expression of PMP70- and CYP2K1/2M1-like proteins in the mummichog. *MAR ENVIRON RES* 54, 297-301.
- Ibabe, A., Bilbao, E., and Cajaraville, M. (2005). Expression of peroxisome proliferator-activated receptors in zebrafish (*Danio rerio*) depending on gender and developmental stage. *HISTOCHEM CELL BIOL* 123, 75-87.
- Krasnov, A., Koskinen, H., Rexroad, C., Afanasyev, S., Molsa, H., and Oikari, A. (2005). Transcriptome responses to carbon tetrachloride and pyrene in the kidney and liver of juvenile rainbow trout (*Oncorhynchus mykiss*). *AQUAT TOXICOL* 74, 70-81.
- Krishnakumar, P., Casillas, E., and Varanasi, U. (1997). Cytochemical responses in the digestive tissue of *Mytilus edulis* complex exposed to microencapsulated PAHs or PCBs. *COMP BIOCHEM PHYS C* 118, 11-18.
- Livingstone, D., Martinez, P., Michel, X., Narbonne, J., Ohara, S., Ribera, D., and Winston, G. (1990). Oxyradical production as a pollution-mediated mechanism of toxicity in the common mussel, *Mytilus-edulis*-L, and other mollusks. *FUNCT ECOL* 4, 415-424.
- Mather-Mihaich, E., and Di Giulio, R. T. (1991). Oxidant, mixed-function oxidase and peroxisomal responses in channel catfish exposed to a bleached kraft mill effluent. *Arch Environ Contam Toxicol* 20, 391-7.
- Orbea, A., Beier, K., Volkl, A., Fahimi, H. D., and Cajaraville, M. P. (1999). Ultrastructural, immunocytochemical and morphometric characterization of liver peroxisomes in gray mullet, *Mugil cephalus*. *Cell and Tissue Research* 297, 493-502.
- Orbea, A., Marigomez, I., Fernandez, C., Tarazona, J. V., Cancio, I., and Cajaraville, M. P. (1999). Structure of peroxisomes and activity of the marker enzyme catalase in digestive epithelial cells in relation to PAH

- content of mussels from two Basque estuaries (Bay of Biscay): Seasonal and site-specific variations. *Archives of Environmental Contamination and Toxicology* 36, 158-166.
- Orbea, A., Ortiz-Zarragoitia, M., and Cajaraville, M. P. (2002). Interactive effects of benzo(a) pyrene and cadmium and effects of di(2-ethylhexyl) phthalate on antioxidant and peroxisomal enzymes and peroxisomal volume density in the digestive gland of mussel *Mytilus galloprovincialis* Lmk. *Biomarkers* 7, 33-48.
- Orbea, A., Ortiz-Zarragoitia, M., Sole, M., Porte, C., and Cajaraville, M. P. (2002). Antioxidant enzymes and peroxisome proliferation in relation to contaminant body burdens of PAHs and PCBs in bivalve molluscs, crabs and fish from the Urdaibai and Plentzia estuaries (Bay of Biscay). *AQUAT TOXICOL* 58, 75-98.
- Ortiz-Zarragoitia, M., and Cajaraville, M. P. (2005). Effects of selected xenoestrogens on liver peroxisomes, vitellogenin levels and spermatogenic cell proliferation in male zebrafish. *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 141, 133-144.
- Oulmi, Y., Negele, R., and Braunbeck, T. (1995). Cytopathology of liver and kidney in Rainbow-trout *oncorhynchus-mykiss* after long-term exposure to sublethal concentrations of linuron. *DIS AQUAT ORGAN* 21, 35-52.
- Oulmi, Y., Negele, R., and Braunbeck, T. (1995). Segment specificity of the cytological response in Rainbow-trout (*Onchorhynchus-mykiss*) renal tubules following prolonged exposure to sublethal concentrations of atrazine. *ECOTOX ENVIRON SAFE* 32, 39-50.
- Pedrajás, J. R., LopezBarea, J., and Peinado, J. (1996). Dieldrin induces peroxisomal enzymes in fish (*Sparus aurata*) liver. *COMP BIOCHEM PHYS C* 115, 125-131.
- Reddy, J. K., Azarnoff, D. L., and Hignite, C. E. (1980). Hypolipidaemic hepatic peroxisome proliferators form a novel class of chemical carcinogens. *Nature* 283, 397-8.
- Rocha, E., Lobo-da-Cunha, A., Monteiro, R. A. F., Silva, M. W., and Oliveira, M. H. (1999). A stereological study along the year on the hepatocytic peroxisomes of brown trout (*Salmo trutta*). *Journal of Submicroscopic Cytology and Pathology* 31, 91-105.
- Yang, J. H., Kostecky, P. T., Calabrese, E. J., and Baldwin, L. A. (1990). Induction of peroxisome proliferation in rainbow trout exposed to ciprofibrate. *Toxicol Appl Pharmacol* 104, 476-82.

Chapter 10 (Acetyl cholinesterase inhibition)

- Akcha, F., Izuel, C., Venier, P., Budzinski, H., Burgeot, T., and Narbonne, J. F. (2000). Enzymatic biomarker measurement and study of DNA adduct formation in benzo[a]pyrene-contaminated mussels, *Mytilus galloprovincialis*. *Aquatic Toxicology* 49, 269-287.
- Barron, M. G., Hansen, J. A., and Lipton, J. (2002). Association between contaminant tissue residues and effects in aquatic organisms. *Rev Environ Contam Toxicol* 173, 1-37.
- Beauvais, S., Jones, S., Brewer, S., and Little, E. (2000). Physiological measures of neurotoxicity of diazinon and malathion to larval rainbow trout (*Oncorhynchus mykiss*) and their correlation with behavioral measures. *ENVIRON TOXICOL CHEM* 19, 1875-1880.
- Bocquene, G., Chantreau, S., Clerendeau, C., Beausir, E., Menard, D., Raffin, B., Minier, C., Burgeot, T., Leszkowicz, A. P., and Narbonne, J. F. (2004). Biological effects of the "Erika" oil spill on the common mussel (*Mytilus edulis*). *Aquatic Living Resources* 17, 309-316.
- Bocquené, G., and Galgani, F. 1998. Biological effects of contaminants: Cholinesterase inhibition by organophosphate and carbamate compounds. *ICES Techniques in Marine Environmental Science*, 23, 12 pp.
- Bocquene, G., Roig, A., and Fournier, D. (1997). Cholinesterases from the common oyster (*Crassostrea gigas*) - Evidence for the presence of a soluble acetylcholinesterase insensitive to organophosphate and carbamate inhibitors. *FEBS LETT* 407, 261-266.
- Bodin, N., Burgeot, T., Stanisiere, J. Y., Bocquene, G., Menard, D., Minier, C., Boutet, I., Amat, A., Cherel, Y., and Budzinski, H. (2004). Seasonal variations of a battery of biomarkers and physiological indices for

- the mussel *Mytilus galloprovincialis* transplanted into the northwest Mediterranean Sea. *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 138, 411-427.
- Cripe, G. M., Goodman, L. R., and Hansen, D. J. (1984). Effect Of Chronic Exposure To Epn And To Guthion On The Critical Swimming Speed And Brain Acetylcholinesterase Activity Of Cyprinodon-Variegatus. *Aquatic Toxicology* 5, 255-266.
- Escartin, E., and Porte, C. (1997). The use of cholinesterase and carboxylesterase activities from *Mytilus galloprovincialis* in pollution monitoring. *ENVIRON TOXICOL CHEM* 16, 2090-2095.
- Flammarion, P., Noury, P., and Garric, J. (2002). The measurement of cholinesterase activities as a biomarker in chub (*Leuciscus cephalus*): the fish length should not be ignored. *ENVIRON POLLUT* 120, 325-330.
- Fulton, M. H., and Key, P. B. (2001). Acetylcholinesterase inhibition in estuarine fish and invertebrates as an indicator of organophosphorus insecticide exposure and effects. *ENVIRON TOXICOL CHEM* 20, 37-45.
- Galgani, F., and Bocequene, G. (1990). Invitro inhibition of acetylcholinesterase from 4 marine species by oragnophosphates and carbamates. *B ENVIRON CONTAM TOX* 45, 243-249.
- Kopecka, J., Rybakowas, A., Barsiene, J., and Pempkowiak, J. (2004). AChE levels in mussels and fish collected off Lithuania and Poland (southern Baltic). *Oceanologia* 46, 405-418.
- LeBris, H., Maffart, P., Bocquene, G., Buchet, V., Galgani, F., and Blanc, G. (1995). Laboratory study on the effect of dichlorvos on two commercial bivalves. *Aquaculture* 138, 139-144.
- Moreira, S., and Guilhermino, L. (2005). The use of *Mytilus galloprovincialis* acetylcholinesterase and glutathione S-transferases activities as biomarkers of environmental contamination along the northwest Portuguese coast. *Environmental Monitoring and Assessment* 105, 309-325.
- Morgan, M. J., Fancey, L. L., and Kiceniuk, J. W. (1990). Response And Recovery Of Brain Acetylcholinesterase Activity In Atlantic Salmon (*Salmo-Salar*) Exposed To Fenitrothion. *Canadian Journal Of Fisheries And Aquatic Sciences* 47, 1652-1654.
- Roche, H., Buet, A., and Ramade, F. (2002). Accumulation of lipophilic microcontaminants and biochemical responses in eels from the Camargue Biosphere Reserve. *Ecotoxicology* 11, 155-164.
- Romeo, M., Mourgaud, Y., Geffard, A., Gnassia-Barelli, M., Amiard, J. C., and Budzinski, H. (2003). Multimarker approach in transplanted mussels for evaluating water quality in Charentes, France, coast areas exposed to different anthropogenic conditions. *Environmental Toxicology* 18, 295-305.
- Sanchez-Hernandez, J. C., Fossi, M. C., Leonzio, C., Focardi, S., Barra, R., Gavilan, J. F., and Parra, O. (1998). Use of biochemical biomarkers as a screening tool to focus the chemical monitoring of organic pollutants in the Biobio river basin (Chile). *Chemosphere* 37, 699-710.
- Sandahl, J. F., Baldwin, D. H., Jenkins, J. J., and Scholz, N. L. (2005). Comparative thresholds for acetylcholinesterase inhibition and behavioral impairment in coho salmon exposed to chlorpyrifos. *ENVIRON TOXICOL CHEM* 24, 136-145.
- VanDolah, R. E., Maier, P. P., Fulton, M. H., and Scott, G. I. (1997). Comparison of azinphosmethyl toxicity to juvenile red drum (*Sciaenops ocellatus*) and the mummichog (*Fundulus heteroclitus*). *ENVIRON TOXICOL CHEM* 16, 1488-1493.
- Williams, A. K., and Sova, C. (1966). Acetylcholinesterase levels in brains of fishes from polluted waters. *Bull. Environ. Contam. Toxicol* 1, 198-204.

Chapter 11 (Vitellogenin)

- Allen Y, Scott AP, Matthiessen P, Haworth S, Thain JE, Feist S (1999) Survey of estrogenic activity in United Kingdom estuarine and coastal waters and its effects on gonadal development of the flounder *Platichthys flesus*. *Environ Toxicol Chem* 18:1791-1800
- Andersen L, Holbech H, Gessbo A, Norrgren L, Petersen GI (2003) Effects of exposure to 17 alpha-ethinylestradiol during early development on sexual differentiation and induction of vitellogenin in zebrafish (*Danio rerio*). *Comparative Biochemistry and Physiology* 134C:365-374

- Arukwe A, Celius T, Walther BT, Goksoyr A (2000) Effects of xenoestrogen treatment on zona radiata protein and vitellogenin expression in Atlantic salmon (*Salmo salar*). *Aquat Toxicol* 49:159-170
- Christensen LJ, Korsgaard B, Bjerregaard P (1999) The effect of 4-nonylphenol on the synthesis of vitellogenin in the flounder *Platichthys flesus*. *Aquat Toxicol* 46:211-219
- Christiansen T, Korsgaard B, Jespersen A (1998) Effects of nonylphenol and 17 beta-oestradiol on vitellogenin synthesis, testicular structure and cytology in male eelpout *Zoarces viviparus*. *Journal of Experimental Biology* 201:179-192
- Hylland, K., Berge, J.A., Goksøyr, A., Pettersen, O., Sætre, T., Efraimsen, H. 1998. Effekter av østrogenlignende stoffer i norske kystfarvann. NIVA-rapport 3668, 71 s. [in Norwegian]
- Kirby MF, Allen YT, Dyer RA, Feist SW, Katsiadaki I, Matthiessen P, Scott AP, Smith A, Stentiford GD, Thain JE, Thomas KV, Tolhurst L, Waldock MJ (2004) Surveys of plasma vitellogenin and intersex in male flounder (*Platichthys flesus*) as measures of endocrine disruption by estrogenic contamination in United Kingdom estuaries: Temporal trends, 1996 to 2001. *Environ Toxicol Chem* 23:748-758
- Madsen LL, Korsgaard B, Bjerregaard P (2003) Estrogenic effects in flounder *Platichthys flesus* orally exposed to 4-tert-octylphenol. *Aquat Toxicol* 64:393-405
- Oppen-Berntsen DO, Hyllner SJ, Haux C, Helvik JV, Walther BT (1992) Eggshell zona radiata-proteins from cod (*Gadus morhua*): Extra-ovarian origin and induction by estradiol-17 beta. *IntJDevBiol* 36:247-254
- Scott, A.P., and Hylland, K. 2002. Biological effects of contaminants: Radioimmunoassay (RIA) and enzyme-linked immunosorbent assay (ELISA) techniques for the measurement of marine fish vitellogenins. *ICES Techniques in Marine Environmental Sciences*. 21pp.
- Scott AP, Katsiadaki I, Witthames PR, Hylland K, Davies IM, McIntosh AD, Thain J (2005) Vitellogenin in the blood plasma of male cod (*Gadus morhua*): A sign of oestrogenic endocrine disruption in the open sea? *Mar Environ Res* in press.
- Scott, AP; Stewart, C; Allen, Y; Matthiessen, P. 2000. 17 beta -oestradiol in male flatfish. *Reproductive Physiology of Fish*. p. 382.
- Vethaak AD, Lahr J, Schrap SM, Belfroid AC, Rijs GBJ, Gerritsen A, de Boer J, Bulder AS, Grinwis GCM, Kuiper RV, Legler J, Murk TAJ, Peijnenburg W, Verhaar HJM, de Voogt P (2005) An integrated assessment of estrogenic contamination and biological effects in the aquatic environment of The Netherlands. *Chemosphere* 59:511-524
- Örn S, Holbech H, Madsen TH, Norrgren L, Petersen GI (2003) Gonad development and vitellogenin production in zebrafish (*Danio rerio*) exposed to ethinylestradiol and methyltestosterone. *Aquat Toxicol* 65:397-411

Chapter 12 (Metallothionein)

- Bebiano MJ, Langston WJ (1991) Metallothionein Induction in *Mytilus-Edulis* Exposed to Cadmium. *Marine Biology* 108:91-96
- Beyer J, Sandvik M, Hylland K, Fjeld E, Egaas E, Aas E, Skare JU, Goksoyr A (1996) Contaminant accumulation and biomarker responses in flounder (*Platichthys flesus* L) and Atlantic cod (*Gadus morhua* L) exposed by caging to polluted sediments in Sorfjorden, Norway. *Aquatic Toxicology* 36:75-98
- Beyer J, Sandvik M, Skare JU, Egaas E, Hylland K, Waagbo R, Goksoyr A (1997) Time- and dose-dependent biomarker responses in flounder (*Platichthys flesus* L) exposed to benzo[a]pyrene, 2,3,3',4,4',5-hexachlorobiphenyl (PCB-156) and cadmium. *Biomarkers* 2:35-44
- Cajaraville MP, Bebianno MJ, Blasco J, Porte C, Sarasquete C, Viarengo A (2000) The use of biomarkers to assess the impact of pollution in coastal environments of the Iberian Peninsula: a practical approach. *Science Of The Total Environment* 247:295-311

- Cheung APL, Lam THJ, Chan KM (2004) Regulation of Tilapia metallothionein gene expression by heavy metal ions. *Marine Environmental Research* 58:389-394
- Dondero F, Piacentini L, Banni M, Rebelo M, Burlando B, Viarengo A (2005) Quantitative PCR analysis of two molluscan metallothionein genes unveils differential expression and regulation. *Gene* 345:259-270
- Engel DW, Roesijadi G (1987) Metallothioneins: a monitoring tool. In: Vernberg WB, Calabrese A, Thurberg FP, Vernberg JF (eds) *Pollution Physiology of Estuarine Organisms*. University of S. Carolina Press, p 421-438
- George S, Gubbins M, MacIntosh A, Reynolds W, Sabine V, Scott A, Thain J (2004) A comparison of pollutant biomarker responses with transcriptional responses in European flounders (*Platichthys flesus*) subjected to estuarine pollution. *Marine Environmental Research* 58:571-575
- George SG, Olsson P-E (1994) Metallothioneins as indicators of trace metal pollution. In: Kramer KJM (ed) *Biomonitoring of coastal waters and estuaries*. CRC Press, Boca Raton, p 151-178
- Haux C, Förlin L (1989) Selected assays for health status in natural fish populations. In: Landner L (ed) *Chemicals in the aquatic environment*. Springer Verlag, Berlin, p 197-215
- Hogstrand C, Gassman NJ, Popova B, Wood CM, Walsh PJ (1996) The physiology of massive zinc accumulation in the liver of female squirrelfish and its relationship to reproduction. *Journal of Experimental Biology* 199:2543-2554
- Hogstrand C, Haux C (1996) Naturally high levels of zinc and metallothionein in liver of several species of the squirrelfish family from Queensland, Australia. *Marine Biology* 125:23-31
- Hogstrand C, Olsson PE, Haux C (1989) A RADIOIMMUNOASSAY FOR METALLOTHIONEIN IN FISH. *Marine Environmental Research* 28:183-186
- Holth, TF (2004) Effekter av miljøgifter på torsk og skrubbe fra indre Oslofjord. MSc Thesis, Department of Biology, University of Oslo. [in Norwegian]
- Hylland K (1999) Biological effects of contaminants: quantification of metallothionein in fish. *TechmarenvironSci*:1-18
- Hylland K, Sandvik M, Skare JU, Beyer J, Egaas E, Goksoyr A (1996) Biomarkers in flounder (*Platichthys flesus*): An evaluation of their use in pollution monitoring. *Marine Environmental Research* 42:223-227
- Kagi JHR, Schaffer A (1988) Biochemistry of Metallothionein. *Biochemistry* 27:8509-8515
- Klaassen CD, Liu J, Choudhuri S (1999) Metallothionein: An intracellular protein to protect against cadmium toxicity. *Annual Review of Pharmacology and Toxicology* 39:267-294
- Lacorn M, Piechotta G, Simat TJ, Kammann U, Wosniok W, Lang T, Muller WEG, Schroder HC, Jenke HS, Steinhart H (2001) Annual cycles of apoptosis, DNA strand breaks, heat shock proteins, and metallothionein isoforms in dab (*Limanda limanda*): influences of natural factors and consequences for biological effect monitoring. *Biomarkers* 6:108-126
- Masters BA, Kelly EJ, Quaipe CJ, Brinster RL, Palmiter RD (1994) Targeted Disruption of Metallothionein-I and Metallothionein-II Genes Increases Sensitivity to Cadmium. *Proceedings of the National Academy of Sciences of the United States of America* 91:584-588
- Mourgaud Y, Martinez E, Geffard A, Andral B, Stanisiere JY, Amiard JC (2002) Metallothionein concentration in the mussel *Mytilus galloprovincialis* as a biomarker of response to metal contamination: validation in the field. *Biomarkers* 7:479-490
- Nissen-Lie, TR (1997) Årstidsvariasjon i spormetallmetabolismen til skrubbe, *Platichthys flesus* L., fra Hvalerområdet. MSc Thesis, Department of Biology, University of Oslo. [in Norwegian]
- Noël-Lambot F (1976) Distribution of cadmium, zinc and copper in the mussel *Mytilus edulis*. Existence of cadmium-binding proteins similar to metallothioneins. *Experientia* 32:324-325
- Olafson RW, Olsson P-E (1991) Electrochemical detection of metallothionein. In: Riordan JF, Vallee BL (eds) *Metallobiochemistry Part B: metallothionein and related molecules*. Academic Press, London, p 205-213

- Roch M, McCarter JA (1984) Metallothionein Induction, Growth and Survival of Chinook Salmon Exposed to Zinc, Copper and Cadmium. *Bull Environ Contam Toxicol* 32:478-485
- Roesijadi G, Unger ME, Morris JE (1988) Immunochemical quantification of metallothioneins of a marine mollusc. *Can J Fish Aquat Sci* 45:1257-1263
- Rotchell JM, Clarke KR, Newton LC, Bird DJ (2001) Hepatic metallothionein as a biomarker for metal contamination: age effects and seasonal variation in European flounders (*Pleuronectes flesus*) from the Severn Estuary and Bristol Channel. *Marine Environmental Research* 52:151-171
- Ruus A, Hylland K, Green NW (2003) Joint Assessment and Monitoring Programme (JAMP). Biological Effects Methods, Norwegian Monitoring 1997-2001. Report No. 4649-2003, NIVA, Oslo
- Sandvik M, Beyer J, Goksoyr A, Hylland K, Egaas E, Skaare JU (1997) Interaction of benzo[a]pyrene, 2,3,3',4,4',5-hexachlorobiphenyl (PCB-156) and cadmium on biomarker responses in flounder (*Platichthys flesus* L). *Biomarkers* 2:153-160
- Sulaiman N, George S, Burke MD (1991) Assessment of sublethal pollutant impact on flounders in an industrialized estuary using hepatic biochemical indices. *Marine Ecology Progress Series* 68:207-212
- Thompson ED, Mayer GD, Balesaria S, Glover CN, Walsh PJ, Hogstrand C (2003) Physiology and endocrinology of zinc accumulation during the female squirrelfish reproductive cycle. *Comparative Biochemistry and Physiology* 134A:819-828
- Tom M, Chen N, Segev M, Herut B, Rinkevich B (2004) Quantifying fish metallothionein transcript by real time PCR for its utilization as an environmental biomarker. *Marine Pollution Bulletin* 48:705-710
- Viarengo A, Burlando B, Dondero F, Marro A, Fabbri R (1999) Metallothionein as a tool in biomonitoring programmes. *Biomarkers* 4:455-466
- Viarengo A, Lafaurie M, Gabrielides GP, Fabbri R, Marro A, Romeo M (2000) Critical evaluation of an intercalibration exercise undertaken in the framework of the MED POL biomonitoring program. *Marine Environmental Research* 49:1-18
- WGBEC (2005). Report of the ICES working group on biological effects of contaminants (WGBEC). <http://www.ices.dk>.
- Zorita I, Strogyloudi E, Buxens A, Mazon LI, Papathanassiou E, Soto M, Cajaraville MP (2005) Application of two SH-based methods for metallothionein determination in mussels and intercalibration of the spectrophotometric method: laboratory and field studies in the Mediterranean Sea. *Biomarkers* 10:342-359

Chapter 13 (δ -aminolevulinic acid dehydratase)

- Addison RF, Fitzpatrick D, Renton KW (1990) Distribution of d-aminolevulinic acid synthetase and d-aminolevulinic acid dehydratase in liver and kidney of rainbow trout (*Salmo gairdneri*). *Comp Biochem Physiol* 95B:317-319
- Burden VM, Sandheinrich MB, Caldwell CA (1998) Effects of lead on the growth and delta-aminolevulinic acid dehydratase activity of juvenile rainbow trout, *Oncorhynchus mykiss*. *Environmental Pollution* 101:285-289
- Dwyer FJ, Schmitt CJ, Finger SE, Mehrle PM (1988) Biochemical changes in longear sunfish, *Lepomis megalotis*, associated with lead, cadmium and zinc from mine tailings. *J Fish Biol* 33:307-317
- Haux C, Förlin L (1988) Biochemical methods for detecting effects of contaminants on fish. *Ambio* 17:376-381
- Haux C, Larsson Å, Lithner G, Sjöbeck M-L (1986) A field study of physiological effects on fish in lead-contaminated lakes. *Environ Toxicol Chem* 5:283-288
- Hodson PV (1976) Delta-Amino Levulinic Acid Dehydratase Activity of Fish Blood as an Indicator of a Harmful Exposure to Lead. *J Fish Res Board Can* 33:268-271
- Hodson PV, Blunt BR, Spry DJ (1978) Chronic toxicity of water-borne and dietary lead to rainbow trout (*Salmo gairdneri*) in Lake Ontario water. *Water Res* 12:869-878

- Hodson PV, Blunt BR, Spry DJ, Austen K (1977) Evaluation of Erythrocyte Delta-Amino Levulinic Acid Dehydratase Activity as a Short-Term Indicator in Fish of a Harmful Exposure to Lead. *J Fish Res Board Can* 34:501-508
- Hodson PV, Dixon DG, Kaiser KLE (1984) Measurement of Median Lethal Dose as a Rapid Indication of Contaminant Toxicity to Fish. *Environ Toxicol Chem* 3:243-254
- Hodson PV, Dixon DG, Spry DJ, Whittle DM, Sprague JB (1982) Effect of Growth-Rate and Size of Fish on Rate of Intoxication by Waterborne Lead. *Can J Fish Aquat Sci* 39:1243-1251
- Holth TF (2004) Effekter av miljøgifter på torsk og skrubbe fra indre Oslofjord. The university of Oslo
- Hylland K (2004) Biological effects of contaminants: quantification of δ -aminolevulinic acid dehydratase (ALA-D) activity in fish blood. *ICES Techniques in Marine Environmental Science* 34:1-9
- Jackim E (1973) Influence of lead and other metals on fish d-aminolevulinic acid dehydratase activity. *J Fish Res Board Can* 30:560-562
- Johansson-Sjöbeck M-L, Larsson Å (1978) The effect of cadmium on the hematology and on the activity of d-aminolevulinic acid dehydratase (ALA-D) in blood and hematopoietic tissues of the flounder, *Pleuronectes flesus* L. *Environ Res* 17:191-204
- Johansson-Sjöbeck M-L, Larsson Å (1979) Effects of inorganic lead on delta-aminolevulinic acid dehydratase activity and hematological variables in the rainbow trout, *Salmo gairdnerii*. *Arch Environ Contam Toxicol* 8:419-431
- Krajnovic-Ozretic M, Ozretic B (1980) The ALA-D activity test in lead-exposed grey mullet *Mugil cephalus*. *Mar Ecol Prog Ser* 3:187-191
- Larsson Å, Haux C, Sjöbeck M-L (1985) Fish Physiology and Metal Pollution: Results and Experiences from Laboratory and Field Studies. *Ecotoxicol Environ Safety* 9:250-281
- Nakagawa H, Sato T, Kubo H (1995) Evaluation of chronic toxicity of water lead for carp *Cyprinus carpio* using its blood 5-aminolevulinic acid dehydratase. *Fisheries Sci* 61:956-959
- Nakagawa H, Tajima T, Sato T, Kuboyama M (1998) A field study of water lead pollution in fresh water areas of northern Kyushu, based on 5-aminolevulinic acid dehydratase activity and lead concentration in the blood of crucian carp, *Carassius auratus langsdorfii*. *J Fac Agr Kyushu U* 43:209-215
- Ruus A, Hylland K, Green NW (2003) Joint Assessment and Monitoring Programme (JAMP). Biological Effects Methods, Norwegian Monitoring 1997-2001. Report No. 4649-2003, NIVA, Oslo
- Schmitt CJ, Dwyer FJ, Finger SE (1984) Bioavailability of Pb and Zn from mine tailings as indicated by erythrocyte d-aminolevulinic acid dehydratase (ALA-D) activity in suckers (Pisces: Catostomidae). *Can J Fish Aquat Sci* 41:1030-1040
- Schmitt CJ, Whyte JJ, Brumbaugh WG, Tillitt DE (2005) Biochemical effects of lead, zinc, and cadmium from mining on fish in the Tri-States District of northeastern Oklahoma, USA. *Environ Toxicol Chem* 24:1483-1495
- Schmitt CJ, Wildhaber ML, Hunn JB, Nash T, Tieger MN, Steadman BL (1993) Biomonitoring of Lead-Contaminated Missouri Streams with an Assay for Erythrocyte Delta-Aminolevulinic-Acid Dehydratase Activity in Fish Blood. *Arch Environ Con Tox* 25:464-475
- Sordyl H, Osterland A (1990) Investigation of factors influencing d-aminolevulinic acid dehydratase activity (ALA-D) of rainbow trout. *Zool Jb Physiol* 94:445-460