

Accepted Manuscript

This is an Accepted Manuscript of the following article:

S. Jannicke Moe, Anne Lyche Solheim, Hanna Soszka, Małgorzata Gołub, Andrzej Hutorowicz, Agnieszka Kolada, Joanna Picińska-Fałtynowicz, Witold Białokoz. Integrated assessment of ecological status and misclassification of lakes: The role of uncertainty and index combination rules. *Ecological Indicators*. Volume 48, 2015, pages 605-615, ISSN 1470-160X.

The article has been published in final form by Elsevier at

<http://dx.doi.org/10.1016/j.ecolind.2014.08.018>

© 2104. This manuscript version is made available under the

CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

It is recommended to use the published version for citation.

Manuscript Number: ECOLIND-3500R1

Title: Integrated assessment of ecological status and misclassification of lakes: the role of uncertainty and index combination rules

Article Type: Research Paper

Keywords: Biological quality element
Integrated lake assessment
Modeling
Uncertainty
Water Framework Directive
Waterbody classification

Corresponding Author: Dr. Jannicke Moe, PhD

Corresponding Author's Institution: Norwegian Institute for Water Research

First Author: Jannicke Moe, PhD

Order of Authors: Jannicke Moe, PhD; Anne Lyche Solheim; Hanna Soszka; Małgorzata Gołub; Andrzej Hutorowicz; Agnieszka Kolada; Joanna Picińska-Fałtynowicz

Abstract: The European Water Framework Directive (WFD) requires that the ecological status of waterbodies is assessed using multiple biological quality elements (BQEs) that are combined into a single status class. The recommended combination rule (the "one-out, all-out" rule; OOA) has been criticized for being unreasonably conservative and for being sensitive to uncertainty. In this study, the objective was to compare the sensitivity to uncertainty of four different combination rules: (1) OOA, (2) OOA with exclusion of one element (3) average and (4) weighted average. Index values for 5 BQEs (phytoplankton, phytobenthos, macrophytes, macroinvertebrates and fish) sampled from 10 lakes in the Wel River catchment in Poland were used to classify the lakes according to the OOA and the three alternative combination rules. Based on the mean and (where possible) standard deviation of these index values, we modelled the risk of misclassification by simulating 10,000 resamples for each BQEs in each lake, classifying each resample and calculating the proportion of misclassified resamples under each combination rule. For individual BQEs, the risk of misclassification increased both with higher uncertainty and with the proximity of the index value to a class boundary. Under the OOA rule, the risk of misclassification was more biased towards worse status ("underclassification") than towards better status. Furthermore, risk of underclassification was more affected by uncertainty under the OOA rule compared with the alternative combination rules. This analysis has demonstrated the weaknesses associated with the OOA rule for integration of BQEs for lake classification. However, the alternative combination rules are associated with other shortcomings, such as the need for subjective judgement, and involve a higher risk of not protecting the most sensitive BQE and thus the whole ecosystem. We recommend that future versions of instructions for WFD implementation consider alternatives to the OOA combination rule, and provide guidelines for weighting of individual BQEs.

Response to Reviewers: All replies to the reviewers are given in a separate document.

Highlights

- The "one-out, all-out" (OOAO) rule for combining assessment results for different biological quality elements is more prone to underestimation of the "correct" ecological status than alternative combination rules
- The OOA rule's tendency of underestimation of ecological status increases with the index values' uncertainty
- Analysis of misclassification is complicated by the fact that the rate of misclassification inherently increases with the index values' proximity to status class boundaries

1 **Title:** Integrated assessment of ecological status and misclassification of lakes: the role of
2 uncertainty and index combination rules

3
4
5
6
7 **Authors:**

8
9 S. Jannicke Moe^{a,*}, Anne Lyche Solheim^a, Hanna Soszka^b, Małgorzata Gołub^b,
10
11 Andrzej Hutorowicz^c, Agnieszka Kolada^b, Joanna Picińska-Fałtynowicz^d, Witold Białokoz^e

12
13
14
15
16
17 ^a *Norwegian Institute for Water Research, Gaustadalléen 23, 0349 OSLO, Norway*

18
19 ^b *Institute of Environmental Protection, National Research Institute, Department of*
20
21 *Freshwater Assessment Methods and Monitoring, Kolektorska 4, 01-692 Warszawa, Poland*

22
23
24 ^c *The Stanisław Sakowicz Inland Fisheries Institute, Department of Hydrobiology,*
25
26 *Oczapowskiego 10, 10-719 Olsztyn, Poland*

27
28
29 ^d *Institute of Meteorology and Water Management, National Research Institute, Wrocław*
30
31 *Branch, Parkowa 30, 52-616 Wrocław, Poland*

32
33
34 ^e *The Stanisław Sakowicz Inland Fisheries Institute, Department of Lake Fisheries, Rajska 2,*
35
36 *11-500 Giżycko, Poland*

37
38
39
40
41 **E-mail addresses:** jmo@niva.no, als@niva.no, hasoszka@ios.edu.pl, mgolub@ios.edu.pl,
42
43 ahut@infish.com.pl, akolada@ios.edu.pl, joanna.faltynowicz@imgw.pl,
44
45
46 wbialokoz@infish.com.pl

47
48
49
50
51 *** Corresponding author:**

52
53 Jannicke Moe, NIVA, Gaustadalléen 21, NO-0349 Oslo, Norway

54
55
56 jmo@niva.no

57
58 Telephone: +47 908 98 108, Fax: +47 22 18 52 00

27 ABSTRACT

28 The European Water Framework Directive (WFD) requires that the ecological status of
29 waterbodies is assessed using multiple biological quality elements (BQEs) that are combined
30 into a single status class. The recommended combination rule (the "one-out, all-out" rule;
31 OOA) has been criticized for being unreasonably conservative and for being sensitive to
32 uncertainty. In this study, the objective was to compare the sensitivity to uncertainty of four
33 different combination rules: (1) OOA, (2) OOA with exclusion of one element (3) average
34 and (4) weighted average. Index values for 5 BQEs (phytoplankton, phytobenthos,
35 macrophytes, macroinvertebrates and fish) sampled from 10 lakes in the Wel River catchment
36 in Poland were used to classify the lakes according to the OOA and the three alternative
37 combination rules. Based on the mean and (where possible) standard deviation of these index
38 values, we modelled the risk of misclassification by simulating 10,000 resamples for each
39 BQEs in each lake, classifying each resample and calculating the proportion of misclassified
40 resamples under each combination rule. For individual BQEs, the risk of misclassification
41 increased both with higher uncertainty and with the proximity of the index value to a class
42 boundary. Under the OOA rule, the risk of misclassification was more biased towards worse
43 status ("underclassification") than towards better status. Furthermore, risk of
44 underclassification was more affected by uncertainty under the OOA rule compared with the
45 alternative combination rules. This analysis has demonstrated the weaknesses associated with
46 the OOA rule for integration of BQEs for lake classification. However, the alternative
47 combination rules are associated with other shortcomings, such as the need for subjective
48 judgement, and involve a higher risk of not protecting the most sensitive BQE and thus the
49 whole ecosystem. We recommend that future versions of instructions for WFD
50 implementation consider alternatives to the OOA combination rule, and provide guidelines
51 for weighting of individual BQEs.

53	<i>Keywords:</i>
54	Biological quality element
55	Integrated lake assessment
56	Modeling
57	Uncertainty
58	Water Framework Directive
59	Waterbody classification
60	
61	<i>Abbreviations:</i>
62	BQE: biological quality element
63	EQR: ecological quality ratio
64	nEQR: normalised ecological quality ratio
65	OOAO: One-out, all-out (combination rule)
66	WFD: Water Framework Directive
67	

68 1. Introduction

69
70 The Water Framework Directive (WFD; EC 2000) of the European Union requires that
71 member states must assess the ecological status of their surface waterbodies, including lakes.
72 Across Europe, WFD-compliant national classification systems have been developed and
73 adapted for assigning waterbodies to one of five classes of ecological status (high, good,
74 moderate, poor and bad) (Hering et al., 2010). The WFD further requires that all waterbodies
75 obtain good ecological status by 2015, and consequently all waterbodies found to be in
76 moderate or worse status must be restored. Moreover, the WFD states that estimates of
77 confidence and precision attained by the monitoring system should be provided in river basin
78 management plans (Annex V, Section 1.3.4). Since restoration measures can be expensive, the
79 uncertainty associated with waterbody classification should be of high interest for water
80 resource management (Højberg et al., 2007; Irvine, 2004). If a lake in good or better status is
81 wrongly classified as having less-than-good status ("underclassified"), money may be wasted
82 on restoration measures that were not strictly needed (Prato et al., 2014). On the other hand, if
83 a lake in less-than-good status is wrongly classified as good or better ("overclassified"), the
84 ecosystem quality and services may be compromised.

85
86 Classification of ecological status of lakes should be based on a set of biological quality
87 elements (BQEs) representing main ecosystem components, i.e. (1) phytoplankton, (2)
88 macrophytes and phytobenthos, (3) benthic invertebrate fauna (here called
89 "macroinvertebrates") and (4) fish (WFD, Annex V, Section 1.2.2). The WFD states that the
90 policy should be based on the precautionary principle (§11); the idea of this principle is that if
91 at least one component of ecosystem is impaired, this indicates that something is wrong in the
92 ecosystem (waterbody) as a whole. Moreover, the WFD requires that the ecological status

93 class for a waterbody "shall be represented by the lower of the values for the biological and
94 physico-chemical monitoring results for the relevant quality elements" (Annex V, Section
95 1.4.2 (i)). This implies that the status is determined by either the combined biological
96 monitoring result or by the physical-chemical monitoring result (the lower of the two).
97 However, the directive does not specify how to combine the values of multiple BQEs into one
98 biological monitoring result. The guidance on classification provided by the Common
99 Implementation Strategy for the WFD (EC 2005) has recommended the method known as
100 "One-out, all-out" (OOAO): the waterbody status is determined by the BQE with the worst
101 status. However, based on comparison with alternative rules for integrating BQEs, such as
102 (weighted) average, median or other weight-of-evidence approaches, several authors have
103 stated that the OOAO tend to result in a stricter classification than what seems reasonable
104 (Alahuhta et al., 2009; Borja and Rodriguez, 2010; Caroni et al., 2013; Gottardo et al., 2011;
105 Hering et al., 2010; Moss et al., 2003; Nõges et al., 2009; Nõges and Nõges, 2006; Prato et al.,
106 2014; Rask et al., 2010; Sutela et al., 2013; Søndergaard et al., 2005). Another concern with
107 the OOAO method is that higher uncertainty in index values tend to result in even stricter
108 classification (Caroni et al., 2013; EC (European Commission), 2005; Nõges et al., 2009;
109 Sandin, 2005).

110
111 Uncertainty in biological index values results from many sources, including natural temporal
112 and spatial variation and sampling variation (see Clarke, 2013). The quantification of sources
113 of uncertainty in index values and their significance for status classification have been
114 addressed in many studies (Carvalho et al., 2013; Clarke and Hering, 2006; Kelly et al.,
115 2009b; Thackeray et al., 2013). Nevertheless, few studies have investigated the role of joint
116 uncertainty of indices when several BQEs are integrated (but see Caroni et al., 2013). There is
117 therefore a need for more research on how the OOAO and other BQE combination rules

118 perform in waterbody classification based on real data under different levels of sampling
119 uncertainty.

120
121 In our study, we have analysed the effects of joint uncertainty for five BQEs (phytoplankton,
122 phytobenthos, macrophytes, macroinvertebrates and fish) sampled from 10 lakes in Poland.
123 The analysis was based on simulations of index values for all BQEs with three levels of
124 uncertainty (section 3.1), and application of four different combination rules (section 2.3) for
125 the resulting BQE status classes. The objective of this paper was to address the following
126 question: How does increasing levels of uncertainty affect the risk of misclassification of
127 lakes under different BQE combination rules? To answer this question, we also investigated
128 how uncertainty in index values affect the risk of misclassification at the BQE level, and how
129 this risk was transferred to the whole-lake level under the different combination rules.

130

131 **2. Materials and methods**

132 133 *2.1. Data*

134
135 The study area is the catchment of the lowland river Wel in central Poland, with a surface area
136 of 822 km². Surface waters in the Wel catchment are affected mainly by eutrophication due to
137 agricultural runoff (app. 60% of areas of extensive agriculture in the catchment) and also by a
138 few point sources of organic pollution. Ten lakes with surface area above 0.5 km² are located
139 in this catchment (Fig. 1, Table 1). The biological data used in this study were collected from
140 all of the ten lakes in 2009 during the Polish-Norwegian project deWELopment (Soszka,
141 2011).

142

143 2.2. *Biological index values and classification system*

1
2 144
3
4
5 145 In this study, each biological quality element (BQE) was represented by one index, as follows.

6
7 146 - Phytoplankton: Phytoplankton Metric for Polish Lakes (Hutorowicz et al., 2011).

8
9 147 - Phytobenthos: Diatom index for lakes (phytobenthos) (Picińska-Fałtynowicz, 2011).

10
11
12 148 - Macrophytes: Ecological State Macrophyte Index (Kolada et al., 2011)

13
14 149 - Macroinvertebrates: Benthic Quality Index based on Chironomid Pupal Exuviae Technique
15
16 (macroinvertebrates; based on Ruse, 2010) (Gołub et al., 2011).

17 150
18
19 151 - Fish: Lake Fish Index N2 (Białokoz and Chybowski, 2011).

20
21
22 152 For each index, the sampling method, calculation, the responses to eutrophication pressure
23
24 153 gradients as well as classification scheme are described in the given references. For

25
26 154 phytoplankton and macrophytes, respectively, a full description of the national assessment

27
28
29 155 methods are given in the Technical Reports from the Intercalibration phase 2 (Phillips et al.,

30
31 156 2014; Portielje et al., 2014). Although the WFD defines phytobenthos and macrophytes as one

32
33
34 157 BQE, the two organism groups are treated as two separate BQEs in this paper. The reason is

35
36 158 that Poland, like most countries in the Central-Baltic region, has chosen to develop separate

37
38
39 159 assessment methods for macrophytes and phytobenthos (Kelly et al., 2009a), and no

40
41 160 integration rules exist at the moment (Portielje et al., 2014, Table 4.4). Moreover, changing

42
43
44 161 environmental conditions may affect macrophytes and phytobenthos indices differently due to

45
46 162 the differences in generation time and dispersal rate; therefore these organism groups may

47
48
49 163 provide different information about ecosystem stability (Schneider et al., 2012).

50
51 164
52
53 165 The ecological classification system used in this study (Soszka, 2011) comprises, for each

54
55
56 166 biological index, a *reference condition* representing the index value assumed for lakes

57
58 167 undisturbed by anthropogenic impact, and *class boundaries* defining the index values on the

168 borders between the five ecological status classes (high, good, moderate, poor and bad). More
169 information on the methods used for setting reference conditions and class boundaries for the
170 Polish classification system is available in the WISER database on national assessment
171 methods (<http://www.wiser.eu/results/method-database>; Birk et al., 2012), for all BQEs
172 except macroinvertebrates. The full ecological classification system includes also physico-
173 chemical variables, which were not included here. For each index, as required by the WFD,
174 the ecological quality ratio (EQR) was calculated as the index value divided by the reference
175 condition value. The resulting indices in EQR scale have range 0-1 (Appendix A, Table A.1).
176 Likewise, the class boundaries for each index were converted to EQR scale (Table A.2) by
177 division by the respective reference condition value. Note that the class boundaries are non-
178 evenly spaced for all BQEs except phytoplankton (Table A.2); this is essential in comparison
179 of the BQE classifications. For example, a BQE with narrow class width for good status (e.g.
180 macrophytes, class width = 0.17) may be more susceptible to bias in the assessment of good
181 status compared to a BQE with a wider class (e.g. fish, class width = 24).

182
183 To facilitate comparison of index values for different BQEs, the EQR values (Table A.1) were
184 normalised (nEQR) by a piecewise linear transformation procedure (Caroni et al., 2013). The
185 normalisation is based on the distance from the index value (in EQR scale) to the nearest class
186 boundaries (Eq. 1):

$$\text{nEQR} = (\text{EQR} - \text{lower_EQR}) * \frac{(\text{upper_nEQR} - \text{lower_nEQR})}{(\text{upper_EQR} - \text{lower_EQR})} + \text{lower_nEQR}, \text{ (Eq. 1)}$$

187
188
189 where lower_EQR and upper_EQR are lower and upper class boundaries in EQR scale for the
190 given index (Table A.2), and lower_nEQR and upper_nEQR lower and upper class

193 boundaries in normalised EQR scale (high/good = 0.8, good/moderate = 0.6, moderate/poor=
194 0.4, poor/bad= 0.2). The transformation to nEQR scale ensures standard class widths and
195 boundaries for all BQEs (see also EC (2011), Fig. 12)¹. This way, one can infer directly from
196 each nEQR value (Table 2) both the status class and the distance to the nearest class
197 boundaries.

2.3. BQE combination rules

201 Four alternative rules for combining the ecological status of multiple BQEs were applied in
202 this study (Table 2).

204 *Rule 1: "OOAO"* (One-out, all-out). The status of the lake was determined by the lowest status
205 of all the BQEs.

206 *Rule 2: "OOAO-E"* (One-out, all-out after exclusion of one BQE). This combination rule is
207 recommended in cases where one BQE has high variability or is for other reasons associated
208 with low confidence (EC 2005). Here, macroinvertebrates were excluded (see section 3.1).

209 *Rule 3: "Avg"* (Average). Following the WISERBUGS method (Clarke, 2013, see section
210 3.1.), the status class for each BQE was converted to an integer (H=1, G=2, M=3, P=4, B=5),
211 and the arithmetic average for all BQEs was calculated. This conversion implies that the
212 proximity of an index value to class boundaries is ignored, which is not ideal. We
213 nevertheless chose to base the average on integers instead of the actual nEQR values, to make
214 our results comparable with other studies using WISERBUGS (e.g., Caroni et al., 2013;
215 Kolada et al., 2013). If the average was halfway between two classes, it was assigned to the
216 worse of the two classes.

¹ A more detailed illustration of normalisation of EQR values can be found in the European Environment Agency's Data Dictionary for Lakes: http://forum.eionet.europa.eu/nrc-eionet-freshwater/library/wise_reporting_2011/biological_reporting/biologydd_20110617jpg

1
2 217 *Rule 4: "Avg-W" (Weighted average).* The status classes were converted to numeric values as
3
4 218 for Rule 3, but the class value for each BQE was multiplied by a weight inversely related to
5
6 219 the uncertainty assumed for the BQE (see section 3.1). In this study macroinvertebrates were
7
8 220 down-weighted (weight = 10%) relative to the other BQEs (each 22.5%).

9
10 221 The notation "OOAO(-E)" will be used when the two rules OOAO and OOAO-E are
11
12 222 considered jointly, and "Avg(-W)" for the two rules Avg and Avg-W considered jointly.

13
14 223

15 16 224 **3. Calculation: the WISERBUGS method for estimating risk of misclassification**

17
18
19 225

20
21 226 For analysing the risk of misclassification, we adopted the WISERBUGS method (Clarke,
22
23 227 2013). The method assumes that index values used for ecological classification of a
24
25 228 waterbody follow a normal distribution that can be specified by the mean and standard
26
27 229 deviation of replicated samples. The standard deviation (SD) then represents the sampling
28
29 230 uncertainty of the index. The estimated mean and SD defines a normal probability
30
31 231 distribution, from which resamples of the index can be simulated by random drawing. This
32
33 232 method recognises that the true status class of a waterbody is unknown, but considers the
34
35 233 assessment based on the measured index value (cf. Table 2) as the "correct" class.

36
37 234 Misclassification of the simulated index values is defined as assignment to any other class
38
39 235 than the "correct" class. The risk of misclassification is thus based on the precision of the
40
41 236 index values, which is measured by standard deviation (SD), but does not consider the
42
43 237 accuracy (the proximity to the unknown correct status) (Clarke, 2013).

44
45
46 238

47 48 239 *3.1. Uncertainty in index values*

49
50
51 240

241 Following the WISERBUGS method, characterisation of the probability distribution and
242 estimation of sampling variation for indices should ideally be based on a large number of
243 properly replicated samples, which are not available in most biological studies including ours.
244 However, our aim was not to predict the exact risk of misclassification, but to compare the
245 relative risk of misclassification for different levels of uncertainty. Therefore a pragmatic
246 approach was taken: where possible, the SDs for each BQE was based on multiple samples
247 from the same lake (taken at different stations, in different seasons or by different personnel),
248 and calculated as pooled SD ("SD1") for all lakes. This uncertainty measure may be
249 considered to include spatial and/or temporal variation in addition to sampling variation. For
250 phytoplankton (SD1 = 0.056) and phytobenthos (SD1 = 0.046), the SD1 was calculated from
251 4 lakes with 2-3 stations sampled once in summer and once in autumn, respectively. For
252 macrophytes (SD1 = 0.051), the SD1 was calculated from 10 lakes surveyed by 2-3 different
253 persons, once in the peak of the growing season. The faunal indices had insufficient samples
254 for calculation of SD. However, since macroinvertebrates often had lower nEQR values than
255 the other BQEs, we were particularly interested in how the exclusion or down-weighting of
256 this BQE would affect the overall assessment and risk of misclassification. The
257 macroinvertebrate index is used in national classification but the assessment system has not
258 yet been intercalibrated with the systems of other Central-Baltic countries (Böhmer et al.,
259 2014), therefore this index was associated with lower confidence than the botanical indices.
260 To reflect this lower confidence, we chose as a pragmatic solution to assign higher uncertainty
261 for the macroinvertebrate index (SD1 = 0.10) than for the other BQEs. For fish, for simplicity,
262 the sampling uncertainty was set to the same level as the botanical elements (SD1 = 0.05).

263

264 *3.2. Resampling and probability of misclassification of BQEs and lakes*

265

266 The modelling approach in this study follows the WISERBUGS method (Clarke, 2013):
267 stochastic simulation of biological index values with sampling uncertainty, and calculation of
268 misclassification under different combination rules. Three levels of uncertainty were applied,
269 denoted SD1, SD2 and SD3. Uncertainty level SD1 corresponds to the estimated or assumed
270 SD for the respective indices, as described above. For level SD2, the SD for each BQE was
271 multiplied by $\sqrt{2}$ (i.e., the variance was doubled). In level SD3, correspondingly, the SDs
272 were multiplied by $\sqrt{3}$. We simulated 10,000 resamples for each BQE in each lake, classified
273 each resample and calculated the proportion of misclassified resamples compared with the
274 "correct" class. The simulation routine was programmed in R version 2.14.1 (R Development
275 Core Team, 2011), and can be summarised in the following steps, for each lake.

- 276
277 1. For each BQE and each SD level, assume that the index values follow a normal probability
278 distribution $N \sim (\text{mean}, \text{SD})$ defined by the mean index value for the lake (in EQR scale;
279 Table A.1) and the pooled SD (section 3.1).
- 280 2. For each BQE and each SD level, simulate 10 000 samples (index values) drawn randomly
281 from their respective probability distributions $N \sim (\text{mean}, \text{SD})$.
- 282 3. For each simulation, assess the status class for each BQE based on their respective index
283 values and class boundaries.
- 284 4. For each simulation and each BQE combination rule, assess the integrated status class for
285 the lake according to the obtained BQE status classes.
- 286 5. For each SD level and each combination rule, calculate the proportion of "correct"
287 classification as the number of simulations with the same class as obtained for the input data
288 with the same combination rule (Table 1) divided by the total number of simulations. The
289 remaining proportion of simulations represents the probability of misclassification (under the
290 given combination rule).

291

1

2 292 For example, for macrophytes in Lake Kiełpińskie, the mean EQR is 0.51 (Table A.1) and the

3

4 293 pooled SD1 is 0.051 (section 3.1). Simulation of 10 000 resamples from the normal

5

6 294 distribution $N \sim (\text{mean}=0.51, \text{SD}=0.051)$ resulted in 60.03% resamples in high class and

7

8 295 39.96% in good class, as displayed in Fig. 2a (leftmost bar). Since the "correct" class in this

9

10 296 case is high (cf. Table 2), the probability of misclassification is 39.96% (Fig. 2b, leftmost

11

12 296 case is high (cf. Table 2), the probability of misclassification is 39.96% (Fig. 2b, leftmost

13

14 297 bar).

15

16 298

17

18 298

19

19 299 *3.3. Cross-lake comparisons of risk of misclassification*

20

21 300

22

23 300

24

24 301 The risk misclassification for the ten lakes combined was analysed by linear models, with the

25

26 302 response variable being the number of misclassified simulated resamples for each lake (as

27

28 303 described in section 3.2, step 5). Significant difference in misclassification among BQEs was

29

30 304 tested both with BQE as a single predictor variable (one-way ANOVA) and with SD as a

31

32 305 continuous co-variable (ANCOVA). The dataset used for this test comprised the number of

33

34 306 misclassifications in the 10 lakes x 5 BQEs x 3 SD levels (in total 150 records). Likewise,

35

36 307 difference in misclassification among combination rules was tested both with combination

37

38 308 rule as single predictor variable and with SD level as a co-variable (dataset: 10 lakes x 4

39

40 309 combination rules x 3 SD levels; in total 120 records). In addition, the number of over- and

41

42 310 underclassifications were also used as alternative response variables. Pairwise comparison of

43

44 311 the number of misclassifications between BQEs and between combination rules was

45

46 312 performed by Tukey's "honestly significant difference" method (using the R function

47

48 313 "TukeyHSD").

49

50 314

51

52 314

53

54 314

55

56 314

57

58 314

59

60 314

61

62 314

63

64 314

65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

315 Ecological status assessment sensu WFD should in principle be applied to waterbodies (and
316 their components), not to larger geographical levels such as catchment. Nevertheless, to
317 describe more general patterns for the whole catchment in this case study, an "aggregated
318 class" for all lakes combined was assigned to each BQE and each combination rule (i.e., the
319 class with the highest proportion of resamples for all lakes combined). For each BQE and
320 combination rule, the number of misclassifications for all lakes combined was calculated as
321 the total number of resamples for all lakes deviating from the correct "aggregated class".

323 **4. Results and Discussion**

324 *4.1. Cross-lake patterns in status classification: effects of BQE combination rule*

326
327 Averaging the status class of individual BQEs generally resulted in higher status than
328 applying the OOAo rule (nine out of ten lakes; Table 2), as expected. Different combination
329 rules for BQE classes have been explored and compared to the OOAo in several other
330 studies, such as average (Caroni et al., 2013; Nõges and Nõges, 2006; Sutela et al., 2013),
331 median (Alahuhta et al., 2009; Caroni et al., 2013; Rask et al., 2010), and weight-of-evidence
332 approaches or decision trees (Borja et al., 2009; Gottardo et al., 2011; Veríssimo et al., 2013).
333 In each case, the alternative rule has given equal or better classification than the OOAo.
334 Many of the authors have expressed concerns that the OOAo seems too conservative,
335 especially when several BQEs are used. For example, the two largest lakes in Estonia
336 (Võrtsjärv and Peipsi) both obtained moderate status, while more subjective expert-based
337 estimates suggest that the status should be good (Nõges and Nõges, 2006).

339 Excluding macroinvertebrates (rule OAO-E) improved the "correct" class compared with
1
2 340 the OAO for only two lakes (Table 2; Lake Grądy and Lake Hartowieckie). In these two
3
4 341 lakes, the macroinvertebrates had the lowest status. Correspondingly, down-weighting
5
6
7 342 macroinvertebrates when averaging the BQEs improved the "correct" class for only two lakes
8
9 343 (Table 2; Lake Dąbrowa Mała and Lake Grądy). For Lake Dąbrowa Mała there was large
10
11 344 disagreement among the BQEs, ranging from high to poor. Therefore, down-weighting one of
12
13
14 345 the two poor BQEs was sufficient to shift the "correct" class from moderate to good.
15
16

17 346

19 347 *4.2. Misclassification of individual BQEs: effects of uncertainty*

22 348

24 349 Higher SD levels generally increased the risk of misclassification at the BQE level (e.g., Lake
25
26 350 Kiełpińskie, Fig. 2b), as could be expected. However, the probability distribution across status
27
28
29 351 classes for the simulated samples (Fig. 2a) was also determined by the proximity of the index
30
31 352 value to a class boundary. The proximity to a class boundary in normalised EQR scale can be
32
33
34 353 inferred from the normalised EQR values in Table 2. For Lake Zarybinek, for example, the
35
36 354 nEQR of phytoplankton and macrophytes (0.39 and 0.37, respectively) were just below the
37
38
39 355 moderate/poor boundary (0.4). This was reflected in the simulated resamples and resulting
40
41 356 classification at the BQE level (Fig. 3a): the two mentioned BQEs had almost equal
42
43
44 357 probability of assessment to moderate or poor class. Consequently, the probability of
45
46 358 misclassification (Fig. 3b) was high (>40%) for these BQEs. For these BQEs where
47
48
49 359 misclassification was already high due to the proximity to a class boundary, higher SD
50
51 360 typically increased this risk only slightly.

53 361

56 362 The importance of uncertainty in index values for risk of misclassification at BQE level, as
57
58 363 demonstrated here, has also been clearly demonstrated in previous studies (Caroni et al.,
59
60
61
62
63
64
65

364 2013; Clarke et al., 2006; Kelly et al., 2009b; Ruse, 2010; Szoszkiewicz et al., 2007). The
1
2 365 estimated or assumed sampling uncertainty for the BQEs in this study (approx. 0.05 - 0.10)
3
4 366 were based on few samples, but correspond well to SD levels estimated for index values in
5
6
7 367 other studies (with similar index scale; 0-1). Examples include invertebrates in rivers (SD
8
9 368 0.058-0.065; Clarke et al., 2006), invertebrates in lakes (SD 0.032-0.094; Caroni et al., 2013),
10
11 369 diatoms in rivers and lakes (temporal variation; SD approx. 0-0.1; Kelly et al., 2009b). The
12
13 370 SD levels in this study can therefore be considered to be within a realistic range for sampling
14
15 371 uncertainty. For monitoring and classification in practice, index values will also be affected
16
17 372 by other sources of uncertainty (e.g. natural temporal variation in the ecosystem). The higher
18
19 373 levels of uncertainty used in the simulations (SD2 up to 0.20) might be considered a
20
21 374 conservative estimate of other uncertainty sources as well.
22
23
24
25
26
27
28

29 375
30 376 The importance of an index value's proximity to class boundaries for the risk of
31
32 377 misclassification of the BQE has also been demonstrated in numerous other studies
33
34 378 (Carstensen, 2007; Clarke and Hering, 2006; Kelly et al., 2009b; Kolada et al., 2013; Ruse,
35
36 379 2010; Szoszkiewicz et al., 2007). However, although the proximity to a class boundary
37
38 380 represents a source of uncertainty for the classification, this factor is not an error that can be
39
40 381 reduced. Thus, instead of defining the status as one class (e.g. poor for phytoplankton in Lake
41
42 382 Zarybinek, Fig. 3a), one might consider the proximity to class boundaries and describe the
43
44 383 status as "poor-to-moderate", or in probabilistic terms (e.g. 60 % poor and 40% moderate). A
45
46 384 more advanced approach - a fuzzy inference system - was used by Gottardo et al. (2011): they
47
48 385 considered also uncertainty in the class boundaries and assigned the membership of each
49
50 386 index to two neighbouring classes, expressed by percentages. If the status of a BQE is defined
51
52 387 as belonging to two classes in such a probabilistic way, the very concept of misclassification
53
54 388 should be reconsidered.
55
56
57
58
59
60
61 389

390 *4.3. Misclassification of individual lakes: effects of BQE combination rule and uncertainty*

391
392 As for individual BQEs, the misclassification of whole lakes increased with the level of SD.
393 However, the effects of SD for individual lakes were confounded by the effects of proximity
394 of index values to class boundaries. For Lake Kiełpińskie, the pattern of misclassification
395 under the OOA(-E) rules (Fig. 2d) reflected the pattern of the worst BQE (macrophytes and
396 macroinvertebrates; Fig. 2b). Under the Avg(-W) rules, in contrast, the risk of
397 misclassification was very low (Fig. 2d), reflecting the fact that most index values were far
398 from the class boundaries (cf. Fig. 2a and Table 2). For Lake Zarybinek, in comparison, the
399 Avg(-W) rules resulted in a high degree of overclassification compared with the OOA(-E)
400 rules (Fig. 3d); this reflects that three of the BQEs were close to an upper class boundary (as
401 described above; Fig 3c). For Lake Dąbrowa Mała, where several BQEs had almost equal
402 probability of two neighbouring classes (Fig. 4a), and thus high risk of misclassification (Fig.
403 4b), the "correct" lake class was altered by changed weighting in the combination rule (Fig.
404 4c). In this case, the large difference in risk of misclassification for Avg-W vs. Avg (Fig. 4d)
405 was due to the shift in the "correct" class.

406
407 The currently recommended modification of the OOA rule - excluding the BQE with
408 highest uncertainty (EC 2005) - will automatically reduce the risk of underclassification.
409 However, there is also a risk that the excluded BQE actually is the most vulnerable
410 component, and that excluding this element will result in e.g. good status when moderate
411 status would be more appropriate, and therefore will fail to protect this BQE. More generally,
412 from a scientific point of view, discarding available information (even if uncertain) is not the
413 best means for obtaining a more reliable result. Moreover, if one BQE is routinely excluded
414 from status classification, it is more likely that it will eventually be excluded from monitoring

1
2 415 programmes (Søndergaard et al., 2005); this loss of information may in the long run increase
3 416 the risk of inappropriate management decisions.

4
5 417
6
7 418 The average combination rules may seem favourable from a statistical point of view, because
8
9 419 they make better use of all available information, and give more robust and balanced results
10
11 420 also under high uncertainty. However, these combination rules will not necessarily ensure the
12
13 421 protection of the whole ecosystem, especially in cases where there is large disagreement
14
15 422 among the BQEs (e.g. Kiełpińskie, Fig. 2). Down-weighting of BQEs with low confidence
16
17 423 may reduce the risk of misclassification (Fig. 6b), but may also fail to protect the most
18
19 424 sensitive BQEs (e.g. Dąbrowa Mała, Fig. 4). Moreover, weighted average or other weight-of-
20
21 425 evidence approaches (e.g., Gottardo et al., 2011) are not straightforward to implement,
22
23 426 because the choices will need to be justified, and there is a risk that the weighting can be
24
25 427 manipulated in order to obtain desired results. Guidelines for weighting of different BQEs,
26
27 428 e.g. based on uncertainty or other measures of confidence, would therefore be useful.
28
29
30
31
32
33

34 429 35 36 430 *4.4. Cross-lake patterns in misclassification: effects of uncertainty under different* 37 38 431 *combination rules*

39
40
41 432
42
43 433 The "aggregated class" of individual BQEs for all lakes combined (see section 3.3) ranged
44
45 434 from poor to good (Fig. 5a), and was not affected by the uncertainty level in index values
46
47 435 (SD1 - SD3). Nevertheless, the uncertainty levels affected the probability distribution across
48
49 436 status classes, and hence the risk of misclassification (Fig. 5b).
50

51 437
52
53 438 The inclusion of uncertainty also revealed a more nuanced picture of the overall effects of
54
55 439 combination rules. Although exclusion of macroinvertebrates generally did not alter the
56
57
58
59
60
61
62
63
64
65

1 440 "correct" lake class compared to the OOAO combination rule (Table 2), it shifted the overall
2 441 aggregated distribution of simulated classes towards higher status (Fig. 5c). Consequently, the
3
4 442 overall rates of misclassification were slightly reduced by this modification of the
5
6
7 443 combination rule (Fig. 5d).

8
9 444
10
11 445 Analysis of among-lake variation in misclassification for individual BQEs showed significant
12
13 446 effects of both BQE ($F_{4,144} = 10.58$, $p < 0.001$) and SD level ($F_{1,144} = 29.23$, $p < 0.001$). The
14
15 447 Tukey HSD test (Fig. 6a) revealed that on average, misclassification was significantly higher
16
17 448 for macroinvertebrates and macrophytes than for phyto**ben**thos and fish, with phytoplankton
18
19 449 in-between. The extra high SD for assumed macroinvertebrates (approximately twice as high
20
21 450 as for the other BQEs) did not result in a correspondingly high rate of misclassification for
22
23 451 this BQE; this indicates that the proximity to class boundaries is an equally important factor
24
25 452 for the risk of misclassification. The bias towards overclassification (Fig. 6a), especially for
26
27 453 macrophytes, reflects that the index values were often close to upper class boundaries.
28
29
30
31
32
33

34 454
35
36 455 The probability of misclassification did not differ significantly among the different
37
38 456 combination rules, according to the ANOVA test (Fig. 6b). However, the combination rules
39
40 457 influenced the numbers of under- or overclassification. The number of underclassifications
41
42 458 were significantly lower under the Avg(-W) rules than under the OOAO rule ($F_{3,116} = 8.68$, p
43
44 459 < 0.001). Conversely, the Avg rule resulted in more overclassifications than OOAO ($F_{3,116} =$
45
46 460 3.30 , $p < 0.03$).

47
48
49 461
50
51 462 The effect of SD level on the number of misclassifications of whole lakes varied among the
52
53 463 combination rules. For the two OOAO(-E) rules, the rate of misclassification increased
54
55 464 significantly with SD ($F_{1,58} = 8.61$, $p < 0.005$). In contrast, under Avg(-W), misclassification
56
57
58
59
60
61
62
63
64
65

465 was not significantly affected by SD ($F_{1,58} = 0.148$, $p < 0.70$). Correspondingly, the number of
1
2 466 underclassifications increased with SD under the OOA(-E) rules ($F_{1,58} = 3.31$, $p = 0.062$),
3
4 467 but was not affected by SD under the Avg(-W) rules ($F_{1,58} = 1.60$, $p = 0.21$). The number of
5
6
7 468 overclassifications was not affected by SD (both $F_{1,58} < 0.73$, $p > 0.39$).
8

9 469
10
11
12 470 A similar pattern was found by Caroni et al. (2013), using the WISERBUGS simulation
13
14 471 approach for Swedish lakes with data on 2-4 BQEs with SD ranging from 0.00001 to 0.25:
15
16
17 472 The proportion of misclassifications, as well as the bias towards "underclassification",
18
19 473 increased more with SD when BQEs were combined by OOA than when BQE classes were
20
21
22 474 averaged.
23

24 475
25
26 476 In conclusion, three tendencies can be inferred from the aggregated distribution of status of all
27
28
29 477 lakes (Fig. 5) and from the statistical testing of percentage misclassification among lakes.
30
31 478 First, the total number of misclassifications is slightly higher under the OOA rule than under
32
33
34 479 the other three combination rules. Second, under the OOA there are considerably more
35
36 480 underclassifications than overclassifications; under the other combination rules these two
37
38
39 481 types of misclassifications are more balanced. Third, higher uncertainty (SD) increases the
40
41 482 percentage of misclassification, and especially the percentage of underclassification, more
42
43
44 483 under the OOA than under the other rules. In other words, because the OOA rule never
45
46 484 gives "the benefit of the doubt", higher levels of doubt (uncertainty) will generally lead to
47
48
49 485 stricter assessments.
50

51 486 52 53 487 *4.5. Implications for water management policy* 54 55

56 488
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

489 The most critical outcome of status classification may be whether a waterbody fails to meet
490 the WFD objective of good ecological status, and therefore will need restoration measures.
491 Classification with the OOA rule resulted in moderate or worse status for all lakes in the
492 Wel catchment; accordingly all of these lakes need restoration. The average combination rules
493 improved the status from moderate to good in two cases; the weighted average even improved
494 the status from poor to good in one case. In such situations, the choice of a strict combination
495 rule such as the OOA can determine management decisions in favour lake restoration, and
496 therefore cause considerable economic costs (Prato et al., 2014). Conversely, selecting a more
497 liberal average-based combination rule e.g. for Kiełpińskie would imply that lake restoration
498 is not needed, even though the moderate status of two BQEs indicated that improvement was
499 needed in this case.

500
501 For lake management in practice, the quantification of uncertainty of index values (as
502 required by the WFD) may be difficult, and estimation of the risk of misclassification will
503 therefore be a challenge. Based on this study, it is not possible to conclude for a given lake
504 that an average-based combination rule will give higher or lower risk of misclassification than
505 the OOA rule. Nevertheless, one can generally expect that the risk of misclassification will
506 be more affected by uncertainty in index values if the OOA combination rule is used
507 compared with an average-based rule. Moreover, using the OOA, one can expect a higher
508 risk of underclassification compared with overclassification if the uncertainty is high.

509
510 The OOA rule for classification of waterbodies was recommended by the EC (2005) as a
511 means implementing the precautionary principle and protecting the whole ecosystem.
512 Moreover, the biological indices based on different taxonomic groups may indicate
513 anthropogenic pressures of different types (e.g., nutrient enrichment vs. habitat degradation)

514 or occurring at different spatial and temporal scales (e.g., local habitats vs. watershed-level)
1
2 515 (Carlisle et al., 2008; Walters et al., 2009). In this respect, the OOA rule makes more sense
3
4
5 516 than other, less conservative rules.
6

7 517
8
9 518 Although the OOA rule is simple to implement in practice, the consequences of using this
10
11
12 519 rule become more complicated when considering the effects of uncertainty, as shown by our
13
14 520 analysis and by Caroni et al. (2013). In future versions of guidelines for WFD
15
16
17 521 implementation, these findings should be considered and alternative combination rules should
18
19 522 be discussed. Like other authors (Alahuhta et al., 2009), we will not conclude by
20
21
22 523 recommending one particular combination rule as the most appropriate, but hope that our
23
24 524 results contribute to a better understanding of the benefits and shortcomings of different
25
26
27 525 combination rules when applied to different ecosystem.
28

29 526
30
31 527 We support the statements that more research is needed on combination rules for integrated
32
33
34 528 waterbody assessment under uncertainty (Caroni et al., 2013; Nõges et al., 2009). The existing
35
36
37 529 datasets in this and other cited studies provide opportunities for more investigation using
38
39 530 simulation approaches such as WISERBUGS, e.g. with different criteria for weighting BQEs
40
41 531 and different uncertainty levels under alternative combination rules. The aim should be to
42
43
44 532 obtain a combination rule that ensures the protection of the whole ecosystem elements under
45
46 533 pressure, while the risk of underclassification and "false alarm" for restoration is acceptable
47
48
49 534 for waterbody management in practice.
50

51 535

53 536 **Acknowledgments**

54
55
56 537

1
2 538 The study was financed by a grant from the Norwegian Financial Mechanism (Project No.
3 539 PNRF – 220 – A I – 1/07) within the project *deWELopment* (Development and validation of
4 540 methods for integrated assessment of ecological status of rivers and lakes to support river
5 541 basin management plans). We are grateful to Hanna Ciecierska and Joanna Ruszczyńska
6
7 542 (University of Warmia and Mazury) for macrophyte data, Agnieszka Pasztaleniec (Institute of
8
9 543 Environmental Protection – NRI), Agnieszka Napiórkowska-Krzebietke, Joanna Hutorowicz
10
11 544 (Inland Fisheries Institute) for phytoplankton data, and Łucjan Chybowski (Inland Fisheries
12
13 545 Institute) for fish data. Sebastian Kutyla is acknowledged for kindly supporting us in the
14
15 546 preparation of Fig. 1. We thank Ralph Clarke and Mike Dunbar for advice on the use of the
16
17 547 WISERBUGS method. Anders Gjørwad Hagen and four anonymous reviewers are
18
19 548 acknowledged for helpful comments to earlier versions of the manuscript.
20
21
22
23
24
25
26
27
28

29

550 **Supplementary data**

551

33
34 552 Supplementary information associated with this article can be found in the online version.
35
36 553 Figs. S1-7: Uncertainty in classification for each of the remaining seven lakes not shown in
37
38 554 Figs. 2-4.
39
40

41

44 556 **References**

45
46 557

47
48 558 Alahuhta, J., Vuori, K.-M., Hellsten, S., Järvinen, M., Olin, M., Rask, M., Palomäki, A.,
49
50 559 2009. Defining the ecological status of small forest lakes using multiple biological quality
51
52 560 elements and palaeolimnological analysis. *Fundamental and Applied Limnology / Archiv für*
53
54 561 *Hydrobiologie* 175, 203-216.
55
56
57
58
59
60
61
62
63
64
65

562 Białokoz, W., Chybowski, Ł., 2011. 5. Ichtyofauna, in: Soszka, H. (Ed.), Ecological status
1
2 563 assessment of the waters in the Wel River catchment. Guidelines for integrated assessment of
3
4 564 ecological status of rivers and lakes to support river basin management plans. Institute of
5
6
7 565 Inland Fisheries, Olsztyn, Poland, pp. 217-234 (in Polish with English summary).
8
9
10 566 Birk, S., Bonne, W., Borja, A., Brucet, S., Courrat, A., Poikane, S., Solimini, A., van de
11
12 567 Bund, W., Zampoukas, N., Hering, D., 2012. Three hundred ways to assess Europe's surface
13
14 568 waters: An almost complete overview of biological methods to implement the Water
15
16
17 569 Framework Directive. Ecological Indicators 18, 31-41.
18
19 570 Borja, A., Bald, J., Franco, J., Larreta, J., Muxika, I., Revilla, M., Rodríguez, J.G., Solaun, O.,
20
21 571 Uriarte, A., Valencia, V., 2009. Using multiple ecosystem components, in assessing
22
23
24 572 ecological status in Spanish (Basque Country) Atlantic marine waters. Mar Pollut Bull 59, 54-
25
26
27 573 64.
28
29 574 Borja, A., Rodriguez, J.G., 2010. Problems associated with the 'one-out, all-out' principle,
30
31 575 when using multiple ecosystem components in assessing the ecological status of marine
32
33
34 576 waters. Mar Pollut Bull 60, 1143-1146.
35
36 577 Böhmer, J., Arbaciauskas, K., Benstead, R., Wim Gabriels, Porst, G., Reeze, B., Timm, H.,
37
38
39 578 2014. Central Baltic Lake Benthic invertebrate ecological assessment methods, in: Poikane, S.
40
41 579 (Ed.), Water Framework Directive Intercalibration Technical Report. European Commission
42
43 580 Joint Research Centre.
44
45
46 581 Carlisle, D.M., Hawkins, C.P., Meador, M.R., Potapova, M., Falcone, J., 2008. Biological
47
48 582 assessment of Appalachian streams based on predictive models for fish, macroinvertebrate,
49
50
51 583 and diatom assemblages. Journal of North American Benthological Society 27, 16-37.
52
53 584 Caroni, R., Bund, W., Clarke, R.T., Johnson, R.K., 2013. Combination of multiple biological
54
55
56 585 quality elements into waterbody assessment of surface waters. Hydrobiologia 704, 437-451.
57
58
59
60
61
62
63
64
65

586 Carstensen, J., 2007. Statistical principles for ecological status classification of Water
1
2 587 Framework Directive monitoring data. *Mar Pollut Bull* 55, 3-15.
3
4 588 Carvalho, L., Poikane, S., Lyche Solheim, A., Phillips, G., Borics, G., Catalan, J., Hoyos, C.,
5
6
7 589 Drakare, S., Dudley, B.J., Järvinen, M., Laplace-Treytore, C., Maileht, K., McDonald, C.,
8
9
10 590 Mischke, U., Moe, J., Morabito, G., Nöges, P., Nöges, T., Ott, I., Pasztaleniec, A., Skjelbred,
11
12 591 B., Thackeray, S.J., 2013. Strength and uncertainty of phytoplankton metrics for assessing
13
14 592 eutrophication impacts in lakes. *Hydrobiologia* 704, 127-140.
15
16 593 Clarke, R.T., 2013. Estimating confidence of European WFD ecological status class and
17
18 594 WISER Bioassessment Uncertainty Guidance Software (WISERBUGS). *Hydrobiologia* 704,
19
20 595 39-56.
21
22 596 Clarke, R.T., Hering, D., 2006. Errors and uncertainty in bioassessment methods – major
23
24 597 results and conclusions from the STAR project and their application using STARBUGS.
25
26 598 *Hydrobiologia* 566, 433-439.
27
28 599 Clarke, R.T., Lorenz, A., Sandin, L., Schmidt-Kloiber, A., Strackbein, J., Kneebone, N.T.,
29
30
31 600 Haase, P., 2006. Effects of sampling and sub-sampling variation using the STAR-AQEM
32
33 601 sampling protocol on the precision of macroinvertebrate metrics. *Hydrobiologia* 566, 441-
34
35 602 459.
36
37
38 603 EC (European Commission), 2000. Directive 2000/60/EC of the European Parliament and of
39
40 604 the council of 23 October 2000 establishing a framework for Community action in the field of
41
42 605 water policy. , Official Journal of the European Communities, Luxembourg.
43
44 606 EC (European Commission), 2005. Overall approach to the classification of ecological status
45
46 607 and ecological potential, Common Implementation Strategy for the Water Framework
47
48 608 Directive (2000/60/CE). Office for Official Publications of the European Communities,
49
50 609 Luxembourg.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

610 EC (European Commission), 2011. Guidance document on the intercalibration process 2008-
1
2 611 2011. Technical Report - 2011 - 045, Common Implementation Strategy for the Water
3
4 612 Framework Directive (2000/60/CE). Office for Official Publications of the European
5
6
7 613 Communities, Luxembourg.
8
9 614 Gołub, M., Prus, P., Koszałka, J., 2011. 4. Macrozoobenthos, in: Soszka, H. (Ed.), Ecological
10
11 615 status assessment of the waters in the Wel River catchment. Guidelines for integrated
12
13 616 assessment of ecological status of rivers and lakes to support river basin management plans.
14
15 617 Institute of Inland Fisheries, Olsztyn, Poland, pp. 203-216 (in Polish with English summary).
16
17 618 Gottardo, S., Semenzin, E., Giove, S., Zabeo, A., Critto, A., de Zwart, D., Ginebreda, A.,
18
19 619 Marcomini, A., 2011. Integrated risk assessment for WFD ecological status classification
20
21 620 applied to Llobregat river basin (Spain). Part I—Fuzzy approach to aggregate biological
22
23 621 indicators. *Science of The Total Environment* 409, 4701-4712.
24
25 622 Hering, D., Borja, A., Carstensen, J., Carvalho, L., Elliott, M., Feld, C.K., Heiskanen, A.-S.,
26
27 623 Johnson, R.K., Moe, J., Pont, D., Solheim, A.L., de Bund, W.v., 2010. The European Water
28
29 624 Framework Directive at the age of 10: A critical review of the achievements with
30
31 625 recommendations for the future. *Science of The Total Environment* 408, 4007-4019.
32
33 626 Hutorowicz, A., Napiórkowska-Krzebietke, A., Pasztaleniec, A., Hutorowicz, J., Lyche
34
35 627 Solheim, A., Skjelbred, B., 2011. 1. Phytoplankton, in: Soszka, H. (Ed.), Ecological status
36
37 628 assessment of the waters in the Wel River catchment. Guidelines for integrated assessment of
38
39 629 ecological status of rivers and lakes to support river basin management plans. Institute of
40
41 630 Inland Fisheries, Olsztyn, Poland, pp. 143-168 (in Polish with English summary).
42
43 631 Højberg, A.L., Refsgaard, J.C., Geer, F., Jørgensen, L.F., Zsuffa, I., 2007. Use of models to
44
45 632 support the monitoring requirements in the Water Framework Directive. *Water Resour*
46
47 633 *Manage* 21, 1649-1672.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

634 Irvine, K., 2004. Classifying ecological status under the European Water Framework
1
2 635 Directive: the need for monitoring to account for natural variability. *Aquatic Conservation:*
3
4 636 *Marine and Freshwater Ecosystems* 14, 107-112.
5
6
7 637 Kelly, M., Bennett, C., Coste, M., Delgado, C., Delmas, F., Denys, L., Ector, L., Fauville, C.,
8
9 638 Ferréol, M., Golub, M., Jarlman, A., Kahlert, M., Lucey, J., Ní Chatháin, B., Pardo, I., Pfister,
10
11 639 P., Picinska-Faltynowicz, J., Rosebery, J., Schranz, C., Schaumburg, J., van Dam, H.,
12
13 640 Vilbaste, S., 2009a. A comparison of national approaches to setting ecological status
14
15 641 boundaries in phytobenthos assessment for the European Water Framework Directive: results
16
17 642 of an intercalibration exercise. *Hydrobiologia* 621, 169-182.
18
19 643 Kelly, M., Bennion, H., Burgess, A., Ellis, J., Juggins, S., Guthrie, R., Jamieson, J.,
20
21 644 Adriaenssens, V., Yallop, M., 2009b. Uncertainty in ecological status assessments of lakes
22
23 645 and rivers using diatoms. *Hydrobiologia* 633, 5-15.
24
25
26 646 Kolada, A., Ciecierska, H., Rusczyńska, J., Dynowski, P., 2013. Sampling techniques and
27
28 647 inter-surveyor variability as sources of uncertainty in Polish macrophyte metric for lake
29
30 648 ecological status assessment. *Hydrobiologia* DOI 10.1007/s10750-013-1591-9.
31
32
33 649 Kolada, A., Ciecierska, H., Rusczyńska, J., Mjelde, M., Dziedzic, J., Dynowski, P.,
34
35 650 Jasnorzewski, A., 2011. 2. Macrophytes, in: Soszka, H. (Ed.), *Ecological status assessment of*
36
37 651 *the waters in the Wel River catchment. Guidelines for integrated assessment of ecological*
38
39 652 *status of rivers and lakes to support river basin management plans. Institute of Inland*
40
41 653 *Fisheries, Olsztyn, Poland, pp. 169-186 (in Polish with English summary).*
42
43 654 Moss, B., Stephen, D., Alvarez, C., Becares, E., Bund, W.V.D., Collings, S.E., Donk, E.V.,
44
45 655 Eyto, E.D., Feldmann, T., Fernández-Aláez, C., Fernández-Aláez, M., Franken, R.J.M.,
46
47 656 García-Criado, F., Gross, E.M., Gyllström, M., Hansson, L.-A., Irvine, K., Järvalt, A., Jensen,
48
49 657 J.-P., Jeppesen, E., Kairesalo, T., Kornijów, R., Krause, T., Künnap, H., Laas, A., Lill, E.,
50
51 658 Lorens, B., Luup, H., Rosa Miracle, M., Nöges, P., Nöges, T., Nykänen, M., Ott, I., Peczula,
52
53
54
55
56
57
58
59
60
61
62
63
64
65

659 W., Peeters, E.T.H.M., Phillips, G., Romo, S., Russell, V., Salujõe, J., Scheffer, M.,
1
2 660 Siewertsen, K., Smal, H., Tesch, C., Timm, H., Tuvikene, L., Tonno, I., Virro, T., Vicente, E.,
3
4 661 Wilson, D., 2003. The determination of ecological status in shallow lakes — a tested system
5
6 662 (ECOFRAME) for implementation of the European Water Framework Directive. *Aquatic*
7
8
9 663 *Conservation: Marine and Freshwater Ecosystems* 13, 507-549.
10
11 664 Nõges, P., Bund, W., Cardoso, A.C., Solimini, A.G., Heiskanen, A.-S., 2009. Assessment of
12
13 665 the ecological status of European surface waters: a work in progress. *Hydrobiologia* 633, 197-
14
15 666 211.
16
17 667 Nõges, P., Nõges, T., 2006. Indicators and criteria to assess ecological status of the large
18
19 668 shallow temperate polymictic lakes Peipsi (Estonia/Russia) and Vortsjarv (Estonia). *Boreal*
20
21 669 *Environ. Res.* 11, 67-80.
22
23 670 Phillips, G., Free, G., Karottki, I., Laplace-Treyture, C., Maileht, K., Mischke, U., Ott, I.,
24
25 671 Pasztaleniec, A., Portielje, R., Søndergaard, M., Trodd, W., Wichelen, J.V., 2014. Central
26
27 672 Baltic Lake Phytoplankton ecological assessment methods, in: Poikane, S. (Ed.), *Water*
28
29 673 *Framework Directive Intercalibration Technical Report*. European Commission Joint
30
31 674 Research Centre.
32
33 675 Picińska-Fałtynowicz, J., 2011. 3. Phytobenthos, in: Soszka, H. (Ed.), *Ecological status*
34
35 676 *assessment of the waters in the Wel River catchment. Guidelines for integrated assessment of*
36
37 677 *ecological status of rivers and lakes to support river basin management plans*. Institute of
38
39 678 *Inland Fisheries*, Olsztyn, Poland, pp. 187-202 (in Polish with English summary).
40
41 679 Portielje, R., Bertrin, V., Denys, L., Grinberga, L., Karottki, I., Kolada, A., Krasovskienė, J.,
42
43 680 Leiputé, G., Maemets, H., Ott, I., Phillips, G., Pot, R., Schaumburg, J., Schranz, C., Soszka,
44
45 681 H., Stelzer, D., Søndergaard, M., Willby, N., 2014. Central Baltic Lake Macrophyte
46
47 682 *ecological assessment methods*, in: Poikane, S. (Ed.), *Water Framework Directive*
48
49 683 *Intercalibration Technical Report*. European Commission Joint Research Centre.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

684 Prato, S., La Valle, P., De Luca, E., Lattanzi, L., Migliore, G., Morgana, J.G., Munari, C.,
1
2 685 Nicoletti, L., Izzo, G., Mistri, M., 2014. The "one-out, all-out" principle entails the risk of
3
4 686 imposing unnecessary restoration costs: a study case in two Mediterranean coastal lakes. *Mar*
5
6
7 687 *Pollut Bull* 80, 30-40.
8
9 688 R Development Core Team, 2011. R: A language and environment for statistical computing.
10
11 689 R Foundation for Statistical Computing, Vienna, Austria.
12
13
14 690 Rask, M., Vuori, K.-M., Hämäläinen, H., Järvinen, M., Hellsten, S., Mykrä, H., Arvola, L.,
15
16 691 Ruuhijärvi, J., Jyväsjärvi, J., Kolari, I., Olin, M., Salonen, E., Valkeajärvi, P., 2010.
17
18 692 Ecological classification of large lakes in Finland: comparison of classification approaches
19
20
21 693 using multiple quality elements. *Hydrobiologia* 660, 37-47.
22
23
24 694 Ruse, L., 2010. Classification of nutrient impact on lakes using the chironomid pupal exuvial
25
26 695 technique. *Ecological Indicators* 10, 594-601.
27
28
29 696 Sandin, L., 2005. Testing the EC Water Framework Directive "one-out, all-out" rule -
30
31 697 simulating different levels of assessment errors along a pollution gradient in Swedish streams.
32
33 698 *Verhandlungen der Internationale Vereinigung für Theoretische und Angewandte Limnologie*
34
35
36 699 29, 334-336.
37
38 700 Schneider, S.C., Lawniczak, A.E., Picińska-Faltynowicz, J., Szoszkiewicz, K., 2012. Do
39
40 701 macrophytes, diatoms and non-diatom benthic algae give redundant information? Results
41
42 702 from a case study in Poland. *Limnologica - Ecology and Management of Inland Waters* 42,
43
44
45 703 204-211.
46
47 704 Soszka, H., 2011. Ecological status assessment of the waters in the Wel River catchment.
48
49 705 Guidelines for integrated assessment of ecological status of rivers and lakes to support river
50
51 706 basin management plans. Institute of Inland Fisheries, Olsztyn, Poland (in Polish with English
52
53
54
55 707 summary).
56
57
58
59
60
61
62
63
64
65

708 Sutela, T., Aroviita, J., Keto, A., 2013. Assessing ecological status of regulated lakes with
1
2 709 littoral macrophyte, macroinvertebrate and fish assemblages. *Ecological Indicators* 24, 185-
3
4
5 710 192.
6
7 711 Szoszkiewicz, K., Jusik, S., Zgola, T., Czechowska, M., Hryc, B., 2007. Uncertainty of
8
9 712 macrophyte-based monitoring for different types of lowland rivers. *Belgian Journal of Botany*
10
11 713 140, 7-16.
12
13
14 714 Søndergaard, M., Jeppesen, E., Peder Jensen, J., Lildal Amsinck, S., 2005. Water Framework
15
16 715 Directive: ecological classification of Danish lakes. *Journal of Applied Ecology* 42, 616-629.
17
18
19 716 Thackeray, S.J., Nõges, P., Dunbar, M.J., Dudley, B.J., Skjelbred, B., Morabito, G., Carvalho,
20
21 717 L., Phillips, G., Mischke, U., Catalan, J., de Hoyos, C., Laplace, C., Austoni, M., Padedda,
22
23 718 B.M., Maileht, K., Pasztaleniec, A., Järvinen, M., Solheim, A.L., Clarke, R.T., 2013.
24
25 719 Quantifying uncertainties in biologically-based water quality assessment: A pan-European
26
27 720 analysis of lake phytoplankton community metrics. *Ecological Indicators* 29, 34-47.
28
29
30
31 721 Veríssimo, H., Lane, M., Patrício, J., Gamito, S., Marques, J.C., 2013. Trends in water quality
32
33 722 and subtidal benthic communities in a temperate estuary: Is the response to restoration efforts
34
35 723 hidden by climate variability and the Estuarine Quality Paradox? *Ecological Indicators* 24, 56-
36
37 724 67.
38
39
40
41 725 Walters, D.M., Roy, A.H., Leigh, D.S., 2009. Environmental indicators of macroinvertebrate
42
43 726 and fish assemblage integrity in urbanizing watersheds. *Ecological Indicators* 9, 1222-1233.
44
45
46 727
47
48
49 728
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

729 **Appendix A**

730

731 **Table A.1.** Mean index values in EQR scale for each BQE.

732

Lake name	Biological quality element (BQE) ^a				
	PP	PB	MP	MI	FI
Dąbrowa Wielka	0.42	0.70	0.67	0.53	0.78
Dąbrowa Mała	0.39	0.82	0.50	0.43	0.94
Rumian	0.31	0.70	0.39	0.38	0.61
Zarybinek	0.39	0.59	0.32	0.36	0.39
Tarczyńskie	0.12	0.56	0.33	0.09	0.08
Grądy	0.30	0.79	0.33	0.06	0.31
Lidzbarskie	0.34	0.67	0.32	0.24	0.36
Kiełpińskie	0.81	0.69	0.44	0.59	0.94
Hartowieckie	0.46	0.78	0.48	0.31	0.33
Zwiniarz	0.18	0.85	0.26	0.28	0.25

733

734 **Table A.2.** Class boundaries in EQR scale used for status classification and for calculation of
 1
 2 735 normalised EQR values, for each biological quality element (BQE).

3
 4
 5 736

7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22	BQE ^a	Class boundaries ^b			
		H/G	G/M	M/P	P/B
11	PP	0.8	0.6	0.4	0.2
14	PB	0.8	0.6	0.4	0.15
17	MP	0.68	0.51	0.34	0.17
19	MI	0.9	0.69	0.45	0.21
21	FI	0.69	0.45	0.25	0.1

23
 24 737 ^a PP = phytoplankton, PB = phytobenthos, MP = macrophytes, MI = macroinvertebrates, FI =
 25
 26 738 fish

27
 28
 29 739 ^b H = high, G = good, M = moderate, P = poor, B = bad
 30

31 740
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65

Figure captions

Fig. 1. Location of the study area. Upper left panel: map of Europe with location of Poland; lower left panel: map of Poland with location of the Wel river catchment; right panel: location of ten lakes within the Wel river catchment in North-Central Poland. The numbers refer to lake names in Table 1.

Fig. 2. Uncertainty in classification of Lake Kiełpińskie for different biological quality elements (BQEs), different BQE combination rules and different uncertainty levels in BQE index values. For more information on the lake, see Table 1. The labels above the bars show the "correct class" (cf. Table 2). For abbreviations and more details, see Table 2. The distribution of status classes based on 10,000 simulated resamples (see section 3.2). (a) Percentage of status classes assessed for each BQE and for each uncertainty level (1, 2, 3). (b) Percentage of resamples of each BQE categorised as underclassification and overclassification, respectively. (c) Percentage of waterbody status classes assessed for each BQE combination rule and for each uncertainty level. (d) Percentage of waterbody status classes categorised as underclassification and overclassification, respectively.

Fig. 3. Uncertainty in classification of Lake Zarybinek for different biological quality elements (BQEs), different BQE combination rules and different uncertainty levels in BQE index values. For abbreviations and more details, see Fig. 2.

Fig. 4. Uncertainty in classification of Lake Dąbrowa Mała for different biological quality elements (BQEs), different BQE combination rules and different uncertainty levels in BQE index values. For abbreviations and more details, see Fig. 2.

Fig. 5. Summarised uncertainty in classification of all ten lakes combined for different biological quality elements (BQEs), different BQE combination rules and different uncertainty levels in BQE index values. In plots (a) and (c), the percentage of resamples for each status class is summed for all lakes. The "correct aggregated status" is the class with the highest proportion of resamples for all lakes combined, for each BQE (a) and each combination rule (c), respectively. For abbreviations and more details, see Fig. 2.

Fig. 6. Outcome of the analysis of variance (ANOVA) in misclassification for the ten lakes by different BQEs, different combination rules and different levels of uncertainty in individual BQE index values. The calculation of misclassification for each simulated resample is described in section 3.2, step 5. The displayed percentage of misclassification represents the average number of misclassified resamples for all lakes. The letters above the bars in plot (a) indicate significant differences between BQEs according to the ANOVA (see method description in section 3.3): pairs of BQEs with significantly different percentage of misclassification have no common letters above the bars. For abbreviations and more details, see Fig. 2.

Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.




Fig. 6.




Table 1

The main characteristics of the ten lakes used in the study. Mixing types: s = stratified, ns = non-stratified. Land use forms: NAT = natural and semi-natural; AGR = agriculture, URB = urban. After Soszka (2011). For geographic location, see Fig. 1.

No.	Lake	Surface area (km ²)	Max depth (m)	Mean depth (m)	Retention time (years)	Mixing type	Catchment area (km ²)	Land use in catchment (%)		
								NAT	AGR	URB
1	Dąbrowa Wielka	6.15	34.7	8.2	2.50	s	95.5	22.6	76.2	1.2
2	Dąbrowa Mała	1.73	34.5	10.0	0.59	s	159.1	25.2	73.8	1.0
3	Rumian	3.06	14.4	6.5	0.36	s	254.4	22.5	76.7	0.8
4	Zarybinek	0.74	7.0	2.4	0.03	ns	270.1	22.7	76.4	0.9
5	Tarczyńskie	1.64	9.2	3.8	0.10	ns	296.2	23.8	75.4	0.8
6	Grądy	1.13	9.1	4.7	0.10	ns	332.9	25.8	73.4	0.8
7	Lidzbarskie	1.22	25.5	10.1	0.12	s	534.3	29.4	69.3	1.3
8	Kiełpińskie	0.61	11.0	6.1	2.50	s	14.3	56.9	42.8	0.3
9	Hartowieckie	0.70	5.2	2.9	1.11	ns	8.7	17.9	82.1	0.0
10	Zwiniarz	0.50	5.8	3.0	1.11	ns	7.9	34.3	65.7	0.0

Table 2

Ecological status of the ten lakes (Table 1) based on data for each biological quality element (BQE) and integration of all BQEs using different combination rules (see section 2.3). The values under BQEs are normalised ecological quality ratios (nEQR; see section 2.2). Ecological status classes correspond to the following intervals of nEQR: 0-0.2 = Bad, 0.2-0.4 = Poor, 0.4-0.6 = Moderate, 0.6-0.8 = Good, 0.8-1 = High.

Lake name	Biological quality element ^a					Combination rule ^b			
	PP	PB	MP	MI	FI	OOAO	OOAO-E	Avg	Avg-W
Dąbrowa Wielka	0.42	0.70	0.79	0.47	0.86	Moderate	Moderate	Good	Good
Dąbrowa Mała	0.39	0.82	0.59	0.38	0.96	Poor	Poor	Moderate	Good
Rumian	0.31	0.70	0.46	0.34	0.73	Poor	Poor	Moderate	Moderate
Zarybinek	0.39	0.59	0.37	0.33	0.54	Poor	Poor	Poor	Poor
Tarczyńskie	0.12	0.56	0.39	0.09	0.16	Bad	Bad	Poor	Poor
Grądy	0.30	0.79	0.38	0.06	0.46	Bad	Poor	Poor	Moderate
Lidzbarskie	0.34	0.67	0.38	0.23	0.51	Poor	Poor	Moderate	Moderate
Kielpińskie	0.83	0.69	0.51	0.52	0.96	Moderate	Moderate	Good	Good
Hartowieckie	0.46	0.78	0.56	0.28	0.48	Poor	Moderate	Moderate	Moderate
Zwiniarz	0.18	0.85	0.30	0.26	0.40	Bad	Bad	Poor	Poor

^a PP = phytoplankton, PB = phytobenthos, MP = macrophytes, MI = macroinvertebrates, FI = fish

^b OOA = one-out, all-out, OOA-E = one-out, all-out after exclusion of MI, Avg = average; Avg-W = average with down-weighting of MI




Fig. S1. Uncertainty in classification of Lake Dąbrowa Wielka for different BQEs, different BQE combination rules and different levels of uncertainty in individual BQE index values. For more details, see Fig. 2.




Fig. S2. Uncertainty in classification of Lake Rumian for different BQEs, different BQE combination rules and different levels of uncertainty in individual BQE index values. For more details, see Fig. 2.




Fig. S3. Uncertainty in classification of Lake Tarczyńskie for different BQEs, different BQE combination rules and different levels of uncertainty in individual BQE index values. For more details, see Fig. 2.




Fig. S4. Uncertainty in classification of Lake Grądy for different BQEs, different BQE combination rules and different levels of uncertainty in individual BQE index values. For more details, see Fig. 2.




Fig. S5. Uncertainty in classification of Lake Lidzbarskie for different BQEs, different BQE combination rules and different uncertainty levels in BQE index values. For more details, see Fig. 2.




Fig. S6. Uncertainty in classification of Lake Hartowieckie for different BQEs, different BQE combination rules and different levels of uncertainty in individual BQE index values. For more details, see Fig. 2.




Fig. S7. Uncertainty in classification of Lake Zwiniarz for different BQEs, different BQE combination rules and different levels of uncertainty in individual BQE index values. For more details, see Fig. 2.