

Original article (Orijinal araştırma)

First record of the egg parasitoids of *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae) in Turkey using DNA barcoding

Türkiye’de *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae)’un yumurta parazitoidlerinin DNA Barkodu kullanılarak belirlenmiş ilk kaydı

Feza CAN CENGİZ^{1*} Kamuran KAYA² Başak ULAŞLI¹ Jérôme MORINIÈRE³

Summary

Chilo partellus (Swinhoe) (Lepidoptera: Crambidae) is an invasive insect species attacking maize (*Zea mays* L.) and other cereal crops causing important yield losses. The occurrence of this insect in Turkey was first reported in maize growing areas of some provinces in the East Mediterranean region of Turkey in 2014. Chemical or other pest control methods do not always provide acceptable control of this pest, so biological control is considered an important alternative. However, for a successful biological control, the first step is to reliably identify the natural enemies of a target pest, which is difficult to achieve using methods based on morphology. Recent developments in molecular techniques allow more reliable identification of insect species and their parasitoids. Therefore, the aim of this study was to identify the egg parasitoids of *C. partellus* by molecular methods. Parasitized eggs were collected from maize fields in Hatay province, Turkey, from September to October 2014 and in September 2015. Eggs were maintained in the laboratory and emerging adult parasitoids were subjected to molecular analysis. Using DNA barcoding, two native natural enemies, *Trichogramma brassicae* Bezdenko (Hymenoptera: Trichogrammatidae) and *Telenomus busseolae* (Gahan) (Hymenoptera: Platygastridae) were identified as egg parasitoids of *C. partellus* for the first time in Turkey.

Keywords: *Trichogramma brassicae*, *Telenomus busseolae*, COI, spotted stem borer

Özet

Chilo partellus (Swinhoe) (Lepidoptera: Crambidae) mısır ve diğer tahıl ürünlerinde zararlı olan ve önemli ürün kayıplarına neden olan istilacı bir türdür. Bu böceğin Türkiye’deki varlığı ilk kez 2014 yılında Türkiye’nin Doğu Akdeniz Bölgesi’nin bazı illerinin mısır alanlarında kaydedilmiştir. Bu zararlı için insektisitler veya diğer zararlı mücadele metodları her zaman yeterli kontrolü sağlamamakta, biyolojik mücadele ise önemli bir alternatif olarak düşünülmektedir. Ancak biyolojik metodların başarılı bir şekilde uygulanması için zararlının doğal düşmanlarının ilk adımda iyi tanımlanmış olması gerekmektedir, Morfolojiye dayanan klasik teşhis metodlarıyla bunu gerçekleştirmek oldukça zordur. Moleküler tekniklerdeki son gelişmeler, böcek türlerinin ve onların parazitoidlerinin daha doğru tanımlanmasına olanak vermektedir. Dolayısıyla bu çalışmanın amacı *C. partellus*’ un yumurta parazitoidlerini, moleküler metodlar kullanarak tanılamaktır. Bu zararlının parazitlenmiş yumurtaları, 2014 yılının eylül ve ekim ayları ile 2015 yılı eylül ayında Türkiye’nin Hatay ilindeki mısır alanlarından toplanmıştır. Yumurtalar daha sonra laboratuvar koşullarında kültüre alınmış ve çıkış yapan erginlerin moleküler analizleri yapılmıştır. DNA barkod ile İki yerel doğal düşman, *Trichogramma brassicae* Bezdenko (Hymenoptera: Trichogrammatidae) ve *Telenomus busseolae* (Gahan) (Hymenoptera: Platygastridae) *C. partellus*’un yumurta parazitoidleri olarak ilk kez kaydedilmiştir.

Anahtar sözcükler: *Trichogramma brassicae*, *Telenomus busseolae*, COI, benekli gövdekurdu

¹ Mustafa Kemal University, Agricultural Faculty, Plant Protection Department, Entomology Section 31040 Hatay-Turkey

² Mustafa Kemal University, Agricultural Research and Application Center 31040 Hatay-Turkey

³ SNSB - Bavarian State Collection of Zoology (ZSM), Münchhausenstrasse 21; 81247 München-Germany

* Corresponding author (Sorumlu yazar) e-mail: cezafan_onurcan@hotmail.com

Received (Alınış): 06.02.2016

Accepted (Kabul ediliş): 06.04.2016

Published Online (Çevrimiçi Yayın Tarihi): 25.05.2016

Introduction

Maize (*Zea mays* L.) is a globally important cereal crop providing staple food in many countries. It takes the third place among the most common cereal crops after wheat and rice (FAO-STAT, 2014), and in Turkey after wheat and barley (TÜİK, 2014). Maize production is the main income source of farmers in developing countries (Tagne et al., 2008). Despite the importance of maize production, a large number of lepidopteran pests continue to cause significant economic loss. The invasive pest, *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae), commonly known as spotted stem borer, is one of the most serious pests of cereal crops, especially maize and sorghum in Asia and South Africa. *Chilo partellus* has a broad host range, including both wild and cultivated plants (Khan et al., 1997; Rebe et al., 2004), but is of most concern as a pest of maize. *Chilo partellus* has been reported to cause severe yield losses in maize throughout its geographical distribution. In some studies, maize yield losses attributable to *C. partellus* ranged from 24 to 75% (Kumar & Mihm, 1995; Kumar, 2002). Farid et al. (2007) reported 10 to 50% damage in the Peshawar Valley, Pakistan. Yield losses caused by stem borers in Africa have been as high as 80 to 88% in maize (Van den Berg, 2009; Kfir, 1990) and as much as 88% in sorghum (Seshu Reddy, 1988). This pest was recorded for the first time in maize fields in some provinces in the East Mediterranean region of Turkey in 2014 and 2015 (Sertkaya et al., 2014; Bayram & Tonğa, 2015).

There are many chemical formulations which have been applied in the field for the management of *C. partellus*. Research has shown that a wide range of insecticide in various formulations is effective against *C. partellus*. However, the use of insecticides for pest control is not only expensive for small farm holders, but also has undesirable consequences such as resistance development, secondary pest outbreaks, environmental pollution and risk to operators. Furthermore, stem borer larvae are difficult spray targets as they are hidden within the plant stem, which reduces the efficiency of insecticides by preventing non-systemic chemicals from reaching the larvae. Moreover, insecticides have a negative impact on beneficial fauna. Therefore, biological control agents are preferable for control of this important pest and different development stages of the pest, such as eggs, should be targeted. A number of parasitoids of stem borer lepidopteran pests in cereal crops have been recorded around the globe. Hymenopteran parasitoids are one of the most species-rich groups of animals, potentially accounting for more than 20% of the world insects (LaSalle & Gauld, 1991). Most of these parasitoid species belong to the families Braconidae, Chalcidoidea, Eulophidae, Ichneumonidae, Platygasteridae and Trichogrammatidae.

Many egg, larval and pupal parasitoids of *C. partellus* have been reported from different countries. *Trichogramma chilonis* Ishii and *Trichogrammatoidea lutea* Girault (Trichogrammatidae) were found to be important egg parasitoids of *C. partellus* (Neupane et al., 1985; Kfir, 1990). Jalali & Singh (2006), reported that the parasitism rates by *T. chilonis* on *C. partellus* eggs on fodder maize were up to 75.2 and 62.6% in the first generation when parasitoids were released at 3 and 5 d intervals, respectively. In the second generation, parasitism rates were 90.4 and 78.4% at 3 and 5 d release intervals, respectively. In the study of Kfir (1990), one egg parasite, seven larval parasites (two of them egg-larval parasite) and two pupal parasites were recorded from parasitized *C. partellus* collected from maize and grain sorghum in South Africa. *Apanteles sesamiae* Cameron (Braconidae) was shown to be the most abundant larval parasite followed by an *Iphiaulax* sp. (Braconidae). *Dentichasmias busseolae* Heinrich (Ichneumonidae) and *Pediobius furvus* (Gahan) (Eulophidae) were the most abundant pupal parasites, while other species were found to be uncommon larval or pupal parasites. Kfir (1990) also reported *Chelonus curvimaculatus* Cameron and an *Chelonus* sp., (Braconidae) as egg-larval parasites of *C. partellus*. In another study, *Cotesia flavipes* Cameron, *Cotesia sesamiae* (Cameron) (Braconidae) and *Psilochalsis soudanensis* (Steffan) (Chalcidoidea) were determined to be larval parasitoids, and *P. furvus* and *D. busseolae* to be pupal parasitoids of *C. partellus* in Uganda (Rwomushana et al. 2005). Divya et al. (2009) recorded two larval parasitoids, one braconid (*C. flavipes*) and one tachinid (*Sturmiopsis inferens* Townsend), and one eulophid (*Tetrastichus* sp.) pupal parasitoid.

Trichogramma spp. (Trichogrammatidae) and *Telenomus* spp. (Platygastridae) are tiny parasitoid wasps that are important biological control agents of lepidopterous insect pests. Some of them have been used successfully in the control of crop pests (Borror et al., 1981). *Trichogramma* spp. are also effective biocontrol agents, because they can control the pest in the egg stage (Bournier, 1982; Somchoudhury & Dutt, 1988) and they are cost effective. This is an ideal choice for releases against crop borers because of its ease of propagation and application (Farid et al., 2007). The species of genus *Telenomus* are eggs parasites of a wide range of hosts, but they preferably attack Lepidoptera and Heteroptera (Masner & Johnson, 1979). *Telenomus* spp. are important in the natural control of insect pest populations (Yuliarti et al., 2002).

A reliable identification of effective parasitoid species is the first and most important step for a successful biological control program. Identification of species of these wasps has been exclusively based on the morphology of the male genitalia (Sorokina, 1993). In the cases where males are not found or are in low numbers, identification becomes even more difficult (Aeschlimann, 1990). However, the small size of these parasitoids (<1 mm long) and lack of clear diagnostic features make them difficult to identify, and this confusion can impede the implementation of effective biological control programs (Smith & Hubbes, 1986; Pinto et al., 1989; Pinto & Stouthamer 1994). Given the difficulties in distinguishing species of the genera, *Trichogramma* and *Telenomus*, molecular studies have gained importance in recent years (Yuliarti et al., 2002; Sümer et al., 2009; Poorjavad et al., 2012; Nasir et al., 2013). Therefore, with the advent of molecular techniques, DNA-based approaches have been used to generate molecular markers that are useful for the characterization of closely related or cryptic species for biological control work (Landry et al., 1993; Hoy et al., 2000; Chang et al., 2001). The general ease of species diagnosis reveals one of the great values of a DNA-based approach for identification. Newly encountered species will be recognized by their genetic divergence from known members of the assemblage (Hebert et al., 2003). Since standard morphological methods are not always sufficiently precise to differentiate micro-hymenopteran to species level (Landry et al., 1993) genetic means have become the tools of choice for the routine identification of *Trichogramma* spp. (Sümer et al., 2009). The use of a standardized DNA region for fast and reliable species identification, such as cytochrome c oxidase I (COI), has proven to be efficient in hyperdiverse groups for which morphological identification is difficult or impossible (Valentini et al., 2009).

The present study aimed to determine and identify *C. partellus* and its egg parasitoids in Turkey using DNA barcoding.

Materials and Methods

Egg masses of *C. partellus* were collected during maize field surveys in Reyhanlı-Hatay Province in September to October 2014 and September 2015. Each egg mass was put into a glass tube with parts of the plant. These egg masses were incubated at $26\pm 1^{\circ}\text{C}$, $70\pm 5\%$ RH, 16L:8D h photoperiod, until larval or pupal formation, or adult emergence of any possible natural enemies (i.e. egg parasitoids) occurred. Any emerged egg parasitoids were collected and preserved in 96% ethanol for molecular analysis. Larvae of *C. partellus* were cultured in plastic containers to obtain adults. Emerged adults of *C. partellus* were identified based on male genitalia by the first author. All specimens of the pest were deposited in the Museum of Mustafa Kemal University, Hatay, Turkey.

Amplification of the 5' COI barcoding region was undertaken by first extracting DNA from dry legs of eight *C. partellus* specimens and from the whole specimen for four micro hymenopterans (three *Trichogramma* and one *Telenomus*). DNA extraction was performed using the DNEasy tissue kit following the manufacturer's protocol (Qiagen, Hilden, Germany).

Polymerase chain reactions were performed following standard protocols of the entomology lab of the Bavarian State Collection of Zoology (http://zsm-entomology.de/wiki/Coleoptera#The_Beetle_DNA_Lab) using the primers dgHco 5'-TAACTTCAGGGTGACCAARAAYCA-3' and mLCOintF 5'-GGWACWGGWTGAACWGTWTAYCCYCC-3' (Leray et al., 2013), which amplify 300- to 350-bp

fragments targeting the center of the 5' region of COI. PCR products were purified and processed for sequencing, using BigDye v3.1 (Applied Biosystems, Foster City, CA, USA). The assembly and editing of the sequences was performed using Sequencher 4.10.1 (Gene Codes, Ann Arbor, MI, USA). In order to screen for pseudogenes, successfully amplified sequences were aligned with reference sequences in MEGA v6.0 (Tamura et al., 2013) and the coding frame was checked for stop codons. Successfully sequenced PCR products were identified using BLAST searches with the identification tool on the BOLD system (<http://www.boldsystems.org>, Ratnasingham & Hebert, 2007). Sequences of *C. partellus* were successfully identified with >98% similarity to published *C. partellus* specimens [genbank accessions: KP233794, KP233796; BIN (Cluster ID) BOLD:AAN5677]. *Trichogramma brassicae* Bezdenko sequences were identified with >99% similarity (KC488653, KC488655 and KC488656; BOLD:AAD6262). The single sequence of *Telenomus busseolae* (Gahan) was identified with >95% similarity (DQ888418).

Results and Discussion

Two native natural enemies, *T. busseolae* belonging to Platygasteridae and *T. brassicae* belonging to Trichogrammatidae were identified from parasitized eggs of *C. partellus* from maize collected from Reyhanlı of Hatay Province of the eastern Mediterranean region of Turkey on 15 September and 15 October 2014, respectively. *C. partellus* was present in both years. *Chilo partellus* was recorded recently for the first time from some provinces in the eastern Mediterranean region of Turkey, but natural enemies of this pest were not determined in these studies (Sertkaya et al., 2014; Bayram & Tonğa, 2015). When new insect species are introduced into a region, some can become invasive and cause significant economic damage. The success or failure of invasion of a species may depend on the its life history parameters, its response to climatic conditions, the competition with native species and the impact of natural enemies (Grabenweger et al., 2010). Natural enemies are considered to be the most important and practical option for keeping populations of invasive insects under pressure, thereby minimizing spread. Indigenous natural enemies may be able to reduce stem borer populations in the field (Bonhof, 2000; Midega et al., 2004). To control the stem borer lepidopteran pest by non-chemical means, it is first necessary to identify the key parasitoids species, which is difficult to achieve by morphological methods. Therefore, the results of this study are important, because they demonstrated that DNA barcode sequencing was successful in identifying *C. partellus* as well as two associated egg parasites, *T. brassicae* and *T. busseolae*, by blasting the barcode sequences using the BOLD identification tool (Table 1).

Telenomus busseolae is a solitary egg parasitoid of several species of noctuid stem borers including *C. partellus*. Parasitism by *T. busseolae* can be high in some areas of the Mediterranean basin (Kornoşor et al., 1994; Sétamou & Schulthess, 1995). However, periodic releases of mass-produced parasitoids can be necessary to increase the rate of parasitism in the field (Alexandri & Tsitsipis, 1990; Kornoşor et al., 1994). Egg parasitoids of the genus *Trichogramma*, including *T. brassicae*, have been successfully used to control mainly lepidopteran pests in biological control worldwide (Li et al., 1994)

However, changes in the rate of parasitism may happen when hosts are switched. Therefore, the ability of a parasitoid species to be reared on a non-target host and still keep a high rate of parasitism of the in the target pest after successive rearing is an important determinant of parasitoid quality and must be investigated (Pomari-Fernandes et al., 2015). Consequently, these species could be considered as appropriate candidates for biological control of the spotted stem borer. In future, mass rearing and evaluation of releases to support its naturally-occurring activity in areas of Turkey where the efficacy of the parasitoid is insufficient could be undertaken. The data obtained in this study are expected to contribute to biological management studies of the pest.

Table 1. Accession numbers of sequences submitted to the GenBank database

Species	Sampling location	BOLD process IDs	Accession numbers
<i>Chilo partellus</i>	Hatay-Turkey	GBMIX2732-16	KU687432
		GBMIX2733-16	KU687434
		GBMIX2734-16	KU687439
		GBMIX2735-16	KU687435
		GBMIX2736-16	KU687440
		GBMIX2737-16	KU687438
		GBMIX2738-16	KU687442
		GBMIX2739-16	KU687437
Parasitoids			
<i>Telenomus busseolae</i>	Hatay-Turkey	GBMIX2741-16	KU687431
<i>Trichogramma brassicae</i>	Hatay-Turkey	GBMIX2742-16	KU687441
		GBMIX2743-16	KU687433
		GBMIX2744-16	KU687436

Acknowledgments

We are grateful to Axel Hausmann [Bavarian State Collection of Zoology (ZSM), Munich, Germany] for help and valuable suggestions, including advice during preparation of this manuscript. We also thank the reviewers for their valuable comments.

References

- Aeschlimann, J. P., 1990. Simultaneous occurrence of thelytoky and bisexuality in hymenopteran species, and its implications for the biological control of pests. *Entomophaga* 35: 3–5.
- Alexandri, M. P. & J. A. Tsitsipis, 1990. Influence of the egg parasitoid *Platytenomus busseolae* (Hym.; Scelionidae) on the population of *Sesamia nonagrioides* (Lep.; Noctuidae) in Central Greece. *Entomophaga*, 35: 16-25.
- Avcı, M., 2003. Parasitism of egg-batches of the cedar processionary moth, *Traumatocampa ispartaensis* in Turkey. *Phytoparasitica* 31 (2): 1-6.
- Bayram, A. & A. Tonga, 2015. First report of *Chilo partellus* in Turkey, a new invasive maize pest for Europe. *Journal of Applied Entomology*, 140 (3): 236-240.
- Bonhof, M. J., 2000. The Impact of Predators on Maize Stem Borers in Coastal Kenya . Wageningen University, PhD thesis, The Netherlands, 181 pp.
- Borror, D. J., D. M. De Long & C. A. Triplehorn, 1981. An Introduction to the Study of Insects. Saunders College Publishing, 827 pp.
- Bournier, J. P., 1982. "Behaviour of various strains of *Trichogramma* towards eggs of *Heliothis armigera* Hbn (Lep. Noctuidae) and *Chilo partellus* Swinhoe (Lep. Pyralidae), 85-96" .Les trichogrammes. Ier Symposium International (10-15 November 1986, Guangzhou-Chine).
- Chang, S. C, N. T. Hu, C. Y. Hsin & C. N. Sun, 2001. Characterization of differences between two *Trichogramma* wasps by molecular markers. *Biological Control* 21 (1): 75–78.
- Divya, K., K. N. Marulasiddesha, K. Krupanidhi & M. Sankar, 2009. Population dynamics of spotted stem borer, *Chilo partellus* (Swinhoe) and its interaction with natural enemies in sorghum. *Indian Journal of Science and Technology*, 3 (1): 70-74.

- Edgar, R. C., 2004. MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, 32 (5): 1792-1797.
- FAO-STAT, 2014. Food and Agriculture Organization of the United States Statistics, (Web page: <http://faostat.fao.org>), (Date accessed: January 2016).
- Farid, A., M. I. N. Khan, A. Khan, S. U. K. Khattak & A. Sattar, 2007. Studies on maize stem borer, *Chilo partellus* in Peshawar Valley. *Pakistan Journal of Zoology*, 39 (2): 127-131.
- Grabenweger, G., P. Kehrl, I. Zweimüller, S. Augustin, N. Avtzis, S. Bacher, J. Freise, S. Girardoz, S. Guichard, W. Heitland, C. Lethmayer, M. Stolz, R. Tomov, L. Volter & M. Kenis, 2010. Temporal and spatial variations in the parasitoid complex of the horse chestnut leafminer during its invasion of Europe. *Biological Invasions*, 12: 2797-2813.
- Hebert, P. D. N., A. Cywinska, S. L. Ball & J. R. Deward, 2003. Biological identifications through DNA barcodes. *Proceedings Royal Society London Biological Sciences*, 270:313-321.
- Hoy, M.A., A. Jeyaprakash, R. Morakote, P. K. C. Lo & R. Nguyen, 2000. Genomic analyses of two populations of *Ageniaspis citricola* (Hymenoptera: Encyrtidae) suggest that a cryptic species may exist. *Biological Control* 17: 1-10.
- Jalali, S. K. & S. P. Singh, 2006. Biological control of *Chilo partellus* using egg parasitoid *Trichogramma chilonis* and *Bacillus thuringiensis*. *Indian Journal of Agricultural Research* 40 (3): 184-189.
- Kfir, R., 1990. Parasites of the spotted stalk borer, *Chilo partellus* (Lep: Pyralidae) in South Africa. *Entomophaga* 35 (3): 403-410.
- Khan, Z. R., P. Chiliswa, K. Ampong-Nyarko, L. E. Smart, A. Polaszek, J. Wandera & M. A. Mulaa, 1997. Utilisation of wild gramineous plants for the management of cereal stemborers in Africa. *International Journal of Tropical Insect Science*, 17 (1): 143-150.
- Kornoşor, S., E. Sertkaya & A. Özpinar, 1994. "Distribution of the egg parasitoid *Platytenomus busseolae* (Gahan) (Hym., Scelionidae) and its effect on the population of *Sesamia nonagrioides* Lef. (Lep., Noctuidae) in the Mediterranean Region of Turkey, 193-200". 4th. International Symposium on *Trichogramma* and Other Egg Parasites (4-7 October, Cairo, Egypt).
- Kumar, H. & J. A. Mihm, 1995. Antibiosis and tolerance to fall armyworm, *Spodoptera frugiperda* (J. E. Smith), southwestern corn borer, *Diatraea grandiosella* Dyar and sugarcane borer, *Diatraea saccharalis* Fabricius in selected maize hybrids and varieties. *Maydica*. 40: 245-251.
- Kumar, H., 2002. Resistance in maize to larger grain borer, *Prospahanus truncates* (Horn) (Coleoptera: Bostrichidae). *Journal of Stored Products Research*, 38: 267-280.
- Landry, B. S., L. Dextraze & G. Boivin, 1993. Random amplified polymorphic DNA markers for DNA fingerprinting and genetic variability assessment of minute parasitic wasp species (Hymenoptera: Mymaridae and Trichogrammatidae) used in biological control programs of phytophagous insects. *Genome*, 36 (3): 580-587.
- LaSalle, J. & I. D. Gauld, 1991. Parasitic Hymenoptera and the biodiversity crisis. *Redia*, 74 (3): 315-334.
- Leray, M., J. Y. Yang., C. P. Meyer, S. C. Mills, N. Agudelo, V. Ranwez, J. T. Boehm & R. J. Machida, 2013. A new versatile primer set targeting a short fragment of the mitochondrial COI region for metabarcoding metazoan diversity: application for characterizing coral reef fish gut contents. *Frontiers in Zoology*, 10: 34.
- Li, S. Y., D. E. Henderson & J. H. Meyer, 1994. Selection of suitable *Trichogramma* species for potential control of the blackheaded fire worm infesting cranberries. *Biocontrol*, 4 (3): 244-248.
- Masner, L. & N. F. Johnson, 1979. A new species of *Telenomus* (Hymenoptera: Scelionidae) an egg parasite of the Ambush bug, *Phymata* sp. (Heteroptera: Phymatidae). *Canadian Entomologist* 111 (10): 1115-1119.
- Midega, C. A. O., C. K. P. O. Ogot & W. A. Overholt, 2004. Effect of agroecosystem diversity on natural enemies of maize stemborers in coastal Kenya. *International Journal of Tropical Insect Science*, 4 (4): 280-286.
- Nasir, M. F., G. Hagedorn, C. Büttner, C. Reichmuth & M. Schöller, 2013. Molecular identification of *Trichogramma* species from Pakistan, using ITS-2 region of rDNA. *BioControl* 58 (4): 483-491.
- Neupane, F. P., H. C. Coppel & R. K. Chapman, 1985. Bionomics of the maize borer, *Chilo partellus* (Swinhoe), in Nepal. *International Journal of Tropical Insect Science*. 6 (4): 547-553.
- Pinto, J. D., R. K. Velten, G. R. Platner & E. R. Oatman, 1989. Phenotypic plasticity and characters in *Trichogramma*. *Annals of the Entomological Society America*, 85: 413-422.

- Pinto, J. D. & R. Stouthamer, 1994. "Systematics of the Trichogrammatidae with Emphasis on *Trichogramma*.1-36". In: Biological Control with Egg Parasitoids (Eds: E. Wajnberg & S. A. Hassan). CAB international, 266 pp.
- Pomari- Fernandes, A., A. F. Bueno, A. P. Queiroz & S. A. De Bortoli, 2015. Biological parameters and parasitism capacity of *Telenomus remus* Nixon (Hymenoptera: Platygastridae) reared on natural and factitious hosts for successive generations. African Journal of Agricultural Research, 10 (33): 3225-3233.
- Poorjavad, N., S. H. Goldansaz, T. Machtelinckx, L. Tirry, R. Stouthamer & T. van Leeuwen, 2012. Iranian *Trichogramma*: ITS2 DNA characterization and natural *Wolbachia* infection. BioControl, 57 (3): 361-374.
- Ratnasingham, S. & P. D. Hebert, 2007. BOLD: The Barcode of Life Data System (<http://www.barcodinglife.org>). Molecular Ecology Notes, 7 (3), 355-364.
- Rebe, M., J. Van Den Berg & M. A. McGeoch, 2004. Colonization of cultivated and indigenous graminaceous host plants by *Busseola fusca* (Fuller) (Lepidoptera: Noctuidae) and *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae) under field conditions. African Entomology, 12 (2): 187-199.
- Rwomushana, I., S. Kyamanywa, A. J. Ngi-Song & C. O. Omwega, 2005. Incidence of larval and pupal parasitoids of Lepidoptera in Uganda with special reference to new associations. International Journal of Tropical Insect Science, 25 (4): 274-280.
- Sertkaya, E., V. Akmeşe & E. Atay, 2014. Türkiye'de mısırdaki yeni bir zararlı, *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae). Türkiye Entomoloji Bülteni, 4 (3): 197-200.
- Seshu Reddy, K. V., 1988. Assessment of on-farm yield losses in sorghum due to insect pests. International Journal of Tropical Insect Science, 9: 679-685.
- Sétamou, M. & F. Schulthess, 1995. The influence of egg parasitoids belonging to *Telenomus busseolae* (Hym. Scelionidae) species complex on *Sesamia calamistis* (Lepidoptera: Noctuidae) population in maize fields in Southern Benin. Biocontrol Science and Technology 5: 69–81.
- Smith, S. M. & M. Hubbes, 1986. Strains of the egg parasitoid *Trichogramma minutum* Riley. Journal of Applied Entomology, 101 (1-5) : 223-239.
- Somchoudhury, A. K. & N. Dutt, 1988. Influence of hosts and host ages on the bionomics of *Trichogramma perkinsi* Girault and *Trichogramma australicum* Girault. Indian Journal of Entomology, 50 (3): 374-379.
- Sorokina, A. P., 1993. Keys to the Species of Genus *Trichogramma* Westw. (Hymenoptera: Trichogrammatidae) of the World Fauna. Kolos Publishing House, Moscow, 77 pp.
- Sümer, F., A. S. Tunçbilek, S. Öztemiz, B. Pintureau, P. Rugman-Jones & R. Stouthamer, 2009. A molecular key to the common species of *Trichogramma* of the Mediterranean region. BioControl, 54: 617-624.
- Tagne, A., T. P. Feujio & C. Sonna, 2008. "Essential oil and plant extracts as potential substitutes to synthetic fungicides in the control of fungi" ENDURE International Conference, Diversifying crop protection (12-15 October 2008, La Grande-Motte-France), 42. Oral presentation.
- Tamura, K., G. Stecher, D. Peterson, A. Filipski & S. Kumar, 2013. MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. Molecular Biology and Evolution, 30 (12): 2725–2729.
- TÜİK, 2014. Türkiye İstatistik Kurumu. (Web page: http://www.tuik.gov.tr/PreTablo.do?alt_id=1001), (Date accessed: January 2016).
- Valentini, A., F. Pompanon & P. Taberlet, 2009. Barcoding for ecologist. Trends in Ecology and Evolution, 24: 110–117.
- Van den Berg, J., A. F. Nur & A. Polaszek, 1998. "Cultural Control, 333–347" In: Cereal Stem Borers in Africa: Economic Importance, Taxonomy, Natural Enemies and Control (Ed. A. Polaszek). International Institute of Entomology. CAB International, 592 pp.
- Van den Berg, J., 2009. "Can Vetiver grass be used to manage insect pests on crops, 254-264". Third International Conference on Vetiver (6-9 October 2003, Guangzhou- China), China Agriculture Press, 572 pp.
- Wakeil-EI, N., N. Gaafar, A. Sallam & C. Volkmar, 2013. Side effects of insecticides on natural enemies and possibility of their integration in plant protection strategies. Insecticides-development of safer and more effective technologies-Chapter 1-January 2013. 56 pp.
- Yuliarti, N., P. Hidayat & D. Buchori, 2002. Molecular identification of egg parasitoid, *Telenomus* spp. (Hym.: Scelionidae) from several locations in Java using RAPD-PCR. Biotropia 19: 57-64.