

RESEARCH ARTICLE

Open Access



Musculoskeletal pain and re-employment among unemployed job seekers: a three-year follow-up study

Chioma A. Nwaru*, Clas-Håkan Nygård and Pekka Virtanen

Abstract

Background: Poor health is a potential risk factor for not finding employment among unemployed individuals. We investigated the associations between localized and multiple-site musculoskeletal pain and re-employment in a three-year follow-up of unemployed job seekers.

Methods: Unemployed people ($n = 539$) from six localities in southern Finland who participated in various active labour market policy measures at baseline in 2002/2003 were recruited into a three-year health service intervention trial. A questionnaire was used to collect data on musculoskeletal health and background characteristics at baseline and on employment status at the end of the follow-up. We conducted a complete case ($n = 284$) and multiple imputation analyses using logistic regression to investigate the association between baseline musculoskeletal pain and re-employment after three years.

Results: Participants with severe pain in the lower back were less likely to become re-employed. This was independent of potential confounding variables. Pain in the hands/upper extremities, neck/shoulders, lower extremities, as well as multiple site were not determinants of re-employment.

Conclusions: Our findings lend some support to the hypothesis that poor health can potentially cause health selection into employment. There is the need to disentangle health problems in order to clearly appreciate their putative impact on employment. This will allow for more targeted interventions for the unemployed.

Keywords: Unemployment, Musculoskeletal pain, Localized pain, Multiple pain sites, Re-employment

Background

Unemployment has a detrimental effect on the health and well-being of individuals [1], their spouses [2], their children [3, 4], and the public at large [5–7]. Prospective studies have shown that re-employment could improve the health of the unemployed. Evidence of such improvement has been demonstrated in both a five-year [8] and a ten-year [9] follow-up study, where a significant improvement in mental health was reported among the unemployed after they re-entered paid employment. Schuring et al. [10] and Carlier et al. [11] also demonstrated that re-employment improved physical health, hence suggested that labour force participation should be considered as a therapeutic measure within the health promotion framework for the unemployed.

Poor health is an important risk factor for not finding employment. According to the health selection theory, unemployed persons with poor health may be less likely hired by prospective employers, thus are at risk of being selected into prolonged spell of unemployment [8, 12]. Many studies have investigated health selection using mental or physical health as determining factors. Findings regarding mental health are inconsistent. In a two-year follow-up study in Norway, mental disorders and physician-diagnosed psychiatric syndromes or personality disorders were risk factors for not regaining employment among long-term (more than 12 weeks) unemployed people [13]. In a five-year follow-up of that study, only the doctor's diagnosis of psychiatric syndromes or personality disorders was however significantly associated with reduced re-employment [8]. In a three-year study in Finland, psychological distress was not associated with re-employment

* Correspondence: Chioma.nwaru@uta.fi
School of Health Sciences, University of Tampere, FI-33014 Tampere, Finland

among registered unemployed persons [14], but a twelve-year follow-up study in Britain reported an increased likelihood of re-employment among unemployed women with psychological distress [15].

Regarding physical health, van de Mheen et al. [16] reported that poor general health, a chronic condition, and health complaints were determinants of re-employment after 4.5 years. Similar findings were reported in the European Household survey, with poor health and chronic conditions as determinants of not entering paid employment in most European countries [17]. Poor general health [18, 19] and decreased work performance due to impaired health [20] have also been shown to reduce likelihood of re-employment. One limitation in these studies is that the indicators of physical health were measured in a general context, i.e. in terms of chronic health problems or general self-rated health, which despite being important and valid measures – do not give indication of the specific roles of the health problems and diseases.

Musculoskeletal pain is a widespread problem among the working population, and it is a known risk factor for poor work ability [21, 22], increased absence due to sickness [23], early retirement [24], and health-related job loss [25]. Musculoskeletal pain may also reduce the possibility of regaining employment, but the evidence emerge from studies conducted among persons with arthritis and musculoskeletal disorders who were unemployed [26] and those of pre-retirement age [27]. Generalizing these findings to the general unemployed population would require further studies among individuals with differential symptom patterns and unemployment histories. In the present study, we investigate whether localized and multiple-site musculoskeletal pain are associated with re-employment in a three-year follow-up of registered unemployed people aged 18 to 59 years in Finland.

Methods


Study design and subjects

The study data originated from the Career Health Care (CHC) project, a three-year intervention trial that was launched in 2002–2003 in Finland with the goal of tackling health problems and risks related to unemployment [28]. Participants in the project were unemployed people ($n = 539$) from six localities in southern Finland who were enrolled in active labour market policy (ALMP) measures. They were recruited at the beginning of the ALMP measures, during which they received oral and written information about the study. This information made it explicit that participation was voluntary and not a condition for participation in the ALMP or access to the associated benefits. Those who consented to the study were randomly allocated to the intervention and control groups. The intervention group ($n = 265$) were recipients of the CHC package (i.e. the extra health services that

targeted the unemployed). The control group ($n = 274$) only used communal health services. Both groups completed the baseline questionnaire during the recruitment exercise. Follow-up data was collected three years after the first encounter and 311 persons responded to this follow-up. The intervention group completed the follow-up questionnaires during the CHC encounter, and the control group returned their questionnaires by post. We excluded a group of respondents ($n = 27$) who were classified as non-job seekers at follow-up from the present study, because they were not at risk for unemployment. This gave rise to a sample of 284 people aged 18–59 years who responded to the three-year follow-up (see Fig. 1).

Measurements

We measured musculoskeletal complaints at baseline using a modified version of the Nordic Musculoskeletal Questionnaire [29]. Respondents were asked to report, on a scale of 0 to 10, whether they had experienced pain or numbness in four locations during the preceding week. The locations were the hands or upper extremities, neck or shoulders, lower back, and the feet or lower extremities. The response for each pain variable was categorized into three groups: 0 = no pain, 1–5 = mild pain, and 6–10 = severe pain. To construct a multiple site



pain measure, mild and severe categories were combined into any pain = 1 and no pain = 0. All four musculoskeletal pain variables were then added up and the summed variable was expressed as the number of sites with pain (from 0 = no pain in any site to 4 = pain in four sites).

Other variables that were measured at baseline and considered as potential confounders included age, gender, educational attainment, marital status, duration of unemployment, alcohol use, smoking, physical activity, somatic diseases, and depression. Age was categorized into three groups: “18–29”, “30–44”, and “45–59”. Educational attainment was classified into three levels: “college/university degree”, “vocational school degree”, and “no occupational degree”. Marital status was categorized as “single”, “married/cohabiting”, or “widowed/divorced”. Duration of unemployment was dichotomized to “less than one year” and “more than one year”. Alcohol use was elicited with the question “how often do you drink beer, wine or other alcoholic drinks?” The response was categorized into three: “never/less often”, “2–4 times a month”, and “2 or more times/week”. Smoking was dichotomized to “smokers” and “none smokers”, and leisure-time physical activity (i.e. frequency of vigorous physical activity for at least 15 to 20 min) was categorized into three: “not at all or only a little”, “moderate” (once per week), and “much” (twice or more per week). General health was assessed with the question “do you have diseases diagnosed by a physician?” A list of 18 different diseases was provided with a dichotomized reply (yes or no). We considered the responses that included one or more of the nine somatic diseases listed, i.e. cardiovascular illnesses, respiratory illnesses, diabetes, etc. (with the exception of musculoskeletal diseases). The sum score of the diseases was dichotomized (yes or no), and those subjects reporting one or more diseases were categorized as having somatic disease(s). Depression was measured with the Beck Depression Inventory [30] and dichotomized to “depressed” and “not depressed”.

Current employment status was determined in the three-year follow-up questionnaire and classified into two categories: “re-employed” and “unemployed”. Subjects were defined as re-employed if they reported being either employed or self-employed. The unemployed group consisted of those who reported not being in any paid job but seeking employment during the follow-up.

Statistical analysis

The description of the subjects’ characteristics are presented as frequencies and percentages, and differences between groups were tested with a chi-squared test for categorical variables. The association between musculoskeletal pain and re-employment was examined with binary logistic regression. Re-employment was coded in

such a manner that an odds ratio > 1 indicated an increased likelihood of re-employment. We conducted both complete-case (i.e. those who participated in both baseline and three-year follow-up) and multiple imputation (i.e. to impute data of the three-year follow-up for those who did not participate in the follow-up) analyses. The complete-case analysis was undertaken using IBM SPSS Statistics for Windows, version 20.0. (Armonk, NY: IBM Corp). In the complete-case analysis, unadjusted and adjusted models were performed. The unadjusted model (Model I) estimated the independent effect of the various localized pains, as well as the number of pain sites. The adjusted models included potential confounders in the model, with Model II simultaneously controlling for age, gender, educational attainment, and marital status. Model III additionally adjusted for the duration of unemployment, alcohol use, smoking, physical activity, somatic diseases, depression and participation in CHC. Although a recent study by Romppainen et al. (2014) did not find any beneficial effect of the CHC on re-employment, we also explored its role as a potential effect-modifying variable by entering an interaction term between musculoskeletal pain and participation in CHC in the adjusted model in relation to re-employment. If the interaction term was significant, we stratified the analysis by participation in CHC and calculated the stratum-specific estimates adjusting for all other confounders.

The multiple imputation (assuming missing at random) was conducted using the Multiple Imputation by Chained Equations (MICE) algorithm in Stata (version 13). A total of 20 imputed datasets were created. All variables that were used in the complete-case analysis, irrespective of whether they had missing or not, were included in the imputation model. After the imputation, we then repeated the Model III logistic regression analysis conducted with the complete-case analysis. An interaction term between musculoskeletal pain and participation in CHC was also investigated in the Model III of the multiple imputation model. Results are presented as odds ratios (OR) with their 95 % confidence intervals (CI), and their statistical significance was defined as the two-sided *p*-value < 0.05.

Results

At the three-year follow-up, 311 of the original 539 participants responded to the questionnaire survey (response rate 58 %). An analysis of non-respondents versus respondents showed a lower response rate among males (47 %) than among females (64 %), among smokers (50 %) than among none smokers (64 %), and among participants in the intervention (49 %) than among the control (66 %) group. Participants who were either widowed or divorced had the lowest response rate (47 %) compared to their counterparts who were single (52 %) or were married or cohabiting (64 %). Differences in other individual

characteristics (age, educational attainment, alcohol use, physical activity, somatic diseases, and depression) as well as musculoskeletal pain were not statistically significant (see Additional file 1: Table S1).

By excluding 27 (9 % of those who completed both questionnaire) ineligible respondents, who consisted of retirees or those receiving disability pension ($n = 9$), those on parental leave ($n = 7$), non-job seekers ($n = 1$), or those excluded for some other reason ($n = 10$), the subsequent analyses included 52 % (284/539) of the original study population. The baseline individual and health characteristics of the 284 respondents are given in Table 1. The participants were predominantly middle-aged (45 %, $n = 127$), with most of them (67 %, $n = 190$) having been unemployed for less than one year at baseline. Twenty-two percent of them had attained a college/university degree. In the week preceding the baseline measurement, 147 (52 %) reported mild-to-severe pain in the hands/upper extremities, 195 (69 %) in the neck/shoulders, 154 (52 %) in the lower back, and 141 (50 %) in the feet/lower extremities. Over half of the respondents (59 %, $n = 168$) had concurrent pain in two or more sites.

Participants with somatic diseases were more likely to report pain compared to those without somatic diseases, regardless of the pain type (Table 2). Reporting pain also increased with decreasing participation in vigorous physical activity although the differences were significant only for low back pain ($p = 0.016$) and lower extremity pain ($p = 0.047$). Other characteristics, such as age, gender, educational attainment, marital status, duration of unemployment, participation in CHC, alcohol use, smoking, and depression were not significant determinants of most musculoskeletal pain. Regarding employment status during the three-year follow-up, over half (55 %, $n = 156$) of the participants were re-employed. The likelihood of re-employment decreased with increasing age and decreasing educational attainment. Participants who were either widowed or divorced (40 %) were less likely to regain employment than those who were either single (49 %) or married/cohabiting (61 %).

Table 3 shows the results of the associations between musculoskeletal pain at baseline and re-employment after three years. Based on the unadjusted result, those with severe pain in the lower back or feet/lower extremities had a reduction of up to 59 % in the likelihood of re-employment. In the adjusted models, the reduced likelihood of re-employment with pain in the lower back (OR 0.37, 95 % CI 0.15–0.92) or feet/lower extremities (OR 0.38, 95 % CI 0.15–0.93) remained unchanged even after controlling for age, gender, educational attainment, marital status, duration of unemployment, participation in CHC, alcohol use, smoking, physical activity, somatic diseases, and depression. A reduced likelihood for re-employment was also found for those participants with

three (OR 0.48, 95 % CI 0.23–0.99) or four (OR 0.51, 95 % CI 0.27–0.99) pain sites, although these associations were not retained when adjustments for confounders were introduced into the model (Table 4). The interaction between participation in CHC and musculoskeletal pain was not significant for most pain types except for low back pain. When we stratified the analysis by participation in CHC, the estimated odds for finding employment was significantly lower for those individuals in the control group who had severe low back (OR 0.18, 95 % CI 0.04–0.77) (see Additional file 2: Table S2).

Results from the complete-case and multiple imputation analyses were generally similar to each other, except that the confidence interval for lower extremity pain included one in the multiple imputation analysis (complete-case: OR 0.38, 95 % CI 0.15–0.93; multiple imputation: OR 0.51, 95 % CI 0.22–1.16). In addition, the significant interaction effect between low back pain and participation in CHC observed in the complete-case analysis was not seen in the multiple imputation analysis, suggesting that the complete-case interaction may be a chance finding.

Discussion

We found that severe pain in the lower back was associated with a reduced likelihood of re-employment after three years among unemployed job seekers. Pain in the hands/upper extremities, neck/shoulders, the lower extremities, as well as multiple site did not influence re-employment. These results were similar both in complete-case and multiple imputation analyses.

We recorded a moderate but acceptable participation rate of 58 % at three-year follow-up, which is similar to those achieved in previous studies [31, 32]. Usually high drop-out rates have been observed for the unemployed [33, 34]. Although differences between participants and non-participant at the three-year follow-up were observed only for sex, marital status, smoking, and participation in the CHC, we undertook multiple imputation analysis to impute missing data for those who did not take part in the follow-up assessment. This provided us with relevant sensitivity analysis to appraise the extent of bias due to follow-up with the complete-case analysis. Our assessment of the subjects' musculoskeletal pain status was based on a self-report, which may introduce information bias, however self-reporting of pain indicators has been noted to be reliable [29] and it is commonly used for pain studies [24, 35, 36]. The time into the past (one week) participants were asked to recall any pain is short and therefore should minimize the risk of recall bias.

There may be the possibility of residual confounding since we could not assess the influence of all potential confounders, particularly body mass index, although previous studies [37] did not find an independent association

Table 1 Distribution of study participants by baseline socio-demographic and health characteristics

| Individual characteristics | Unemployed job-seekers (N = 284) n (%) |
|---|--|
| Age (years) | |
| 18–29 | 68 (23.9) |
| 30–44 | 127 (44.7) |
| 45–59 | 80 (28.2) |
| Missing | 9 (3.2) |
| Gender | |
| Male | 89 (31.3) |
| Female | 194 (68.3) |
| Missing | 1 (0.4) |
| Educational attainment | |
| No occupational education | 93 (32.7) |
| Vocational school | 120 (42.3) |
| College/university | 64 (22.5) |
| Missing | 7 (2.5) |
| Marital status | |
| Single | 82 (28.9) |
| Married/cohabiting | 170 (59.9) |
| Widowed/divorced | 30 (10.6) |
| Missing | 2 (0.7) |
| Duration of unemployment | |
| Less than one year | 190 (66.9) |
| More than one year | 94 (33.1) |
| Participation in CHC | |
| Intervention group | 119 (41.9) |
| Control group | 165 (58.1) |
| Lifestyle/health characteristics | |
| Alcohol use | |
| Never/less often | 113 (39.8) |
| 2–4 times/month | 128 (45.1) |
| 2 or more times/week | 43 (15.1) |
| Smoker | |
| No | 179 (63.0) |
| Yes | 105 (37.0) |
| Physical activity | |
| Much | 91 (32.0) |
| Moderate | 70 (24.6) |
| Not at all or only a little | 111 (39.1) |
| Missing | 12 (4.2) |
| Somatic diseases | |
| No | 153 (53.9) |
| Yes | 110 (38.7) |
| Missing | 21 (7.4) |

Table 1 Distribution of study participants by baseline socio-demographic and health characteristics (*Continued*)

| | |
|---|------------|
| Depression | |
| No | 253 (89.1) |
| Yes | 17 (6.0) |
| Missing | 14 (4.9) |
| Hands/upper extremity pain | |
| None | 120 (42.3) |
| Mild | 90 (31.7) |
| Severe | 57 (20.1) |
| Missing | 17 (6.0) |
| Neck/shoulder pain | |
| None | 75 (26.4) |
| Mild | 119 (41.9) |
| Severe | 76 (26.8) |
| Missing | 14 (4.9) |
| Low back pain | |
| None | 106 (37.3) |
| Mild | 106 (37.3) |
| Severe | 48 (16.9) |
| Missing | 24 (8.5) |
| Feet/lower extremity pain | |
| None | 126 (44.4) |
| Mild | 96 (33.8) |
| Severe | 45 (15.8) |
| Missing | 17 (6.0) |
| Number of musculoskeletal pain sites | |
| 0 | 74 (26.1) |
| 1 | 42 (14.8) |
| 2 | 46 (16.2) |
| 3 | 50 (17.6) |
| 4 | 72 (25.4) |

between body mass index and re-employment. The generalizability of our findings is equally limited owing to the fact that our data were based on unemployed persons who actively participated in various labour market policy measures. Hence, they constituted a relatively unique group that may not be representative of the unemployed population as a whole. Nonetheless, the findings of this study reflect evidence from unemployed people who still belong to the labour force. Vesalainen and Vuori [14] showed that the level of job-seeking activities might influence an individual's probability of finding a job. It is also possible that the level of job-seeking activities will vary among members of different unemployment groups. Our study excluded those in other unemployment groups such as retirees, those receiving disability pensions, those on

Table 2 Individual characteristics of participants by baseline musculoskeletal pain and re-employment at three-year follow-up

| | Baseline musculoskeletal pain | | | | | | | | | |
|--------------------------|---|---------|--|---------|--------------------------------------|---------|--|---------|--|---------|
| | % with no hands/upper extremity pain (n = 120) | p-value | % with no neck/shoulder pain (n = 75) | p-value | % with no low back pain (n = 106) | p-value | % with no feet/lower extremity pain (n = 126) | p-value | % re-employed at 3-year follow-up (n = 156) | p-value |
| Age (years) | | 0.079 | | 0.410 | | 0.420 | | 0.605 | | 0.001 |
| 18–29 | 54.5 | | 25.8 | | 43.3 | | 53.0 | | 67.6 | |
| 30–44 | 46.3 | | 30.3 | | 44.4 | | 49.6 | | 59.1 | |
| 45–59 | 37.5 | | 27.4 | | 35.3 | | 40.3 | | 38.8 | |
| Gender | | 0.819 | | 0.001 | | 0.286 | | 0.342 | | 0.222 |
| Male | 44.8 | | 42.4 | | 47.6 | | 41.4 | | 49.4 | |
| Female | 45.3 | | 20.7 | | 37.7 | | 50.3 | | 57.2 | |
| Educational attainment | | 0.098 | | 0.001 | | 0.167 | | 0.119 | | 0.051 |
| No occupational educ. | 40.9 | | 25.8 | | 38.8 | | 44.3 | | 46.2 | |
| Vocational educ. | 43.6 | | 24.3 | | 36.8 | | 45.5 | | 56.7 | |
| College/university | 54.0 | | 35.9 | | 54.8 | | 58.1 | | 65.6 | |
| Marital status | | 0.023 | | 0.702 | | 0.957 | | 0.174 | | 0.046 |
| Single | 38.0 | | 27.5 | | 42.3 | | 39.2 | | 48.8 | |
| Married/cohabiting | 50.3 | | 26.5 | | 39.4 | | 53.1 | | 60.6 | |
| Widowed/divorced | 35.7 | | 34.6 | | 46.2 | | 37.0 | | 40.0 | |
| Duration of unemployment | | 0.970 | | 0.081 | | 0.616 | | 0.745 | | 0.093 |
| Less than one year | 45.2 | | 24.2 | | 39.4 | | 47.2 | | 58.4 | |
| More than one year | 44.4 | | 35.2 | | 43.5 | | 47.2 | | 47.9 | |
| Participation in CHC | | 0.645 | | 0.898 | | 0.124 | | 0.163 | | 0.929 |
| Intervention group | 43.0 | | 26.3 | | 38.8 | | 40.9 | | 54.6 | |
| Control group | 46.4 | | 28.8 | | 41.4 | | 52.0 | | 55.2 | |
| Alcohol use | | 0.692 | | 0.363 | | 0.432 | | 0.456 | | 0.879 |
| Never/less often | 48.1 | | 25.7 | | 45.6 | | 43.4 | | 53.1 | |
| 2–4 times/month | 43.8 | | 25.4 | | 34.5 | | 46.7 | | 56.3 | |
| 2 or more times/week | 40.5 | | 39.5 | | 46.3 | | 58.5 | | 55.8 | |
| Smoker | | 0.819 | | 0.965 | | 0.442 | | 0.028 | | 0.679 |
| No | 43.9 | | 27.2 | | 38.0 | | 44.8 | | 55.9 | |
| Yes | 46.6 | | 28.7 | | 45.4 | | 51.0 | | 53.3 | |
| Physical activity | | 0.128 | | 0.068 | | 0.016 | | 0.047 | | 0.774 |
| Much | 55.4 | | 35.6 | | 53.6 | | 55.3 | | 54.9 | |

Table 2 Individual characteristics of participants by baseline musculoskeletal pain and re-employment at three-year follow-up (Continued)

| | | | | | | | | | |
|--------------------------|------|-------|------|-------|------|-------|------|-------|-------|
| Moderate | 41.8 | | 24.3 | | 36.4 | | 50.7 | | 58.6 |
| Not at all/only a little | 39.0 | | 20.0 | | 32.3 | | 36.5 | | 53.2 |
| Somatic diseases | | 0.006 | | 0.057 | | 0.002 | | 0.001 | 0.118 |
| No | 52.8 | | 32.7 | | 50.0 | | 54.5 | | 58.8 |
| Yes | 32.7 | | 19.2 | | 29.3 | | 35.2 | | 49.1 |
| Depression | | 0.059 | | 0.063 | | 0.220 | | 0.007 | 0.082 |
| No | 47.1 | | 29.3 | | 43.0 | | 49.8 | | 56.9 |
| Yes | 17.6 | | 17.6 | | 23.5 | | 17.6 | | 35.3 |

P-value by χ^2 tests

Table 3 Associations between localized pain at baseline and re-employment at three-year follow-up

| Localized musculoskeletal pain | Re-employment at 3-year follow-up | | | |
|--------------------------------|-----------------------------------|-----------------------|------------------------|--|
| | OR (95 % CI) | | | |
| | Model I ^a | Model II ^b | Model III ^c | Multiple imputation model ^d |
| Hands/upper extremity | | | | |
| None | 1.00 | 1.00 | 1.00 | 1.00 |
| Mild | 0.84 (0.48–1.47) | 1.45 (0.76–2.73) | 1.40 (0.69–2.87) | 1.22 (0.67–2.20) |
| Severe | 0.54 (0.28–1.02) | 0.63 (0.31–1.27) | 0.63 (0.28–1.38) | 0.54 (0.27–1.09) |
| Neck/shoulder | | | | |
| None | 1.00 | 1.00 | 1.00 | 1.00 |
| Mild | 0.93 (0.51–1.66) | 1.01 (0.52–1.94) | 0.87 (0.42–1.81) | 1.05 (0.50–2.23) |
| Severe | 0.78 (0.41–1.49) | 0.72 (0.35–1.49) | 0.99 (0.44–2.24) | 0.72 (0.41–2.32) |
| Low back | | | | |
| None | 1.00 | 1.00 | 1.00 | 1.00 |
| Mild | 0.92 (0.53–1.60) | 1.11 (0.61–2.04) | 0.96 (0.48–1.90) | 1.07 (0.50–2.29) |
| Severe | 0.41 (0.21–0.83) | 0.40 (0.18–0.88) | 0.37 (0.15–0.92) | 0.35 (0.16–0.78) |
| Feet/lower extremity | | | | |
| None | 1.00 | 1.00 | 1.00 | 1.00 |
| Mild | 0.73 (0.42–1.25) | 1.10 (0.60–2.01) | 1.20 (0.60–2.40) | 1.05 (0.48–2.29) |
| Severe | 0.41 (0.20–0.82) | 0.46 (0.21–0.98) | 0.38 (0.15–0.93) | 0.51 (0.22–1.16) |

^aUnadjusted model^bAdjusted for age, gender, educational attainment, and marital status^cAdjusted Model II + duration of unemployment, participation in CHC, alcohol use, smoking, physical activity, somatic diseases and depression

Models I, II, and III are based on complete-case analysis (N = 284)

^dAdjusted for age, gender, educational attainment, marital status, duration of unemployment, participation in CHC, alcohol use, smoking, physical activity, somatic diseases and depression (N = 539)

parental leave, non-job seekers, and those in other situations who are likely to adopt passive job-seeking behaviour.

Our findings of reduced re-employment among participants with severe lower back pain supports those of Straaton et al. [26], Yelin, Trupin & Sebasta [27], and Virtanen, Janlert, & Hammarstöm [37], who all showed that musculoskeletal pain was a determinant factor in regaining re-employment. The contribution of the present study is that it distinguished pain in local sites from that

in multiple sites, and provided insight into their respective roles in the relationship between health and employment. This is necessary considering that the differences in the risk factors and prognosis of the various pain types require different interventional measures.

A potential explanation why pain in the lower back was associated with a reduced likelihood of re-employment while pain in the other body regions (hands/upper extremities, neck/shoulders, and lower extremities) was not may

Table 4 Associations between number of musculoskeletal pain sites at baseline and re-employment at three-year follow-up

| Number of musculoskeletal pain sites | Re-employment at 3-year follow-up | | | |
|--------------------------------------|-----------------------------------|-----------------------|------------------------|--|
| | OR (95 % CI) | | | |
| | Model I ^a | Model II ^b | Model III ^c | Multiple imputation model ^d |
| 0 | 1.00 | 1.00 | 1.00 | 1.00 |
| 1 | 0.99 (0.45–2.16) | 0.82 (0.36–1.88) | 0.85 (0.35–2.10) | 0.86 (0.40–1.84) |
| 2 | 1.04 (0.48–2.22) | 1.26 (0.55–2.88) | 1.57 (0.61–4.02) | 1.05 (0.48–2.27) |
| 3 | 0.48 (0.23–0.99) | 0.58 (0.26–1.29) | 0.86 (0.35–2.09) | 0.68 (0.31–1.47) |
| 4 | 0.51 (0.27–0.99) | 0.72 (0.35–1.49) | 0.69 (0.29–1.61) | 0.66 (0.33–1.32) |

^aUnadjusted model^bAdjusted for age, gender, educational attainment, and marital status^cAdjusted Model II + duration of unemployment, participation in CHC, alcohol use, smoking, physical activity, somatic diseases and depression

Models I, II, and III are based on complete-case analysis (N = 284)

^dAdjusted for age, gender, educational attainment, marital status, duration of unemployment, participation in CHC, alcohol use, smoking, physical activity, somatic diseases and depression (N = 539)

be that low back pain may have persisted during periods of unemployment and thus, discouraged the motivation for finding employment. The occurrence of low back pain is not only associated with work-related factors, but also with psychological (anxiety, depression, emotional instability) and lifestyle-related (smoking and excess body weight) factors [38], which are prevalent among unemployed individuals [39, 40]. In addition, empirical evidence has shown that pain in the lower back is highly recurrent and rarely resolves [38], with some studies showing that low back pain may be associated with activity restriction [41]. It could be that these characteristics of low back pain may limit job search activities among individuals suffering from severe low back pain.

It was a surprising finding that the number of pain sites was not associated with re-employment considering the deleterious impact of pain on work and productivity [24, 42]. It is possible that pain in multiple sites is less burdensome during periods of unemployment due to reduced exposure to occupational factors that are considered major predisposing agents for pain in multiple sites [43].

Conclusion

In this study, we find that severe low back pain is a significant determinant of re-employment among unemployed job-seekers. This finding demonstrates the need to disentangle health problems in order to clearly appreciate their putative impact on employment. This is of paramount importance, especially for those health problems that may be modifiable. In further research, it would be helpful to understand whether similar associations may exist for chronic versus acute musculoskeletal pain.

Additional files

Additional file 1: Table S1. Relation of background characteristics to respondents to the questionnaire survey at 3-year follow-up. (DOCX 18 kb)

Additional file 2: Table S2. Modification of the effect of low back pain on re-employment by participation in CHC. (DOCX 16 kb)

Abbreviations

CHC, Career Health Care; ALMP, Active labour market policy; MICE, Multiple Imputation by Chained Equations

Acknowledgements

We would like to thank Dr Bright Nwaru for his helpful comments and statistical guidance.

Funding

This study was supported by grants from the Academy of Finland (grant no 207515), the Finnish Ministry of Employment and the Economy, the Finnish Ministry of Social Affairs and Health, and the Juho Vainio Foundation.

Availability of data and materials

The dataset that was used in this article is available from Dr Pekka Virtanen (pekka.j.virtanen@uta.fi) on request.

Authors' contributions

PV conceived this study and is the principal investigator of the CHC project; he prepared and delivered the data for the study and contributed to the analyses and writing of the manuscript. CN participated in the data analysis and was chiefly responsible for writing the manuscript drafts. CHN participated in the planning and implementation of the CHC project and contributed to the writing of the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors have no competing interests to declare.

Consent for publication

All authors have read and approved the final version of the paper being submitted for publication.

Ethics approval and consent to participate

At the time of the planning and implementation of the study, the Medical Research Act dealing with Ethics Committees had not yet come into force in Finland. There were Ethics Boards, which, however, were oriented narrowly to biomedical experiments, and this kind of study on health promotion services was not subjected to external ethical assessment. However, the Ethics Committee of Pirkanmaa University Hospital District assessed the study plan retrospectively, and stated that a study with a corresponding design would be approvable (ETL-code R13024). The study had a steering group consisting of representatives from the Ministry of Employment and the Economy and the Ministry of Social Affairs and Health.

Received: 17 November 2015 Accepted: 8 June 2016

Published online: 08 July 2016

References

- McKee-Ryan F, Wanberg CR, Song Z, Kinicki A. Psychological and physical well-being during unemployment: a meta-analytic study. *J Appl Psychol*. 2005;90(1):53–76.
- Marcus J. The effect of unemployment on the mental health of spouses—Evidence from plant closures in Germany. *J Health Econ*. 2013;32:546–58.
- Raatikainen K, Heiskanen N, Heinonen S. Does unemployment in family affect pregnancy outcome in conditions of high quality maternity care? *BMC Public Health*. 2006;6:46.
- Sleskova M, Salonna F, Madarasova A, Deckova M, et al. Does parental unemployment affect adolescents' health? *J Adolesc Health*. 2006;38:527–35.
- Kuhn A, Lalive R, Zweimuller J. The public health costs of job loss. *J Health Econ*. 2009;28:1099–115.
- Räisänen S, Kramer MR, Gissler M, Saari J, Heinonen S. Unemployment at municipality level is associated with an increased risk of small for gestational age births— a multilevel analysis of all singleton births during 2005–2010 in Finland. *Int J Equity Health*. 2014;13:95.
- Maruthappu M, Watkins J, Taylor A, Williams C, Ali R, Zeltner T, Atun R. Unemployment and prostate cancer mortality in the OECD, 1990–2009. *ecancer*. 2015;9:538.
- Claussen B. Health and re-employment in a five-year follow-up of long term unemployed. *Scand J Public Health*. 1999;27(2):94–100.
- Thomas C, Benzeval M, Stansfeld S. Psychological distress after employment transitions: the role of subjective financial position as a mediator. *J Epidemiol Community Health*. 2007;61:48–52.
- Schuring M, Mackenbach J, Voorham T, Burdorf A. The effect of re-employment on perceived health. *J Epidemiol Community Health*. 2011;65:639–44.
- Carlier BE, Schuring M, Lötters FJB, Bakker B, Borgers N, Burdorf A. The influence of re-employment on quality of life and self-rated health, a longitudinal study among unemployed persons in the Netherlands. *BMC Public Health*. 2013;13:503.
- Stewart JM. The impact of health status on the duration of unemployment spells and the implications for studies of the impact of unemployment on health status. *J Health Econ*. 2001;781–796.
- Claussen B, Bjorndal A, Hjort PF. Health and re-employment in a two year follow up of long term unemployed. *J Epidemiol Community Health*. 1993;47:14–8.
- Vesalainen J, Vuori J. Job-seeking, adaptation and re-employment experiences of the unemployed: a 3-year follow-up. *J Community Appl Soc Psychol*. 1999;9:383–94.

15. García-Gómez P, Jones AM, Rice N. Health effects on labor market exits and entries. *Labour Econ.* 2010;17:62–76.
16. van de Mheen H, Stronks K, Schrijvers CTM, Mackenbach JP. The influence of adult ill health on occupational mobility and mobility out of and into employment in The Netherlands. *Soc Sci Med.* 1999;49:509–18.
17. Schuring M, Mackenbach J, Voorham T, Burdorf A. The effects of ill health on entering and maintaining paid employment: evidence in European countries. *J Epidemiol Community Health.* 2007;61:597–604.
18. Schuring M, Robroek SJM, Otten Ferdy WJ, Arts CH, Burdorf A. The effect of ill health and socioeconomic status on labor force exit and re-employment: a prospective study with ten years follow-up in the Netherlands. *Scand J Work Environ Health.* 2013;39(2):134–43.
19. Lötters F, Carlier B, Bakker B, Borgers N, Schuring M, Burdorf A. The influence of perceived health on labor participation among long term unemployed. *J Occup Rehabil.* 2013;23:300–8.
20. Wagenaar AF, Kompier MAJ, Houtman ILD, van den Bossche SNJ, Taris TW. Who gets fired, who gets hired: the role of workers' contract, age, health, work ability, performance, work satisfaction and employee investments. *Int Arch Occup Environ Health.* 2015;88:321–34.
21. Neupane S, Miranda H, Virtanen P, Siukola A, Nygård CH. Multi-site pain and work ability among industrial population. *Occup Med.* 2011;61(8):563–9.
22. Lindegård A, Larsman P, Hadzibajramovic E, Ahlborg Jr G. The influence of perceived stress and musculoskeletal pain on work performance and work ability in Swedish health care workers. *Int Arch Occup Environ Health.* 2014;87:373–9.
23. Morken T, Riise T, Moen B, et al. Low back pain and widespread pain predict sickness absence among industrial workers. *BMC Musculoskelet Disord.* 2003;4:21.
24. Miranda H, Kaila-kangas L, Heliövaara M, et al. Musculoskeletal pain at multiple sites and its effects on work ability in a general working population. *Occup Environ Med.* 2010;67:449–55.
25. Solomon C, Poole J, Palmer KT, Coggon D. Health-related job loss: findings from a community-based survey. *Occup Environ Med.* 2007;64:144–9.
26. Straaton KV, Maisiak R, Wrigley JM, White MB, Johnson P, Fine PR. Barriers to return to work among persons unemployed due to arthritis and musculoskeletal disorders. *Arthritis Rheum.* 1996;101–109.
27. Yelin ED, Trupin LS, Sebesta DS. Transitions in employment, morbidity, and disability among persons ages 51–61 with musculoskeletal and non-musculoskeletal conditions in the US, 1992–1994. *Arthritis Rheum.* 1999;42(4):769–79.
28. Romppainen K, Saloniemä A, Kinnunen U, Liukkonen V, Virtanen P. Does provision of targeted health care for the unemployed enhance re-employment? *BMC Public Health.* 2014;14:1200.
29. Kuorinka I, Jonsson B, Kilbom A, et al. Standardized Nordic Questionnaires for the analysis of musculoskeletal symptoms. *Appl Ergon.* 1987;18:233–7.
30. Beck AT, Wald CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry.* 1961;4:53–63.
31. Skärlund M, Åhs A, Westerling R. Health-related and social factors predicting non-reemployment amongst newly unemployed. *BMS Public Health.* 2012;12:893.
32. Carlier BE, Schuring M, van Lenthe FJ, Burdorf A. Influence of health on job search behavior and re-employment: the role of job-search cognitions and coping resources. *J Occup Rehabil.* 2014;24:670–9.
33. Caetano R, Ramisetty-Mikler S, McGrath C. Characteristics of non-respondents in a US national longitudinal survey on drinking and intimate partner violence. *Addiction.* 2003;98:791–7.
34. Drivsholm T, Falgaard EL, Davidsen M, Jørgensen TI, Hollnagel H, Borch-Johnsen K. Representativeness in population-based studies: a detailed description of non-response in a Danish cohort study. *Scand J Public Health.* 2006;34(6):623–31.
35. Natvig B, Eriksen W, Bruusgaard D. Low back pain as a predictor of long-term work disability. *Scand J Public Health.* 2002;30(4):288–92.
36. Neupane S, Virtanen P, Leino-Arjas P, Miranda H, Siukola A, Nygård C-H. Multi-site pain and working conditions as predictors of work ability in a 4-year follow-up among industry employees. *Eur J Pain.* 2013;17(3):444–51.
37. Virtanen P, Janlert U, Hammarström A. Health status and health behavior as predictors of the occurrence of unemployment and prolonged unemployment. *Public Health.* 2013;127:46–52.
38. Woolf AD, Pfleger B. Burden of major musculoskeletal conditions. *Bull World Health Organ.* 2003;81(9):646–56.
39. Theodossiou I. The effects of low-pay and unemployment on psychological well-being: a logistic regression approach. *J Health Econ.* 1998;17:85–104.
40. Bartley M. Unemployment and ill health: understanding the relationship. *J Epidemiol Community Health.* 1994;48:333–7.
41. Gupta G, Nandini N. Prevalence of low back pain in non-working rural housewives of Kampur, India. *Int J Occup Med Environ Health.* 2015;28(2):313–20.
42. Haukka E, Kaila-Kangas L, Ojajarvi A, Miranda H, Karppinen J, Viikari-Juntura E, et al. Pain in multiple sites and sickness absence trajectories: a prospective study among Finns. *Pain.* 2013;154(2):306–12.
43. Solidaki E, Chatzi L, Bitsios P, Coggon D, Palmer K, Kogevinas M. Risk factors for persistent multisite pain in three occupational groups: CUPID study in Crete. *Occup Environ Med.* 2011;68:A8. doi:10.1136/oemed-2011-100382.23.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

