

Title: Genotype and diet affect resistance, survival, and fecundity but not fecundity tolerance

Author(s): Kutzer, M. A. M.; Kurtz, J.; Armitage, S. A. O.

Document type: Postprint

right to use is granted. This document is intended solely for personal, non-commercial use.

Citation: This is the peer reviewed version of the following article:
Kutzer, M. A. M., Kurtz, J., & Armitage, S. A. O. (2017). Genotype and diet affect resistance, survival, and fecundity but not fecundity tolerance. *Journal of Evolutionary Biology*, 31(1), 159–171. <https://doi.org/10.1111/jeb.13211>,
which has been published in final form at <https://doi.org/10.1111/jeb.13211>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

1 Genotype and diet affect resistance, survival, and fecundity but not
2 fecundity tolerance

3

4 Megan A. M. Kutzer^{1,2}, Joachim Kurtz¹ & Sophie A. O. Armitage^{1,3}

5

6 ¹ Institute for Evolution and Biodiversity, Hüfferstrasse 1, University of Münster, 48149

7 Münster, Germany

8 ² Current address: IST Austria (Institute of Science and Technology Austria), Am Campus 1,

9 3400 Klosterneuburg, Austria

10 ³ Current address: Institute of Biology, Freie Universität Berlin, Königin-Luise-Str. 1-3,

11 14195 Berlin, Germany

12

13 Corresponding author:

14 E-mail: sophie.armitage@fu-berlin.de

15 Telephone: -

16 Fax: +49 30 838461356

17 Running headline: Genotype affects resistance not tolerance

18 **Abstract**

19 Insects are exposed to a variety of potential pathogens in their environment, many of which
20 can severely impact fitness and health. Consequently, hosts have evolved resistance and
21 tolerance strategies to suppress or cope with infections. Hosts utilising resistance improve
22 fitness by clearing or reducing pathogen loads and hosts utilising tolerance reduce harmful
23 fitness effects per pathogen load. To understand variation in, and selective pressures on
24 resistance and tolerance we asked to what degree they are shaped by host genetic
25 background, whether plasticity in these responses depends upon dietary environment, and
26 whether there are interactions between these two factors. Females from ten wild-type
27 *Drosophila melanogaster* genotypes were kept on high or low protein (yeast) diets, and
28 infected with one of two opportunistic bacterial pathogens, *Lactococcus lactis* or
29 *Pseudomonas entomophila*. We measured host resistance as the inverse of bacterial load in
30 the early infection phase. The slope linking fly fecundity and individual-level bacteria load
31 provided our fecundity tolerance measure. Genotype and dietary yeast determined host
32 fecundity and strongly affected survival after infection with pathogenic *P. entomophila*.
33 There was considerable genetic variation in host resistance, a commonly found phenomenon
34 resulting from e.g. varying resistance costs or frequency-dependent selection. Despite this
35 variation and the reproductive cost of higher *P. entomophila* loads, the slopes linking bacteria
36 load and fecundity did not vary across genotypes. Absence of genetic variation in tolerance
37 may suggest that at this early infection stage fecundity tolerance is fixed or that any evolved
38 tolerance mechanisms are not expressed under these infection conditions.

39

40 **Keywords:** diet, DGRP, ecological immunology, fecundity tolerance, fitness, *Lactococcus*
41 *lactis*, pathogen, *Pseudomonas entomophila*, resistance, yeast.

42 **Introduction**

43 The composition of a hosts' microbial community is in part determined by how a host
44 responds towards invading microbes. Such host reactions towards microbes are composed of
45 resistance and tolerance, two disparate strategies whose deployment may ultimately depend
46 on a combination of intrinsic, innate factors and external, environmental factors (Råberg *et*
47 *al.*, 2009; Graham *et al.*, 2011; Kutzer & Armitage, 2016a). Once an infection becomes
48 established within a host, a host can actively resist the pathogen by clearance or by targeting
49 pathogen replication rate, which can aid host recovery time, but often comes at a cost to host
50 fitness (Kraaijeveld *et al.*, 2002). In contrast, host tolerance limits the deleterious fitness and
51 health effects of a pathogenic infection without targeting pathogen load (Roy & Kirchner,
52 2000; Råberg *et al.*, 2007). Both strategies can have far reaching impacts on host-pathogen
53 co-evolutionary trajectories (Best *et al.*, 2014).

54

55 The co-evolution of host resistance and pathogen virulence has been well characterised (e.g.
56 Masri *et al.*, 2015; Woolhouse *et al.*, 2002). Briefly, when a host resists a pathogenic
57 infection, it reduces pathogen prevalence in a population. After the pathogen counter-adapts
58 to circumvent the host resistance mechanisms, pathogen frequency increases in the host
59 population, resulting in a negative feedback loop and antagonistic co-evolution in both the
60 host and pathogen populations (Roy & Kirchner, 2000). Resistance mechanisms can be highly
61 host-pathogen specific or they can be more general. For example, resistance can result from
62 allelic variation in only a few loci (Luijckx *et al.*, 2013), the same antimicrobial peptide
63 (AMP) can increase in expression to a range of different pathogens (e.g. Lemaitre, Reichhart
64 & Hoffmann 1997), and different AMPs can act synergistically against one pathogen (Marxer
65 *et al.*, 2016).

66

67 Models predict that the evolution of host tolerance can act in two ways in a population,
68 depending upon whether hosts show fecundity- or mortality-tolerance (Best *et al.*, 2010,
69 2014). Mortality-tolerance is the ability to reduce the negative effect of infection on host
70 survival, and is important for pathogen prevalence. If an infected host lives longer, then the
71 pathogen also has a greater chance of being transmitted among hosts, which could lead to
72 disease reservoir expansion, and greater mortality in the host population (Roy & Kirchner,
73 2000; Miller *et al.*, 2006; Best *et al.*, 2008; Vale *et al.*, 2011). Fecundity tolerance, which we
74 test here and is the ability to reduce the negative effect of infection on host fecundity, should
75 be neutral to pathogen prevalence because the pathogen's infectious period is neither
76 prolonged nor shortened (Best *et al.*, 2010). However, if fecundity tolerance comes at a cost
77 to host lifespan, the pathogen's infectious period will be reduced along with host lifespan,
78 which can lead to a negative feedback and potentially, genetic variation in fecundity tolerance
79 (Best *et al.*, 2008).

80

81 Numerous mechanisms can lead to disease tolerance in animals, and these seem to be
82 dependent on pathogen and host type (Ayres & Schneider, 2012). Both hosts and pathogens
83 have optimal fitness strategies (Råberg, 2014), but these need not be fixed in their respective
84 populations (Best *et al.*, 2008). Host tolerance can be, but is not necessarily, genetically
85 determined (e.g. Råberg *et al.* 2007; Blanchet, Rey & Loot 2010; Sternberg *et al.* 2013;
86 Howick & Lazzaro 2014; Parker, Garcia & Gerardo 2014), and it can also be a plastic
87 response, where its expression is determined by the host environment, for example
88 concentration of dietary glucose (Howick & Lazzaro, 2014) or yeast (Kutzer & Armitage
89 2016b). Thereby, variation in defense strategies within and between populations can be
90 attributed to genetic (G) and environmental (E) factors or a combination of both (i.e. G x E
91 interactions), but studies exploring how different populations express resistance and tolerance

92 in response to changing environmental factors are under-represented (but see Howick &
93 Lazzaro 2014). Genetic variation in host immune function in particular can be maintained and
94 selected for by fluctuations in the host environment (Mitchell *et al.*, 2005; Lazzaro & Little,
95 2009; Hawley & Altizer, 2011; Sadd, 2011).

96

97 Resource availability and acquisition are important for mounting and maintaining an effective
98 immune response. Hosts can mask the deleterious effects of infection by increasing their
99 resource intake (Ayres & Schneider, 2009; Bashir-Tanoli & Tinsley, 2014), therefore
100 manipulating dietary components like protein or carbohydrates may uncover trade-offs or
101 costs that are not present under *ad libitum* conditions (Moret & Schmid-Hempel 2000;
102 Sternberg *et al.* 2012; Howick & Lazzaro 2014; Kutzer & Armitage 2016b). Such
103 physiological trade-offs (i.e. immune function versus fitness) are central to life history theory,
104 and can be either genetically fixed or variable, which will ultimately determine if the trade-off
105 is selected for in a population (Flatt *et al.*, 2011). In *Drosophila melanogaster*, experimental
106 dietary manipulation has mixed effects on the immune response, giving weight to the idea that
107 these relationships are largely context dependent (Vale *et al.*, 2011). For example, dietary
108 yeast restriction uncovered pathogen dependent, intra-genotypic variation in host tolerance
109 but not resistance in a single population of flies infected with *Escherichia coli* (Kutzer &
110 Armitage, 2016b), but in a separate study, resistance to *E. coli* was improved in flies with *ad*
111 *libitum* access to food compared with their counterparts on standard medium (McKean &
112 Nunney, 2005). Infections can impose considerable costs on hosts by competing for host
113 resources, decreasing host reproductive output, and causing host death (Stearns, 1992; Hurd,
114 2009), so hosts may use different immune strategies depending on an infection's pathology.
115 That is, fecundity compensation or reduction may be caused by infection or it may be a host
116 strategy (Hurd, 2001), which should be intimately connected to host defense strategies like

117 resistance and tolerance. For example a pathogenic infection may result in a host allocating
118 resources away from resistance to reproduction, appearing tolerant in the short-term (Vale &
119 Little, 2012; Leventhal *et al.*, 2014).

120

121 Resistance and tolerance can be plastic responses, changing over the course of an infection
122 (Howick & Lazzaro, 2014; Lough *et al.*, 2015; Kutzer & Armitage, 2016b; Louie *et al.*,
123 2016), but we were curious to know to what extent these responses show environmental
124 plasticity and genetic variability. Therefore, our novel approach was to test whether dietary
125 restriction through yeast (protein) limitation affects resistance and tolerance, and examine the
126 environmental interaction with genotype by testing ten wild-type *D. melanogaster* genotypes.
127 We infected flies with one of two opportunistic bacterial pathogens,
128 *Pseudomonas entomophila* and *Lactococcus lactis*, with different infection progressions and
129 contrasting short-term pathogenicity, and examined acute-phase resistance and tolerance to
130 infection to explore the extent to which these strategies are affected by genotype and the
131 environment. Here we defined acute phase infections as early stage infections occurring
132 between 0 and 72 hours post infection when pathogen levels are at their peak (e.g. Howick &
133 Lazzaro 2014). We measured range tolerance (Little *et al.*, 2010), where the slope of the
134 regression that results from the pathogen load and fecundity for every individual in the group,
135 describes tolerance for each treatment group (Råberg *et al.*, 2007; Graham *et al.*, 2011;
136 Lefèvre *et al.*, 2011). This provides more information than a single mean value for bacterial
137 load because host tolerance is measured over a range of pathogen loads. A group with a
138 steeper negative slope is less tolerant than a group with a flatter slope, because the former
139 loses their fitness more rapidly as pathogen load increases. Our use of range tolerance
140 contrasts with other studies on *D. melanogaster* (except Kutzer & Armitage 2016b), which
141 used means per group or different individuals for estimates of fecundity and bacteria load

142 (Corby-Harris *et al.*, 2007; Ayres & Schneider, 2008; Howick & Lazzaro, 2014). Because the
143 expression of defense strategies could be determined by a combination of infection pathology,
144 resource availability, and genetic factors, we predicted that 1. there is genetic variation for
145 resistance and tolerance, 2. dietary restriction would uncover costs manifested as reduced
146 fecundity tolerance in response to infection with a pathogenic bacterium, *P. entomophila*, and
147 3. dietary restriction may uncover trade-offs between resistance and tolerance. We find that
148 while fecundity and survival are determined by host genotype and dietary yeast, resistance is
149 largely genetically determined, and fecundity tolerance is unaffected by either genotype or
150 environment.

151

152 **Methods**

153 *Drosophila melanogaster* culture conditions

154 We used ten wild-type populations. The locally-collected population used in this study
155 (1_4WS; Kutzer & Armitage 2016b) was maintained in a population cage with overlapping
156 generations. Nine populations with variable fecundity (Ral208, Ral350, Ral367, Ral373,
157 Ral375, Ral379, Ral406, Ral509, Ral765) from the *Drosophila* Genetic Reference Panel
158 (DGRP) originating from North Carolina, USA (Mackay *et al.*, 2012) were maintained in
159 vials and placed onto new food every two weeks. For the purposes of this study we consider
160 each of these populations as being a distinct genotype (e.g. Mackay *et al.*, 2012), but we note
161 that the populations will inevitably differ from one another not only in their genetics, but also
162 in factors such as the microbiota that they contain. Therefore, we use ‘genotype’ in a broader
163 sense. All stocks were kept at 25 °C, 70 % relative humidity on a 12-12 hour light-dark cycle,
164 and were reared on a standard sugar, yeast, agar medium (SYA medium: 1.5 % agar, 5 %
165 sugar, 10 % brewer’s yeast [inactive *Saccharomyces cerevisiae* that is approximately 45 %

166 protein], 3 % nipagin, 0.3 % propionic acid) (Bass *et al.*, 2007). The procedures described
167 below were repeated independently to give a total of seven experimental replicates.

168

169 *Experimental animals and dietary treatments*

170 The individuals used in the experiment, as well as their parents, were reared at constant larval
171 density following protocols described in Kutzer & Armitage (2016b) with the following
172 modifications. Between 300 – 500 flies from each DGRP genotype were placed in embryo
173 collection cages to generate the F1 generation for each of the seven replicates. We collected
174 approximately 400 to 500 larvae of each of the ten genotypes for both the F1 and F2
175 generations. After the F2 generation eclosed, virgin females were allocated in groups of 20 to
176 one of the two dietary treatments, SYA or reduced yeast (RY) medium. RY medium
177 contained 25 % of the yeast contained in the SYA medium (Kutzer & Armitage, 2016b).
178 Males were kept in groups of 20 on SYA medium until mating.

179

180 *Mating assay and diet treatments*

181 Five to six days after adult eclosion we performed group mating assays at room temperature.
182 Beginning at 9:00 am, 10 male flies were placed into vials with 10 virgin females and allowed
183 30 minutes to mate. Limiting the time to 30 minutes decreased the chance of remating, which
184 could affect the immune response (Short *et al.*, 2012). Female and male flies were separated
185 by brief CO₂ anaesthetization. Males were discarded and females were individualised on a RY
186 diet or on an added yeast (AY) diet. The AY diet was SYA medium supplemented with active
187 baker's yeast granules, giving *ad libitum* access to yeast.

188

189 *Bacterial preparation and infections*

190 We chose two infective bacteria species with distinct infection dynamics. *L. lactis* does not
191 cause significant host death between 0 and 24 hours post-injection (hpi) when injected with
192 the dose we use below, but replicates rapidly in the host from 0 to 24 hpi (Kutzer & Armitage,
193 2016b). *P. entomophila* is comparatively more pathogenic hence the lower injection dose
194 used below, and can cause host death beginning approximately 20 - 22 hpi (personal
195 observation). Our *L. lactis* strain (gift from Brian Lazzaro) was isolated from a wild caught *D.*
196 *melanogaster* in State College, Pennsylvania (Lazzaro, 2002). The *P. entomophila* strain was
197 isolated from a wild caught fruit fly in Guadeloupe (gift from Bruno Lemaitre) (Vodovar *et*
198 *al.*, 2005; Vallet-Gely *et al.*, 2008). Both are opportunistic pathogens of *D. melanogaster*.
199 Aliquots of *L. lactis* and *P. entomophila* were stored in 34.4 % glycerol at -80 °C. *L. lactis*
200 was plated on lysogeny broth (LB) agar and *P. entomophila* was plated on LB agar containing
201 1 % milk to select for protease positive clones (Neyen *et al.*, 2012), after which, bacterial
202 preparation and infections were carried out following Kutzer & Armitage (2016b) using a
203 randomized block design of 60 total treatment groups (10 genotypes x 2 diets x 3 infection
204 treatments). In each experimental replicate we processed 3 flies per treatment group, giving
205 21 flies per genotype x diet x infection treatment, i.e. 1260 flies in total. A volume of 18.4 nL
206 of bacterial or a control solution was injected into the lateral side of the thorax using a fine
207 glass capillary attached to a Nanoject II™ (Drummond). For *P. entomophila* we injected 18.4
208 nL of a 5×10^6 cells mL⁻¹ bacterial solution where the bacteria was suspended in *Drosophila*
209 Ringer's solution (Werner *et al.*, 2000), which was equivalent to approximately 92 bacteria
210 per individual. In preliminary experiments with the 1_4WS genotype we found that this dose
211 resulted in about 10 % mortality 24 hpi. Flies infected with *L. lactis* were injected with 18.4
212 nL of a 1×10^8 cells mL⁻¹ bacterial solution, which was equivalent to 1840 bacteria per fly.
213 Control flies were injected with 18.4 nL of *Drosophila* Ringer's solution. Females were
214 returned to 25 °C, 70 % relative humidity after infection. We diluted the leftover injection

215 bacteria aliquots to 1×10^3 cells/ mL⁻¹ and plated 50 μ l of each on LB plates, which should
216 have yielded 50 CFUs. Bacterial counts from each aliquot ranged from 30 to 76 CFUs for *L.*
217 *lactis* and 23 and 67 for *P. entomophila*. We found no evidence of contamination for any
218 replicate.

219

220 *Fitness measure*

221 We measured pre-infection fitness as the total number of adult offspring produced by females
222 in the ~26 hours between mating and injections. Infected fitness was the total number of adult
223 offspring produced by each individual female in the 24 hpi. After we had removed females
224 from their vials for the bacterial load assay (below), the vials were kept at 25 °C until the
225 offspring had completed development and eclosed. Flies on the AY medium were given 12
226 days to complete development and those on the RY medium were given 17 days. The vials
227 were then turned upside down, frozen, and the offspring were counted after the experiment
228 ended.

229

230 *Standardising fecundity for fecundity tolerance*

231 There were considerable genotypic differences in the number of adult offspring produced by
232 uninfected flies (general vigour), which will partly determine fly fecundity when infected.
233 Following the example of Graham *et al.* (2011) we therefore assessed fecundity as the cost of
234 infection. We standardised the values by calculating the percent change in adult offspring
235 number relative to uninfected Ringer's controls as our response variable. The calculations
236 were performed using the mean fecundity of the Ringer's group for each genotype/diet
237 combination. The percent change for each individual was therefore calculated as $((\text{individual}$
238 $\text{infected fecundity } \omega_i - \text{mean Ringer's group fecundity } \omega_0) / \omega_0) \times 100$. For statistical reasons,

239 we standardized the change in fitness in this way to make the fecundity values more
240 comparable across the two dietary treatments (Kutzer & Armitage 2016).

241

242 *Bacterial load assay*

243 We assayed bacterial load at 24 hpi. The inverse of load determines the resistance of
244 individual flies (methods as described in Kutzer & Armitage 2016b). In brief, after surface
245 sterilisation, we serially diluted homogenates of whole flies infected with *L. lactis* in LB
246 medium at 1:1, 1:100 and 1:1000 and homogenates of flies infected with *P. entomophila* were
247 diluted 1:1 and 1:50 for each replicate. We plated 50 µl of each dilution onto LB agar and
248 incubated the plates at 30 °C for 20 hours and then counted bacterial colony forming units
249 (CFUs). We did not homogenize control flies injected with *Drosophila* Ringer's solution
250 because we found from previous work that these were usually negative for bacterial growth
251 (e.g. Kutzer & Armitage, 2016b, 7 % of all treatment groups had colony morphology that was
252 inconsistent with the injected bacteria). If a plate contained too many CFUs to count at the
253 highest dilution (~2%), we assigned the value as the greatest number of CFUs counted in the
254 genotype/treatment group (e.g. Vincent & Sharp, 2014).

255

256 *Statistical analyses*

257 Statistical analyses were performed in R version 3.3.1 (R Core Team, 2016). The statistical
258 models are detailed in Appendix S1, and model parameter estimates and standard errors are in
259 Tables S1-S4. Because of the substantial mortality in flies infected with *P. entomophila* on
260 RY medium we removed this group from all analyses except for tests on survival.

261

262 *Genome wide association study for resistance to P. entomophila infection*

263 We took advantage of the availability of whole genome sequences for the 9 DGRP genotypes,
264 and performed an exploratory genome wide association study to test for associations between
265 SNPs/INDELs and resistance to *P. entomophila*, using median bacterial load per genotype for
266 the AY environment. We used *P. entomophila* load because we reasoned that it showed the
267 strongest phenotypic differences across genotypes (see Results).

268

269 **Results**

270 *Effect of diet and genotype on survival*

271 Survival 24 hpi with Ringer's or *L. lactis* was high (mean % survival \pm SE: Ringer's: $98.8 \pm$
272 0.47 %; *L. lactis*: 98.6 ± 0.61 %) and unaffected by either diet or fly genotype (Figs 1A and
273 B; Table 1). However, after infection with *P. entomophila*, the genotypes differed in the
274 degree to which diet reduced their survival (Fig. 1C, interaction between genotype and diet in
275 Table 1). A reduced yeast diet strongly reduced survival, and there were significant
276 differences in how well the genotypes could survive infection over this short time (Fig. 1C,
277 Table 1).

278

279 *Effect of diet and genotype on bacterial load and bacterial load correlations*

280 Fly genotypes varied in their *L. lactis* bacterial loads (Fig. 2A, Table 2). However, we found
281 no evidence for a dietary effect or genotype-by-diet interaction on *L. lactis* load. There was a
282 marked difference in *P. entomophila* load across genotypes (Fig. 2B, Table 2). We were not
283 able to test whether there was a dietary effect on bacterial load because of the high mortality
284 we observed in the RY treatment (Fig. 1C). There was no relationship between *L. lactis* and
285 *P. entomophila* load across the ten genotypes using mean (Spearman's rank correlation, $\rho =$
286 -0.55 , $p = 0.10$) or median load ($\rho = 0.17$, $p = 0.65$).

287

288 *Effect of diet, infection status and genotype on fecundity*

289 Post-injection fecundity, i.e. adult offspring, was unaffected by infection treatment (Ringer's,
290 *L. lactis*, *P. entomophila*) in flies on AY media (Table 3, Model 3a). However, fecundity
291 varied significantly across genotypes (Figs 3A, B and C; Table 3) and there was a strong
292 positive correlation between pre- and post-infection fecundity (Table 3). We found
293 interactions between diet x genotype and genotype x infection status when comparing post-
294 infection fecundity among dietary treatments in flies injected with Ringer's solution or *L.*
295 *lactis* (Figs 3A, B, D and E, Table 3, Model 3b). Diet, genotype and pre-infection fecundity
296 were also significant predictors of post-infection fecundity in this model. A number of
297 females did not produce adult offspring, which may have been due to protein restriction
298 and/or the possibility that the flies did not mate during the group mating assay. However,
299 female fecundity after group matings in this experiment was comparable to fecundity after
300 observations of single pair matings: in this experiment 17 % of 1_4WS females on RY
301 medium produced zero offspring, which is similar to a previous experiment in which we
302 observed single pair matings (22 % with zero offspring, genotype 1_4WS on RY medium,
303 (Kutzer & Armitage, 2016b)

304

305 *Effect of diet and genotype on fecundity tolerance*

306 We found no effect of bacterial load on fecundity tolerance to *L. lactis*. Tolerance towards *L.*
307 *lactis* did not vary by genotype or diet, or a combination of both, which is illustrated by the
308 lack of significant interactions between these factors and bacterial load (Fig. 4A, Table 4).
309 However, genotype and diet affected percent change in adult offspring (Table 4).

310

311 Fecundity tolerance to *P. entomophila* tended to decrease as bacteria load increased, which
312 was independent of genotype (Fig. 4B). We observed no effect of genotype on fecundity
313 tolerance to a *P. entomophila* infection (Table 4).

314

315 *Genome wide association study for resistance to P. entomophila infection*

316 We found no significant associations between median *P. entomophila* load per genotype and
317 any of the SNPs or INDELS present in those genotypes (Fig. S1), which is most likely due to
318 a lack of power from only nine genotypes. Therefore, we do not discuss this analysis further.

319

320 **Discussion**

321 We tested the degree to which host genotype, dietary environment and G x E interactions
322 influence survival, fecundity, resistance, and fecundity tolerance to two acute phase bacterial
323 infections in ten wild-type *D. melanogaster* populations. Host genotype strongly predicted
324 variation in fecundity and resistance, but not tolerance, after infection with both *L. lactis* and
325 *P. entomophila*. As expected, a lower dietary yeast environment reduced fecundity, however,
326 it did not affect host resistance or tolerance to *L. lactis*. In contrast, lower dietary yeast
327 markedly reduced survival after a *P. entomophila* infection, which in combination with
328 genotype and a G x E effect on survival suggests that dietary environment and genetic
329 background play central roles in host defense during the early stages of a pathogenic
330 infection.

331

332 *Host genotype and diet determine survival after P. entomophila infection*

333 The importance of nutrition on traits including immune function, reproduction, and lifespan
334 cannot be disputed (Moret & Schmid-Hempel, 2000; Siva-Jothy & Thompson, 2002; McKean
335 *et al.*, 2008; Ayres & Schneider, 2009; Sadd, 2011; Sternberg *et al.*, 2012; Stahlschmidt *et al.*,

336 2013; Howick & Lazzaro, 2014; Kutzer & Armitage, 2016b). In our study, adult dietary yeast
337 restriction had a stronger negative effect on the survival of some genotypes infected with *P.*
338 *entomophila*, indicating a G x E interaction for this phenotype, which may suggest genotypic
339 variation for adult nutritional acquisition or energy storage (e.g. Bashir-Tanoli & Tinsley,
340 2014; Unckless *et al.*, 2015a, b). Furthermore, lower yeast availability consistently reduced
341 the survival of *P. entomophila*-infected flies across the ten genotypes tested, suggesting that
342 resource acquisition and/or availability is an important determinant of infection outcome in
343 this host-pathogen interaction and highlights the importance of considering the environmental
344 context in immune studies. It is interesting that survival after systemic infection of *P.*
345 *entomophila* in eight of the DGRP genotypes found in this study is positively correlated
346 (Spearman's rank correlation: $\rho = 0.84$, $p = 0.009$) with survival three days after an oral *P.*
347 *entomophila* infection using the same genotypes in another study (Sleiman *et al.*, 2015,
348 Ral367 was not used by these authors), despite the fact that the studies used different
349 infection routes and were done in different laboratories. Survival in the *L. lactis* and Ringer's
350 treatments was high and unaffected by any of our experimental factors.

351

352 *Genotypic variation in resistance to acute infections*

353 Evolutionary models predict that individuals within a host population will vary in their ability
354 to ward off infection (Miller *et al.*, 2007; Duffy & Forde, 2009; Boots *et al.*, 2012). In
355 addition, environmental heterogeneity can alter the expression of host susceptibility to
356 pathogens and resistance. However, adult dietary manipulation did not result in diet-induced
357 variability in resistance in the *L. lactis* infected groups, or G x E effects, and we were unable
358 to test this hypothesis explicitly in the *P. entomophila* infected groups because of high
359 mortality on the reduced yeast diet. The *L. lactis* observations were consistent with the results
360 of a previous study, where reduced access to protein did not affect bacterial load within a

361 single wild-type genotype (Kutzer & Armitage, 2016b). We found genotypic variation for
362 resistance (the inverse of bacteria load) in response to acute stage infection with both bacteria
363 species. Resistance is predicted to vary among genotypes or populations (Miller *et al.*, 2007;
364 Duffy & Forde, 2009; Boots *et al.*, 2012; Vale & Little, 2012), so it is not unexpected that
365 these 10 genotypes exhibit variation in their capacity for resistance. The genotypic variation
366 in resistance to a *P. entomophila* infection was considerable, highlighting the importance of
367 testing infections across different host genetic backgrounds and the difficulty in making
368 generalisations from single genotypes, as suggested by Sleiman *et al.* (2015). The five-fold
369 change in median *P. entomophila* load from the most to the least resistant genotype could
370 result from both variation in host immune responses and, potentially, the ability of the
371 bacteria to grow inside the host, for example if resources available for bacterial growth vary
372 across hosts. Genotypic variation in resistance is pervasive, it has been found across *D.*
373 *melanogaster* genotypes (Lazzaro *et al.*, 2006; Magwire *et al.*, 2012; Hotson & Schneider,
374 2015; Unckless *et al.*, 2015b), including those orally infected with *P. entomophila* (Sleiman *et*
375 *al.*, 2015), or injected with *L. lactis* (Lazzaro *et al.*, 2006). Lazzaro, Sackton & Clarke (2006)
376 found that many genotypes infected with *L. lactis* displayed a narrow phenotypic distribution,
377 which they suggest is driven by high bacteria loads. In the present study, we observed high
378 loads and a flat distribution at the maximum load, and there seemed to be an infection ceiling
379 of approximately 1×10^8 bacteria per fly, which may indicate that *L. lactis* reaches a growth
380 plateau within the fly independent of genetic variation. Furthermore, the individual bacterial
381 loads of *L. lactis* infected flies were surprisingly variable, especially given that the DGRP
382 genotypes were initially inbred for 20 generations. Individual variation for resistance
383 therefore seems surprising, but it is noteworthy that considerable phenotypic variation within
384 DGRP genotypes has been observed for traits such as antiviral resistance and susceptibility,

385 sleep, and food acquisition (Magwire *et al.*, 2012; Harbison *et al.*, 2013; Garlapow *et al.*,
386 2015).

387

388 *Genotype and diet, but not infection, influence post-infection fecundity*

389 Hosts must balance the costs of mounting an immune response, infection clearance, and
390 repairing infection induced- or self- damage, with life history traits such as reproduction.

391 Insects are sensitive to changes in their dietary environment and rapidly adjust egg production
392 accordingly (e.g. Kutzer & Armitage 2016a), therefore we predicted that adult offspring

393 numbers would decrease in response to protein restriction and that there would be variation
394 across genotypes, which was the case. We observed a marginal effect of infection and

395 genotype on host fecundity but this may have been driven by the addition of diet in the model,

396 as we found no interaction effect when we compared fecundity within the *ad libitum* yeast

397 groups. This may have been due to the ability of the flies to compensate for the effects of

398 infection with a pathogenic bacterium like *P. entomophila*, which reduces survival (e.g.

399 McKean *et al.* 2008). Interestingly, we also found that the degree of reduction in adult

400 offspring after yeast restriction varied across genotypes.

401

402 *No evidence for variation in host tolerance*

403 The expression of host tolerance within populations should be dependent upon disease

404 pathology and host immunopathology. Pathogen infection dynamics vary according to

405 infective dose, the route of infection, and infection outcome (Schmid-Hempel, 2011;

406 Schneider, 2011). Considering this, we used two bacteria with different infection dynamics

407 and differing degrees of virulence to explore the relationship among host fitness, diet, and

408 bacteria load to estimate fecundity tolerance.

409

410 We predicted genetic variation for tolerance phenotypes in response to infection and
411 environment with at least one bacteria species, given that a number of studies have found such
412 effects (e.g. Råberg *et al.*, 2007; Blanchet *et al.*, 2010; Graham *et al.*, 2011; Adelman *et al.*,
413 2013; Sternberg *et al.*, 2013; Parker *et al.*, 2014). For example, dietary manipulation of
414 glucose lead to a genotype-by-environment effect for *D. melanogaster* fecundity tolerance
415 that was most pronounced early during a *Providencia rettgeri* infection (Howick & Lazzaro,
416 2014). However, we found no inter-genotypic differences in fecundity tolerance in response
417 to infection with either bacteria species, and fecundity tolerance to *L. lactis* was not affected
418 by changes in host diet. The latter result confirms a previous study, where it was shown that
419 under similar experimental conditions the 1_4WS genotype does not show environmentally
420 induced variation for tolerance in response to infection with *L. lactis* (Kutzer & Armitage,
421 2016b). We also note that the considerable variation in our response variables, fecundity and
422 bacteria load, makes it potentially difficult to detect relationships between these two
423 variables.

424

425 Despite striking differences in bacterial load across genotypes, all genotypes showed similar
426 reductions in fitness with increasing *P. entomophila* load, indicating a general reproductive
427 cost to an increasing bacteria load. It is possible that assaying fecundity at an earlier infection
428 time point would have uncovered genotypic variation in host tolerance (e.g. Howick &
429 Lazzaro, 2014; Kutzer & Armitage, 2016b), or that fecundity tolerance is somehow fixed or
430 has reached saturation (e.g. Miller *et al.*, 2006). It is also possible that the co-expression of
431 host immune strategies shows temporal variation in this system, with flies surviving infection
432 by *P. entomophila* expressing resistance early on and then expressing tolerance later in the
433 infection compared to non-survivors (e.g. Lough *et al.*, 2015), which may explain the
434 considerable variation in resistance we see across these ten genotypes.

435

436 Eco-immunological studies have found support for genetic variation in host tolerance but
437 such variation is not the rule. For instance mortality tolerance declined in families of monarch
438 butterflies as *Ophryocystis elektroscirrha* inoculation dose increased, but was unaffected by
439 genotype (Lefèvre *et al.*, 2011), and aphid genotypes displayed variation in fecundity
440 tolerance but not in mortality tolerance (Parker *et al.*, 2014). A common theme that emerges
441 from these studies is the importance in the choice of the fitness or health measure and its
442 relationship to the pathogen in question, i.e. the relationship between fitness and disease
443 pathology. We suggest that fitness measures should be carefully considered in light of disease
444 pathology and infection dynamics.

445

446 *Conclusion*

447 Here we tested the effects of dietary environment and genotype on resistance and tolerance to
448 two acute phase bacterial infections. Genotype and dietary environment were strong
449 predictors of mortality from an infection with *P. entomophila* as well as predictors of
450 fecundity. We show that there is considerable genetic variability in resistance to infection,
451 while tolerance does not vary among genotypes or according to environment in the ten tested
452 genotypes. Context dependence is a recurring theme in the resistance and tolerance literature.
453 The expression of these immune strategies depends on genetics, environment and the unique
454 infection trajectories of the pathogens. Plotting detailed infection trajectories that cover an
455 entire course of infection whether at the individual or genotype level, as well as successively
456 quantifying within host damage, will help to tease apart the mechanisms governing the
457 expression of resistance and tolerance.

458

459 **Acknowledgements**

460 We would like to thank (acknowledgments redacted for blind review).

461

462 **Data Accessibility**

463 Data from the manuscript will be made accessible upon acceptance in the Dryad database.

464 **References**

- 465 Adelman, J.S., Kirkpatrick, L., Grodio, J.L. & Hawley, D.M. 2013. House finch populations
466 differ in early inflammatory signaling and pathogen tolerance at the peak of
467 *Mycoplasma gallisepticum* infection. *Am. Nat.* **181**: 674–89.
- 468 Ayres, J.S. & Schneider, D.S. 2008. A signaling protease required for melanization in
469 *Drosophila* affects resistance and tolerance of infections. *PLoS Biol.* **6**: 2764–2773.
- 470 Ayres, J.S. & Schneider, D.S. 2009. The role of anorexia in resistance and tolerance to
471 infections in *Drosophila*. *PLoS Biol.* **7**: e1000150.
- 472 Ayres, J.S. & Schneider, D.S. 2012. Tolerance of Infections. *Annu. Rev. Immunol.* **30**: 271–
473 294.
- 474 Bashir-Tanoli, S. & Tinsley, M.C. 2014. Immune response costs are associated with changes
475 in resource acquisition and not resource reallocation. *Funct. Ecol.* **28**: 1011–1019.
- 476 Bass, T.M., Grandison, R.C., Wong, R., Martinez, P., Partridge, L. & Piper, M.D.W. 2007.
477 Optimization of dietary restriction protocols in *Drosophila*. *J. Gerontol.* **62A**: 1071–
478 1081.
- 479 Best, A., White, A. & Boots, M. 2008. Maintenance of host variation in tolerance to
480 pathogens and parasites. *Proc. Natl. Acad. Sci. U. S. A.* **105**: 20786–20791.
- 481 Best, A., White, A. & Boots, M. 2010. Resistance is futile but tolerance can explain why
482 parasites do not always castrate their hosts. *Evolution.* **64**: 348–357.
- 483 Best, A., White, A. & Boots, M. 2014. The coevolutionary implications of host tolerance.
484 *Evolution.* **68**: 1426–1435.
- 485 Blanchet, S., Rey, O. & Loot, G. 2010. Evidence for host variation in parasite tolerance in a
486 wild fish population. *Evol. Ecol.* **24**: 1129–1139.
- 487 Boots, M., White, A., Best, A. & Bowers, R. 2012. The importance of who infects whom: the
488 evolution of diversity in host resistance to infectious disease. *Ecol. Lett.* **15**: 1104–11.

489 Carr, D.E., Murphy, J.F. & Eubanks, M.D. 2006. Genetic variation and covariation for
490 resistance and tolerance to Cucumber mosaic virus in *Mimulus guttatus* (Phrymaceae): a
491 test for costs and constraints. *Heredity*. **96**: 29–38.

492 Corby-Harris, V., Habel, K.E., Ali, F.G. & Promislow, D.E.L. 2007. Alternative measures of
493 response to *Pseudomonas aeruginosa* infection in *Drosophila melanogaster*. *J. Evol.*
494 *Biol.* **20**: 526–533.

495 Duffy, M.A. & Forde, S.E. 2009. Ecological feedbacks and the evolution of resistance.

496 Fineblum, W.L. & Rausher, M.D. 1995. Tradeoff between resistance and tolerance to
497 herbivore damage in a morning glory. *Nature* **377**: 517–520.

498 Flatt, T., Heyland, A. & Stearns, S.C. 2011. What mechanistic insights can or cannot
499 contribute to life history evolution. In: *Mechanisms of Life History Evolution: The*
500 *Genetics and Physiology of Life History Traits and Trade-Offs* (T. Flatt & A. Heyland,
501 eds), pp. 375–379. Oxford University Press, Oxford.

502 Garlapow, M.E., Huang, W., Yarboro, M.T., Peterson, K.R. & Mackay, T.F.C. 2015.
503 Quantitative genetics of food intake in *Drosophila melanogaster*. *PLoS One* **10**.

504 Graham, A.L., Shuker, D.M., Pollitt, L.C., Auld, S.K.J.R., Wilson, A.J. & Little, T.J. 2011.
505 Fitness consequences of immune responses: Strengthening the empirical framework for
506 ecoimmunology. *Funct. Ecol.* **25**: 5–17.

507 Harbison, S.T., McCoy, L.J. & Mackay, T.F.C. 2013. Genome-wide association study of
508 sleep in *Drosophila melanogaster*. *BMC Genomics* **14**: 281.

509 Hawley, D.M. & Altizer, S.M. 2011. Disease ecology meets ecological immunology:
510 Understanding the links between organismal immunity and infection dynamics in
511 natural populations. *Funct. Ecol.* **25**: 48–60.

512 Hotson, A.G. & Schneider, D.S. 2015. *Drosophila melanogaster* Natural Variation Affects
513 Growth Dynamics of Infecting *Listeria monocytogenes*. *G3 Genes|Genomes|Genetics* **5**:

514 2593–2600.

515 Howick, V.M. & Lazzaro, B.P. 2014. Genotype and diet shape resistance and tolerance
516 across distinct phases of bacterial infection. *BMC Evol. Biol.* **14**: 56.

517 Hurd, H. 2009. Chapter 4 Evolutionary Drivers of Parasite-Induced Changes in Insect Life-
518 History Traits. From Theory to Underlying Mechanisms. *Adv. Parasitol.* **68**: 85–110.

519 Hurd, H. 2001. Host fecundity reduction: A strategy for damage limitation? *Trends Parasitol.*
520 **17**: 363–368.

521 Kraaijeveld, A. R., Ferrari, J. & Godfray, H.C.J. 2002. Costs of resistance in insect-parasite
522 and insect-parasitoid interactions. *Parasitology* **125 Suppl**: S71–S82.

523 Kutzer, M.A.M. & Armitage, S.A.O. 2016a. Maximising fitness in the face of parasites: a
524 review of host tolerance. *Zoology* **119**: 281–289.

525 Kutzer, M.A.M. & Armitage, S.A.O. 2016b. The effect of diet and time after bacterial
526 infection on fecundity, resistance, and tolerance in *Drosophila melanogaster*. *Ecol. Evol.*
527 **6**: 4229–42.

528 Lazzaro, B.P. 2002. A Population and Quantitative Genetic Analysis of the *Drosophila*
529 *melanogaster* Antibacterial Immune Response. The Pennsylvania State University.

530 Lazzaro, B.P. & Little, T.J. 2009. Immunity in a variable world. *Philos. Trans. R. Soc. Lond.*
531 *B. Biol. Sci.* **364**: 15–26.

532 Lazzaro, B.P. & Rolff, J. 2011. Danger, microbes, and homeostasis. *Science.* **332**: 43–44.

533 Lazzaro, B.P., Sackton, T.B. & Clark, A.G. 2006. Genetic variation in *Drosophila*
534 *melanogaster* resistance to infection: A comparison across bacteria. *Genetics* **174**: 1539–
535 1554.

536 Lefèvre, T., Williams, A.J. & de Roode, J.C. 2011. Genetic variation in resistance, but not
537 tolerance, to a protozoan parasite in the monarch butterfly. *Proc. R. Soc. B* **278**: 751–
538 759.

539 Lemaitre, B., Reichhart, J.M. & Hoffmann, J.A. 1997. *Drosophila* host defense: differential
540 induction of antimicrobial peptide genes after infection by various classes of
541 microorganisms. *Proc. Natl. Acad. Sci. U. S. A.* **94**: 14614–9.

542 Leventhal, G.E., Dünner, R.P. & Barribeau, S.M. 2014. Delayed Virulence and Limited Costs
543 Promote Fecundity Compensation upon Infection. *Am. Nat.* **183**: 480–93.

544 Little, T.J., Shuker, D.M., Colegrave, N., Day, T. & Graham, A.L. 2010. The coevolution of
545 virulence: tolerance in perspective. *PLoS Pathog.* **6**: e1001006.

546 Lough, G., Kyriazakis, I., Bergman, S., Lengeling, A. & Doeschl-Wilson, A.B. 2015. Health
547 trajectories reveal the dynamic contributions of host genetic resistance and tolerance to
548 infection outcome. *Proc. R. Soc. B-Biological Sci.* **282**: 20152151.

549 Louie, A., Song, K.H., Hotson, A., Thomas Tate, A. & Schneider, D.S. 2016. How Many
550 Parameters Does It Take to Describe Disease Tolerance? *PLoS Biol.* **14**.

551 Luijckx, P., Fienberg, H., Duneau, D. & Ebert, D. 2013. A matching-allele model explains
552 host resistance to parasites. *Curr. Biol.* **23**: 1085–1088.

553 Mackay, T.F.C., Richards, S., Stone, E.A., Barbadilla, A., Ayroles, J.F., Zhu, D., *et al.* 2012.
554 The *Drosophila melanogaster* Genetic Reference Panel. *Nature* **482**: 173–178. Nature
555 Publishing Group, a division of Macmillan Publishers Limited. All Rights Reserved.

556 Magwire, M.M., Fabian, D.K., Schweyen, H., Cao, C., Longdon, B., Bayer, F., *et al.* 2012.
557 Genome-Wide Association Studies Reveal a Simple Genetic Basis of Resistance to
558 Naturally Coevolving Viruses in *Drosophila melanogaster*. *PLoS Genet* **8**: e1003057.

559 Marxer, M., Vollenweider, V. & Schmid-Hempel, P. 2016. Insect antimicrobial peptides act
560 synergistically to inhibit a trypanosome parasite. *Phil. Trans. R. Soc. B* **371**: 20150302.

561 Mauricio, R., Rausher, M.D. & Burdick, D.S. 1997. Variation in the defense strategies of
562 plants: Are resistance and tolerance mutually exclusive?

563 McKean, K.A. & Nunnery, L. 2005. Bateman’s principle and immunity: phenotypically

564 plastic reproductive strategies predict changes in immunological sex differences.
565 *Evolution* **59**: 1510–1517.

566 McKean, K.A., Yourth, C.P., Lazzaro, B.P. & Clark, A.G. 2008. The evolutionary costs of
567 immunological maintenance and deployment. *BMC Evol. Biol.* **8**: 76.

568 Miller, M.R., White, A. & Boots, M. 2007. Host life span and the evolution of resistance
569 characteristics. *Evolution.* **61**: 2–14.

570 Miller, M.R., White, A. & Boots, M. 2006. The evolution of parasites in response to
571 tolerance in their hosts: the good, the bad, and apparent commensalism. *Evolution* **60**:
572 945–956.

573 Mitchell, S.E., Rogers, E.S., Little, T.J. & Read, A.F. 2005. Host-Parasite and Genotype-By-
574 Environment Interactions: Temperature Modifies Potential for Selection By a Sterilizing
575 Pathogen. *Evolution.* **59**: 70–80.

576 Moreno-García, M., Condé, R., Bello-Bedoy, R. & Lanz-Mendoza, H. 2014. The damage
577 threshold hypothesis and the immune strategies of insects. *Infect. Genet. Evol.* **24**: 25–
578 33.

579 Moret, Y. & Schmid-Hempel, P. 2000. Survival for immunity: the price of immune system
580 activation for bumblebee workers. *Science.* **290**: 1166–1168.

581 Neyen, C., Poidevin, M., Roussel, A. & Lemaitre, B. 2012. Tissue- and ligand-specific
582 sensing of gram-negative infection in *Drosophila* by PGRP-LC isoforms and PGRP-LE.
583 *J. Immunol.* **189**: 1886–97.

584 Parker, B.J., Garcia, J.R. & Gerardo, N.M. 2014. Genetic variation in resistance and
585 fecundity tolerance in a natural host-pathogen interaction. *Evolution.* **68**: 2421–2429.

586 Råberg, L. 2014. How to Live with the Enemy: Understanding Tolerance to Parasites. *PLoS*
587 *Biol.* **12**: e1001989.

588 Råberg, L., Graham, A.L. & Read, A.F. 2009. Decomposing health: tolerance and resistance

589 to parasites in animals. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **364**: 37–49.

590 Råberg, L., Sim, D. & Read, A.F. 2007. Disentangling genetic variation for resistance and
591 tolerance to infectious diseases in animals. *Science*. **318**: 812–814.

592 R Core Team. 2016. R: A Language and Environment for Statistical Computing. R
593 Foundation for Statistical Computing, Vienna, Austria.

594 Restif, O. & Koella, J.C. 2004. Concurrent evolution of resistance and tolerance to
595 pathogens. *Am. Nat.* **164**: E90–E102.

596 Roy, B.A. & Kirchner, J.W. 2000. Evolutionary dynamics of pathogen resistance and
597 tolerance. *Evolution*. **54**: 51–63.

598 Sadd, B.M. 2011. Food-environment mediates the outcome of specific interactions between a
599 bumblebee and its trypanosome parasite. *Evolution*. **65**: 2995–3001.

600 Schmid-Hempel, P. 2011. *Evolutionary Parasitology: The Integrated Study of Infections,*
601 *Immunology, Ecology, and Genetics*, 2nd ed. Oxford University Press, Oxford.

602 Schneider, D.S. 2011. Tracing personalized health curves during infections. *PLoS Biol.* **9**.

603 Short, S.M., Wolfner, M.F. & Lazzaro, B.P. 2012. Female *Drosophila melanogaster* suffer
604 reduced defense against infection due to seminal fluid components. *J. Insect Physiol.* **58**:
605 1192–1201.

606 Siva-Jothy, M.T. & Thompson, J.J.W. 2002. Short-term nutrient deprivation affects immune
607 function. *Physiol. Entomol.* **27**: 206–212.

608 Sleiman, M.S., Osman, D., Massouras, A., Hoffmann, A.A., Lemaitre, B. & Deplancke, B.
609 2015. Genetic, molecular and physiological basis of variation in *Drosophila* gut
610 immunocompetence. *Nat Commun* **6**: 7829.

611 Stahlschmidt, Z.R., Rollinson, N., Acker, M. & Adamo, S. a. 2013. Are all eggs created
612 equal? Food availability and the fitness trade-off between reproduction and immunity.
613 *Funct. Ecol.* **27**: 800–806.

- 614 Stearns, S.C. 1992. *The Evolution of Life Histories*. Oxford University Press, London.
- 615 Sternberg, E.D., Lefèvre, T., Li, J., de Castillejo, C.L.F., Li, H., Hunter, M.D., *et al.* 2012.
- 616 Food plant derived disease tolerance and resistance in a natural butterfly-plant-parasite
- 617 interactions. *Evolution*. **66**: 3367–3376.
- 618 Sternberg, E.D., Li, H., Wang, R., Gowler, C. & de Roode, J.C. 2013. Patterns of host-
- 619 parasite adaptation in three populations of monarch butterflies infected with a naturally
- 620 occurring protozoan disease: virulence, resistance, and tolerance. *Am. Nat.* **182**: E235–
- 621 E248.
- 622 Unckless, R.L., Rottschaefer, S.M. & Lazzaro, B.P. 2015a. A Genome-Wide Association
- 623 Study for Nutritional Indices in *Drosophila*. *G3 Genes|Genomes|Genetics*
- 624 g3.114.016477.
- 625 Unckless, R.L., Rottschaefer, S.M. & Lazzaro, B.P. 2015b. The Complex Contributions of
- 626 Genetics and Nutrition to Immunity in *Drosophila melanogaster*. *PLoS Genet.* **11**:
- 627 e1005030.
- 628 Vale, P.F. & Little, T.J. 2012. Fecundity compensation and tolerance to a sterilizing pathogen
- 629 in *Daphnia*. *J. Evol. Biol.* **25**: 1888–1896.
- 630 Vale, P.F., Wilson, A.J., Best, A., Boots, M. & Little, T.J. 2011. Epidemiological,
- 631 evolutionary, and coevolutionary implications of context-dependent parasitism. *Am. Nat.*
- 632 **177**: 510–21.
- 633 Vallet-Gely, I., Lemaitre, B. & Boccard, F. 2008. Bacterial strategies to overcome insect
- 634 defences. *Nat. Rev. Microbiol.* **6**: 302–313.
- 635 Vincent, C.M. & Sharp, N.P. 2014. Sexual antagonism for resistance and tolerance to
- 636 infection in *Drosophila melanogaster*. *Proc. R. Soc. B* **281**: 20140987.
- 637 Vodovar, N., Vinals, M., Liehl, P., Basset, A., Degrouard, J., Spellman, P., *et al.* 2005.
- 638 *Drosophila* host defense after oral infection by an entomopathogenic *Pseudomonas*

- 639 species. *Proc. Natl. Acad. Sci. U. S. A.* **102**: 11414–9.
- 640 Werner, T., Liu, G., Kang, D., Ekengren, S., Steiner, H. & Hultmark, D. 2000. A family of
641 peptidoglycan recognition proteins in the fruit fly *Drosophila melanogaster*. *Proc. Natl.*
642 *Acad. Sci. U. S. A.* **97**: 13772–7.
- 643 Woolhouse, M.E.J., Webster, J.P., Domingo, E., Charlesworth, B. & Levin, B.R. (2002)
644 Biological and biomedical implications of the co-evolution of pathogens and their hosts.
645 *Nat. Genet.* **32**, 569-577.
- 646
- 647 **Online Supporting Information**
- 648 Appendix S1. Statistical analyses.
- 649 Figure S1. P-value plots of GWAS.
- 650 Table S1-S4. Parameter estimates for statistical models.

651 **Tables**

652 **Table 1.** The effect of diet and genotype on fly survival 24 hours after injection with either
 653 Ringer's, *L. lactis* or *P. entomophila*.

	<i>Tested effect</i>	<i>df</i>	<i>Dev</i>	<i>Resid.df</i>	<i>Resid.dev</i>	<i>P</i>
Model 1a: Ringer's	Diet	1	0.2	138	34.85	0.65
	Genotype	9	6.9	129	27.96	0.65
	Genotype x Diet	9	6.85	120	21.12	0.65
Model 1b: <i>L. lactis</i>	Diet	1	0.69	138	39.29	0.41
	Genotype	9	9.02	129	30.27	0.44
	Genotype x Diet	9	7.81	120	22.45	0.55
Model 1c: <i>P. entomophila</i>	Diet	1	202.99	138	214.49	<0.0001
	Genotype	9	69.05	129	145.45	<0.0001
	Genotype x Diet	9	22.94	120	122.51	0.006

654

655

656 **Table 2.** The effect of diet and genotype on bacteria load 24 hours post infection. The dashes
 657 (-) indicate that diet and diet x genotype could not be tested.

<i>Tested effect</i>	Model 2a: <i>L. lactis</i>				Model 2b: <i>P. entomophila</i>			
	<i>numDF</i>	<i>denDF</i>	<i>F</i>	<i>P</i>	<i>numDF</i>	<i>denDF</i>	<i>F</i>	<i>P</i>
Diet	1	369	0.58	0.446	-	-	-	-
Genotype	9	369	4.57	<0.0001	9	150	19.25	<0.0001
Diet x Genotype	9	369	0.50	0.874	-	-	-	-

658

659 **Table 3.** The effect of diet, genotype and infection status on post-infection fecundity,
660 measured as adult offspring number. Pre-infection fecundity was included as a covariate.
661 High mortality in the *P. entomophila* group on reduced yeast (RY) precluded testing one fully
662 factorial fecundity model. Therefore Model 3a tests fecundity of the three infection groups,
663 i.e., injection of Ringer's, *L. lactis* or *P. entomophila*, where flies had only *ad libitum* (AY)
664 access to yeast. The dashes (-) indicate that diet and interactions with diet could not be tested.
665 Model 3b tests the fecundity of only Ringer's and *L. lactis* injected flies, on either AY or RY.

<i>Tested effect</i>	Model 3a: AY fecundity			Model 3b: AY vs RY fecundity		
	<i>df</i>	<i>F</i>	<i>P</i>	<i>df</i>	<i>F</i>	<i>P</i>
Diet	-	-	-	1	63	<0.0001
Genotype	9	9.22	<0.0001	9	12.006	<0.0001
Infection status	2	0.13	0.88	1	1.569	0.211
Diet x Genotype	-	-	-	9	9.46	<0.0001
Diet x Infection status	-	-	-	1	1.441	0.23
Genotype x Infection status	18	0.69	0.82	9	1.931	0.045
Pre-infection fecundity	1	233.23	<0.0001	1	46.51	<0.0001

666

667

668 **Table 4.** The effects of bacteria load (CFU), diet and genotype on post-infection fecundity
669 (measured as % adult offspring produced relative to the uninfected Ringer’s control) after
670 infection with *L. lactis* or *P. entomophila*. Significant interactions between CFU and either or
671 both of the other factors would indicate variation in fecundity tolerance. The dashes (-)
672 indicate that diet and interactions with diet could not be tested.

<i>Tested effect</i>	Model 4a: <i>L. lactis</i>				Model 4b: <i>P. entomophila</i>			
	<i>numDF</i>	<i>denDF</i>	<i>F</i>	<i>P</i>	<i>numDF</i>	<i>denDF</i>	<i>F</i>	<i>P</i>
CFU	1	349	2.77	0.097	1	140	11.1	0.001
Diet	1	349	19.62	<0.0001	-	-	-	-
Genotype	9	349	4.83	<0.0001	9	140	1.63	0.114
CFU x Diet	1	349	0.01	0.907	-	-	-	-
CFU x Genotype	9	349	0.96	0.475	9	140	0.57	0.821
Diet x Genotype	9	349	1.39	0.191	-	-	-	-
CFU x Diet x Genotype	9	349	1.24	0.271	-	-	-	-

673

674 **Figure legends**

675 **Figure 1. *D. melanogaster* survival 24 hours post injection.** Females were kept on a diet of
676 *ad libitum* yeast (AY) or reduced yeast (RY) and injected with (A) Ringer's solution as an
677 injection control, or one of two bacteria, (B) *L. lactis*, or (C) *P. entomophila*. Each line
678 represents the reaction norm of one of ten genotypes. Some reactions norms overlap. Each
679 dot represents the proportion of 21 flies that survived.

680

681 **Figure 2. Bacteria load for each genotype 24 hours post infection.** Females were injected
682 with either (A) *L. lactis* or (B) *P. entomophila*. Bacteria load was quantified as the number of
683 colony forming units (CFUs) counted on agar plates containing individual whole fly
684 homogenates. There was no effect of diet on *L. lactis* loads, so AY and RY individuals are
685 combined in (A). Genotypes are arranged in ascending order and diamonds represent medians
686 calculated from between 37 and 42 female flies in (A) and between 15 and 21 female flies in
687 (B). To visualise the data on a log scale, we added 1 to all CFU counts. The dotted lines
688 indicate the approximate infection doses.

689

690 **Figure 3. Fecundity measured as the number of adult offspring produced in the 24 hours**
691 **post injection.** Female flies were subjected to one of the following injection treatments and
692 diet combinations: (A) Ringer's injected AY medium, (B) *L. lactis* injected AY medium, (C)
693 *P. entomophila* injected AY medium, (D) Ringer's injected RY medium, (E) *L. lactis* injected
694 RY medium. Diamonds represent medians calculated from between 15 and 21 female flies per
695 treatment group. Genotypes are arranged in descending order of offspring numbers. Data
696 points with darker shades of grey indicate where the values of more than one individual
697 overlap.

698

699 **Figure 4. Fecundity tolerance after infection with *L. lactis* and *P. entomophila*.** (A)
700 Fecundity tolerance after *L. lactis* infection is unaffected by diet or genotype; (B) Fitness
701 decreases in response to increasing *P. entomophila* load regardless of genotype. The natural
702 log of bacterial load (CFU) is plotted against the percent change in adult offspring number.
703 Each data point represents the bacteria load and fitness of one female fly. Reaction norms are
704 plotted for AY (solid lines) and RY (dashed lines; *L. lactis* only) for each genotype 24 hours
705 post injection. Data points with darker shades of grey indicate where the values of more than
706 one individual overlap. Mortality was too high in *P. entomophila* infected individuals on the
707 RY medium to include them in the analysis.
708