

OPEN

Heading choices of flying *Drosophila* under changing angles of polarized light

Thomas F. Mathejczyk & Mathias F. Wernet*

Many navigating insects include the celestial polarization pattern as an additional visual cue to orient their travels. Spontaneous orientation responses of both walking and flying fruit flies (*Drosophila melanogaster*) to linearly polarized light have previously been demonstrated. Using newly designed modular flight arenas consisting entirely of off-the-shelf parts and 3D-printed components we present individual flying flies with a slow and continuous rotational change in the incident angle of linear polarization. Under such open-loop conditions, single flies choose arbitrary headings with respect to the angle of polarized light and show a clear tendency to maintain those chosen headings for several minutes, thereby adjusting their course to the slow rotation of the incident stimulus. Importantly, flies show the tendency to maintain a chosen heading even when two individual test periods under a linearly polarized stimulus are interrupted by an epoch of unpolarized light lasting several minutes. Finally, we show that these behavioral responses are wavelength-specific, existing under polarized UV stimulus while being absent under polarized green light. Taken together, these findings provide further evidence supporting *Drosophila*'s abilities to use celestial cues for visually guided navigation and course correction.

Like many other animals, insects have developed the ability to efficiently navigate the most complex environments. Over several decades, evidence has accumulated showing that different insect species combine a multitude of visual stimuli in order to take fast and reliable navigational decisions (reviewed in¹). Amongst these cues, the celestial polarization pattern serves as a robust visual stimulus informing the heading choices of many navigating insects^{1–3}. Since Karl von Frisch first described the ability of honeybees to orient their waggle dances using merely a small patch of sky that did not include the sun as a landmark, many insects have also been shown to integrate the directional information provided by the skylight polarization pattern into their repertoire of visual cues⁴. Importantly, this ability is not restricted to central-place foragers like bees or desert ants that rely on visual cues to find their way back to their hive or nest^{5,6}. For example, both diurnal and nocturnal ball-rolling dung beetles have been shown to use the celestial polarization pattern to set a straight path away from the food source where both predators and competitors may aggregate^{7,8}. In this case, dung beetles show the tendency to maintain the same heading over repeated trials⁶. The tendency of other walking insects to set and maintain heading choices under a linearly polarized stimulus remains less well characterized. Although spontaneous behavioral responses to rotating polarization filters (polarotaxis) were demonstrated for crickets and flies when walking on air-suspended balls under laboratory settings^{9–11}, clear characterizations of angular heading choices are missing for these experiments. Similarly, behavioral data for flying insects (other than honeybees), especially when using virtual flight arenas remains relatively scarce^{1,2}. Oriented flights of suspended monarch butterflies under a polarized stimulus have been demonstrated, yet its ethological significance remains somewhat controversial, due to conflicting reports^{12–14}. Probably the most valuable recent progress comes from the fly *Drosophila melanogaster*: spontaneous responses of flying *Drosophila* to linearly polarized light using virtual flight arenas have been demonstrated, both under the natural sky, as well as using an artificial stimulus generated in the laboratory using commercially available polarization filters^{15–18} (reviewed in¹⁹). Most importantly, flies were shown to choose arbitrary angular headings with respect to the orientation of the e-vector of the polarized stimulus and showed the tendency to maintain this navigational decision over several minutes, even when the stimulus presentation was perturbed for several minutes¹⁸. Nevertheless, fairly little is known about the navigational capabilities of free-living fruit flies (reviewed in²⁰). Catch-and-release experiments from a fixed point in the desert suggested that *Drosophila (melanogaster)*

Freie Universität Berlin, Fachbereich Biologie, Chemie und Pharmazie, Institut für Biologie – Neurobiologie, Königin-Luise Strasse 1-3, 14195, Berlin, Germany. *email: mathias.wernet@fu-berlin.de

and *pseudoobscura*) disperse into all directions equally and are able to keep straight headings over extended periods of time, while flying in environments which provide few visual landmarks^{21,22}. For a better quantitative understanding of the mechanisms underlying such processes, skylight navigation experiments using virtual flight arenas therefore serve as an attractive platform for the study of the navigation skills of wild type insects, thereby providing the platform for testing transgenic specimens harboring well-defined circuit perturbations^{23,24}.

The retinal basis of celestial polarization vision across insects is well understood: in virtually all cases, specialized ommatidia located in the ‘dorsal rim area’ (DRA) of the adult eye are morphologically and molecularly specialized for this task²⁵. In flies, DRA inner photoreceptors R7 and R8 express the same UV Rhodopsin Rh3 which is localized within untwisted light-sensing rhabdomeres, resulting in high polarization sensitivity^{23,26–33}. Since rhabdomeres of DRA R7 and R8 of a given ommatidium are oriented orthogonally to each other, these two cells form an opponent analyzer pair^{23,31}. Like in other insects, analyzer directions of DRA ommatidia change gradually along the DRA, forming a ‘fan-shaped array’ of polarization detectors³⁴. Due to the monochromatic Rhodopsin expression in DRA R7 and R8, navigational decisions of *Drosophila* in response to linearly polarized light should be limited to the UV range of the spectrum^{23,25}, whereas light of longer wavelengths should not elicit orientation responses to this stimulus. Interestingly, different insect species express blue-sensitive Rhodopsins in their polarization-sensitive DRA photoreceptors (crickets, locusts)^{35,36}, and in some cases green-sensitive Rhodopsins were reported (cock chafers)³⁷. Although these different Rhodopsin choices seem to reflect adaptations to different ecological niches, the exact ethological reason for these differences remain incompletely understood^{38,39}. Increasing evidence also points towards many insects (including flies) being capable to detect linearly polarized light through a DRA-independent channel (reviewed in⁴⁰). Experiments from *Drosophila* have shown that these polarotactic behaviors are not UV-specific, since behavioral responses can be elicited using polarized green light presented to the ventral half of the retina^{16,23}. Although incompletely understood, these behaviors could be indicative of a so-far poorly understood system in which retinal detectors are used to detect linearly polarized reflections. Such reflections could be used by insects to seek out or avoid water surfaces, evaluate oviposition sites, or even detect prey^{2,40}.

We use virtual flight arenas to test heading decisions of individual flies flying under ‘open-loop’ conditions under a slowly rotating polarization filter. In agreement with previous studies, we find that flies initially choose a heading with respect to the orientation of the incident polarized light that varies between individuals and shows no preference for certain headings over the entire population tested (arbitrary headings). In this configuration, the rotation of the polarization filter therefore forces the fly to constantly adjust its heading in order to hold its original heading decision constant. By quantifying the fly’s ability to adjust its heading relative to the changing e-vector over time we show that the behavioral performance varies greatly within a population, yet a similar behavior is never observed under unpolarized UV light, or linearly polarized green light. Importantly, flies show the tendency to maintain this heading over several minutes: we show that flies that perform well in following the e-vector within a 5-minute experiment show a high tendency to choose a similar heading in a second experiment, even when interrupted by a 5-minute interval of unpolarized light. These experiments underscore the usefulness of the experimental setups presented here and serve as an ‘open source’ platform for the development of new assays optimized for different visual behaviors, in flies as well as other species of flying insects.

Results

The aim of this study was a quantitative analysis of heading choices recorded from single flies flying under a slowly rotating polarization filter. We reasoned that flies that commit to a specific heading angle with respect to the incident angle of polarization would show a tendency to hold this angle constant and therefore correct for the slow rotational drift of the stimulus.

Virtual flight arenas for testing the heading choices of tethered flies under a rotating polarized stimulus.

In order to quantify the heading choices of flies (glued to a metal pin) under a constantly rotating e-vector of linearly polarized light presented dorsally, we used custom built virtual flight arenas assembled from 3D printed and off-the-shelf hardware (Fig. 1a, Supplemental Figs S1 and S2). Similar to what was previously described⁴¹, the magnetic field created by two magnets kept the steel pins (and therefore the flies) vertical and in the center of the dorsally presented stimulus (see below), while allowing individual flies to rotate around their yaw axis, thereby enabling them to freely choose their headings (for 3D-printing and detailed assembly instructions, see <https://doi.org/10.1101/527945> and www.flygen.org/skylight-navigation). On top of the setup, switchable LED light sources (UV or green) were attached to two vertical beams, equipped with a matching set of highly collimated optics (Mightex Inc, see methods), in order to minimize off axis illumination of the polarization filter. The polarization state of the dorsally presented stimulus could be altered by shining light through a switchable filter ‘sandwich’ consisting of diffuser paper and a linear polarizer with either the polarizer or the diffuser facing the fly, as previously demonstrated^{10,11,23}. This allowed for switching between two experimental conditions in which the degree of polarization was either ~0% (unpolarized) and almost 100% linearly polarized while keeping light intensity between trials constant (Fig. 1b, Supplemental Fig. S2A,B). The filter sandwich was placed inside a removable filter cassette housed inside a 3D-printed filter wheel that could be rotated using an Arduino-controlled servo motor (Dynamixel MX-28T, see methods) via a 3D printed gear system, all of which was also attached to the vertical beams (Fig. 1a). The upper magnet holding the tethered fly in place was magnetically attached (via a second magnet) to a UV fused silica window mounted in the light path (Fig. 1a, Supplemental Figs S1 and S2). Similar to what had previously been reported⁴¹, a sapphire bearing for minimizing friction and to keep the steel pins in place, was placed on a small matte white disc (to avoid intensity artefacts) which was glued to the bottom side of the upper magnet. Due to this placement and size of the upper magnet (for holding the fly in place), the stimulus extended over a 17° wide concentric ring in the flies’ dorsal field of view (Fig. 1c, Supplemental Fig. S2C). Keeping this angle as small as possible was crucial in order to minimize transmission artifacts (i.e. intensity fluctuations)




Figure 1. Modular assay for studying skylight navigation in individual flying flies. **(a)** Schematic drawing of a fully assembled virtual flight arena for the quantitative study of skylight navigation in flying *Drosophila*. Commercially available parts include: (1) a 30 cm \times 30 cm double density optical breadboard (Thorlabs); (2) compatible metal beams (Thorlabs); (3) swappable, magnetically attached high-power LED light sources with collimated optics (Mightex); (4) a robotics-grade servo motor for rotating the polarization filter (Dynamixel MX-28T, Robotis), positioned behind the light path (when viewed from the front); (5) an infrared camera shielded from the fly's view, filming its body axis (Firefly MV, Point Grey). Custom-designed 3D-printed parts (see supplemental materials for printing instructions) include: (6) attachment to the vertical bars; (7) filter holder including gear system (see <https://doi.org/10.1101/527945> and www.flygen.org/skylight-navigation for details); (8) horizontal platform holding the UV fused silica plate and top magnet for attaching the magnetotethered fly; (9) infrared illumination (enclosed LED's) and bottom plate with ring magnet, mounted onto the infrared camera; (10) backlit cylinder for reducing linearly polarized reflection artifacts, which can be lifted all the way to the horizontal platform, when flies are tethered. **(b)** Two polarization filter / diffuser orientations can be chosen within the filter holder: Unpolarized (diffused; left), or polarized (right). See Fig. 3b for polarimetric characterization. **(c)** Drawing of the fly's field of view inside the apparatus. $\alpha = 17^\circ$. **(d)** Camera image for the extraction of the fly's body axis (shown in blue).

that can arise from the polarization filter, especially at larger viewing angles¹⁵. Additionally, the suspended fly was placed at the center of a matte, backlit cylinder that was 3D-printed from white material, in order to minimize polarized reflections from the walls surrounding the fly, as well as to shield it from additional visual cues (#10, in Fig. 1a, and Supplemental Fig. S2C). The flies were illuminated with near-infrared LED's and filmed from below with 60hz (Firefly MV, Point Grey) through a pinhole placed on top of the camera in order to avoid reflections off its lens. Using the camera footage, the flies' body axis angles were extracted over time using custom made image processing code so that their heading choices in response to the rotating e-vector could be quantified (Fig. 1d).

Flying *Drosophila* follow a slowly rotating e-vector at an arbitrary angular distance. In order to achieve a more precise quantitative measure for the quality of polarotactic responses over more extended time intervals, as opposed to rapid changes in e-vector orientation¹⁷ (see <https://doi.org/10.1101/527945>), we introduced a new stimulus: flies flying within the virtual flight arena were presented a linearly polarized stimulus rotating slowly with constant angular velocity ($\sim 6^\circ/s$). Given that our test subjects were flying, this speed was chosen $\sim 3x$ faster than what was previously published for walking crickets and houseflies⁹⁻¹¹. In these experiments the 5-minute recording session per trial was split up into 30×10 s windows. For each of these 30 windows the mean angular velocity of each fly was then calculated. If the difference between this angular velocity and the fly's angular velocity was smaller than $3^\circ/s$, the particular time window was categorized as polarotactic behavior (i.e. above threshold; areas shaded blue). For each of these periods the fly's chosen heading was then calculated as the mean angular difference between the fly's body axis and the incident e-vector (blue bar plots: above threshold; grey bar plots: below threshold). A representative fly (Fig. 2a) flying under a constantly rotating e-vector adjusted its heading in about one third of the recorded 10 s time windows (10/30). The number and length of observed interruptions without polarotaxis varied from fly to fly, resulting in a wide spread of behavioral performance quality (as defined by number of polarotactic 10 sec time windows) when integrating over the




Figure 2. Flying *Drosophila* follow a slowly rotating e-vector at an arbitrary angle. **(a)** Top: Flight heading (blue line) of a single fly orienting to a slowly rotating linearly polarized UV stimulus (PolUV1; orange line). 10 s intervals with above threshold polarotactic behavior (see methods) are shaded blue. Bottom: plot of above threshold intervals and circular plot of heading chosen by the animal (blue: above threshold; grey: below threshold). **(b)** Same analysis as above, using an unpolarized UV stimulus (UVunpol); same fly. **(c)** Same analysis as above, using a polarized green stimulus (PolGreen); same fly. **(d)** Same analysis as above, using a re-polarized UV stimulus (PolUV2); same fly.

entire 5 minutes tested. Importantly, the calculated preferred heading of a given fly falls within a narrow angular range when compared across polarotactic periods (Fly in Fig. 2a: mean heading 63.2°, SD = 12°), despite interspersed periods of non-polarotactic behavior (during which the ‘preferred orientation’ varies greatly). This indicates that flies attempt to keep a preferred heading with respect to the celestial e-vector pattern over short periods of time. As expected, virtually no polarotactic periods were detected when the same fly flew under unpolarized UV light (UVunpol), but otherwise unchanged conditions (Fig. 2b). Similarly, when the fly was flying under linearly polarized green light (PolGreen), virtually no polarotaxis was detected (Fig. 2c). However, upon presenting the fly re-polarized UV light again (PolUV2), polarotactic time periods were restored, in some cases even more pronounced than in the first UV trial (Fig. 2d). Pooled data from all tested flies flying under a constantly




Figure 3. Variation of heading choices across individuals tested. **(a)** Summary plot of % time spent following the slowly rotating e-vector, comparing the four conditions from above. Bonferroni-corrected two-tailed Mann-Whitney U test, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. N (left to right) = 66, 22, 38, 43. **(b)** Direct comparison of male versus female polarotaxis from PolUV1 and PolUV2 experiments reveals no significant difference. Two-tailed Mann-Whitney U test. N (left to right) = f (30), m (36), f (19), m (24).

rotating filter under different lighting conditions (PolUV1, UVunpol, PolGreen, PolUV2) reveals that polarotaxis occurs exclusively when using a linearly polarized UV stimulus (Fig. 3a). Flies flying under a slowly rotating linearly polarized UV stimulus spent significantly more time following the e-vector, compared to flying under unpolarized UV or polarized green light, respectively. Interestingly, flies that underwent a second flight under re-polarized UV light spent even more time following the e-vector. Finally, analysis of behavioral responses of female and male flies reveal no significant differences between genders in the first as well as in a second trial under polarized UV light (PolUV1 and PolUV2).

Chosen headings are arbitrary, while behavioral performance varies between flies. By comparing the behavior across many individuals (N = 66) we investigated the spread of preferred headings when single




Figure 4. Chosen headings are arbitrary and different between flies. Top: Flight heading (blue line) of four single flies orienting to a slowly rotating linearly polarized UV stimulus (PolUV1; orange line). 10 s intervals with above threshold polarotactic behavior (see methods) are shaded blue. Bottom: Plots of above threshold intervals over time and circular histograms of headings chosen by the animals (blue: above threshold; grey: below threshold).

flies were flying under a slowly rotating polarization filter. The goal was to investigate whether, in this particular kind of virtual flight arena, certain headings are naturally preferred or avoided, or whether the choice of preferred heading is arbitrary and therefore different between individual flies. The strategy is exemplified by the direct comparison of four representative traces of individual flies in response to a slowly rotating polarized UV stimulus (Fig. 4). Quality of behavioral performance (polarotaxis intervals) and preferred heading were quantified as described above. It appears that in these trials each fly chooses a different preferred heading with respect to the incident angle of polarization (Fig. 4, circular plots). Taken together, the preferred heading angles of all tested flies during their first linearly polarized UV trial (PolUV1) were distributed over the whole angular range (Fig. 5a,b). Importantly, although the quality of behavioral performance (number of polarotaxis intervals, i.e. time spent following the rotating e-vector) varied greatly between individuals, it did not correlate with the angular heading choice of the animals (Fig. 5c). Hence, unlike previous studies on walking fly populations²³, no tendency for a fixed preferred angular heading choice was found across flying individuals.




Figure 5. Variability and robustness of consecutive heading choices. **(a)** Summary of heading angles (angular difference to the polarization filter) chosen by 66 individual flies, under the PolUV1 stimulus. **(b)** Same data plotted on circular coordinates reveals a wide distribution. **(c)** Same data as in B where behavioral performance (%Pol = percentage of time following the e-vector) is represented in false color.




Figure 6. Arbitrarily chosen headings are maintained between trials. **(a)** Plot depicting the angular difference between headings chosen in two consecutive trials interrupted by a period of unpolarized light (PolUV1 vs PolUV2) as a function of the quality of polarotaxis during the PolUV1 period (% of time spent following e-vector). **(b)** The change of polarotaxis probability over the 5 min experimental time window, plotted for all four experimental conditions (PolUV1, UVunpol, PolGreen, and PolUV2).

Arbitrarily chosen headings are maintained between trials. Finally, we tested whether the amount of time that the flies spent following the e-vector (number of polarotaxis intervals, i.e. quality of behavioral performance) within the first linearly polarized UV trial (PolUV1) correlates with the tendency of the flies to choose a similar preferred heading in a second consecutive trial (PolUV2) that was separated from the first by an interruption (5 min of UVunpol). We found that the better the flies' performance within the first trial (more time spent following the e-vector in PolUV1), the higher the likelihood of them choosing a similar heading in the second trial (Fig. 6a). During both PolUV1 and PolUV2 intervals, the probability of tested flies for following the rotating e-vector increased during the 5 min trial (Fig. 6b). In contrast, the overall lower polarotactic values obtained in control conditions (UVunpol and PolGreen) showed no similar increase over time.

Discussion

Navigating insects rely on the detection and integration of a wide variety of visual cues, like celestial bodies (sun, moon, milky way), intensity gradients, and chromatic gradients¹. In addition, the celestial pattern of linearly polarized light serves as an attractive orientation cue that many insects use^{2,42,43}. Spontaneous behavioral responses of both walking and flying *Drosophila* to linearly polarized light ('polarotaxis') have been demonstrated in the past, using both population assays, as well as single fly assays^{16–18,23,34,44,45}. In all these experiments, much care was given to the control and avoidance of intensity artifacts that can result in behavioral decisions that are in fact independent of the linearly polarized component of the stimulus (reviewed in¹⁵). The virtual flight arenas used here have been designed with the dual goal of providing relatively cheap, robust setups that can easily be assembled, while at the same time minimizing intensity/reflection artifacts. The codes, templates and building instructions for the virtual flight arenas are freely available for download to anyone (for a detailed description,

see <https://doi.org/10.1101/527945> and www.flygen.org/skylight-navigation). Due to their modular design, they can be modified for studying behavioral responses to moving stimuli^{46–49}, shapes⁵⁰, colors⁵¹, or celestial bodies²⁴.

Using these new virtual flight arenas, we show that individual flies choose arbitrary headings under a linearly polarized stimulus, and when summed over all individuals tested, all chosen headings appear to spread randomly. This finding is in good agreement with recently published studies, although these were using a rather different kind of stimulation system¹⁸. In further agreement with these past studies, we also find that flies show a clear tendency to keep their chosen heading over several minutes, which indicates that any given fly attempts to maintain its chosen heading. Given the considerable gap in knowledge about the ethology of *Drosophila*, these data provide further support for a potential role of polarization vision in guiding long-range navigation behaviors that have been reported for flies^{20–22}. In contrast, we show that single flies flying under a linearly polarized green stimulus displayed no comparable polarotaxis, which was to be expected due to the fact that orientation responses to polarized stimuli presented dorsally were shown to be mediated exclusively by the DRA²³, whose polarization-sensitive R7 and R8 photoreceptors in the fly both express the UV-sensitive Rhodopsin Rh3^{25,28–30}. Nevertheless, behavioral responses to linearly polarized green light stimuli have also been reported in the past^{16,22}, but only when presented ventrally (far beyond the retinal regions that were illuminated in this study) and the retinal detectors responsible are not well understood^{23,40}.

Our experiments show that well-performing flies show a clear tendency to maintain their chosen heading, even when interrupted by a period of unpolarized stimulation. These data again provide independent support for previous studies¹⁸ and reinforce the idea that a generalist fly like *Drosophila melanogaster* is indeed capable of using skylight polarization for maintaining a chosen course over longer times, which is crucial for achieving more complex navigational tasks^{20–22}. Like previous studies, we aimed at quantifying the quality of behavioral responses, since we expected that behavioral performance of individual flies to be greatly variable due to the strong influence of environmental conditions as well as internal states of the animal(s). For this study, we introduced a simple new stimulus, where flies are suspended under a slowly rotating polarization filter (~6°/s) under 'open loop' conditions. Quantifying the quality of a behavioral response by chopping any given 5-minute experiment into 30 × 10 sec polarotactic periods serves as an attractive new strategy for producing statistically significant data in a reasonable amount of time. Using this method, our experiments indeed revealed that behavioral performance is variable across all individuals tested. Even after tight control of food quality, rearing conditions, temperature, and humidity, the flies' cooperation in these experiments remains unpredictable. Importantly, the distribution of polarotactic 10 s intervals across any given 5 min experiment did not reveal any obvious pattern, except the trend that probability of polarotaxis tended to improve over time (as summarized in Fig. 6). How much this variability could depend on the fly's motivational state or navigational decision making remains to be investigated. Interestingly, even within a given 5 min recording, flies do not necessarily follow the rotating e-vector permanently, but may transition into and out of polarotactic periods (see Fig. 2a). This demonstrates the usefulness of this experimental setup for further studies on the dynamics and modulation of polarotactic behavior (and potentially underlying decision making processes), for instance in response to different internal states. Finally, our experiments reveal no significant differences in the behavioral performance of male versus female flies. This was to be expected since catch-and-release experiments did not reveal any sex differences in *Drosophila's* tendency to disperse^{21,22}. Furthermore, the size and structure of DRA ommatidia does not differ in a systematic way, between sexes. Although male-specific, Fruitless-expressing neurons have been characterized in the *Drosophila* brain, none of them appear to be clustered in the dorsal periphery of the visual system^{52,53}.

Many insect species use the celestial polarization pattern in conjunction with other visual stimuli like celestial bodies, intensity gradients, chromatic gradients, and landmarks¹. The hierarchy in which these stimuli are combined might differ between species as well as depending on context. One recent study reported that single *Drosophila* flying in a virtual flight arena are also able to use an artificially generated celestial body (the sun) as a reference to choose a heading (menotaxis)²⁴, a behavior that requires 'compass neurons' in a central brain region known as the central complex⁵⁴. This function is therefore in good agreement with physiological properties described for these neurons in locusts⁵⁵. Classic data from larger insects⁵, as well as more recent studies from *Drosophila*⁵⁶ are beginning to elucidate the neural circuitry of the 'compass pathway', along which menotactic and polarotactic information are being integrated by the insect brain, resulting in time-compensated compass information in the central complex^{5,57}. Despite important similarities, it remains unclear whether flies use their anatomical compass pathway for performing exactly the same computations for skylight navigation^{24,56,58–62}. The experiments presented here therefore serve as an important new platform for the efficient combination of *Drosophila* molecular genetic tools for the cell-type specific manipulation of neuronal function with quantitative behavior assays for testing skylight navigation.

Methods

Fly rearing. Wild type Oregon R flies (isogenized) were reared at 25 °C and 60% relative humidity on standard cornmeal agarose food under a 12h-light/12h-dark cycle. Care was taken to keep population densities low within fly vials by flipping flies on a daily basis.

Fly preparation. Experiments were performed at 25 °C and 50% relative humidity during the flies' evening activity peaks up until one hour after the light period within the respective rearing incubators would have ended. Flies were glued to 10 mm long, 100 µm diameter steel pins (ENTO SPHINX s.r.o., Czech Republic), so that when positioned vertically, they held the flies at a natural flying angle (about 60° from horizontal) and were allowed to recover for at least 20 minutes from the gluing procedure before being tested. To prevent the flies from flying during the recovery phase, small pieces of Kimwipes were transferred to their tarsi. Initial flight behavior was triggered by inducing a little air puff towards the fly from below. This was also quickly done when flies stopped

flying during the experiments, but not more than 3 times per experiment without excluding those flies from data analysis.

Flight simulator setup. *Virtual flight arenas.* Detailed building instructions of the virtual flight arenas used in this study including 3D printing and step-by-step assembly instructions and the codes necessary for their operation are freely available (<https://doi.org/10.1101/527945> and www.flygen.org/skylight-navigation). Furthermore, the experimental setup, including stimulus properties (normalized intensities of the LED light sources and polarimetric characterization of the stimuli used) are described in greater detail in Supplemental Figures S1 and S2.

Stimulus delivery. Above the fly a switchable cassette holding a 50 mm × 50 mm sheet linear polarizer (OUV5050, Knight Optical, UK) and 13 layers of thin, non-fluorescent diffuser paper (80 g/sqm, Max Bringman KG) was inserted into a motorized rotatable cassette holder. Light from a collimated UV or green LED (365 nm: LCS-0365-13-B, 530 nm: LCS-0530-15-B, Mightex) coming from above and passing through the filter cassette was polarized (pol filter at bottom) or depolarized (diffuser paper at bottom) depending on the orientation of the filter cassette and presented to the dorsal part of the flies' eyes. Using a spectrometer (Flame, Ocean Optics) the intensity of the two LEDs was set approximately isoquantally at 2×10^{12} photons/s/cm². By rotating the cassette holder, it was possible to precisely control the angle of the polarized stimulus' e-vector. The recording protocol consisted of rotating the E-vector with constant angular velocity (5.97 deg/s) for 5 minutes while synchronously recording the flies' behavior.

Extraction of flight heading. The heading of each fly in each acquired video frame was extracted using a custom-written macro script for the open-source software Fiji⁶³. In short, it binarizes each video and fits an ellipse around the fly's body to extract its heading in a range from 0° to 180°, in accordance with the directional ambiguity of the presented e-vector. The Fiji tracking results were analyzed using MatLab. Circular statistics were used. In short, a fly's heading changes were quantified as polarotactic behavior if the mean difference between the fly's angular velocity and the e-vector's angular velocity was smaller than 3°/s for a given 10 s time window. By calculating the fly's heading relative to the e-vector during such polarotactic episodes it was possible to calculate a measure of the 'preferred e-vector' for each fly. The custom Fiji- and Matlab-scripts used in this study, as well as setup building instructions and documentation is freely available under www.flygen.org/skylight-navigation.

Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Received: 30 May 2019; Accepted: 30 October 2019;

Published online: 14 November 2019

References

1. Heinze, S. Unraveling the neural basis of insect navigation. *Curr Opin Insect Sci* **24**, 58–67, <https://doi.org/10.1016/j.cois.2017.09.001> (2017).
2. Mathejczyk, T. F. & Wernet, M. F. Sensing Polarized Light in Insects. Oxford Encyclopedia of Neuroscience. Online publication date: Sep 2017, <https://doi.org/10.1093/acrefore/9780190264086.013.10> (2017).
3. Nilsson, D. E. & Warrant, E. J. Visual discrimination: Seeing the third quality of light. *Curr Biol* **9**, R535–537 (1999).
4. von Frisch, K. Die Polarisation des Himmelslichts als orientierender Faktor bei den Tänzern der Bienen. *Experientia* **5**, 142–148 (1949).
5. Homberg, U. Sky Compass Orientation in Desert Locusts-Evidence from Field and Laboratory Studies. *Front Behav Neurosci* **9**, 346, <https://doi.org/10.3389/fnbeh.2015.00346> (2015).
6. El Jundi, B., Baird, E., Byrne, M. J. & Dacke, M. The brain behind straight-line orientation in dung beetles. *J Exp Biol* **222**, <https://doi.org/10.1242/jeb.192450> (2019).
7. Foster, J. J. *et al.* Stellar performance: mechanisms underlying Milky Way orientation in dung beetles. *Philos Trans R Soc Lond B Biol Sci* **372**, <https://doi.org/10.1098/rstb.2016.0079> (2017).
8. El Jundi, B. *et al.* A Snapshot-Based Mechanism for Celestial Orientation. *Curr Biol* **26**, 1456–1462, <https://doi.org/10.1016/j.cub.2016.03.030> (2016).
9. Vonphilipsborn, A. & Labhart, T. A Behavioral-Study of Polarization Vision in the Fly, *Musca-Domestica*. *Journal of Comparative Physiology a-Sensory Neural and Behavioral Physiology* **167**, 737–743 (1990).
10. Brunner, D. & Labhart, T. Behavioral Evidence for Polarization Vision in Crickets. *Physiol Entomol* **12**, 1–10, <https://doi.org/10.1111/j.1365-3032.1987.tb00718.x> (1987).
11. Henze, M. J. & Labhart, T. Haze, clouds and limited sky visibility: polarotactic orientation of crickets under difficult stimulus conditions. *J Exp Biol* **210**, 3266–3276, <https://doi.org/10.1242/jeb.007831> (2007).
12. Stalleicken, J. *et al.* Do monarch butterflies use polarized skylight for migratory orientation? *J Exp Biol* **208**, 2399–2408, <https://doi.org/10.1242/jeb.01613> (2005).
13. Sauman, I. *et al.* Connecting the navigational clock to sun compass input in monarch butterfly brain. *Neuron* **46**, 457–467, <https://doi.org/10.1016/j.neuron.2005.03.014> (2005).
14. Reppert, S. M., Zhu, H. & White, R. H. Polarized light helps monarch butterflies navigate. *Curr Biol* **14**, 155–158 (2004).
15. Foster, J. J. *et al.* Polarisation vision: overcoming challenges of working with a property of light we barely see. *Naturwissenschaften* **105**, 27, <https://doi.org/10.1007/s00114-018-1551-3> (2018).
16. Wolf, R., Gebhardt, B., Gademann, R. & Heisenberg, M. Polarization Sensitivity of Course Control in *Drosophila-Melanogaster*. *J Comp Physiol* **139**, 177–191 (1980).
17. Weir, P. T. & Dickinson, M. H. Flying *Drosophila* Orient to Sky Polarization. *Current Biology* **22**, 21–27 (2012).
18. Warren, T. L., Weir, P. T. & Dickinson, M. H. Flying *Drosophilamelanogaster* maintain arbitrary but stable headings relative to the angle of polarized light. *J Exp Biol* **221**, <https://doi.org/10.1242/jeb.177550> (2018).

19. Warren, T. L., Giraldo, Y. M. & Dickinson, M. H. Celestial navigation in *Drosophila*. *J Exp Biol* **222**, <https://doi.org/10.1242/jeb.186148> (2019).
20. Dickinson, M. H. Death Valley, *Drosophila*, and the Devonian toolkit. *Annu Rev Entomol* **59**, 51–72, <https://doi.org/10.1146/annurev-ento-011613-162041> (2014).
21. Coyne, J. A., Bryant, S. H. & Turelli, M. Long-distance migration of *Drosophila*. 2. Presence in desolate sites and dispersal near a desert oasis. *Am. Nat.*, 847–861 (1987).
22. Coyne, J. A. *et al.* Long-distance migration of *Drosophila*. *Am. Nat.*, 589–595 (1982).
23. Wernet, M. F. *et al.* Genetic dissection reveals two separate retinal substrates for polarization vision in *Drosophila*. *Curr Biol* **22**, 12–20, <https://doi.org/10.1016/j.cub.2011.11.028> (2012).
24. Giraldo, Y. M. *et al.* Sun Navigation Requires Compass Neurons in *Drosophila*. *Curr Biol* **28**, 2845–2852 e2844, <https://doi.org/10.1016/j.cub.2018.07.002> (2018).
25. Labhart, T. & Meyer, E. P. Detectors for polarized skylight in insects: a survey of ommatidial specializations in the dorsal rim area of the compound eye. *Microsc Res Tech* **47**, 368–379 (1999). 10.1002/(SICI)1097-0029(19991215)47:6<368::AID-JEMT2>3.0.CO;2-Q
26. Wada, S. Special Marginal Ommatidia of Flies (Diptera-Brachycera) - Architecture and Distribution in Compound Eyes. *Z Morphol Tiere* **77**, 87–125 (1974).
27. Wada, S. Special Rhabdomic Type in Compound Eye of Flies. *Experientia* **27**, 1237–& (1971).
28. Fortini, M. E. & Rubin, G. M. The Optic Lobe Projection Pattern of Polarization-Sensitive Photoreceptor Cells in *Drosophila-Melanogaster*. *Cell and Tissue Research* **265**, 185–191 (1991).
29. Fortini, M. E. & Rubin, G. M. Analysis of Cis-Acting Requirements of the Rh3 and Rh4 Genes Reveals a Bipartite Organization to Rhodopsin Promoters in *Drosophila-Melanogaster*. *Gene Dev* **4**, 444–463 (1990).
30. Wernet, M. F. *et al.* Homothorax switches function of *Drosophila* photoreceptors from color to polarized light sensors. *Cell* **115**, 267–279 (2003).
31. Wunderer, H. & Smola, U. Fine-Structure of Ommatidia at the Dorsal Eye Margin of Calliphora-Erythrocephala Meigen (Diptera, Calliphoridae) - an Eye Region Specialized for the Detection of Polarized-Light. *Int J Insect Morphol* **11**, 25–38 (1982).
32. Smola, U. & Wunderer, H. Fly Rhabdomeres Twist Invivo. *J Comp Physiol* **142**, 43–49 (1981).
33. Tomlinson, A. Patterning the peripheral retina of the fly: decoding a gradient. *Dev Cell* **5**, 799–809 (2003).
34. Weir, P. T. *et al.* Anatomical Reconstruction and Functional Imaging Reveal an Ordered Array of Skylight Polarization Detectors in *Drosophila*. *J Neurosci* **36**, 5397–5404, <https://doi.org/10.1523/JNEUROSCI.0310-16.2016> (2016).
35. Henze, M. J., Dannenhauer, K., Kohler, M., Labhart, T. & Gesemann, M. Opsin evolution and expression in arthropod compound eyes and ocelli: insights from the cricket *Gryllus bimaculatus*. *BMC Evol Biol* **12**, 163, <https://doi.org/10.1186/1471-2148-12-163> (2012).
36. Schmeling, F. *et al.* Opsin expression, physiological characterization and identification of photoreceptor cells in the dorsal rim area and main retina of the desert locust, *Schistocerca gregaria*. *J Exp Biol* **217**, 3557–3568, <https://doi.org/10.1242/jeb.108514> (2014).
37. Labhart, T., Meyer, E. P. & Schenker, L. Specialized ommatidia for polarization vision in the compound eye of cockchafers, *Melolontha melolontha* (Coleoptera, Scarabaeidae). *Cell Tissue Res* **268**, 419–429 (1992).
38. Hegedus, R., Horvath, A. & Horvath, G. Why do dusk-active cockchafers detect polarization in the green? The polarization vision in *Melolontha melolontha* is tuned to the high polarized intensity of downwelling light under canopies during sunset. *J Theor Biol* **238**, 230–244, <https://doi.org/10.1016/j.jtbi.2005.05.033> (2006).
39. Barta, A. & Horvath, G. Why is it advantageous for animals to detect celestial polarization in the ultraviolet? Skylight polarization under clouds and canopies is strongest in the UV. *Journal of Theoretical Biology* **226**, 429–437 (2004).
40. Heinloth, T., Uhlhorn, J. & Wernet, M. F. Insect Responses to Linearly Polarized Reflections: Orphan Behaviors Without Neural Circuits. *Front Cell Neurosci* **12**, 50, <https://doi.org/10.3389/fncel.2018.00050> (2018).
41. Bender, J. A. & Dickinson, M. H. Visual stimulation of saccades in magnetically tethered *Drosophila*. *J Exp Biol* **209**, 3170–3182, <https://doi.org/10.1242/jeb.02369> (2006).
42. Labhart, T. & Wehner, R. In *Invertebrate Vision* (eds Warrant, E. J. & Nilsson, D. E.) 291–348 (Cambridge University Press, 2006).
43. Wehner, R. Polarization vision—a uniform sensory capacity? *J Exp Biol* **204**, 2589–2596 (2001).
44. Velez, M. M., Gohl, D., Clandinin, T. R. & Wernet, M. F. Differences in neural circuitry guiding behavioral responses to polarized light presented to either the dorsal or ventral retina in *Drosophila*. *J Neurogenet* **28**, 348–360, <https://doi.org/10.3109/01677063.2014.922556> (2014).
45. Velez, M. M., Wernet, M. F., Clark, D. A. & Clandinin, T. R. Walking *Drosophila* align with the e-vector of linearly polarized light through directed modulation of angular acceleration. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **200**, 603–614, <https://doi.org/10.1007/s00359-014-0910-6> (2014).
46. Rister, J. *et al.* Dissection of the peripheral motion channel in the visual system of *Drosophila melanogaster*. *Neuron* **56**, 155–170, <https://doi.org/10.1016/j.neuron.2007.09.014> (2007).
47. Clark, D. A., Bursztyn, L., Horowitz, M. A., Schnitzer, M. J. & Clandinin, T. R. Defining the computational structure of the motion detector in *Drosophila*. *Neuron* **70**, 1165–1177, <https://doi.org/10.1016/j.neuron.2011.05.023> (2011).
48. Maisak, M. S. *et al.* A directional tuning map of *Drosophila* elementary motion detectors. *Nature* **500**, 212–216, <https://doi.org/10.1038/nature12320> (2013).
49. Tuthill, J. C., Nern, A., Holtz, S. L., Rubin, G. M. & Reiser, M. B. Contributions of the 12 neuron classes in the fly lamina to motion vision. *Neuron* **79**, 128–140, <https://doi.org/10.1016/j.neuron.2013.05.024> (2013).
50. Liu, G. *et al.* Distinct memory traces for two visual features in the *Drosophila* brain. *Nature* **439**, 551–556, <https://doi.org/10.1038/nature04381> (2006).
51. Melnattur, K. V. *et al.* Multiple redundant medulla projection neurons mediate color vision in *Drosophila*. *J Neurogenet* **28**, 374–388, <https://doi.org/10.3109/01677063.2014.891590> (2014).
52. Manoli, D. S. *et al.* Male-specific fruitless specifies the neural substrates of *Drosophila* courtship behaviour. *Nature* **436**, 395–400, <https://doi.org/10.1038/nature03859> (2005).
53. Stockinger, P., Kvitsiani, D., Rotkopf, S., Tirian, L. & Dickson, B. J. Neural circuitry that governs *Drosophila* male courtship behavior. *Cell* **121**, 795–807, <https://doi.org/10.1016/j.cell.2005.04.026> (2005).
54. Turner-Evans, D. B. & Jayaraman, V. The insect central complex. *Curr Biol* **26**, R453–457, <https://doi.org/10.1016/j.cub.2016.04.006> (2016).
55. Heinze, S. & Homberg, U. Maplike representation of celestial E-vector orientations in the brain of an insect. *Science* **315**, 995–997, <https://doi.org/10.1126/science.1135531> (2007).
56. Franconville, R., Beron, C. & Jayaraman, V. Building a functional connectome of the *Drosophila* central complex. *Elife* **7**, <https://doi.org/10.7554/eLife.37017> (2018).
57. el Jundi, B., Pfeiffer, K., Heinze, S. & Homberg, U. Integration of polarization and chromatic cues in the insect sky compass. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **200**, 575–589, <https://doi.org/10.1007/s00359-014-0890-6> (2014).
58. Omoto, J. J. *et al.* Visual Input to the *Drosophila* Central Complex by Developmentally and Functionally Distinct Neuronal Populations. *Curr Biol* **27**, 1098–1110, <https://doi.org/10.1016/j.cub.2017.02.063> (2017).
59. Sun, Y. *et al.* Neural signatures of dynamic stimulus selection in *Drosophila*. *Nat Neurosci* **20**, 1104–1113, <https://doi.org/10.1038/nn.4581> (2017).

60. Kim, S. S., Rouault, H., Druckmann, S. & Jayaraman, V. Ring attractor dynamics in the *Drosophila* central brain. *Science* **356**, 849–853, <https://doi.org/10.1126/science.aal4835> (2017).
61. Seelig, J. D. & Jayaraman, V. Feature detection and orientation tuning in the *Drosophila* central complex. *Nature* **503**, 262–266, <https://doi.org/10.1038/nature12601> (2013).
62. Seelig, J. D. & Jayaraman, V. Studying sensorimotor processing with physiology in behaving *Drosophila*. *Int Rev Neurobiol* **99**, 169–189, <https://doi.org/10.1016/B978-0-12-387003-2.00007-0> (2011).
63. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676–682, <https://doi.org/10.1038/nmeth.2019> (2012).

Acknowledgements

The authors thank Leigh Moss, Tanja Heinloth, Lena Naber and Nurelhoda Abdel Muti for experimental support, Gregor Belušič and Marco Ilić for help with the polarometry, Gerit Linneweber for statistical advice, as well as all members of the Wernet and Hiesinger groups for their input. This manuscript also benefited from the helpful comments by two anonymous reviewers. This work was supported by the Deutsche Forschungsgemeinschaft through grants WE 5761/2-1 and SFB958 (Teilprojekt A23), through AFOSR grant FA9550-19-1-7005, through the Berlin Excellency Cluster NeuroCure, with support from the Fachbereich Biologie, Chemie & Pharmazie of the Freie Universität Berlin, as well as the Division of Neurobiology at Freie Universität Berlin (support of FU Berlin and the National Institute of Health to Robin Hiesinger).

Author contributions

T.F.M. and M.F.W. planned the experiments. T.F.M. built the assay and performed all experiments. T.F.M. and M.F.W. designed the figures. M.F.W. wrote the manuscript, M.F.W. and T.F.M. finalized the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-019-53330-y>.

Correspondence and requests for materials should be addressed to M.F.W.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019