

10-29-2019

Nucleus pulposus primary cilia alter their length in response to changes in extracellular osmolarity but do not control TonEBP-mediated osmoregulation.

Hyowon Choi
Thomas Jefferson University

Vedavathi Madhu
Thomas Jefferson University

Irving M. Shapiro
Thomas Jefferson University

Makarand V. Risbud
Thomas Jefferson University
Follow this and additional works at: <https://jdc.jefferson.edu/orthofp>



Part of the [Orthopedics Commons](#), and the [Surgery Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Choi, Hyowon; Madhu, Vedavathi; Shapiro, Irving M.; and Risbud, Makarand V., "Nucleus pulposus primary cilia alter their length in response to changes in extracellular osmolarity but do not control TonEBP-mediated osmoregulation." (2019). *Department of Orthopaedic Surgery Faculty Papers*. Paper 129.

<https://jdc.jefferson.edu/orthofp/129>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Orthopaedic Surgery Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

OPEN

Nucleus pulposus primary cilia alter their length in response to changes in extracellular osmolarity but do not control TonEBP-mediated osmoregulation

Hyowon Choi^{1,2}, Vedavathi Madhu¹, Irving M. Shapiro^{1,2} & Makarand V. Risbud^{1,2*}

The nucleus pulposus (NP) cells adapt to their physiologically hyperosmotic microenvironment through Tonicity-responsive enhancer binding protein (TonEBP/nuclear factor of activated T-cell5 [NFAT5])-mediated osmoregulation. Primary cilia in different organs serve diverse roles including osmosensing, but its contribution to NP cell osmoadaptive response is unknown. A high percentage of cultured primary NP cells possessed primary cilia that changed length in response to osmotic stimuli. Stable silencing of *Intraflagellar Transport 88* (*Ift88*) or *Kinesin Family Member 3 A* (*Kif3a*) to inhibit the formation of primary cilia did not affect hyperosmotic upregulation of TonEBP. While *ShKif3a* blocked hyperosmotic increase of TonEBP-Transactivation Domain (TAD) activity, overall the knockdown of either gene did not alter the hyperosmotic status of proximal promoter activities and transcription of key TonEBP targets. On the other hand, a small decrease in TonEBP level under hypoosmotic condition was attenuated by *Ift88* or *Kif3a* knockdown. Noteworthy, none of the TonEBP target genes were responsive to hypoosmotic stimulus in control and *Ift88* or *Kif3a* knockdown cells, suggesting the primary role of TonEBP in the hyperosmotic adaptation of NP cells. Similarly, in *Kif3a* null mouse embryonic fibroblasts (MEFs), the overall TonEBP-dependent hyperosmotic responses were preserved. Unlike NP cells, TonEBP targets were responsive to hypoosmolarity in wild-type MEFs, and these responses remained intact in *Kif3a* null MEFs. Together, these results suggest that primary cilia are dispensable for TonEBP-dependent osmoadaptive response.

The nucleus pulposus (NP) is a gelatinous center-most structure of the intervertebral disc. It is surrounded by concentric layers of annulus fibrosus and sandwiched between cartilaginous endplates. The extracellular matrix of NP consists of collagens and abundant proteoglycans that are bound by sulfated glycosaminoglycan chains¹. The high negative charge density of glycosaminoglycan molecules draws water and cations, in particular, Na⁺, into the NP tissue. This contributes to the elevated osmotic swelling pressure within the tissue compartment and provides the tissue with its ability to resist daily spinal loadings^{1,2}. The NP matrix is, therefore, hyperosmotic with estimated tissue osmolarity ranging from 430 to 496 mOsm/kg H₂O^{3,4}. With aging and degeneration of the tissue, the aberrant changes in cell phenotype lead to altered extracellular matrix composition, decreased osmotic pressurization of the NP, and compromised the mechanical function of the tissue^{1,5,6}. Therefore, adaptation to their osmodynamic niche for survival and function is crucial for the resident NP cells.

Tonicity-responsive enhancer binding protein (TonEBP/nuclear factor of activated T-cell5 [NFAT5]) is a Rel homology transcription factor that has been well characterized for its osmolarity-dependent function in mammalian cells including NP cells⁷⁻¹⁴. In response to a hyperosmotic stimulus, TonEBP transcriptionally controls the expression of several osmoprotective genes including taurine transporter (TauT), betaine-GABA transporter (BGT-1), sodium/myo-inositol co-transporter (SMIT), and aldose reductase (AR), all of which are essential regulators of intracellular levels of non-ionic osmolytes¹⁵⁻¹⁸. Additionally, TonEBP transcriptionally increases the

¹Department of Orthopaedic Surgery, Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, PA, USA. ²Graduate Program in Cell Biology and Regenerative Medicine, Thomas Jefferson University, Philadelphia, PA, USA. *email: makarand.risbud@jefferson.edu

expression of heat shock protein-70 (HSP-70) to maintain proper protein folding, cellular trafficking, and degradation of misfolded proteins under hyperosmotic conditions^{19,20}. It was recently shown that TonEBP is highly expressed in NP as well as notochord cells of developing mouse embryo, supporting its importance in NP development and maintenance²¹.

Primary cilia found in most types of mammalian cells have diverse roles, including modulation of key signaling pathways^{22–28}. They can also function as a sensory organelle that relays extracellular stimuli, such as changes in osmolarity, to intracellular signaling pathways^{29–38}. In *Caenorhabditis elegans*, OSM-9, a homolog of mammalian transient receptor potential channel vanilloid subfamily (TRPV), and OCR-2, OSM-9/capsaicin receptor related TRPV channel, that are localized in primary cilia are important for sensing extracellular hyperosmotic stimulus³⁵. Similarly, primary cilia of mammalian cholangiocytes and articular chondrocytes are shown to link extracellular hypoosmotic stimulus to intracellular calcium signaling pathway^{36,37}. In a recent report, Siroky *et al.* showed that renal epithelial cells without primary cilia have weakened induction of TonEBP target genes *AR/Akr1b3* as well as *Bgt1* under hyperosmotic conditions³⁸. Although the role of TonEBP in modulating osmoresponse in NP cells has been well studied, it is unknown whether primary cilia contribute to this process. The objective of this study was to investigate if primary cilia function as osmosensory organelles in NP cells. Specifically, we examined if primary cilia control TonEBP-mediated osmoadaptive response through loss-of-function studies measuring the expression of TonEBP and its target genes after inhibition of primary cilia formation. Furthermore, we confirmed our findings in NP cells using *Kif3a* null mouse embryonic fibroblasts (MEFs) that are completely devoid of primary cilia.

Results

The length of primary cilia in NP cells is responsive to changes in extracellular osmolarity.

Primary cilia were visualized in cultured primary rat NP cells by co-immunostaining acetylated α -tubulin and γ -tubulin, labeling ciliary axoneme and basal bodies, respectively (Fig. 1a,b). Previous studies showed that the length of primary cilia in different types of cells changed in response to extracellular stimuli^{39–41}. To examine if primary cilia in NP cells respond to extracellular osmotic stimulus, we cultured NP cells under different osmotic conditions and measured the length of the cilia. The average length of primary cilia was significantly shorter under hypoosmotic condition (200 mOsm/kg H₂O) compared to isoosmotic (330 mOsm/kg H₂O) condition (Fig. 1c,d; $p < 0.0001$ for both 200 mOsm/kg H₂O and 450 mOsm/kg H₂O groups). On the other hand, the length of primary cilia increased under hyperosmotic condition (450 mOsm/kg H₂O), suggesting that the primary cilia in NP cells are sensitive to changes in extracellular osmolarity.

Inhibition of primary cilia formation in NP cells does not affect TonEBP expression in hyperosmotic condition.

The function of primary cilia as an osmosensor has been observed in *C. elegans* as well as in some types of mammalian cells, including renal tubular epithelial cells, articular chondrocytes, and cholangiocytes^{35–38}. NP cells reside in an osmotically active microenvironment due to high proteoglycan content of the NP matrix and dynamic loading of the spine. We examined if primary cilia of the NP cells play a role in sensing extracellular osmolarity and mediating cellular osmotic response. We inhibited formation of primary cilia in NP cells by performing stable knockdown of *Ift88*, a component of complex B of the intraflagellar transport (IFT) particles^{42–45} (Fig. 2a–c; Supp. Fig. S1–1), or *Kif3a*, a motor subunit of kinesin 2⁴⁶ (Fig. 2d–f), both of which are necessary for ciliogenesis. Lentiviral transduction of NP cells with shRNA against *Ift88* or *Kif3a* resulted in a significant decrease in the transcript and protein levels of IFT88 (Fig. 2a–c; $p < 0.0001$ for all statistical comparisons) or KIF3A (Fig. 2d–f; $p < 0.0001$ for Sh*Kif3a* #1 and #2 isoosmotic groups in Fig. 2d; $p = 0.0002$ for Sh*Kif3a* #1 isoosmotic group, $p = 0.346$ for Sh*Kif3a* #2 isoosmotic group in Fig. 2f; Supp. Fig. S1–1), respectively. Stable silencing of either gene resulted in a decreased number of cells with primary cilia (Fig. 2g). Quantification of the number of cells with primary cilia confirmed this result (Fig. 2h; $p < 0.0001$ for all groups). Overall, the lengths of primary cilia that remained after the stable silencing of *Ift88* or *Kif3a* were not significantly different from that of the control cells (Fig. 2i; $p = 0.0334$ for Sh*Ift88* #2, all other groups were statistically not significant).

To determine if inhibition of primary cilia formation resulted in dysregulation of cellular osmoresponse, we stably silenced *Ift88* or *Kif3a* and measured the expression of TonEBP/NFAT5, a transcription factor crucial for osmoregulation in NP cells^{8–11}. While there was a trend of increase in *TonEBP/Nfat5* transcript levels under hyperosmotic conditions (550 mOsm/kg H₂O), this increase did not reach a statistical significance in both control and *Ift88* knockdown cells (Fig. 3a; all analyses were statistically not significant). Hyperosmotic increase in TonEBP protein levels was unaffected by *Ift88* knockdown (Fig. 3b,c; $p = 0.0586$ for Sh*Ctrl* 550 mOsm/kg H₂O, $p = 0.0012$ for Sh*Ift88* #1 550 mOsm/kg H₂O, $p = 0.0141$ for Sh*Ift88* #2 550 mOsm/kg H₂O; Supp. Fig. S1–1). Similarly, when *Kif3a* was stably silenced, the trend of increase in *TonEBP/Nfat5* transcript levels and the upregulation of TonEBP protein under hyperosmotic condition were maintained (Fig. 3d–f; Statistical non-significance for all analyses in Fig. 3d; $p = 0.023$ for Sh*Ctrl* 550 mOsm/kg H₂O, $p = 0.059$ for Sh*Kif3a* #1 550 mOsm/kg H₂O; Supp. Fig. S1–1). Taken together, our data suggest that inhibition of primary cilia formation in NP cells does not affect their TonEBP expression.

Primary cilia do not modulate TonEBP activity and TonEBP target gene expression in NP cells under hyperosmotic conditions.

Hyperosmotic stimulus leads to the increased transcriptional activity of TonEBP in the nucleus^{9,11,47}. We investigated if primary cilia controlled TonEBP activity without affecting its expression. Changes in proximal promoter activities of well-characterized TonEBP target genes including, *AR* (aldose reductase), *HSP-70* and *TauT* (taurine transporter) were measured in NP cells transfected with shRNA against *Ift88* or *Kif3a* (Fig. 4a–c). These promoters contain highly conserved TonEBP-binding sites that are active in NP cells^{10,15,19,20,47,48}. Hyperosmotic (550 mOsm/kg H₂O) increase in the *AR* (Fig. 4a; $p < 0.0001$) and *TauT* (Fig. 4c; $p < 0.0001$) promoter activities was not affected by either Sh*Ift88* ($p = 0.0004$ for clone #1, $p = 0.0012$




Figure 1. NP cell primary cilia modulate their lengths in response to changes in extracellular osmolarity. (a) Immunofluorescence staining of acetylated α -tubulin (green) and γ -tubulin (red) to mark primary cilia axoneme and basal bodies, respectively, in primary rat NP cells. Scale bar = 30 μm . (b) Zoomed-in image of a primary cilium from the area demarcated by the white square in panel (a). (c,d) Primary cilia of rat NP cells cultured under different osmotic conditions for 24 h were visualized by immunofluorescence staining of acetylated α -tubulin. (c) The lengths of primary cilia increase in response to increased osmolarity (450 mOsm/kg H_2O) compared to isoosmotic control (330 mOsm/kg H_2O) conditions, whereas they appear shorter under hypoosmotic conditions (200 mOsm/kg H_2O). White arrows mark primary cilia. Scale bar = 50 μm . (d) Quantification of primary cilium length was done using ImageJ software. (n = 3 experiments; at least 150 cells/group) Data are represented as scatter plots (mean \pm SEM). One-way ANOVA with Dunnett's multiple comparison test was used to determine statistical significance.

for clone #2 for AR-luc; $p < 0.0001$ for both clones for TauT-luc) or ShKif3a ($p = 0.001$ for clone #1, $p = 0.0003$ for clone #2 for AR-luc; $p = 0.0009$ for clone #1 and $p < 0.0001$ for clone #2 for TauT-luc). Similarly, the HSP-70 promoter activity increased under hyperosmotic conditions with or without ShIft88 or ShKif3a. While one of the ShIft88 and ShKif3a clone groups did not reach a statistical significance, the trend of hyperosmotic induction of HSP-70 promoter activity was maintained (Fig. 4b; $p = 0.0011$ for ShIft88 #2, $p = 0.0357$ for ShKif3a #2). In addition, we measured the activity of the TonEBP transactivation domain (TAD) in response to the hyperosmotic stimulus (550 mOsm/kg H_2O) with or without ShIft88 and ShKif3a. While ShIft88 #2 inhibited the hyperosmotic increase of TonEBP-TAD activity, the other clone had no effect (Fig. 4d; $p = 0.0259$ for ShIft88 #1). Both clones of ShKif3a prevented a further increase of TonEBP-TAD activity in response to the hyperosmotic stimulus. However, the average level of TonEBP-TAD activity under hyperosmotic conditions was similar or higher in knockdown cells when compared to ShCtr group, suggesting that the overall TonEBP-TAD activity was unaffected by ShIft88 or ShKif3a (Fig. 4d).

To confirm that TonEBP-mediated hyperosmotic response was independent of primary cilia, we measured mRNA levels of TonEBP targets in NP cells with Ift88 or Kif3a knockdown. AR/Akr1b1, TauT/Slc6a6 as well as sodium/myo-inositol co-transporter (SMIT/Slc5a3) are osmolarity-dependent target genes of TonEBP^{15–18}. Ift88 knockdown did not affect the hyperosmotic (550 mOsm/kg H_2O) increase in the expression of AR/Akr1b1 and SMIT/Slc5a3 genes (Fig. 5a,b; $p = 0.0065$ for ShCtr, $p = 0.0444$ for ShIft88 #1, $p = 0.0105$ for ShIft88 #2 in Fig. 5a; $p = 0.0024$ for ShCtr, $p = 0.0357$ for ShIft88 #1, $p = 0.0031$ for ShIft88 #2 in Fig. 5b). While the increase in TauT/Slc6a6 levels in response to hyperosmotic stimulus did not reach a statistical significance, the trend of increase




Figure 2. Stable knockdown of *Ift88* or *Kif3a* inhibits formation of NP cell primary cilia. **(a)** *Ift88* mRNA levels in NP cells transduced with control (ShCtr) or two different Sh*Ift88* clones were measured by qRT-PCR to confirm the knockdown ($n \geq 5$). **(b)** Western blot image showing significant reduction of IFT88 protein levels after the knockdown of *Ift88*. **(c)** Densitometry analyses of Western blots confirm significant knockdown of IFT88 ($n \geq 5$). **(d–f)** qRT-PCR, Western blot, and corresponding densitometry analyses, show significant downregulation of KIF3A after stable knockdown using two different Sh*Kif3a* clones ($n \geq 4$). **(g)** Acetylated α -tubulin immunofluorescence staining after lentiviral transduction of Sh*Ift88* or Sh*Kif3a* shows inhibition of primary cilia formation in majority of rat NP cells. Scale bar = 75 μ m. White arrowheads point to primary cilia. **(h,i)** Quantitation of percentage of NP cells with primary cilia and primary cilium length after stable silencing of *Ift88* or *Kif3a* ($n = 3$; at least 150 cells/group). Data are represented as scatter plots (mean \pm SEM). ns = not significant. One-way ANOVA or Kruskal-Wallis test with Sidak's, Holm-Sidak's, or Dunn's multiple comparison test was used based on the distribution of the data to determine statistical significance. For statistical comparison of the percentages of NP cells with primary cilia, Fisher's exact test was used. Western blot images were cropped and acquired under same experimental conditions. See Supplementary Fig. S1–1 for un-cropped Western blot images.




Figure 3. Knockdown of *Ift88* or *Kif3a* in NP cells does not affect hyperosmotic upregulation of TonEBP. (a) *TonEBP/Nfat5* mRNA levels in NP cells with *Ift88* knockdown ($n \geq 5$). (b) Western blot image showing increased TonEBP expression in response to hyperosmolarity (550 mOsm/kg H₂O) independently of *Ift88* knockdown. (c) Densitometry analyses of TonEBP with *Ift88* knockdown ($n \geq 4$). (d) *TonEBP/Nfat5* mRNA levels in NP cells with *Kif3a* knockdown ($n \geq 3$). (e) Western blot image showing that hyperosmotic induction of TonEBP is maintained after *Kif3a* knockdown. (f) Densitometry analyses of TonEBP after *Kif3a* knockdown ($n \geq 4$). Data are represented as scatter plots (mean \pm SEM). ns = not significant. One-way ANOVA or Kruskal-Wallis test with Sidak's or Dunn's multiple comparison test was used based on the distribution of the data to determine statistical significance. Western blot images were cropped and acquired under same experimental conditions. See Supplementary Fig. S1-1 for un-cropped Western blot images.

was unaffected by *Ift88* knockdown (Fig. 5c). Likewise, stable silencing of *Kif3a* did not affect the induction of *SMIT/Slc5a3* and *AR/Akr1b1* in response to hyperosmolarity (550 mOsm/kg H₂O) (Fig. 5d, e; $p = 0.0766$ for ShCtr, $p = 0.0777$ for Sh*Kif3a* #1, $p < 0.0001$ for Sh*Kif3a* #2 in Fig. 5d; $p = 0.0015$ for ShCtr, $p = 0.0559$ for Sh*Kif3a* #1, $p = 0.0214$ for Sh*Kif3a* #2 in Fig. 5e). In addition, the hyperosmotic increase of *TauT/Slc6a6* expression was preserved with Sh*Kif3a*, except one of the clones (Fig. 5f; $p = 0.0459$ for ShCtr, $p = 0.008$ for Sh*Kif3a* #1). Taken together, the gene expression data showed that inhibition of primary cilia formation in NP cells did not affect the overall TonEBP transcriptional activity or the target gene expression under hyperosmotic conditions.

The absence of primary cilia has a minor effect on hypoosmotic downregulation of TonEBP and its targets in NP cells.

The tissue osmotic conditions of the NP compartment undergo diurnal changes due to dynamic loading and unloading of the spine. During disc degeneration, the breakdown of the proteoglycan matrix by proteases decreases the tissue osmolarity^{49–51}. Consequently, NP cells need to maintain their function and integrity under both hyper- and hypo-osmotic conditions^{52,53}. We, therefore, assessed the effects of inhibiting primary cilia formation in NP cells on TonEBP expression under hypoosmotic conditions (200 mOsm/kg H₂O). NP cells with *Ift88* (Fig. 6a–c; $p < 0.0001$ for all analyses in Fig. 6a,c; Supp. Figs S1 and 2) or *Kif3a* (Fig. 6f–h; $p = 0.0019$ for Sh*Kif3a* #1, $p = 0.0075$ for Sh*Kif3a* #2 in Fig. 6f; $p = 0.0139$ for Sh*Kif3a* #1, $p = 0.026$ for Sh*Kif3a* #2 in Fig. 6h) knockdown showed significant hypoosmotic decrease in *TonEBP/Nfat5* transcript, compared to control cells that showed similar decreasing trend (Fig. 6d,i; $p = 0.0043$ for Sh*Ift88* #1, $p = 0.0029$ for Sh*Ift88* #2 in Fig. 6d; $p = 0.0089$ for Sh*Kif3a* #2 in Fig. 6i). On the other hand, a small but significant decrease in TonEBP protein levels seen in control cells under the hypoosmotic condition was attenuated following knockdown of *Ift88* (Fig. 6b,e; $p = 0.0266$ for ShCtr, $p = 0.0761$ for Sh*Ift88* #1) or *Kif3a* (Fig. 6g,j; $p = 0.0515$ for ShCtr; Supp. Figs S1 and 2). Although the changes in mRNA and protein levels in response to hypoosmotic stimuli were somewhat




Figure 4. *ShIfi88* or *ShKif3a* does not affect the promoter activities of TonEBP target genes under hyperosmotic conditions. (a–c) Proximal promoter activities of *AR* (a), *HSP-70* (b), and *TauT* (c) in response to hyperosmotic stimuli (550 mOsm/kg H₂O) with or without *ShIfi88* or *ShKif3a*. The promoter activities of TonEBP target genes are mostly unaffected by *ShIfi88* or *ShKif3a*. (d) *ShKif3a* as well as one of the *ShIfi88* clones prevent further upregulation of TonEBP-TAD (Ton-TAD) activity in response to hyperosmolarity. Interestingly, the levels of TonEBP-TAD activity under isoosmotic conditions are increased with *ShIfi88* or *ShKif3a*. Data are represented as scatter plots (mean ± SEM) (n ≥ 3 with 3 technical replicates per biological replicate). ns = not significant. One-way ANOVA or Kruskal-Wallis test with Sidak's or Dunn's multiple comparison test was used based on the distribution of the data to determine statistical significance.

opposing in cells lacking primary cilia, these results suggested that primary cilia may play a minor role in maintaining TonEBP levels in NP cells under hypoosmotic conditions.

We then measured mRNA levels of TonEBP targets to investigate if the small changes in TonEBP levels in cells without primary cilia had functional implications in overall hypoosmotic cellular response. *AR/Ark1b1*, *SMIT/Slc5a3*, and *TauT/Slc6a6* were not affected by stable knockdown of *Ifi88* or *Kif3a* (Fig. 7a–f; all statistical analyses were not significant in Fig. 7a,e,f; $p = 0.0046$ for *ShIfi88* #1 in Fig. 7b; non-significant for *ShCtr* and *ShIfi88* #1, $p = 0.0139$ for *ShIfi88* #2 in Fig. 7c; non-significant for *ShCtr* and *ShKif3a* #2, $p = 0.0088$ for *ShKif3a* #1 in Fig. 7d). Interestingly, none of these osmotic targets showed a significant decrease in their expression under hypoosmotic conditions, and this trend was not affected by either *Ifi88* or *Kif3a* knockdown (Fig. 7a–f). Taken together, our results showed that primary cilia may have a small role in modulating hypoosmotic levels of TonEBP, but this did not lead to any appreciable effect on target gene expression.

TonEBP expression and activity are not mediated by primary cilia in mouse embryonic fibroblasts. Similar to NP cells, mouse embryonic fibroblasts (MEFs) induce TonEBP-dependent osmotic response under hyperosmotic conditions⁸. Parallel experiments were performed using *Kif3a* null MEFs that completely lack primary cilia to further confirm that primary cilia do not control TonEBP-dependent osmotic response⁵⁴. As reported in other studies, TonEBP mRNA (Fig. 8a; $p = 0.0085$) and protein (Fig. 8b,c; $p = 0.0005$) levels were significantly upregulated in wild-type MEFs under hyperosmotic conditions (600 mOsm/kg H₂O). On the other




Figure 5. Hyperosmotic upregulation of TonEBP target gene expression is not affected by inhibition of primary cilia formation in NP cells. (a–c) mRNA levels of TonEBP targets, *AR/Akr1b1*, *SMIT/Slc5a3* and *TauT/Slc6a6*, showing that stable silencing of *Ift88* does not affect their response to hyperosmotic stimuli (550 mOsm/kg H₂O) ($n \geq 5$). (d–f) mRNA levels of *AR/Akr1b1*, *SMIT/Slc5a3* and *TauT/Slc6a6* demonstrate that *Kif3a* knockdown also does not influence hyperosmotic upregulation of these genes ($n \geq 3$). Data are represented as scatter plots (mean \pm SEM). ns = not significant. One-way ANOVA or Kruskal-Wallis test with Sidak's or Dunn's multiple comparison test was used based on the distribution of the data to determine statistical significance.

hand, the increase of TonEBP expression in *Kif3a* null MEFs under hyperosmotic conditions did not reach a statistical significance (Fig. 8a–c). In response to hypoosmotic conditions (200 mOsm/kg H₂O), both wild-type and *Kif3a* null MEFs downregulated TonEBP transcript and protein levels (Fig. 8a–c; $p = 0.0081$ for wild-type MEFs, $p = 0.0015$ for *Kif3a* null MEFs in Fig. 8a; $p = 0.0308$ for wild-type MEFs, $p = 0.0042$ for *Kif3a* null MEFs in Fig. 8c; Supp. Figs S1 and 3).

To investigate if cilia controlled osmolarity-dependent TonEBP activity in MEFs, we measured the activation status of TonEBP-TAD in wild-type and *Kif3a* null MEFs under different osmotic conditions. Both wild-type and *Kif3a* null MEFs increased and decreased TonEBP-TAD activities under hyperosmotic conditions (600 mOsm/kg H₂O) (Fig. 8d; $p = 0.0126$ for 500 mOsm/kg H₂O, $p = 0.0002$ for 600 mOsm/kg H₂O in wild-type MEFs; $p < 0.0001$ for both hyperosmotic conditions in *Kif3a* null MEFs), and hypoosmotic conditions (200 mOsm/kg H₂O) (Fig. 8e; $p < 0.0001$ for both wild-type and *Kif3a* null MEFs), respectively. These results suggested that the effect of osmolarity on TonEBP-TAD activity remained unaffected in the absence of primary cilia.

We then measured the activities of TonEBP responsive reporters and the level of TonEBP target genes in wild-type and *Kif3a* null MEFs to determine if the presence of primary cilia affected TonEBP transcriptional




Figure 6. Inhibition of primary cilia formation in NP cells have minimal effect on hypoosmotic downregulation of TonEBP levels. (a–c) qRT-PCR, Western blot and densitometry analyses demonstrating IFT88 levels under hypoosmotic conditions (200 mOsm/kg H₂O) after stable knockdown of *Ift88*. (d) *TonEBP/Nfat5* expression in NP cells with *Ift88* knockdown under hypoosmotic conditions ($n \geq 3$). (d) Hypoosmotic decrease in TonEBP protein levels is attenuated by *Ift88* knockdown ($n \geq 4$). (f,g) KIF3A mRNA and protein levels in NP cells with *Kif3a* knockdown. (h) *TonEBP/Nfat5* mRNA expression in NP cells with *Kif3a* knockdown under hypoosmotic conditions ($n \geq 3$). (i) Hypoosmotic downregulation of TonEBP is blunted after *Kif3a* knockdown ($n \geq 4$). Data are represented as scatter plots (mean \pm SEM). ns = not significant. One-way ANOVA or Kruskal-Wallis test with Sidak's or Dunn's multiple comparison test was used based on the distribution of the data to determine statistical significance. See Supplementary Figs S1 and 2 for un-cropped Western blot images.

activity. Hyperosmotic stimulus (600 mOsm/kg H₂O) significantly increased *AR* and *HSP-70* promoter activities in both wild-type and *Kif3a* null MEFs (Fig. 9a,b; $p = 0.0003$ for 500 mOsm/kg H₂O, $p = 0.0001$ for 600 mOsm/kg H₂O for wild-type MEFs, $p = 0.0237$ for 500 mOsm/kg H₂O, $p < 0.0001$ for 600 mOsm/kg H₂O for *Kif3a* null MEFs in Fig. 9a; $p = 0.0596$ for 500 mOsm/kg H₂O, $p = 0.0017$ for 600 mOsm/kg H₂O for wild-type MEFs, $p < 0.0001$ for hyperosmolarity in *Kif3a* null MEFs in Fig. 9b). Likewise, the temporal response of *TauT* promoter to the hyperosmotic stimulus was similar in both wild-type and *Kif3a* null MEFs, where a robust activation at 16 h post-treatment was observed (Fig. 9c; $p < 0.0001$ for 16 h and 24 h in both wild-type and *Kif3a*




Figure 7. TonEBP target gene expression under hypoosmotic conditions is unaffected by inhibition of primary cilia formation in NP cells. (a–f) mRNA levels of *AR/Akr1b1*, *SMIT/Slc5a3* and *TauT/Slc6a6* under hypoosmotic conditions (200 mOsm/kg H₂O) following knockdown of *Ifi88* (a–c) or *Kif3a* (d–f). Stable silencing of either *Ifi88* or *Kif3a* does not affect the trend of downregulation in *SMIT/Slc5a3* and *TauT/Slc6a6* under hypoosmotic conditions ($n \geq 3$). Data are represented as scatter plots (mean \pm SEM). ns = not significant. One-way ANOVA or Kruskal-Wallis test with Sidak's or Dunn's multiple comparison test was used based on the distribution of the data to determine statistical significance.

null MEFs). Interestingly, the *AR* promoter activity was insensitive to the hypoosmotic stimulus (200 mOsm/kg H₂O) in wild-type cells, while it was significantly downregulated in *Kif3a* null MEFs (Fig. 9d; $p < 0.0001$). Both wild-type and *Kif3a* null MEFs showed a hypoosmotic reduction in *HSP70* promoter activity (Fig. 9e; $p = 0.0262$ for wild-type, $p < 0.0001$ for *Kif3a* null MEFs). A gradual and time-dependent decrease in *TauT* promoter activity was also observed under hypoosmotic conditions (Fig. 9f; $p < 0.0001$).

A qRT-PCR analysis was performed to measure the expression of select TonEBP target genes in MEFs under different osmotic conditions. *AR/Akr1b1* was significantly upregulated in both wild type and *Kif3a* null MEFs under hyperosmotic conditions (600 mOsm/kg H₂O) (Fig. 9g; $p = 0.0015$ for wild-type, $p < 0.0001$ for null MEFs). In agreement with its promoter activity, *AR/Akr1b1* expression was not changed in wild-type MEFs but was significantly decreased in *Kif3a* null MEFs in response to the hypoosmotic stimulus (200 mOsm/kg H₂O) (Fig. 9g; $p = 0.021$). *SMIT/Slc5a3* expression in both wild-type and *Kif3a* null MEFs was significantly increased and decreased under hyperosmotic and hypoosmotic conditions, respectively (Fig. 9h; $p < 0.0001$ for hyperosmotic condition in wild-type and *Kif3a* null MEFs, $p = 0.0008$ for hypoosmotic condition in wild-type MEFs,




Figure 8. The absence of primary cilia in *Kif3a* null MEFs has a small effect on hyperosmotic induction of TonEBP without influencing its TAD activity (a) *TonEBP/Nfat5* gene expression in wild-type and *Kif3a* null (*Kif3a*^{-/-}) MEFs under hyper- (600 mOsm/kg H₂O) and hypoosmotic (200 mOsm/kg H₂O) conditions. (b,c) Western blot image and corresponding densitometry analyses showing TonEBP levels under different osmotic conditions in wild-type and *Kif3a* null MEFs. *Kif3a* null MEFs show slightly attenuated hyperosmotic increase but unaffected hypoosmotic decrease in TonEBP expression. (d,e) TonEBP-TAD (Ton-TAD) activity in wild-type and *Kif3a* null MEFs under hyperosmotic (d) and hypoosmotic (e) conditions (n = 4). Similar to wild-type cells, *Kif3a* null MEFs show increase and decrease in TonEBP-TAD activity under hyperosmotic and hypoosmotic conditions, respectively (n ≥ 3 with 3 technical replicates per biological replicate). Data are represented as scatter plots (mean ± SEM). ns = not significant. One-way ANOVA or Kruskal-Wallis test with Sidak's or Dunn's multiple comparison test was used based on the distribution of the data to determine statistical significance. Western blot images were cropped and acquired under same experimental conditions. See Supplementary Figs S1 and 3 for un-cropped Western blot images.

$p < 0.0001$ for hypoosmotic condition in *Kif3a* null MEFs). Interestingly, unlike in wild-type MEFs, *TauT/Slc6a6* mRNA levels were not induced by hyperosmolarity in *Kif3a* null MEFs (Fig. 9i; $p = 0.0105$ in wild-type MEFs). The hypoosmotic decrease in *TauT/Slc6a6* levels was maintained in both wild-type and *Kif3a* null MEFs (Fig. 9j; $p < 0.0001$ in wild-type, $p = 0.0001$ in null MEFs). Taken together, the osmotic regulation of TonEBP activity in MEFs remained largely unchanged in the absence of primary cilia.

Discussion

Primary cilia serve diverse physiological functions that are well-conserved from *C. elegans* to vertebrates. Their sensory function in various cell types and organ systems is essential in multiple biological processes such as development and maintenance of tissue homeostasis. There are only a few studies examining the presence of primary cilia in the disc tissues⁵⁵. However, few studies have investigated the role of primary cilia in the intervertebral disc⁵⁶. A few studies have suggested that primary cilia can function as osmosensors in mammalian cells^{36–38}. In rat cholangiocytes or porcine articular chondrocytes, removal of primary cilia inhibits intracellular Ca²⁺ increase




Figure 9. The absence of primary cilia does not affect TonEBP activity and target gene expression in MEFs (a–c) Proximal promoter activities of AR, HSP-70 and *TauT* in wild-type and *Kif3a* null MEFs show that hyperosmotic induction (600 mOsm/kg H₂O) of these promoter activities are intact in null MEFs. (d–f) AR promoter activity is unaffected in wild-type MEFs but decreased in *Kif3a* null MEFs under hypoosmotic conditions (200 mOsm/kg H₂O). Hypoosmotic decrease in the promoter activities of HSP-70 and *TauT* are maintained in *Kif3a* null MEFs. (n ≥ 3 with 3 technical replicates per biological replicate). (g,h) The changes in expression of AR/*Akr1b1* (g) and *SMIT/Slc5a3* (h) under different osmotic conditions are similar between wild-type and *Kif3a* null MEFs (n = 4). (i) Hyperosmotic induction of *TauT/Slc6a6* gene expression observed in wild-type MEFs is absent in *Kif3a* null MEFs. *TauT/Slc6a6* gene expression was downregulated in both wild-type and *Kif3a* null MEFs under hypoosmotic conditions (n = 4). Data are represented as scatter plots (mean ± SEM). ns = not significant. One-way ANOVA or Kruskal-Wallis test with Sidak's or Dunn's multiple comparison test was used based on the distribution of the data to determine statistical significance.

in response to hypoosmotic stimuli^{36,37}. In addition, renal epithelial cells depleted of primary cilia cannot induce TonEBP target genes, *AR/Akr1b3*, and *Bgt1*, under hyperosmotic conditions³⁸. These cell types are frequently exposed to dynamic changes in extracellular osmolarity, similar to those experienced by the NP cells. Importantly, since NP tissue is physiologically hyperosmotic, osmoadaptation is a key survival mechanism for the resident cells. We, therefore, investigated whether primary cilia in NP cells have an osmosensory function and if the activity of TonEBP, a key osmoregulatory transcription factor, is dependent on primary cilia. We show, for the first time, that NP cell primary cilia do not have the osmoregulatory function, although they alter their lengths in response to changes in extracellular osmolarity.

In many cellular pathways that are mediated by primary cilia, the localization of signaling molecules to primary cilia is important for activating downstream signaling pathways^{22,23,57}. Therefore, the changes in the length of primary cilia are thought to be reflective of their functional modification. Indeed, the length of osteocyte primary cilia has been shown to correlate with their sensitivity to the mechanical stimuli⁵⁸. In addition, several known human ciliopathies present with primary cilia of abnormal lengths⁵⁹. In mouse femoral chondrocytes, primary cilia shorten in response to changes in extracellular osmolarity⁴⁰. Likewise, in kidney epithelial cells as well as bone mesenchymal cells, fluid shear-mediated bending of primary cilia results in decreased intracellular cAMP, which in turn causes shortening of primary cilia⁴¹. It is, therefore, possible that the changes in the length of primary cilia in NP cells under different osmotic conditions indicate altered intracellular signaling pathways. However, the pathway mediating the length of cilia in response to osmotic stimuli may be completely independent of the TonEBP-mediated osmotic response.

Our *Ift88* and *Kif3a* knockdown studies showed that the expression of TonEBP and its target genes was unaffected by the absence of primary cilia in NP cells. Under hyperosmotic conditions, TonEBP inhibits excessive sodium influx by increasing the levels of intracellular non-ionic osmolytes including taurine, sodium/myo-inositol, and betaine by inducing transcription of *TauT/Slc6a6*, *Bgt1*, *SMIT/Slc5a3*, and *AR/Akr1b1*^{15–18}. Therefore, disruption of TonEBP activity under hyperosmotic condition can compromise cell survival. The fact that both *Ift88* and *Kif3a* knockdown did not affect the hyperosmotic induction of TonEBP and its target genes suggests that this was the result of the absence of primary cilia rather than a specific effect of the knockdown of either gene. Interestingly, despite the decrease in TonEBP protein levels, the expression of its transcriptional targets, *TauT/Slc6a6*, *SMIT/Slc5a3*, and *AR/Akr1b1* was not significantly affected under hypoosmotic conditions. This may be due to the fact that the osmoregulatory role of TonEBP is more important under hyperosmotic rather than hypoosmotic condition. Nonetheless, neither *Ift88* nor *Kif3a* silencing significantly affected how these genes were regulated in response to hypoosmotic stimuli, strongly suggesting the lack of primary cilia's involvement in this process. In addition, the proximal promoter activities of select TonEBP targets, as well as TonEBP-TAD activity, were mostly unaffected by Sh*Ift88* or Sh*Kif3a* regardless of the osmotic status, suggesting that primary cilia play a minimal role, if any, in osmotic response of NP cells.

Our parallel studies using MEFs further confirmed that primary cilia do not control TonEBP-dependent osmotic response. The use of *Kif3a* null MEFs was advantageous mainly because they completely lack primary cilia, and because MEFs have TonEBP-dependent osmoregulatory pathways similar to that of NP cells^{8,54}. Interestingly, unlike NP cells, *Kif3a* null MEFs were unable to upregulate *TauT* mRNA in response to the hyperosmotic stimulus. This was intriguing since *TauT* promoter activity was not affected by the absence of primary cilia in *Kif3a* null MEFs. This is perhaps due to the high sensitivity of luciferase reporter assay and the fact that it measures the activities of shorter promoter fragments. We cannot exclude the possibility of involvement of either a regulatory sequence outside the analyzed promoter fragment or a primary cilia-associated post-transcriptional regulation. Nonetheless, other osmoreponsive TonEBP targets, *SMIT/Slc5a3* and *AR/Akr1b1*, were upregulated under the hyperosmotic condition in *Kif3a* null MEFs, indicating that the overall cellular osmotic response to hyperosmotic stimuli is intact. Furthermore, we did not observe any obvious increase in *Kif3a* null MEF cell death in response to hyper- or hypo-osmotic stimuli, indicating that these cells were able to maintain their cellular homeostasis without primary cilia. The results of our *Kif3a* null MEFs are particularly interesting because the importance of primary cilia in modulating cell signaling during embryonic development is well-known.

Our data clearly demonstrate that TonEBP-mediated osmoregulation is not controlled by primary cilia in both NP cells and MEFs. The previously reported osmoregulatory and osmosensing function of primary cilia in cholangiocytes³⁶ and kidney epithelial cells³⁸ are therefore likely cell-type specific. Although the *in vitro* culture system is a simplification of the complex *in vivo* extracellular environment, the osmoregulatory role of primary cilia in other cell types was shown to be preserved in *in vitro* settings, suggesting that primary cilia are indispensable in these cell types. On the other hand, the results of our studies indicate that NP cells are capable of initiating TonEBP-mediated cellular osmoreponse without primary cilia. A recent study showed that mouse NP tissue cultured in an *ex vivo* setting decreased their TRPV4 expression in response to cyclic hyperosmolarity⁶⁰. Future studies examining the localization and function of ion channels such as TRPV4, TRM3, or PC-1 and PC-2 that have been associated with cilia osmosensing would be helpful in better understanding how NP cells sense osmotic stimuli.

Materials and Methods

Reagents and plasmids. Lentiviral Sh*Ift88* #1 (TRCN0000178064), Sh*Ift88* #2 (TRCN0000182620), Sh*Kif3a* #1 (TRCN0000339512), Sh*Kif3a* #2 (TRCN0000339514), and control shRNA plasmids were purchased from Sigma. psPAX2 (12260) and pMD2.G (12259) developed by Dr. Didier Trono were obtained from Addgene. Luciferase reporter plasmids were kindly provided by Dr. Takashi Ito, Osaka University (wild type and TonE-mutant TauT-Luc)¹⁵, Dr. H Moo Kwon, University of Maryland (wild type and TonE-mutant HSP70-Luc)²⁰, and Dr. Joan D. Ferraris, NIH (GAL4dbd-548-1531, GAL4dbd, and AR-Luc)^{47,48}. Backbone GAL4dbd contains no TAD but only expresses the GAL4dbd. pFR-Luc reporter (Stratagene) contains the yeast GAL4-binding site, upstream of a minimal promoter driving the firefly luciferase gene.

Cell culture and treatments. All procedures regarding collection of animal tissues was performed as per approved protocols by Institutional Animal Care and Use Committee (IACUC) of the Thomas Jefferson University, in accordance with the IACUC's relevant guidelines and regulations. Rat NP cells were isolated using a method described by Risbud *et al.*⁶¹. After isolation, cells were maintained in Dulbecco's Modified Eagles Medium (DMEM) (Corning, 10-013-CV) with 10% fetal bovine serum (FBS) (Sigma-Aldrich, F6178) supplemented with antibiotics in flask in normoxia (20.9% pO₂). NP cells isolated from two rats were pulled into one T25 flask (P0) and cultured till confluency before passaging (P1). From this point, cells were either plated directly into experiment-specific plates (P2) or passaged. Approximately total of 16 rats were used for the study. Cells from different isolations were used for replication experiments. Cells up to P4 were used for the experiments. Wild type and *Kif3a*^{-/-} MEFs⁵⁴ developed by Dr. Pao-Tien Chuang were kindly provided by Dr. Natalia Riobo, Thomas Jefferson University. For hyper- or hypoosmotic treatment, cells were cultured in DMEM with 10% FBS, containing either no additional osmolytes (330 mOsm/kg H₂O), additional NaCl (60–135 mM to final osmolarity ranging 450–600 mOsm/kg H₂O) or additional dH₂O (to final osmolarity of 200 mOsm/kg H₂O) for 4–24 h. For the hyperosmolarity experiments using NP cells, osmolarity ranging 450 to 550 mOsm/kg H₂O have been used. While the estimated osmolarity of NP tissue *in vivo* ranges from 430 to 496 mOsm/kg H₂O^{3,4}, the above osmolarity range was chosen based on the previous reports on hyperosmotic TonEBP activation in NP cells^{8,11,62}. Similarly, 600 mOsm/kg H₂O was used for MEFs experiments based on the previous report on TonEBP/NFAT5 activity in MEFs, where the peak TonEBP-responsive promoter activity was observed at ~610 mOsm/kg H₂O⁶³.

Real-time RT-PCR analysis. Total RNA was extracted from NP cells using RNeasy mini columns (Qiagen). The purified, DNA-free RNA was converted to cDNA using RNA to cDNA EcoDry™ Premix (Clontech). Template cDNA and gene specific primers were added to SYBR green master mix (Applied Biosystems) and mRNA expression was quantified using the Step One Plus Real-Time PCR System (Applied Biosystems). *Hprt1* was used to normalize gene expression. All the primers used were synthesized by Integrated DNA Technologies, Inc. (Coralville, IA).

Transfections and dual luciferase assay. Cells were plated on 48-well plates (2 × 10⁴ cells/well) one day before transfection. Cells were transfected with 250 ng of wild type TauT, mutant TauT, wild type HSP70, mutant HSP70, or AR reporter plasmid and 250 ng of pRL-TK plasmid. For measuring TonEBP-TAD activity, cells were transfected with 200 ng of pFR-Luc, 200 ng of GAL4dbd-548-1531 (Ton-TAD) and 100 ng of pRL-TK plasmids. For each transfection, plasmids were premixed with the transfection reagent Lipofectamine 2000 (Invitrogen). The treatments were done so that the cells were lysed 48 h after the transfection. Dual-Luciferase™ reporter assay system (Promega) was used for sequential measurements of firefly and Renilla luciferase activities using TECAN Infinite200 Pro microplate reader (TECAN). At least three independent transfections were performed and all analyses were carried out in triplicate.

Immunofluorescence microscopy. NP cells were plated on poly-L-lysine-coated glass coverslips. After treatments, cells were fixed and permeabilized with 4% paraformaldehyde at room temperature for 15 minutes, washed with PBS and then blocked with 5% normal goat serum in PBS with 0.3% Triton X-100 (Sigma Aldrich, T8787) for 1 h at room temperature. Cells on coverslip were then incubated with anti-acetyl- α -tubulin (Lys40) (D20G3) XP[®] antibody (Cell Signaling Technology, #5335, 1:700), or anti-acetylated- α -tubulin (Sigma, T6793, 1:300) and anti- γ -tubulin antibody (Abcam, ab11317, 1:1000) for co-staining in blocking buffer at 4°C overnight, washed with PBS, and then incubated with Alexa Fluor[®]-594 conjugated anti-rabbit, or Alexa Fluor[®]-594 conjugated anti-mouse, or Alexa Fluor[®]-488 conjugated anti-rabbit secondary antibody (Jackson ImmunoResearch Lab, Inc.), at a dilution of 1:700 for 1 h at room temperature in dark. Then the coverslips were washed with PBS and mounted with ProLong[®] Gold Antifade Mountant with DAPI (Thermo Fisher Scientific, P36934). Mounted slides were visualized using a Zeiss AxioImager A2 (Carl Zeiss, Germany), or Zeiss LSM510 confocal microscope (Carl Zeiss, Germany). Three independent experiments were performed for quantitative analysis of cilia length and incidence using ImageJ software (<http://rsb.info.nih.gov/ij/>). Briefly, each cilium was traced using Segmented Lines of the Line Selection Tools, and then the length of the tracing was measured using Measure function. The lengths of at least 150 primary cilia per treatment group were measured.

Protein extraction and western blotting. Following treatment, cells were immediately placed on ice and washed with ice-cold PBS. All the wash buffers and the final cell lysis/re-suspension buffers included 1X complete™ Mini Protease Inhibitor Cocktail (Roche, 11836153001), NaF (5 mM) (Sigma Aldrich, 201154) and Na₃VO₄ (200 μ M) (Sigma Aldrich, S6508). Total cell proteins were resolved by electrophoresis on 8–12% SDS-polyacrylamide gels and transferred by electroblotting to PVDF membranes (EMD Millipore, IPVH00010). Dual-color Lane marker (Bio-Rad, 161-0394) was used as a protein ladder to guide identification of the band sizes. The membranes were blocked with 5% non-fat dry milk in TBST (1% Tween 20 in TBS) and incubated overnight at 4°C in 5% non-fat dry milk in TBST with the antibodies against IFT88 (Proteintech, #13967-1-AP, 1:500), KIF3A (D7G3) (Cell Signaling Technology, #8507, 1:1000), TonEBP/NFAT5 (Novus Biologicals, NB120-3446, 1:1000), or GAPDH (Novus Biologicals, NB300-221, 1:3000). Immunolabeling was detected using the Amersham™ ECL™ Prime Western Blotting Detection Reagent (Thermo Fisher Scientific, 45-002-401). All Western blot experiments were performed at least three independent times.

Lentiviral particle production and viral transduction. HEK 293 T cells (ATCC, CRL-3216) were plated in 10 cm plates (5 × 10⁶ cells/plate) in DMEM with 10% heat-inactivated FBS one day before transfection. Cells were transfected with 9 μ g of Sh*Ctrl*, Sh*Ift88*, or Sh*Kif3a* plasmids along with 6 μ g psPAX2 and 3 μ g pMD2.G using Lipofectamine 2000 (Invitrogen). After 6 h, transfection medium was replaced with DMEM with 10% heat-inactivated FBS and penicillin-streptomycin. Lentiviral medium was harvested at 48 to 60 h

post-transfection, and mixed with 7% PEG 6000 (Sigma Aldrich, 81253) solution and incubated overnight at 4 °C to precipitate virus particles. PEG solution was removed from virus medium before transduction by centrifugation at 1,500 × g for 30 min to pellet virus particles. NP cells were plated in DMEM with 10% heat-inactivated FBS one day before transduction. Cells were transduced with fresh DMEM with 10% heat-inactivated FBS containing viral particles along with 8 µg/ml polybrene (Sigma Aldrich, H9268). After 16 h, the medium was removed and replaced with DMEM with 10% FBS. Cells were harvested for protein extraction 4–5 days after transduction to ensure maximum knockdown efficiency without affecting cell viability. At least three independent experiments were performed.

Statistical analysis. All experiments were performed at least three independent times. Data are presented as the mean ± SE. Differences between multiple groups were assessed by one-way ANOVA or Kruskal-Wallis test depending on the distribution of the data with Sidak's, Holm-Sidak's, or Dunn's multiple comparison test for post-hoc analyses using Prism7 (GraphPad Software). Some data were log-transformed before statistical analyses to account for non-Gaussian distribution. $P < 0.05$ was considered statistically significant.

Received: 25 January 2019; Accepted: 7 October 2019;

Published online: 29 October 2019

References

- Silagi, E. S., Shapiro, I. M. & Risbud, M. V. Glycosaminoglycan synthesis in the nucleus pulposus: Dysregulation and the pathogenesis of disc degeneration. *Matrix Biol.* **71–72**, 368–379 (2018).
- Johnson, Z. I., Shapiro, I. M. & Risbud, M. V. Extracellular osmolarity regulates matrix homeostasis in the intervertebral disc and articular cartilage: Evolving role of TonEBP. *Matrix Biol.* **40**, 10–16 (2014).
- Ishihara, H., Warenso, K., Roberts, S. & Urban, J. P. Proteoglycan synthesis in the intervertebral disk nucleus: the role of extracellular osmolality. *Am. J. Physiol.* **272**, C1499–506 (1997).
- van Dijk, B., Potier, E. & Ito, K. Culturing bovine nucleus pulposus explants by balancing medium osmolarity. *Tissue Eng. Part C. Methods* **17**, 1089–96 (2011).
- Choi, H. *et al.* A novel mouse model of intervertebral disc degeneration shows altered cell fate and matrix homeostasis. *Matrix Biol.* **70**, 102–122 (2018).
- Zhang, Y. *et al.* Early onset of disc degeneration in SM/J mice is associated with changes in ion transport systems and fibrotic events. *Matrix Biol.* **70**, 123–139 (2018).
- Hiyama, A. *et al.* Activation of TonEBP by calcium controls β 1,3-glucuronosyltransferase-I expression, a key regulator of glycosaminoglycan synthesis in cells of the intervertebral disc. *J. Biol. Chem.* **284**, 9824–34 (2009).
- Gajghate, S. *et al.* Osmolarity and intracellular calcium regulate aquaporin2 expression through TonEBP in nucleus pulposus cells of the intervertebral disc. *J. Bone Miner. Res.* **24**, 992–1001 (2009).
- Tsai, T.-T. *et al.* TonEBP/OREBP is a regulator of nucleus pulposus cell function and survival in the intervertebral disc. *J. Biol. Chem.* **281**, 25416–24 (2006).
- Tsai, T.-T. *et al.* MEK/ERK signaling controls osmoregulation of nucleus pulposus cells of the intervertebral disc by transactivation of TonEBP/OREBP. *J. Bone Miner. Res.* **22**, 965–74 (2007).
- Johnson, Z. I., Shapiro, I. M. & Risbud, M. V. RNA Sequencing Reveals a Role of TonEBP Transcription Factor in Regulation of Pro-inflammatory Genes in Response to Hyperosmolarity in Healthy Nucleus Pulposus Cells. *J. Biol. Chem.* **291**, 26686–26697 (2016).
- Choi, H. *et al.* COX-2 expression mediated by calcium-TonEBP signaling axis under hyperosmotic conditions serves osmoprotective function in nucleus pulposus cells. *J. Biol. Chem.* **293**, 8969–8981 (2018).
- Lee, S., Kwon, M. & Woo, S. TonEBP transcriptional activator in the cellular response to increased osmolality. *Pflugers Arch. Eur. J. Physiol.* **444**, 579–585 (2002).
- Libert, S. *et al.* Involvement of TonEBP/NFAT5 in osmoadaptive response of human retinal pigmented epithelial cells to hyperosmolar stress. *Mol. Vis.* **22**, 100–15 (2016).
- Ito, T. *et al.* Expression of taurine transporter is regulated through the TonE (tonicity-responsive element)/TonEBP (TonE-binding protein) pathway and contributes to cytoprotection in HepG2 cells. *Biochem. J.* **382**, 177–82 (2004).
- Miyakawa, H. *et al.* Cis- and trans-acting factors regulating transcription of the BGT1 gene in response to hypertonicity. *Am. J. Physiol.* **274**, F753–61 (1998).
- Rim, J. S. *et al.* Transcription of the sodium/myo-inositol cotransporter gene is regulated by multiple tonicity-responsive enhancers spread over 50 kilobase pairs in the 5'-flanking region. *J. Biol. Chem.* **273**, 20615–21 (1998).
- Ko, B. C., Ruepp, B., Bohren, K. M., Gabbay, K. H. & Chung, S. S. Identification and characterization of multiple osmotic response sequences in the human aldose reductase gene. *J. Biol. Chem.* **272**, 16431–16437 (1997).
- Gogate, S. S., Fujita, N., Skubutyte, R., Shapiro, I. M. & Risbud, M. V. Tonicity enhancer binding protein (TonEBP) and hypoxia-inducible factor (HIF) coordinate heat shock protein 70 (Hsp70) expression in hypoxic nucleus pulposus cells: Role of Hsp70 in HIF-1 α degradation. *J. Bone Miner. Res.* **27**, 1106–1117 (2012).
- Woo, S. K., Lee, S. D., Na, K. Y., Park, W. K. & Kwon, H. M. TonEBP/NFAT5 stimulates transcription of HSP70 in response to hypertonicity. *Mol. Cell. Biol.* **22**, 5753–60 (2002).
- Tessier, S., Madhu, V., Johnson, Z. I., Shapiro, I. M. & Risbud, M. V. NFAT5/TonEBP controls early acquisition of notochord phenotypic markers, collagen composition, and sonic hedgehog signaling during mouse intervertebral disc embryogenesis. *Dev. Biol.*, <https://doi.org/10.1016/j.ydbio.2019.07.004> (2019).
- Rohatgi, R., Milenkovic, L. & Scott, M. P. Patched1 Regulates Hedgehog Signaling at the Primary Cilium. *Science (80-)*. **317**, 372–376 (2007).
- Dorn, K. V., Hughes, C. E. & Rohatgi, R. A Smoothed-Evc2 complex transduces the Hedgehog signal at primary cilia. *Dev. Cell* **23**, 823–35 (2012).
- Huangfu, D. *et al.* Hedgehog signalling in the mouse requires intraflagellar transport proteins. *Nature* **426**, 83–87 (2003).
- Huangfu, D. & Anderson, K. V. Cilia and Hedgehog responsiveness in the mouse. *Proc. Natl. Acad. Sci. USA* **102**, 11325–30 (2005).
- Haycraft, C. J. *et al.* Gli2 and Gli3 localize to cilia and require the intraflagellar transport protein polaris for processing and function. *PLoS Genet.* **1**, e53 (2005).
- Park, T. J., Haigo, S. L. & Wallingford, J. B. Ciliogenesis defects in embryos lacking returned or fuzzy function are associated with failure of planar cell polarity and Hedgehog signaling. *Nat. Genet.* **38**, 303–11 (2006).
- Simons, M. *et al.* Inversin, the gene product mutated in nephronophthisis type II, functions as a molecular switch between Wnt signaling pathways. *Nat. Genet.* **37**, 537–43 (2005).
- McEwen, D. P. *et al.* Hypomorphic CEP290/NPHP6 mutations result in anosmia caused by the selective loss of G proteins in cilia of olfactory sensory neurons. *Proc. Natl. Acad. Sci. USA* **104**, 15917–22 (2007).

30. Bronshtein, A. A. & Minor, A. V. Regeneration of olfactory flagella and restoration of the electroolfactogram following application of triton X-100 to the olfactory mucosa of frogs. *Tsitologia* **19**, 33–9 (1977).
31. Menco, B. P. M. The fine-structural distribution of G-protein receptor kinase 3, beta-arrestin-2, Ca²⁺/calmodulin-dependent protein kinase II and phosphodiesterase PDE1C2, and a Cl⁽⁻⁾-cotransporter in rodent olfactory epithelia. *J. Neurocytol.* **34**, 11–36 (2005).
32. Narita, K., Kawate, T., Kakinuma, N. & Takeda, S. Multiple primary cilia modulate the fluid transcytosis in choroid plexus epithelium. *Traffic* **11**, 287–301 (2010).
33. Elias, R. V., Sezate, S. S., Cao, W. & McGinnis, J. F. Temporal kinetics of the light/dark translocation and compartmentation of arrestin and alpha-transducin in mouse photoreceptor cells. *Mol. Vis.* **10**, 672–81 (2004).
34. Strissel, K. J. *et al.* Recoverin undergoes light-dependent intracellular translocation in rod photoreceptors. *J. Biol. Chem.* **280**, 29250–5 (2005).
35. Tobin, D. M. *et al.* Combinatorial expression of TRPV channel proteins defines their sensory functions and subcellular localization in *C. elegans* neurons. *Neuron* **35**, 307–318 (2002).
36. Gradilone, S. A. *et al.* Cholangiocyte cilia express TRPV4 and detect changes in luminal tonicity inducing bicarbonate secretion. *Proc. Natl. Acad. Sci. USA* **104**, 19138–43 (2007).
37. Phan, M. N. *et al.* Functional characterization of TRPV4 as an osmotically sensitive ion channel in porcine articular chondrocytes. *Arthritis Rheum.* **60**, 3028–3037 (2009).
38. Siroky, B. J. *et al.* Primary cilia regulate the osmotic stress response of renal epithelial cells through TRPM3. *Am. J. Physiol. Physiol.* **312**, F791–F805 (2017).
39. McGlashan, S. R. *et al.* Mechanical loading modulates chondrocyte primary cilia incidence and length. *Cell Biol. Int.* **34**, 441–446 (2010).
40. Rich, D. R. & Clark, A. L. Chondrocyte primary cilia shorten in response to osmotic challenge and are sites for endocytosis. *Osteoarthritis Cartilage* **20**, 923–30 (2012).
41. Besschetnova, T. Y. *et al.* Identification of Signaling Pathways Regulating Primary Cilium Length and Flow-Mediated Adaptation. *Curr. Biol.* **20**, 182–187 (2010).
42. Pazour, G. J. *et al.* Chlamydomonas IFT 88 and Its Mouse Homologue, Polycystic Kidney Disease Gene Tg 737, Are Required for Assembly of Cilia and Flagella. *J. Cell Biol.* **151**, 709–718 (2000).
43. Murcia, N. S. *et al.* The Oak Ridge Polycystic Kidney (orpk) disease gene is required for left-right axis determination. *Development* **127**, 2347–2355 (2000).
44. Yoder, B. K. *et al.* Polaris, a protein disrupted in orpk mutant mice, is required for assembly of renal cilium. *Am. J. Physiol. Renal Physiol.* **282**, F541–52 (2002).
45. Haycraft, C. J. *et al.* Intraflagellar transport is essential for endochondral bone formation. *Development* **134**, 307–16 (2007).
46. Lin, F. *et al.* Kidney-specific inactivation of the KIF3A subunit of kinesin-II inhibits renal ciliogenesis and produces polycystic kidney disease. *Proc. Natl. Acad. Sci. USA* **100**, 5286–91 (2003).
47. Ferraris, J. D. *et al.* Activity of the TonEBP/OREBP transactivation domain varies directly with extracellular NaCl concentration. *Proc. Natl. Acad. Sci. USA* **99**, 739–44 (2002).
48. Ferraris, J. D., Williams, C. K., Martin, B. M., Burg, M. B. & García-Pérez, A. Cloning, genomic organization, and osmotic response of the aldose reductase gene. *Proc. Natl. Acad. Sci. USA* **91**, 10742–6 (1994).
49. Borthakur, A. *et al.* T1ρ magnetic resonance imaging and discography pressure as novel biomarkers for disc degeneration and low back pain. *Spine (Phila. Pa. 1976)* **36**, 2190–6 (2011).
50. Tian, Y. *et al.* Inflammatory Cytokines Associated with Degenerative Disc Disease Control Aggrecanase-1 (ADAMTS-4) Expression in Nucleus Pulposus Cells through MAPK and NF-κB. *Am. J. Pathol.* **182**, 2310–2321 (2013).
51. Wang, J. *et al.* TNF-α and IL-1β promote a disintegrin-like and metalloprotease with thrombospondin type I motif-5-mediated aggrecan degradation through syndecan-4 in intervertebral disc. *J. Biol. Chem.* **286**, 39738–49 (2011).
52. Wilke, H. J., Neef, P., Caimi, M., Hoogland, T. & Claes, L. E. New *in vivo* measurements of pressures in the intervertebral disc in daily life. *Spine (Phila. Pa. 1976)* **24**, 755–62 (1999).
53. Boos, N., Wallin, A., Gbedegbegnon, T., Aebi, M. & Boesch, C. Quantitative MR imaging of lumbar intervertebral disks and vertebral bodies: influence of diurnal water content variations. *Radiology* **188**, 351–4 (1993).
54. Chen, M.-H. *et al.* Cilium-independent regulation of Gli protein function by Sufu in Hedgehog signaling is evolutionarily conserved. *Genes Dev.* **23**, 1910–28 (2009).
55. Donnelly, E., Williams, R. & Farnum, C. The primary cilium of connective tissue cells: imaging by multiphoton microscopy. *Anat. Rec. (Hoboken)* **291**, 1062–73 (2008).
56. Zheng, L. *et al.* Ciliary parathyroid hormone signaling activates transforming growth factor-β to maintain intervertebral disc homeostasis during aging. *Bone Res.* **6**, 21 (2018).
57. Piperi, C. & Basdra, E. K. Polycystins and mechanotransduction: From physiology to disease. *World J. Exp. Med.* **5**, 200 (2015).
58. Spasic, M. & Jacobs, C. R. Lengthening primary cilia enhances cellular mechanosensitivity. *Eur. Cell. Mater.* **33**, 158–168 (2017).
59. McIntyre, J. C., Williams, C. L. & Martens, J. R. Smelling the roses and seeing the light: Gene therapy for ciliopathies. *Trends Biotechnol.* **31**, 355–363 (2013).
60. Palacio-Mancheno, P. E., Evashwick-Rogler, T. W., Laudier, D. M., Purmessur, D. & Iatridis, J. C. Hyperosmolarity induces notochordal cell differentiation with aquaporin3 upregulation and reduced N-cadherin expression. *J. Orthop. Res.* **36**, 788–798 (2017).
61. Risbud, M. V. *et al.* Nucleus pulposus cells express HIF-1 alpha under normoxic culture conditions: a metabolic adaptation to the intervertebral disc microenvironment. *J. Cell. Biochem.* **98**, 152–9 (2006).
62. Liu, C. *et al.* Lack of evidence for involvement of TonEBP and hyperosmotic stimulus in induction of autophagy in the nucleus pulposus. *Sci. Rep.* **7**, 4543 (2017).
63. Go, W. Y., Liu, X., Roti, M. A., Liu, F. & Ho, S. N. NFAT5/TonEBP mutant mice define osmotic stress as a critical feature of the lymphoid microenvironment. *Proc. Natl. Acad. Sci. USA* **101**, 10673–8 (2004).

Acknowledgements

This study is supported by grants from the National Institutes of Health R01 AR055655, R01 AR064733 and R01AR074813.

Author contributions

H.C. conducted the experiments, analyzed data, and wrote the manuscript. V.M. conducted the experiments and analyzed data. M.V.R. and I.M.S. conceived the study, secured funding, and contributed to interpretation of the results. All authors reviewed the results, contributed to editing the manuscript, and approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-019-51939-7>.

Correspondence and requests for materials should be addressed to M.V.R.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019