

## Why do we find so many meteorites on the Nansen blue ice field and where else could we look?

Harry Zekollari<sup>1,2</sup>, Steven Goderis<sup>3</sup>, Vinciane Debaille<sup>4</sup>, Matthias van Ginneken<sup>5</sup>, Jérôme Gattacceca<sup>6</sup>, A J Timothy Jull<sup>7</sup>, Jan Lenaerts<sup>8</sup>, Akira Yamaguchi<sup>9,10</sup>, Naoya Imae<sup>9</sup>, Philippe Huybrechts<sup>3</sup>, Veronica Tollenaar<sup>1</sup>, Stef Lhermitte<sup>1</sup> and Philippe Claeys<sup>3</sup>

<sup>1</sup>*Department of Geoscience and Remote Sensing, Delft University of Technology, Delft, Netherlands*

<sup>2</sup>*Laboratoire de Glaciologie, Université libre de Bruxelles, Brussels, Belgium*

<sup>3</sup>*Earth System Science, Vrije Universiteit Brussel, Brussels, Belgium*

<sup>4</sup>*Laboratoire G-Time, Université Libre de Bruxelles, Brussels, Belgium*

<sup>5</sup>*Centre for astrophysics and planetary science, University of Kent, Kent, United Kingdom*

<sup>6</sup>*Aix Marseille Univ, CNRS, IRD, Coll France, CEREGE, Aix-en-Provence, France*

<sup>7</sup>*Department of Geosciences, University of Arizona, Tucson, United States of America*

<sup>8</sup>*Department of Atmospheric and Oceanic Sciences, University of Colorado, Boulder, United States of America*

<sup>9</sup>*National Institute of Polar Research, Tokyo, Japan*

<sup>10</sup>*The Graduate University for Advanced Studies (SOKENDAI), Tokyo, Japan*

Antarctic blue ice zones are the most productive locations for meteorite recovery on Earth. However, the mechanisms behind these meteorite traps remain a topic of ongoing debate. Here, we propose an interdisciplinary approach combining surface blue ice isotopic analysis, terrestrial age dating of meteorites, remote sensing observations and ice flow and surface mass balance modelling to improve our understanding of the Nansen blue ice field meteorite trap (Dronning Maud Land, East Antarctica). This ice field (2600-3100 m above sea level) was first explored by the 29th Japanese Antarctic Research Expedition (JARE-29) (1987-1989), during which more than 1582 meteorites were retrieved (Naraoka et al., 1990; Yanai et al., 1993). The northern part of the ice field (Nansen A) was subsequently revisited during the austral summer of 2010-11 (Belgian Antarctic Research Expedition, BELARE 2010-2011), resulting in more than 200 newly collected meteorites (Goderis et al., 2011). Two years later, during the austral summer of 2012-13, a meteorite search expedition (JARE-54, BELARE 2012-2013) was organized on the southwestern ice field (Nansen B) and the northern part of the eastern ice field (Debaille et al., 2013; Imae et al., 2013, 2015). During the 2012-13 expedition, more than 400 meteorites were collected, together with 185 surface blue ice samples (Imae et al., 2015; Zekollari et al., 2019).

The surface blue ice samples show one of the largest observed spatial patterns in oxygen isotopic variation to date. Relying on meteorites for which the terrestrial ages are determined using <sup>14</sup>C and <sup>36</sup>Cl, this surface ice is interpreted to date from the Last Interglacial up to the present-day. By combining state-of-the-art satellite derived surface velocities, surface mass balance modelling and ice flow modelling, we estimate that about 75% to 85% of the meteorites found on the ice field were supplied by ice flow after entering the ice sheet in an accumulation area of a few hundred square kilometres located south (upstream) of the ice field. Less than 0.4 new meteorites per year are supplied to the ice field through ice flow, suggesting that the hundreds of meteorites found 25 years after the first visit to this ice field mostly represent meteorites that were previously not found, rather than newly supplied meteorites. By combining these findings, the infall rate of meteorites from space is estimated, which is in line with values from the literature, but situated at the higher end of the range. This study introduces new concepts that can be used to better characterize, understand and identify other meteorite traps. In this presentation, we show some first results of our attempt to identify yet unexplored meteorite traps by combining remote sensing products and various datasets in a big data approach.

### References

- Debaille, V., Imae, N., Yamaguchi, A., Goderis, S., Mikouchi, T., Debouge, W., Hublet, G., Van Roosbroeck, N., Zekollari, H., Kojima, H. and Claeys, P.: The 2012-2013 joint field campaign for collecting meteorites in Antarctica: an efficient collaboration between Japan and Belgium, *Antarct. Meteorites XXXVI*, 11–12, 2013.
- Goderis, S., Kaiden, H., Debaille, V., Kojima, H. and Claeys, P.: Belgian-Japanese search for Antarctic meteorites during the 2010-2011 field season, in *Antarctic Meteorites XXXIV*, p. 12, Tokyo., 2011.
- Imae, N., Akada, Y., Berclaz, C., Claeys, P., Debaille, V., Goderis, S., Hublet, G., Kojima, H., Mikouchi, T., Van Roosbroeck, N., Debouge, W., Yamaguchi, A. and Zekollari, H.: The search for Antarctic meteorites in the Nansen Ice Field by the joint team of Japan and Belgium, *Proc. Japan Geosci. Union Meet.*, PPS24-P05, 2013.

- Imae, N., Debaille, V., Akada, Y., Debouge, W., Goderis, S., Hublet, G., Mikouchi, T., Van Roosbroek, N., Yamaguchi, A., Zekollari, H., Claeys, P. and Kojima, H.: Report of the JARE-54 and BELARE 2012-2013 joint expedition to collect meteorites on the Nansen Ice Field, Antarctica, *Antarct. Rec.*, 59(1), 38–72, 2015.
- Naraoka, H., Yanai, K. and Fujita, S.: Report on Antarctic meteorites search around the Sør Rondane Mountains, *Antarct. Rec.*, 34, 216–224, 1990.
- Yanai, K., Kojima, H. and Naraoka, H.: The Asuka-87 and Asuka-88 collections of Antarctic meteorites: search, discoveries, initial processing, and preliminary identification and classification, *Proc. NIPR Symp. Antarct. Meteorites*, 6, 137–147, 1993.
- Zekollari, H., Goderis, S., van Ginneken, M., Gattacceca, J., ASTER Team, Jull, T. A. J., Yamaguchi, A., Huybrechts, P., Debaille, V. and Claeys, P.: Unravelling the high-altitude Nansen blue ice field meteorite trap (East Antarctica) and implications for regional palaeo-conditions, *Geochim. Cosmochim. Acta*, 248, 289–310, doi:10.1016/j.gca.2018.12.035, 2019.