

(2)

Sclerosing Hepatic Carcinoma

Young Nyun Park, Chanil Park

Department of Pathology, Yonsei University College of Medicine

BRIEF HISTORY

41 가 가
 . 5
 , AST/
 ALT 14/42 IU/L, alkaline phosphatase 91 IU/L,
 total bilirubin 1.8 mg/dL, -fetoprotein 183 ng/
 mL, HBsAg (+), anti-HBs (+) .

GROSS FINDING

8×4×4 cm, 60 gm ,
 10cm
 가 .
 ,
 0.5 cm
 .

MICROSCOPIC FEATURE

. 1)
 ,
 가
 , 2)
 , 3)

2-3 ,

Figure. 1 Gross finding of sclerosing hepatic carcinoma. The cut section showing a well defined, whitish brown, fibrotic mass of a spreading infiltrative growth pattern and several satellite small nodules. The surrounding liver revealing a relatively normal appearance without cirrhotic change.

가

가

AE1

가 cytokeratin 19,

가

• high
 molecular cytokeratin (AE3) alpha-fetoprotein
 low molecular cytokeratin
 (AE1), cytokeratin 19, carcinoembry- onic antigen
 (CEA) epithelial membrane antigen (EMA)
 . Cytokeratin AE1 19
 , CEA EMA

1. ;
 1994;28:636- 642
2. Omata M, Peters RL, Tatter D. Sclerosing
 hepatic carcinoma: relationship to hypercal-
 cemia. Liver 1981;1:33-49