

(Table 1).

1. 약물치료의 위험

가 가

1) , 2) (behavioral teratogenesis)

, 3)

(teratogenicity)

12

5) 가 8)

(teratogen) 가

2 4% 가

12% 6,7) 2

(

) 가

가

가 가

(perinatal syndrome) 2. 정신질환과 연관된 위험

20

가

Table 1. U.S. food and drug administration (FDA) use-in-pregnancy ratings

Category	Interpretation
A	Controlled studies show no risk : Adequate, well controlled studies in pregnant women have failed to demonstrate risk to the fetus.
B	No evidence of risk in humans : Either animal findings show risk, but human findings do not ; or, if no adequate human studies have been done, animal findings are negative.
C	Risk cannot be ruled out : Human studies are lacking, and animal studies are either positive for fetal risk or lacking as well. However, potential benefits may justify the potential risk.
D	Positive evidence of risk : Investigational or postmarketing data show risk to the fetus. Nevertheless, potential benefits may outweigh risks.
X	Contraindicated in pregnancy : Studies in animals or humans, or investigational or postmarketing reports, have shown fetal risk that clearly outweighs any possible benefit to the patient.

Source. Physicians' Desk Reference, 50th Edition. Montvale, NJ, Medical Economics, 1996.

Table 2. Risk factor of psychotropic drugs

Generic name	Risk category ^a	American academy of pediatrics rating ^b	Generic name	Risk category ^a	American academy of pediatrics rating ^b
Antipsychotics			Other antidepressants		
Chlorpromazine	C	Unknown, but of concern	Bupropion	B _m	N/A
Clozapine	B _m	N/A	Mirtazapine	C	N/A
Fluphenazine	C	N/A	Nefazodone	C	N/A
Haloperidol	C _m	Unknown, but of concern	Trazodone	C _m	Unknown, but of concern
Loxapine	C	N/A	Venlafaxine ^c	C _m	N/A
Mesoridazine	C	Unknown, but of concern	Benzodiazepines		
Molindone	C	N/A	Alprazolam	D _m	N/A
Olanzapine	C	N/A	Chlordiazepoxide	D	N/A
Perphenazine	C	Unknown, but of concern	Clonazepam	C	N/A
Pimozide ^c	C	N/A	Clorazepate	D	N/A
Risperidone ^c	C	N/A	Diazepam	D	Unknown, but of concern
Thioridazine	C	N/A	Halazepam ^c	N/A	N/A
Thiothixene	C	N/A	Lorazepam	D _m	Unknown, but of concern
Trifluoperazine	C	N/A	Oxazepam	D	N/A
Medications for antipsychotic side effects			Prazepam ^c	D	Unknown, but of concern
Amantadine	C _m	N/A	Benzodiazepines for insomnia		
Benzotropine	C	N/A	Estazolam ^c	X	N/A
Diphenhydramine	C	N/A	Flurazepam	X _m	N/A
Propranolol	C _m	Compatible	Quazepam ^c	X	Unknown, but of concern
Trihexyphenidyl	C	N/A	Temazepam	X _m	Unknown, but of concern
Heterocyclic antidepressants			Triazolam	X _m	Unknown, but of concern
Anitriptyline	D	Unknown, but of concern	Nonbenzodiazepine anxiolytics and hypnotics		
Amoxapine	C _m	Unknown, but of concern	Buspirone ^c	B	N/A
Clonipramine	C _m	Unknown, but of concern	Chloral hydrate	C _m	Compatible
Desipramine	C	Unknown, but of concern	Zolpidem ^c	C	N/A
Doxepin	C	Unknown, but of concern	Mood-stabilizing medications		
Imipramine	D	Unknown, but of concern	Carbamazepine	C _m	Compatible
Nortriptyline	D	N/A	Clonazepam	C	N/A
Maprotyline	B _m	N/A	Gabapentin	C	N/A
Monoamine oxidase inhibitors			Lamotrigine	C	N/A
Isocarboxazid	C	N/A	Valproic acid	D	Compatible Contraindicated le
Phenelzine	C	N/A	Lithium	D	
Tranylcypromine	C	N/A			
Selective serotonin reuptake inhibitors					
Fluoxetine	C	Unknown, but of concern			
Fluvoxamine ^c	C	Unknown, but of concern			
Paroxetine ^c	C	N/A			
Sertraline	C	N/A			

N/A = not applicable

^aRisk category adapted from Briggs 1994 ; " m " subscript is for data taken from the manufacturer's package insert

^bAmerican Academy of Pediatrics 1994

^cNot listed in Briggs et al. 1994. Risk category taken from Physicians' Desk Reference 1992, 1993, 1994, 1996

임신 중 개별 항정신약물 사용의 영향

10 가 phenothiazine 가 haloperidol cloz - 가 apine 가 haloperidol 가

가 가 가

Table 2

1. 항정신병약물

haloperidol 10 가

⁹⁾ 가 가

가 Rumeau - Rouquette ¹⁰⁾ 가 가

315 3.5% 가 phenothiazines chl - IQ 가 5

orpromazine (metaanalysis) 가

가 ¹¹⁾ Benztropine trihexyphenidyl 가

Slone ¹²⁾ phen - othiazine 1309 50,282 가 diphenhydramine 가

가 prochlorperazine 가

Milkovich Van den Berg ¹³⁾ 6 2. 항우울제 10%

Amphetamine 가 , 가
 amphetamine 가 . floppy baby 가 .
 18,19)

3. 기분 안정제
 1) 리튬 가 , 가 . 70 . Schou²⁶⁾ 5
 가 20,21) , 60 가 ,
 Ebstein's anomaly 가 36 2) 항경련제 가 .
 가 36 . 22) 가 .
 23,24)

Ebstein's anomaly Carbamazepine 1% Valproic acid
 Ebstein's anomaly 1 3% 가 가
 가 가 가 가
 25) Ebste - (orofacial cleft)
 in's anomaly 10 (minor malformation)(, rotated
 20 ears, flat nasal bridge, fingernail hypoplasia)
 20,000 1 ,
 2000 1(0.05%) 1000 1(0.1%)
 Ebstein's anomaly

가 . , .

가 . 가 .

. Desipramine

nortriptyline

fluoxetine

가 . SSRI 가 .

ECT가

Sertraline paroxetine

가 . , .

가 sert -

2. 기분장애

1) 주요 우울증

raline paroxetine 가 가

(SSRI)

SSRI

가 , , ,

28,29) 30)

가

fluoxetine

가 .

MAOIs

가

venlafaxine, nefazodone bupropion,

가

가

가

가

가

risk category가 (metabolites)

FDA () 가

가

3) 용량의 결정 요 약

가 가

가 가

가 가

(가) 가 phenothiazine, lithium, benzodiazepines, valproic acid, carbamazepine

4) 필요시 소아과 의사와 협의 가

가 가 TCAs SSRIs

가 가

모유 수유

가 가

가

가 ,
가

중심 단어 :

참고 문헌

- 1) Kasilo O, Romero M, Bonati M, Tognoni G(1988) : *Information on drug use in pregnancy from the Viewpoint Regional Drug Information Center. Eur J Clin Pharmacol* 35:447-53
- 2) Stowe ZA, Strader JR, Nemeroff CB(1998) : *Psychopharmacology during pregnancy and lactation. In: Textbook of Psychopharmacology. Vol 2, Ed by Schatzberg AF and Nemeroff CB, Washington DC, American Psychiatric Press, pp 979-96*
- 3) Rayburn WF, Andresen BD(1982) : *Principles of perinatal pharmacology. In: Drug Therapy in Obstetrics and Gynecology. Ed by Rayburn WF, Zuspan F, Norwalk CT, Appleton-Century-Crofts, pp1-8*
- 4) Welch R, Findlay J(1981) : *Excretion of drugs in human breast milk. Drug Metab Rev* 12:261-77
- 5) Dicke JM(1989) : *Teratology: Principles and practice. Med Clin North Am* 73:567-81
- 6) Kiely M(1991) : *Reproductive and Perinatal Epidemiology. Boca Raton, FL, CRC Press, pp31-9*
- 7) Riccardi VM(1977) : *The Genetic Approach to Human Disease. New York, Oxford University Press, pp71-98*
- 8) Coyle I, Wayner M, Singer G(1976) : *Behavioral teratogenesis: A critical evaluation. Pharmacol Biochem Behav* 4:191 - 200
- 9) Kopelman AE, McCullar FW, Heggeness L(1975) : *Limb malformations following maternal use of haloperidol. JAMA* 231:62-4
- 10) Rumeau-Rougutte C, Goujard J, Huel G(1977) : *Possible teratogenic effect of phenothiazines in human beings. Teratology* 15:57-64
- 11) Altshuler LL, Cohen LS, Szuba MP, Burt VK, Gitlin M, Mintz J(1996) : *Pharmacological management of psychiatric illness in pregnancy: Dilemmas and guidelines. Am J Psychiatry* 153:592-606
- 12) Slone D, Siskind V, Heinonen OP, Monson RR, Kaufman DW, Shapiro S(1977) : *Antenatal exposure to the phenothiazines in relation to congenital malformations, perinatal mortality rate, birth weight, and intelligence quotient score. Am Obstet Gynecol* 128:486-8
- 13) Milkovich L, Van den Berg BJ(1976) : *An evaluation of the teratogenicity of certain antiemetic drugs. Am J Obstet Gynecol* 125:244-8
- 14) McBride WG(1972) : *Limb deformities associated with ininodi-benzyl hydrochloride [letter]. Med J Aust* 1:492
- 15) Pastuszak A, Schick-Boschetto B, Zuber C, Feldkamp M, Pinelli M, Sihn S, Donnenfeld A, McCormack M, Leen-Mitchell M, Woodland C, Gardner A, Hom M, Koren G(1993) : *Pregnancy outcome following first-trimester exposure to fluoxetine (Prozac). JAMA* 269:2246-8
- 16) Cooper GL(1989) : *The safety of fluoxetine. An update. Br J Psychiatry* 153:77-86
- 17) Heinonen O, Sloan D, Shapiro S(1977) : *Birth Defects and Drugs in Pregnancy. Littleton, Mass, Publishing Services Group*
- 18) Nora J, Vargo T, Nora A(1970) : *A possible environmental trigger in cardiovascular malformations. Lancet* 1:1290-1
- 19) Milkovich L, van den Berg BJ(1977) : *Effects of antenatal exposure to anorectic drugs. Am J Obstet Gynecol* 121:637-42
- 20) Weinstein MR(1976) : *The The International Register of Lithium Babies. Drug Inf J* 50:81-6
- 21) Schou M, Goldfield MD, Weinstein MR, Villaneuve A(1973) : *Lithium and pregnancy, I: Report from the register of lithium babies. Br Med J* 2:135-6
- 22) Kirklin JW, Barratt-Boyes BG(1986) : *Ebstein's malformation. In: Cardiac surgery, Ed by Kirklin JW, Barratt-Boyes BG, New York, John Wiley*
- 23) Kallen B, Tandberg A(1983) : *Lithium and pregnancy: A cohort study on manic-depressive women. Acta Psychiatr Scand* 68:134-9
- 24) Jacobson SJ, Jones K, Johnson K, Ceolin L, Kaur P, Sahn D, Donnenfeld AE, Rieder M, Santelli R, Smythe J, Pastuszak A, Einarson T, Koren G(1992) : *Prospective multicentre study of pregnancy outcome after lithium exposure during first trimester. Lancet* 339:530-3
- 25) Cohen LS, Friedman JM, Jefferson JW, Johnson EM, Weiner ML(1994) : *A reevaluation of risk of in utero exposure to lithium. JAMA* 271:146-50
- 26) Schou M(1976) : *What happened later to the lithium babies: A follow-up study of children born without malformations. Acta Psychiatr Scand* 54:193-7
- 27) Sclonik D, Nulman I, Rovet J, Gladstone D, Czuchta D, Garder HA, Gladstone R, Ashby P, Weksberg R, Einarson T, Koren G(1994) : *Neurodevelopment of children exposed in utero to phenytoin and carbamazepine monotherapy. JAMA* 271:767-70
- 28) Zuckerman B, Bauchner H, Parker S, Cabral H(1990) : *Maternal depressive symptoms during pregnancy, and newborn irritability. J Dev Behav Pediatr* 11:190-4
- 29) Cohn JF, Tronick E(1989) : *Specificity of infants' response to mothers's affective behavior. J Am Acad Child Adolesc Psychiatry* 28:242-8
- 30) Stoukides JA, Stoukides CA(1991) : *Extrapyramidal symptoms upon discontinuation of fluoxetine[letter]. Am J Psychiatry* 148:1263
- 31) Cohen LS, Heller VL, Rosenbaum JF(1989) : *Treatment guidelines for psychotropic drug use in pregnancy. Psychosomatics* 30:25-33
- 32) Omtzigt JGC, Los FJ, Grobbee DE, Pijpers L, Jahoda MGJ, Brandenburg H, Stewart PA, Gaillard HLJ, Sachs ES, Wladimiroff JW, Lindhout D(1992) : *The risk of spina bifida aperta after firsttrimester exposure to valproate in a prenatal cohort. Neurology* 42:119-25
- 33) MRC Vitamine Study Research Group(1991) : *Prevention of neural-tube defects: Results of the medical research council vitamin study. Lancet* 338:131-7
- 34) Ingram IM(1961) : *Obsessional illness in mental hospital patients. J Ment Sci* 107:382-402