

Stephen F. Austin State University
SFA ScholarWorks

Faculty Publications

Mathematics and Statistics

2014

Winter Movements of Louisiana Pine Snakes (*Pituophis ruthveni*) in Texas and Louisiana

Josh B. Pierce
USDA Forest Service, Southern Research Station

D. Craig Rudolph
USDA Forest Service, Southern Research Station

Shirley J. Burgdorf
US Fish and Wildlife Service

Richard R. Schaefer
USDA Forest Service, Southern Research Station

Richard N. Conner
USDA Forest Service, Southern Research Station

See next page for additional authors

Follow this and additional works at: https://scholarworks.sfasu.edu/mathandstats_facultypubs



Part of the [Applied Mathematics Commons](#), and the [Forest Biology Commons](#)

[Tell us](#) how this article helped you.

Repository Citation

Pierce, Josh B.; Rudolph, D. Craig; Burgdorf, Shirley J.; Schaefer, Richard R.; Conner, Richard N.; Himes, John G.; Duran, C. Mike; Hardy, Laurence M.; and Fleet, Robert R., "Winter Movements of Louisiana Pine Snakes (*Pituophis ruthveni*) in Texas and Louisiana" (2014). *Faculty Publications*. 16.
https://scholarworks.sfasu.edu/mathandstats_facultypubs/16

This Article is brought to you for free and open access by the Mathematics and Statistics at SFA ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SFA ScholarWorks. For more information, please contact cdsscholarworks@sfasu.edu.

Authors

Josh B. Pierce, D. Craig Rudolph, Shirley J. Burgdorf, Richard R. Schaefer, Richard N. Conner, John G. Himes, C. Mike Duran, Laurence M. Hardy, and Robert R. Fleet

Winter Movements of Louisiana Pine Snakes (*Pituophis ruthveni*) in Texas and Louisiana

Josh B. Pierce^{1,*}, D. Craig Rudolph¹, Shirley J. Burgdorf², Richard R. Schaefer¹,
Richard N. Conner¹, John G. Himes³, C. Mike Duran⁴, Laurence M. Hardy⁵,
and Robert R. Fleet⁶

Abstract - Despite concerns that the Louisiana Pine Snake (*Pituophis ruthveni*) has been extirpated from large portions of its historic range, only a limited number of studies on their movement patterns have been published. Winter movement patterns are of particular interest since it has been hypothesized that impacts of management practices would be reduced during the winter. Using radiotelemetry, we determined winter movement patterns of Louisiana Pine Snakes (11 males, 8 females) in 5 study areas (2 in Louisiana and 3 in Texas). Movements during winter (November–February) were greatly curtailed compared to the remainder of the year; however, snakes occasionally undertook substantial movements. Relocations were typically within the snake’s previous active-season home range, and movements were more frequent in the early portion of winter. All hibernation sites were within Baird’s Pocket Gopher (*Geomys breviceps*) burrow systems at depths ranging from 13–25 cm. Louisiana Pine Snakes did not use communal hibernacula, nor did individual snakes return to previously used sites in successive years.

Introduction

Snakes of the genus *Pituophis* (Holbrook) are large, terrestrial constrictors that feed primarily on mammals in open habitats (Rodriguez-Robles 2002, Sweet and Parker 1991), and are widely distributed in North America (Sweet and Parker 1991). *Pituophis ruthveni* Stull (Louisiana Pine Snake) is a narrowly distributed species found in eastern Texas and west-central Louisiana (Reichling 1995, Rudolph et al. 2006, Sweet and Parker 1991). The species is thought to have been extirpated from large portions of its historical range, and extant populations are currently known from a limited number of small and fragmented localities (Reichling 1995, Rudolph et al. 2006). The Louisiana Pine Snake is listed as threatened by the Texas Parks and Wildlife Department and is classified by the US Fish and Wildlife Service (USFWS) as a candidate species for listing as threatened or endangered under the Endangered Species Act (USFWS 2012).

Few studies on the ecology of the Louisiana Pine Snake have been published. Recent ecological research has demonstrated that the Louisiana Pine Snake is a

¹USDA Forest Service, Southern Research Station, 506 Hayter Street, Nacogdoches, TX 75965. ²US Fish and Wildlife Service, 510 Desmond Drive, Suite 102, Lacy, WA 98503. ³Florida Fish and Wildlife Conservation Commission, 3911 Highway 2321, Panama City, FL 32409. ⁴The Nature Conservancy, 200 E. Grayson Street #202, San Antonio, TX 78215. ⁵Museum of Life Sciences, Louisiana State University in Shreveport, Shreveport, LA 71115. ⁶Department of Mathematics and Statistics, Stephen F. Austin State University, Nacogdoches, TX 75962. *Corresponding author - jbpierce@fs.fed.us.

diurnal species (Ealy et al. 2004) primarily associated with open pine forests on sandy soils with abundant herbaceous vegetation (Himes et al. 2006a, Rudolph and Burgdorf 1997). Within the snake's historical range, this habitat is maintained by frequent fire (Conner et al. 2001, Frost 1993). The Louisiana Pine Snake preys primarily on *Geomys breviceps* Baird (Baird's Pocket Gopher; Rudolph et al. 2002, 2012) and makes extensive use of pocket gopher burrow systems for shelter, hibernation, and to escape from fires (Rudolph and Burgdorf 1997; Rudolph et al. 1998, 2007). The reduction in pocket gopher populations, resulting from the loss of herbaceous vegetation density and diversity due to fire suppression, is thought to be a factor in the decline of the Louisiana Pine Snake (Rudolph and Burgdorf 1997, Rudolph et al. 2006).

Louisiana Pine Snakes are most active during March–May and September–November, possibly because above-ground temperatures are optimum for movement (Himes et al. 2006a, b). For this reason, land managers have proposed to conduct forestry activities, such as timber harvest, during the winter months when Louisiana Pine Snakes are thought to be dormant (Himes et al. 2006a). These management practices should mitigate incidental take of Louisiana Pine Snakes; however, knowledge of winter activity patterns of Louisiana Pine Snakes is restricted to populations in Bienville Parish, LA, where Himes et al. (2006a) found that snakes remained underground and inactive during the winter months (December–February).

Our objective was to augment our knowledge of winter movement patterns of Louisiana Pine Snakes throughout their currently known range. Better understanding of winter movement patterns should aid land managers in determining the time of year when Louisiana Pine Snakes are least likely to be adversely affected by forestry practices.

Materials and Methods

We captured snakes (11 males, 8 females) in 5 study areas: private land in Bienville Parish, LA (Himes et al. 2006a); Ft. Polk Military Reservation in Louisiana; privately owned Scrappin' Valley in Newton County, TX; Sabine National Forest in Sabine County, TX (Ealy et al. 2004); and Angelina National Forest in Angelina and Jasper counties, TX. All sites had soils with a high sand content, a diverse herbaceous flora dominated by *Schizacharium scoparium* (Michx.) Nash (Little Bluestem) and *Pteridium aquilinum* (L.) Kuhn (Bracken Fern), and an overstory dominated by pines, primarily *Pinus palustris* P. Mill. (Longleaf Pine). All sites have gently rolling topography intersected by intermittent and small permanent streams.

We captured the snakes by hand ($n = 6$) or with drift fence and funnel trap arrays ($n = 13$; Burgdorf et al. 2005) between 1993 and 1997. Treatment of captured individuals was as follows: for each snake we determined weight to the nearest gram, snout–vent length (SVL) to the nearest centimeter, and sex by probing for hemipenes (Schaefer 1934). We then implanted all snakes in Bienville Parish with Holohil SI-2T transmitters (44 x 10 mm, 29-cm whip antennae, weight 12 g; Holohil Systems Ltd., Carp, ON, Canada) intraperitoneally following Reinart and Cundall (1982), except for 1 juvenile snake, in which we similarly implanted a

2.5-g transmitter (constructed by P. Blackburn, Stephen F. Austin State University, Nacogdoches, TX). We implanted the remaining snakes subcutaneously following the general procedures of Weatherhead and Anderka (1984) with Holohil SI-2T transmitters. We anesthetized snakes using ketamine hydrochloride or halothane. Transmitters weighed <2.5% of snake body mass. Transmitter life span was approximately 18–24 months, and maximum transmission range was approximately 1200 m. After surgery, we kept snakes in the laboratory and monitored them for at least 5 days, then released them at their capture location. We replaced transmitters as necessary, generally every 18 months.

We relocated snakes using either an H antenna or a 3-element Yagi antenna and a R2100 receiver (Advanced Telemetry Systems, Inc., Isanti, MN). Generally, we tracked snakes less often and more irregularly during winter (November–February; mean = 2.7 relocations/snake/month) than during active season months (March–October; mean = 6.5 relocations/snake/month).

We used a Trimble GPS Professional™ unit to record relocation site coordinates (any location of a telemetered snake after surgery and subsequent release) and corrected the values using post-processed differential correction. Snakes were not disturbed during this portion of data collection. We considered snake locations ≥ 5 m from a previous location as movement, and collected new coordinates (Himes et al. 2006a).

We excavated seven of 19 Louisiana Pine Snakes from gopher burrows during winter to determine the distance (cm) of the snake from the presumed entrance and the depth (cm) of the snake within the burrow system. For a complete description of this process, see Rudolph et al. 2007.

We calculated monthly movement frequencies by dividing the total number of movements by the total number of relocations for all snakes, across all years. We divided monthly frequencies into seasons (active season: $n = 8$ months; winter: $n = 4$ months). We compared seasonal movement frequencies using a 2-sample *t*-test at an alpha level of 0.05.

We chose each snake's winter dormancy location based on the amount of time the snake spent in its winter locations, attempting to choose the single point that best represented the site of winter dormancy. We calculated the distance from the previous year's winter dormancy location using ArcGIS version 9.3 (Environmental Systems Research Institute, Redlands, CA).

To assess whether snakes were spatially distributed differently between winter and the active season, we measured the distances of winter dormancy locations from the edge of the active-season (March–October) 100% minimum convex polygon (MCP) home ranges (J.B. Pierce, unpubl. data). We calculated home ranges with Home Range Tools for ArcGIS (Rodgers et al. 2007) in ArcGIS version 9.3.

Results

During 1993–1998, we tracked 19 snakes for at least one consecutive active season and winter season. The number of winters tracked (range = 1–4) varied across

individuals, resulting in a total of 37 snake winters (Table 1). We located snakes during winter on a total of 283 occasions, of which only 89 were unique locations because individuals often remained in the same location for one or more subsequent relocations and occasionally returned to previously used locations.

All winter locations were within Baird's Pocket Gopher burrow systems. Rudolph and colleagues (2007) presented data on Louisiana Pine Snake hibernacula using the same snakes as we followed in this study. They excavated 7 snakes from burrows at depths ranging from 13–25 cm (mean = 19.0 ± 4.9 cm; Rudolph et al. 2007). These snakes were <1 m from the presumed point of entrance into the burrow system (Rudolph et al. 2007). Winter refuge placement of all snakes was similar to the positions occupied by snakes at other seasons when using pocket gopher burrows for foraging and refuge (Rudolph et al. 2007).

Because snakes were tracked less often during winter than the other seasons, we might have underestimated the amount of winter movement if snakes moved undetected but returned to previously used sites in the time between our tracking efforts. Snake movement frequencies (mean = 24.0%) were significantly lower during winter when compared to the other seasons (mean = 66.2%; $t = 6.93$, $df = 10$, $P = 0.002$; Fig. 1). Movement frequencies began to decline in September, remaining at $\approx 59\%$ through October. In November, movement frequencies drastically declined and remained low (<37%) until March, at which point snakes returned to a 58% movement frequency (Fig. 1).

Table 1. Snout–vent length (SVL; cm), study site (ANF = Angelina National Forest, BP = Bienville Parish, FP = Ft. Polk, PR = Peason Ridge, SNF = Sabine National Forest, SV = Scrappin' Valley), and number of relocations for each winter Louisiana Pine Snakes were tracked. Dashes (-) indicate years that snakes were not tracked. Asterisks (*) indicate snakes used in Himes et al. 2006a.

| Snake ID # | SVL | Study site | 1993–1994 | 1994–1995 | 1995–1996 | 1996–1997 | 1997–1998 |
|------------|-----|------------|-----------|-----------|-----------|-----------|-----------|
| Male 1 | 136 | ANF | 14 | 9 | 7 | 2 | - |
| Male 2 | 123 | ANF | - | - | 7 | 2 | 4 |
| Male 3 | 132 | SV | - | - | 3 | - | - |
| Male 4 | 131 | SV | - | - | - | 4 | - |
| Male 5 | 105 | FP | - | - | - | 11 | 7 |
| Male 6 | 115 | FP | 2 | - | - | - | - |
| Male 7* | 112 | BP | - | - | 7 | 30 | 9 |
| Male 8* | 135 | BP | - | - | 6 | 10 | - |
| Male 9 | 105 | BP | - | - | 7 | 9 | - |
| Male 10* | 116 | BP | - | - | - | - | 10 |
| Male 11 | 113 | FP | - | - | - | - | 8 |
| Female 1 | 130 | SV | - | - | - | 3 | - |
| Female 2 | 131 | SV | - | - | 3 | - | - |
| Female 3 | 115 | SNF | - | - | 6 | 2 | - |
| Female 4 | 130 | SNF | 14 | 4 | 6 | 1 | - |
| Female 5 | 116 | SNF | - | 9 | 3 | 2 | - |
| Female 6 | 113 | FP | - | - | - | 10 | - |
| Female 7* | 110 | BP | - | - | 7 | 30 | 6 |
| Female 8* | 80 | BP | - | - | - | - | 9 |

Although movements were greatly curtailed compared to the remainder of the year, snakes occasionally undertook substantial movements during winter ($n = 50$, range = 5 to 841 m, mean = 103.2 m). We recorded data for 12 successive winter dormancy locations of individual snakes, which ranged from 112 to 1406 m (mean = 533.5 ± 127.6 SE) from the previous year's winter dormancy location. We did not detect any individuals using the same winter dormancy location in successive years, or multiple individuals using the same dormancy location simultaneously.

Relocations were typically within the previous active-season MCP home range (68 of 89 unique locations). The 21 unique locations outside of the MCP ranged from 2–118 m away from the respective home range (mean = 45.9 ± 8.1 SE). We did not develop minimum convex polygons to determine winter home ranges because sample sizes of unique winter relocations (mean = 4.7) precluded statistical analysis of winter home-range use.

A male individual from the Angelina National Forest displayed the most extreme winter movement among all snakes. During the winter of 1993–1994, it was relocated 14 times in the same winter refuge (Point A; Fig. 2). The moves before and after the winter season were relatively short (43 and 35 m, respectively; Fig. 2). The following winter (1994–1995), the snake was relocated 9 times (7 unique locations), during which it moved over 1000 m. Two relocations accounted for most of this movement (Fig. 2). The snake was returned to the lab on 31 January 1995 for transmitter replacement. During the winter of 1995–1996, we relocated the snake 7 times (4 unique locations). It moved 133 m in mid-November to an area where it remained until late February 1996, at which time it moved 841 m to the west, beginning its post-winter movements (Fig. 2).




Figure 1. Mean monthly and seasonal movement frequency (%) by Louisiana Pine Snakes in Texas and Louisiana, 1993–1997.

Discussion

During the colder months (November–February), Louisiana Pine Snakes greatly curtailed their movements and remained in the burrows of pocket gophers at relatively shallow depths (13–25 cm; Rudolph et al. 2007). A tendency for movements to be more frequent in early winter was observed, suggesting that the behavioral and physiological transition towards less activity is a gradual process. Presumably, Louisiana Pine Snakes also curtailed their feeding activities during the colder months, based on the reduced movement between burrow systems, and their presence in inactive portions of pocket gopher burrow systems (Rudolph et al. 2007). We did not detect Louisiana Pine Snakes using the same pocket gopher burrow system for winter dormancy in successive years. The mean distance from the site of previous year's winter dormancy location was <600 m. Louisiana Pine Snakes did not make directed movements to wintering sites at the end of the fall, but simply curtailed movements at the burrow system occupied at the end of the fall. This strategy may decrease the frequency of relatively long surface moves, in turn potentially reducing predation risk during a return to a permanent or traditional location.

There are some differences between our findings and those of Himes et al. (2006a) regarding winter activity patterns of Louisiana Pine Snakes. While Himes et al. (2006a) found that Louisiana Pine Snakes are most active during




Figure 2. Winter movements of Louisiana Pine Snake Male 1 during 3 successive winters in the Angelina National Forest, TX. Arrows indicate direction of movement.

March–May and September–November, we found movement frequencies to be the highest during April–August. Himes et al. (2006a) found no snakes in the open during the winter months (December–February), while we found snakes were comparatively active (mean = 24%) during that part of the year. Our study includes the same snakes used by Himes et al. (2006a); however differences in the calculation of activity exist between studies. Himes used percentage of observations of snakes located in the open as an indicator of activity. In contrast, we used movement frequency as an indicator of activity, defined as the frequency of geographically different (≥ 5 m apart) subsequent relocations. The disparity between the winter activity patterns of Louisiana Pine Snakes in these studies is likely a combination of the differences between the definition of activity, and the increased sample size of snakes in our study.

There is a considerable amount of variation in the details of winter dormancy within *Pituophis*. Use of previously occupied sites in successive years by multiple individuals is typically reported in more northern populations (Burger et al. 1988, Kapfer et al. 2008, Parker and Brown 1980, Schroder 1950). These sites are relatively long lasting and allow access to greater depths. More southern populations tend to take refuge individually in more temporary (e.g., downed logs and gopher burrows) and superficial sites (Duran 1998, Franz 2005, Gerald et al. 2006, Gregory 1984, Rudolph et al. 2007). This pattern is presumably a response to winter temperatures, with snakes in colder climates requiring more reliable and deeper hibernation sites, and those in milder climates able to use generally more abundant and superficial sites (Rudolph et al. 2007, Sexton and Hunt 1980).

Rudolph and colleagues (1998) observed the behavior of 3 snakes during the course of 2 prescribed burns in February and March. These snakes simply retreated underground as the fire approached them, and were then insulated from the effects of the passing fire. Six other radio-tagged snakes were known to have survived exposure to the prescribed burns without any apparent damage. They concluded that prescribed fire is not a serious threat to the survival of Louisiana Pine Snakes in fire-dependent climax pine communities.

In contrast, Louisiana Pine Snakes have been found dead on the surface during logging operations during active season months, presumably due to mortality caused by logging machinery (D.C. Rudolph, unpubl. data). Thus, it has been hypothesized that impacts of management practices would be less severe during the winter, when snakes are more often underground. Although snakes move less frequently (and presumably shorter distances) during winter, the impacts of management are still unknown during this time period. While snakes may remain underground more often during winter, impacts from heavy machinery may still be detrimental to snakes occurring only 13–25 cm underground, especially because then they are less capable of moving away from potential threats. Therefore, management practices that involve subsurface soil disturbance in areas known or suspected to support populations of Louisiana Pine Snakes should be carefully evaluated before being undertaken.

Acknowledgments

We thank B. Thatcher and two anonymous reviewers for valuable comments on earlier drafts of the manuscript. We are grateful to R. Carrie, T. Trees, J. Helvey, C. Melder, J. Tull, S. Shively, R. Johnson, D. Baggett, P. Taylor, T. Johnson, W. Ledbetter, K. Moore, K. Munderdorf, E. Keith, C. Collins, and R. Maxey for their assistance with this research. Access to study areas was provided by International Paper Company, Temple-Inland, Inc., Champion International, The Nature Conservancy, the Department of Defense, and Mill Creek Ranch. Partial funding was provided by US Fish and Wildlife Service, Texas Parks and Wildlife Department, Louisiana Department of Wildlife and Fisheries, and Temple-Inland, Inc. The USDA Forest Service's Joint Fire Science Program provided additional funding through a grant to R. Rummer, K. Outcalt, D.C. Rudolph, and D. Brockway. Texas Parks and Wildlife Department and Louisiana Department of Wildlife and Fisheries provided the necessary permits. The use of trade, equipment, or firm names in this publication is for reader information only and does not imply endorsement by the US Department of Agriculture of any product or service. All appropriate animal care guidelines were followed (American Society of Ichthyologists and Herpetologists 2004).

Literature Cited

- American Society of Ichthyologists and Herpetologists. 2004. Guidelines for use of live amphibians and reptiles in field and laboratory research, 2nd Edition. American Society of Ichthyologists and Herpetologists. Available online at <http://www.asih.org/files/hacc-final.pdf>. Accessed 23 March 2011.
- Burgdorf, S.J., D.C. Rudolph, R.N. Conner, D. Saenz, and R.R. Schaefer. 2005. A successful trap design for capturing large terrestrial snakes. *Herpetological Review* 36:421–424.
- Burger, J., R.T. Zappalorti, M. Gochfeld, W.J. Boarman, M. Caffrey, V. Doig, S.D. Garber, B. Lauro, M. Mikovsky, C. Safina, and J. Saliva. 1988. Hibernacula and summer den sites of Pine Snakes (*Pituophis melanoleucus*) in the New Jersey Pine Barrens. *Journal of Herpetology* 22:425–433.
- Conner, R.N., D.C. Rudolph, and J.R. Walters. 2001. The Red-cockaded Woodpecker: Surviving in a Fire-maintained Ecosystem. University of Texas Press, Austin, TX. 363 pp.
- Duran, C.M. 1998. Radio-telemetric study of the Black Pine Snake (*Pituophis melanoleucus lodingi*) on the Camp Shelby Training Site. Mississippi Museum of Natural Sciences Technical Report #59. Jackson, MS. 44 pp.
- Ealy, M.J., R.R. Fleet, and D.C. Rudolph. 2004. Diel activity patterns of the Louisiana Pine Snake (*Pituophis ruthveni*) in eastern Texas. *Texas Journal of Science* 56:383–394.
- Franz, R.D. 2005. Up close and personal: A glimpse into the life of the Florida Pine Snake in a North Florida sand hill. Pp. 120–131 *In* W.E. Meshaka, Jr., and K.J. Babbitt (Eds.). *Amphibians and Reptiles: Status and Conservation in Florida*. Kreiger Publishing Company, Malabar, FL. 334 pp.
- Frost, C.C. 1993. Four centuries of changing landscape patterns in the Longleaf Pine ecosystem. *Proceedings of the Tall Timbers Fire Ecology Conference* 18:17–43.
- Gerald, G.W., M.A. Bailey, and J.N. Holmes. 2006. Movements and activity range sizes of Northern Pine Snakes (*Pituophis melanoleucus melanoleucus*) in middle Tennessee. *Journal of Herpetology* 40:503–510.
- Gregory, P.T. 1984. Communal denning in snakes. Pp. 57–75, *In* R.A. Seigel, L.E. Hunt, J.L. Knight, L. Malaret, and N.L. Zuschlag (Eds.). *Vertebrate Ecology and Systematics: A Tribute to Henry S. Fitch*. University of Kansas Publications of the Museum of Natural History, Special Publication 10, Lawrence, KS. 278 pp.

- Himes, J.G., L.M. Hardy, D.C. Rudolph, and S.J. Burgdorf. 2006a. Movement patterns and habitat selection by native and repatriated Louisiana Pine Snakes (*Pituophis ruthveni*): Implications for conservation. *Herpetological Natural History* 9:103–116.
- Himes, J.G., L.M. Hardy, D.C. Rudolph, and S.J. Burgdorf. 2006b. Body temperature variations of the Louisiana Pine Snake (*Pituophis ruthveni*) in a Longleaf Pine ecosystem. *Herpetological Natural History* 9 (2):117–126.
- Kapfer, J.M., J.R. Coggins, and R. Hay. 2008. Spatial ecology and habitat selection of Bullsnakes (*Pituophis catenifer sayi*) at the northern periphery of their geographic range. *Copeia* 2008:815–826.
- Parker, W.S., and W.S. Brown. 1980. Comparative ecology of two colubrid snakes, *Masticophis taeniatus taeniatus* and *Pituophis melanoleucus deserticola*, in northern Utah. *Milwaukee Public Museum Publications in Biology and Geography* 7:1–104.
- Reichling, S.B. 1995. The taxonomic status of the Louisiana Pine Snake (*Pituophis melanoleucus ruthveni*) and its relevance to the evolutionary species concept. *Journal of Herpetology* 29:186–198.
- Reinert, H.K., and D. Cundall. 1982. An improved surgical implantation method for radio-tracking snakes. *Copeia* 1982:702–705.
- Rodgers, A.R., A.P. Carr, H.L. Beyer, L. Smith, and J.G. Kie. 2007. HRT: Home Range Tools for ArcGIS. Version 1.1. Ontario Ministry of Natural Resources, Centre for Northern Forest Ecosystem Research, Thunder Bay, ON, Canada.
- Rodriguez-Robles, J.A. 2002. Feeding ecology of North American Gopher Snakes (*Pituophis catenifer*, Colubridae). *Biological Journal of the Linnean Society* 77:165–183.
- Rudolph, D.C., and S.J. Burgdorf. 1997. Timber Rattlesnakes and Louisiana Pine Snakes: Hypotheses of decline. *Texas Journal of Science* 49:111–122.
- Rudolph, D.C., S.J. Burgdorf, J. Tull, M. Ealy, R.N. Conner, R.R. Schaefer, and R.R. Fleet. 1998. Avoidance of fire by Louisiana Pine Snakes, *Pituophis melanoleucus ruthveni*. *Herpetological Review* 29:146–148.
- Rudolph, D.C., S.J. Burgdorf, R.N. Conner, C.S. Collins, D. Saenz, R.R. Schaefer, T. Trees, C.M. Duran, M. Ealy, and J.G. Himes. 2002. Prey handling and diet of Louisiana Pine Snakes (*Pituophis ruthveni*) and Black Pine Snakes (*P. melanoleucus lodingi*), with comparisons to other selected Colubrid snakes. *Herpetological Natural History* 9:57–62.
- Rudolph, D.C., S.J. Burgdorf, R.R. Schaefer, R.N. Conner, and R.W. Maxey. 2006. Status of the Louisiana Pine Snake, *Pituophis ruthveni*. *Southeastern Naturalist* 5:463–472.
- Rudolph, D.C., R.R. Schaefer, S.J. Burgdorf, M. Duran, and R.N. Conner. 2007. Pine snake (*Pituophis ruthveni* and *Pituophis melanoleucus lodingi*) hibernacula. *Journal of Herpetology* 41:560–565.
- Rudolph, D.C., C.A. Melder, J.B. Pierce, R.R. Schaefer, and B. Gregory. 2012. Diet of the Louisiana Pine Snake (*Pituophis ruthveni*). *Herpetological Review* 43:243–245.
- Schaefer, W.H. 1934. Diagnosis of sex in snakes. *Copeia* 1934:181.
- Schroder, R.C. 1950. Hibernation of Blue Racers and Bull Snakes in western Illinois. *Chicago Academy of Sciences Natural History Miscellanea* 75:1–2.
- Sexton, O.J., and S.R. Hunt. 1980. Temperature relationships and movements of snakes (*Elaphe obsoleta*, *Coluber constrictor*) in a cave hibernaculum. *Herpetologica* 36:20–26.
- Sweet, S.S., and W.S. Parker. 1991. *Pituophis melanoleucus*. *Catalogue of American Amphibians and Reptiles* 474:1–8.
- US Fish and Wildlife Service (USFWS). 2012. Animals proposed for review. *Federal Register*, Washington, DC. 77(225):70,017–70,018.
- Weatherhead, P.J., and F.W. Anderka. 1984. An improved radio transmitter and implantation technique for snakes. *Journal of Herpetology* 18:264–269.