



University of Dundee

Translational control of gene expression via interacting feedback loops

Wang, Liang; Romano, M. Carmen; Davidson, Fordyce A.

Published in:

Physical Review E: Statistical, Nonlinear, and Soft Matter Physics

DOI:

[10.1103/PhysRevE.100.050402](https://doi.org/10.1103/PhysRevE.100.050402)

Publication date:

2019

Document Version

Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Wang, L., Romano, M. C., & Davidson, F. A. (2019). Translational control of gene expression via interacting feedback loops. *Physical Review E: Statistical, Nonlinear, and Soft Matter Physics*, 100(5), [050402]. <https://doi.org/10.1103/PhysRevE.100.050402>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Translational control of gene expression via interacting feedback loops

Liang Wang

Division of Mathematics, University of Dundee, DD1 4HN, UK

M. Carmen Romano

*SUPA, Institute for Complex Systems and Mathematical Biology,
Department of Physics, Aberdeen AB24 3UE, UK and*

Institute of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen AB24 3FX, UK

Fordyce A. Davidson

Division of Mathematics, University of Dundee, DD1 4HN

*Corresponding Author**

(Dated: October 10, 2019)

Translation is a key step in the synthesis of proteins. Accordingly, cells have evolved an intricate array of control mechanisms to regulate this process. By constructing a multi-component mathematical framework we uncover how translation may be controlled via interacting feedback loops. Our results reveal that this interplay gives rise to a remarkable range of protein synthesis dynamics, including oscillations, step-change and bistability. This suggests that cells may have recourse to a much richer set of control mechanisms than was previously understood.

Keywords: translation, feedback, TASEP, oscillation, bistability

Control of gene expression refers to the processes by which the production of proteins is regulated by the cell. This is at the heart of the functioning of all living organisms and it allows cells to adapt to their environment. Control of gene expression can occur at multiple levels [1, 2]. In this paper we focus on translational control.

Translation is the process by which a protein is made from a messenger RNA (mRNA) molecule. An mRNA consists of a sequence of codons, each coding for a certain amino acid. Translation is performed by molecular machines called ribosomes, which bind to the beginning of the mRNA (5' UTR region), scan it for the start codon and hop from one codon to the next, thereby producing the chain of amino acids which form the protein. When the ribosome reaches the stop codon, the protein is complete, is released into the cytoplasm and the ribosome binds off the mRNA.

Recent years have witnessed an explosion of information about how translational mechanisms regulate protein levels [3]. Prominent examples include translational control during cell stress [4], switching in the mechanism responsible for translation initiation during the cell cycle [5], and translational repression by microRNAs [6–9]. One of the main advantages of translational control is that allows for a rapid cell response [10].

In this paper, we focus on one important case of translational control that has remained unexplored within this research framework, namely the interplay between positive and negative regulatory mechanisms.

Proteins that bind to the 5'UTR region of mRNAs and hinder translation play an important role in regulation of gene expression [18–20]. Examples include proteins involved in neural differentiation and plasticity [21], cognitive problems [22], cell cycle and differentiation of mammalian cells [23, 24] and DNA repair [25]. On the other hand, virtually all mRNAs are subject to positive feedback via ribosome recycling due to their pseudo-circular structure [26, 27]. A particularly pertinent example is the Poly(A) binding protein (PABP) [28–32]. PABP binds to the 3' end (poly(A) tail) of all mRNAs and strongly affects both translational efficiency and stability of all transcripts. Moreover, PABPs interact with the initiation factors bound to the cap of the mRNA, facilitating the circularisation of the mRNA and consequent recycling of ribosomes [26, 33]. Finally, PABPs protect the mRNA from degradation [34]. Therefore, PABPs exert control on protein synthesis via a two-fold mechanism, and as consequence, regulation of PABP availability provides a global mode of translational regulation.

Here we show that the interplay between negative translational feedback and ribosome recycling gives rise to a novel range of dynamical behaviour in protein synthesis, including oscillations, step-change and bistability. The existence of at least one negative and one positive feedback loop has been proved to be a necessary condition for oscillatory behaviour and multistability, respectively [11–15], but examples of this dynamical behaviour have been studied mainly in transcriptional regulation [16] or coupling between transcription and translation [17]. To our knowledge, the generation of oscillatory and bistable behaviour by purely translational regulatory mechanisms has not been studied before.

Modelling framework.—Our mathematical framework is a multi-component model that accounts for transla-

* f.a.davidson@dundee.ac.uk




FIG. 1. Schematic of translation model with ribosome recycling and auto-negative feedback. Competitive recycling (dashed line), non-competitive recycling (solid line). Each site can be occupied by no more than one particle, so that at any time t the state at site i given by $S_i(t)$ is either $S_i(t) = 0$ or $S_i(t) = 1$, with $i = 1, \dots, L$, where L is the lattice length. Particles bind to the first site of the lattice at rate $\tilde{\alpha}$, then hop from one site to the next at rate γ (usually rescaled to one and done so here) and finally leave the lattice from the last site at rate β , marking the point where the associated protein synthesis is completed. See text for further details.

tion, protein complex formation and binding of protein complexes and ribosomes at the 5'UTR (see Fig. 1). At its core is a stochastic model of one-dimensional transport extensively studied in non-equilibrium statistical physics - the Totally Asymmetric Simple Exclusion Process (TASEP) [35]. Here we augment this standard model with processes that explicitly account for protein synthesis and associated positive and negative feedbacks. This allows us to consider both steady state and dynamic behaviours. **Our model predicts the average number of particles per site ρ and the consequent production of proteins. Ribosomes are assumed to attach at a rate $\tilde{\alpha}$ and move along the mRNA at a rate γ . On reaching the final site, ribosomes leave with rate β and are recycled at rate k . At this stage a protein is produced. The protein number N is controlled by a balance of synthesis and degradation with rate constant r . Critical to the model is that both proteins and ribosomes feed back to affect the loading rate $\tilde{\alpha}$ (see Fig. 1).** We consider lattices representing realistic mRNA lengths of 500 codons and simulation times that are of the order of the half-life of typical prokaryotic mRNAs (see Supp. Mat.).

Translation. Ribosomes are represented by particles that hop stochastically along the sites of a one dimensional lattice corresponding to the codons of the mRNA [36]. Note that in the standard case (setting recycling and feedback to zero here) steady state traffic on the lattice can be classified into three main phases: *low density* (LD) ($\alpha < \beta$, $\alpha < 1/2$), *high density* (HD) ($\beta < \alpha$, $\beta < 1/2$) and the *maximal current* (MC) ($\alpha, \beta \geq 1/2$). Each of these phases has an associated average density ρ and current J (average flux of particles), which in the limit of an infinitely long lattice are given by: $\rho_{LD} = \alpha$, $\rho_{HD} = 1 - \beta$, $\rho_{MC} = 1/2$

and $J_p = \rho_p(1 - \rho_p)$, $p \in \{LD, HD, MC\}$ [37, 38]. Corresponding expressions for the modified TASEP considered here will be presented below.

Translational negative feedback. A protein can bind (often in multimeric form) to its own mRNA, thereby blocking the loading of ribosomes. Since protein binding/unbinding to the mRNA is generally much faster than ribosome loading [39], the probability of the start codon being free for ribosome loading can be described by a Hill-function $f(N) = 1/(1 + (N/\theta)^n)$, where N is the protein copy number, θ measures the protein level that induces half maximal ribosome binding rate and n measures cooperativity of the protein multimer (see Supp. Mat.). Thus, the intrinsic initiation rate is modified from α (the standard constant rate) to $\alpha f(N)$.

Translational positive feedback. The two ends of the mRNA can interact leading to a pseudo-circular structure [40], which together with the recycling complex Rli1p [41] promote terminating ribosomes to start a new round of translation on the same mRNA [42]. Following [43], a ribosome on site $i = L$ is assumed to either detach at rate β and enter the reservoir of free ribosomes or move directly onto site $i = 1$ at a *recycling rate* k (if $S_1(t) = 0$) to re-initiate the translation process.

Protein degradation. Once synthesised, proteins enter the intra-cellular pool, where they are subjected to degradation. This can be a complex process [44, 45]. However, as a detailed description is not critical to the work presented, we adopt the widely used approach of modelling removal as a Poisson process with resultant removal rate rN .

Finally, note that if the processes detail above are in steady state, then $J = rN$ and hence $N \equiv N^* = J/r$ and hence we can write $f = 1/(1 + (4IJ)^n)$, where we have introduced the reciprocal factor $I := 1/(4\theta r)$ that measures *feedback intensity* (the factor of 4 is for algebraic convenience).

Models for interacting feedback loops. Experimental results suggest that recycled ribosomes are channelled downstream of the normal *de novo* initiation site and thus may evade the blocking effect of the protein complex [46]. This is the case discussed here and referred to as *non-competitive recycling*. However, the relative position of the protein complex binding site and the recycled ribosome initiation site is not clear. Hence, an alternative is that both recycled ribosomes and *de novo* initiation are blocked by the protein complex, (*competitive recycling*) (see Fig. 1). We comment on this alternative case below.

In the *non-competitive recycling* case a system of ordinary differential equations that determine the dynamics

of the average occupancies ρ_i of the lattice sites is:

$$\begin{aligned} \frac{d\rho_1}{dt} &= \underbrace{\alpha f(N)(1-\rho_1)}_{de\ novo} + \underbrace{k\rho_L(1-\rho_1)}_{recycled} - \rho_1(1-\rho_2), \\ \frac{d\rho_i}{dt} &= \rho_{i-1}(1-\rho_i) - \rho_i(1-\rho_{i+1}), \quad i = 2, \dots, L-1, \\ \frac{d\rho_L}{dt} &= \rho_{L-1}(1-\rho_L) - \beta\rho_L - \underbrace{k(1-\rho_1)\rho_L}_{recycled}. \end{aligned} \quad (1)$$

By direct comparison with the corresponding system for the standard TASEP, *effective* entry and exit rates can be defined as follows

$$\alpha_{eff} := \alpha f(N) + k\rho_L, \quad \beta_{eff} := \beta + k(1-\rho_1). \quad (2)$$

Effective rates for the *competitive recycling* case can be defined equivalently (see Supp. Mat.). However, the behaviour of these two cases is similar in almost all parameter regimes and we present only the results for the non-competitive case highlighting where any differences in the two cases arise.

Steady-state analysis of protein production and ribosome density.— Setting the right hand side of System 1 to zero and applying a mean-field approach leads to conditions that partition steady states of the system into phases aligned with those for the standard case (LD, HD and MC). These conditions are defined by direct substitution of α and β with α_{eff} and β_{eff} , respectively, in the standard conditions stated above. In turn and after some lengthy analysis, expressions for these phases and their boundaries in terms of the system parameter are as follows - note these definitions collapse to the standard cases on setting $k = I = 0$ (see Supp. Mat.).

Maximal Current Phase

$$\frac{\alpha}{2\alpha + k(1+I^n)} \leq \begin{cases} \beta & \leq \frac{\alpha k}{1+I^n - 2\alpha} & \alpha < \frac{1}{2}(1+I^n), \\ \beta & & \text{otherwise.} \end{cases} \quad (3)$$

Within this region,

$$\begin{aligned} \alpha_{eff} &= \frac{\alpha}{1+I^n} + \frac{k}{2\left(\beta + \sqrt{\beta\left(\beta + \frac{k}{\alpha}(1+I^n)\right)}\right)} \quad \text{and} \\ \beta_{eff} &= \frac{1}{2}\left(\beta + \sqrt{\beta\left(\beta + \frac{k}{\alpha}(1+I^n)\right)}\right). \end{aligned}$$

Low Density Phase

$$\alpha < \begin{cases} \beta \left(1 + \left(\frac{4I(\beta+k)(1-\beta)}{(1+k)^2}\right)^n\right) & 2\beta + k < 1, \\ \frac{\beta}{2\beta+k}(1+I^n) & \text{otherwise.} \end{cases} \quad (4)$$

Within this region, there exists a unique, positive expression for α_{eff} that yields unique, positive expressions for β_{eff} and N^* .

High Density Phase

$$\alpha > \begin{cases} \beta \left(1 + \left(\frac{4I(\beta+k)(1-\beta)}{(1+k)^2}\right)^n\right) & 2\beta + k < 1, \\ \frac{k\beta}{1-2\beta}(1+I^n) & \text{otherwise.} \end{cases} \quad (5)$$

Within this region, there exist either one, two or three positive solutions, β_{eff} , that yield positive expressions for α_{eff} and N^* .

For $k, I > 0$ and for a large range of parameter space, long run Monte-Carlo simulations of the modified TASEP reveal steady state phases that are well-characterised by Eqs. (3)-(5) (see Figs. S1 and S2). However, a deeper analysis reveals that the complex interplay of positive and negative feedback can generate distinct and novel *dynamical* behaviour.

Negative feedback and ribosome recycling induce oscillations in cellular protein level.—

Monte-Carlo simulations reveal periodic oscillations in the number of proteins $N(t)$ within the initiation limited regime (LD phase). The stochastic nature of the individual simulations makes it difficult to systematically differentiate periodic oscillations from random fluctuations by visual inspection. However, a power spectrum analysis provides a clear demarcation: a tight, single-peaked spectrum is associated with periodic oscillations (Figs. 2A,C) whereas a broad band response is obtained in the case of stochastic fluctuations (Figs. 2B,D).

The time needed for a ribosome to transit the mRNA induces a delay between initiation and completion of protein synthesis. This generates a delay in the action of the negative feedback - a mechanism known to generate oscillatory behaviour [47]. A simplified model for the protein copy number $N(t)$ in the LD regime is:

$$\begin{aligned} \frac{dN(t)}{dt} &= J(t) - rN(t) \\ &= \alpha_{eff}(t-T)(1-\alpha_{eff}(t-T)) - rN(t), \end{aligned} \quad (6)$$

where T denotes the translational delay time [48]. Appealing to Eqs. 2 and setting $\rho_L = J(t)/\beta_{eff}$, it follows that

$$\alpha_{eff}(t) = \frac{\alpha(\beta+k)}{\alpha k + \beta \left(1 + (4IrN(t))^n\right)}. \quad (7)$$

Substituting Eq. 7 into Eq. 6 results in a delay differential equation for N . The translational delay can be estimated as $T = L/(1-\rho^*)$, where $\rho^* := \alpha_{eff}(N^*)$. This simplified model reproduces the amplitude and period of the stochastic simulations (cf. Fig. 2 A and Fig. S3D). It can be shown that on increasing α the steady state of (6) can be driven unstable via a Hopf bifurcation with




FIG. 2. Simulation and power spectrum of protein level in low density phase. (A,B) Protein number as a function of time. Averages over 5,000 realizations of stochastic simulations (black lines), 20 example realizations (grey lines), single example trajectory (green line). Red dashed lines are mean field N^* as computed using the steady state theory. Typical simulation (A) and the corresponding power spectrum (C) from the region where oscillation are predicted to exist ($\alpha = 0.8$; note $\alpha_{eff} = 0.17 < 0.5$ therefore corresponding to the system being in LD phase). Typical simulation (B) and the corresponding power spectrum (D) from the region where oscillation are not predicted to occur ($\alpha = 0.05$). In all cases $\beta = 0.5$, $k = 0.2$, $I = 5/2$, $r = 0.002$, $n = 5$, $L = 500$. **1 time unit = 1/22 secs.**

the Hopf locus given by

$$B \cos(\sqrt{B^2 - r^2} T) + r = 0, \quad (8)$$

where B is a function of the system parameters (see Supp. Mat.). This locus forms a curve in the $\alpha - \beta$ -plane (see Fig. 3). After some algebra, it can be shown that necessary conditions for the existence of the Hopf locus are $n > 1$ and $I > F(\alpha, \beta, k, n)$ for some positive function F (see Supp. Mat.). The condition $n > 1$ indicates that cooperativity in protein binding is necessary for the onset of oscillations [49]. The second condition indicates that the onset of oscillations occurs when the feedback intensity is sufficiently strong.

As one would intuitively expect, increasing feedback intensity induces onset of oscillations at lower values of the intrinsic loading rate α (see Fig. 3A). Interestingly, the Hopf locus also shifts left on increase the recycling rate (see Fig. 3B). Hence, counterintuitively, ribosome recycling - a positive feedback mechanism - also *enhances*




FIG. 3. The Hopf locus and the effect of varying recycling and feedback. Hopf bifurcation locus in the $\alpha - \beta$ plane computed using Eq. (8). The effect of varying feedback intensity (A) and recycling (B) is shown. Black arrows indicate direction of increasing effect. (A) $I = 25/14$ (red), $I = 25/12$ (green), $I = 25/10$ (blue) with $k = 0.2$ (B) $k = 0.2$ (red), $k = 0.4$ (green), $k = 0.6$ (blue) with $I = 25/10$. $L = 500$ sites, $r = 0.002$, $n = 5$.

the onset of oscillations [50].

Interplay between recycling and negative feedback induces bistability in protein production. -In the MC and LD phases, the current J is uniquely defined for any given parameter set. On the contrary, in the HD phase J can be multiply defined: after some algebra it can be shown that β_{eff} is the solution to the following $2n + 1$ degree equation (see Supp. Mat.):

$$4^n k I^n \beta \beta_{eff}^n (1 - \beta_{eff})^{n+1} - (k\beta + \alpha)\beta_{eff} + \beta(\alpha + k) = 0. \quad (9)$$

For $k = 0$, Eq. (9) has the unique solution $\beta_{eff} = \beta$. For $I = 0$, β_{eff} is uniquely defined by $\beta_{eff} = \beta(\alpha + k)/(k\beta + \alpha)$ [43]. However, when both $k, I > 0$, Eq. (9) can have three admissible solutions, depending on the value of α . Thus, for suitably chosen parameters, there exists an interval of values of α for which three steady state values of $N^* = J/r$ co-exist. Figure 4A shows N^* as function of α , so that on increasing α from zero to one, the model transits from the LD to the HD phase (at $\alpha \approx 0.5$).

With values of α selected from the co-existence interval shown in Fig. 4A, simulations reveal that the number of proteins remains centred on high or low states for a time scale orders of magnitude larger than stochastic fluctuations (4B). Rapid switching events between the favoured state are accompanied by a brief hiatus at an intermediate state. The mean locations of these favoured and intermediate states are well-approximated by the an-




FIG. 4. (A) Protein number N^* as a function of the initiation rate α as predicted by the steady state theory. N^* is first monotonically increasing (LD phase) then decreasing (HD phase) [43]. N^* then passes through two fold bifurcations, leading first into, and then out of the interval of co-existent states - with the upper and lower branches (solid black lines) separated by an intermediate state indicated (dashed grey line). (B) Monte-Carlo simulation of protein number a function of time (solid black line) with the mean field solutions N^* (red dashed lines) from A ($\alpha = 0.77$). In both cases $\beta = 0.015$, $k = 0.8$, $I = 6$, $r = 0.002$, $n = 2$, $L = 500$. **1 time unit = 1/22 secs.**

alytic expressions for the steady states obtained from Eq. 9 (Fig. 4A). Frequency histograms reveal the effect of varying α across the bistable region and together with dwell-time histograms, indicate this to be a memory-less stochastic switching process (see Figs. S5 and S6)

Fixing k (resp. I) and increasing I (resp. k) increases the interval of values of α for which the fold exists (fold width - see Fig. S7). Interestingly, the location of the fold is also an increasing function of I and k . Indeed, somewhat counter-intuitively, for a fixed value of α , increasing the intensity of the negative feedback I can force the system from a low to high N^* state. To understand this recall that within the HD regime, any change of parameters leading to a decrease in the ribosome density leads to an increase in the ribosomal current: increasing I , decreases α_{eff} , thereby decreasing the density of ribosomes on the mRNA. Finally, we note that bimodality in protein production rate is a result of a careful balance between the negative and positive feedback loops and tuning one or the other can drive the system both into and out of a bimodal response (see Fig. S7). Indeed, in the case of competitive recycling no




FIG. 5. Step-change in steady state protein levels. (A) Signal-response curve for protein number N^* as a function of the initiation rate α (black curves). Inadmissible solutions for (9) red dotted curve. The LD-HD boundary vertical grey dashed line. (B) Time series of the number of proteins: $\alpha = 0.28$ then switched to $\alpha = 0.3$ at time $t = 4.5 \times 10^6$. In each case $\beta = 0.015$, $k = 0.263$, $I = 6$, $r = 0.002$, $n = 2$, $L = 500$. **1 time unit = 1/22 secs.**

bistability occurs. Following similar calculations, it can be shown that the equation corresponding to Eq. 9 has a unique solution. It appears that feedback dominates and locks the system into a unique steady state.

Feedback interplay can induce step-changes in protein production—If k is fixed to ensure bistability, a critical value of I exists at which the right boundary of the bistable region coincides precisely with the LD-HD boundary. In this case, as the initiation rate α passes through the LD-HD boundary, a discontinuity in the number of proteins occurs (Fig. 5A). [Qualitatively similar behaviour is obtained on keeping I fixed and varying k .] This step-change in the number of proteins can be large, suggesting that small changes in the ribosome initiation rate α can result in a significant shift in protein levels. Simulations confirm this theoretical prediction. On increasing α dynamically during a simulation, a step change (around 75% reduction) was clearly induced on transept of the LD/HD critical value (Fig. 5B). This cliff-edge response is another unique feature of resulting from the interplay between feedback and recycling.

Conclusions.—Our model shows the rich dynamical behaviour caused by the interplay of negative and

positive feedback loops in translation, and it provides a general mathematical framework to analyse other kinds of gene expression regulation, such as regulation exerted by miRNAs [51]. It suggests that this interplay could provide cells with a versatile mechanism to adapt their protein levels according to the environment. The centrally important protein PABP is subject to ribosome recycling and is known to exhibit translational negative feedback. Interestingly, PABP has also been implicated

in circadian oscillations [52]. Finally, disturbances of poly(A) tail length have been linked to a number of physiological and pathological processes. Therefore, a better understanding of the interplay of ribosome recycling and translational negative feedback has far reaching consequences.

Acknowledgements. LW was supported by the Northern Research Partnership

-
- [1] J. Hausser, L. Keren, and U. Alon, *Nature Communications* **10** (2019).
- [2] U. Alon, *An introduction to systems biology: design principles of biological circuits* (Boca Raton, FL, Chapman & Hall/CRC, 2007).
- [3] J. Hershey, N. Sonenberg, and M. Mathews, *Cold Spring Harb Perspect Biol.* **4**, a011528 (2012).
- [4] K. Spriggs, M. Bushell, and A. Willis, *Mol. Cell* **40**, 228 (2014).
- [5] S. Pyronnet, J. Dostie, and N. Sonenberg, *Genes Dev.* **15**, 2083 (2001).
- [6] M. Selbach, B. Schwanhussner, N. Thierfelder, Z. Fang, R. Khanin, and N. Rajewsky, *Nature* **455**, 58 EP (2008).
- [7] A. Zinovyev, N. Morozova, A. N. Gorban, and A. Harel-Belan, “Mathematical modeling of microrna-mediated mechanisms of translation repression,” in *MicroRNA Cancer Regulation: Advanced Concepts, Bioinformatics and Systems Biology Tools*, edited by U. Schmitz, O. Wolkenhauer, and J. Vera (Springer Netherlands, Dordrecht, 2013) pp. 189–224.
- [8] S. Nikolov, J. V. Gonzalez, M. Nenov, and O. Wolkenhauer, *Biotechnology & Biotechnological Equipment* **26**, 3315 (2012).
- [9] A. Zinovyev, N. Morozova, N. Nonne, E. Barillot, A. Harel-Bellan, and A. N. Gorban, *BMC Systems Biology* **4** (2010), 10.1186/1752-0509-4-13.
- [10] J. W. Hershey, N. Sonenberg, and M. B. Mathews, *Cold Spring Harbor Perspectives in Biology* (2018), 10.1101/cshperspect.a032607.
- [11] J. L. Gouz, *Journal of Biological Systems* **06(1)**, 11 (1998).
- [12] E.-H. Snoussi, *Journal of Biological Systems* **06(1)**, 3 (1998).
- [13] P. François and V. Hakim, *Proceedings of the National Academy of Sciences* **101**, 580 (2004), <https://www.pnas.org/content/101/2/580.full.pdf>.
- [14] L. Potvin-Trottier, N. Lord, G. Vinnicombe, and J. Paulsson, *Nature* **538**, 514?517 (2016).
- [15] E. Ullner, A. Koseska, J. Kurths, E. Volkov, H. Kantz, and J. García-Ojalvo, *Phys. Rev. E* **78**, 031904 (2008).
- [16] R. Hermsen, B. Ursem, and P. R. ten Wolde, *PLOS Computational Biology* **6**, 1 (2010).
- [17] Y. Zarai and T. Tuller, *PLOS Computational Biology* **14**, 1 (2018).
- [18] K. Abdelmohsen, *Modulation of Gene Expression by RNA Binding Proteins: mRNA Stability and Translation Fundamental Principles of Optical Lithography* (InTech Open, 2012).
- [19] O. Levi and Y. Arava, *PLOS Biology* **17**, 1 (2019).
- [20] R. Betney, J. de Silva, and I. Stansfield, *RNA-A* **16**, 655 (2010).
- [21] H. M. N and L. H, *Cell Mol Life Sci* **65(20)**, 3168 (2008).
- [22] G. J. Bassell and S. T. Warren, *Neuron* **60**, 201 (2008).
- [23] C. K. Damgaard and J. Lykke-Andersen, *Genes & Development* **25**, 2057 (2011).
- [24] V. Dormoy-Raclet, J. Markovits, A. Jacquemin-Sablon, and H. Jacquemin-Sablon, *RNA Biology* **2**, 112 (2005).
- [25] T. Fukuda, M. Ashizuka, T. Nakamura, K. Shibahara, K. Maeda, H. Izumi, K. Kohno, M. Kuwano, and T. Uchiyumi, *Nucleic Acids Research* **32**, 611 (2004).
- [26] S. Wells, P. Hillner, R. Vale, and A. Sachs, *Mol. Cell* **2**, 135 (1998).
- [27] Z. A. Afonina, A. G. Myasnikov, V. A. Shirokov, B. P. Klaholz, and A. S. Spirin, *Nucleic Acids Research* **42**, 9461 (2014), <http://oup.prod.sis.lan/nar/article-pdf/42/14/9461/14123029/gku599.pdf>.
- [28] E. Hornstein, H. Harel, G. Levy, and O. Meyuhis, *FEBS Letters* **457**, 209 (1999).
- [29] O. Neto, N. Standart, and C. Desa, *Nucleic Acids Research* **23**, 2198 (1995).
- [30] J. Bag, *J. Biol. Chem.* **276**, 47352 (2001).
- [31] G. Patel, S. Ma, and J. Bag, *Nucleic Acids Res.* **33(22)**, 7074?7089 (2005).
- [32] B. Gorgoni and N. Gray, *BRIEFINGS IN FUNCTIONAL GENOMICS AND PROTEOMICS* **3**, 125?141 (2004).
- [33] A. Kahvejian, Y. V. Svitkin, R. Sukarieh, M. N. M’Boutchou, and N. Sonenberg, *Genes & Development* **19**, 104 (2005).
- [34] D. Cao and R. Parker, *RNA* **7**, 1192 (2001).
- [35] B. Derrida, E. Domany, and D. Mukamel, *J. Stat. Phys.* **69**, 667 (1992).
- [36] C. MacDonald, J. Gibbs, and A. Pipkin, *Biopolymers* **6**, 1 (1968).
- [37] B. Derrida, M. Evans, C. Hakim, and V. Pasquier, *J. Phys. A: Math. Gen.* **26**, 1493 (1993).
- [38] G. Schütz and E. Domany, *J. Stat. Phys.* **72**, 277 (1993).
- [39] L. Chen, R. Wang, T. Kobayashi, and K. Aihara, *Phys. Rev. E* **70**, 011909 (2004).
- [40] D. Barthelme, S. Dinkelaker, S.-V. Albers, P. Londei, U. Ermler, and R. Tampe, *Proc. Natl Acad. Sci.* **108**, 3228 (2011).
- [41] C. Shoemaker and R. Green, *Proc Natl Acad. Sci.* **108**, 1392 (2011).
- [42] N. Amrani, S. Ghosh, D. Mangus, and A. Jacobson, *Nature* **435**, 1276 (2008).
- [43] E. Marshall, I. Stansfield, and M. Romano, *J. R. Soc. Interface* **11**, 104 (2014).
- [44] D. A. Rothenberg, M. J. Taliaferro, S. M. Huber, T. J. Begley, P. C. Dedon, and F. M. White, *iScience* **9**, 367 (2010).

- (2018).
- [45] R. Golan-Lavi, C. Giacomelli, G. Fuks, A. Zeisel, J. Sontag, S. Sinha, W. Kstler, S. Wiemann, U. Korf, Y. Yarden, and E. Domany, *Cell Reports* **18**, 3129 (2017).
- [46] L. Rajkowitsch, C. Vilela, K. Berthelot, C. Ramirez, and J. McCarthy, *J. Mol. Biol.* **335**, 71 (2004).
- [47] S. Pigolotti, S. Krishna, and M. Jensen, *Proc Natl Acad. Sci.* **104**, 6533 (2007).
- [48] Note that in agreement with numerical simulations, we only expect oscillations within the LD phase, since in both HD and MC, the current J is independent of the delay between loading and exit (see Supp. Mat.).
- [49] M. Elowitz and S. Leibler, *Nature* **403**, 335 (2000).
- [50] B. Ananthasubramaniam and H. Herzog, *PLoS ONE* **2014**, e104761 (2014).
- [51] S. Oliveto, M. Mancino, N. Manfrini, and S. Biffo, *World J Biol Chem* **8**, 45756 (2017).
- [52] S. Kojima, E. Sher-Chen, and C. Green, *Gene Dev.* **26**, 2724 (2012).