

Acta Biologica Plantarum Agriensis 7: 42–54 (2019)
ISSN 2061-6716 (Print), 2063-6725 (Online)
<http://abpa.ektf.hu/>

DOI:10.21406/abpa.2019.7.42
Research article

ONNIA TRIQUETRA (PERS.) IMAZEKI, A PINE ASSOCIATED POLYPORE SPECIES REPORTED FOR THE FIRST TIME FROM HUNGARY

Balázs Palla^{1*}, Imre Borsicki², Zoltán Lukács³ & Viktor Papp¹

¹Department of Botany and Soroksár Botanical Garden, Faculty of Horticulture,
Szent István University, H-1118 Budapest, Ménesi str. 44. Hungary; ²2500
Esztergom, Bajcsy-Zsilinszky str. 26. Hungary; ³1071 Budapest, Damjanich str. 54.
Hungary; *E-mail: lihelih1@gmail.com

Abstract: *Onnia triquetra*, a *Pinus* specialist polypore species in Europe, is reported for the first time in Hungary. Historical and recent collections of almost 60 years were examined. The macro- and micromorphological characteristics are given, along with ecological inferences of the species and its hosts in Hungary.

Keywords: Hymenochaetaceae, Europe, native, new record

INTRODUCTION

Hungarian forests are composed mainly of deciduous tree species, covering 89.5% of the forest area. The most common species are oaks, poplars, beech and hornbeam, among aliens the black locust. Conifers represent only 10.5% of the forests, with a significant decline in their actual area in the last two decades (from 243 500 hectares in 2000 to 195 000 hectares in 2016). The most widespread conifer species are Scots pine (*Pinus sylvestris* L.) and Austrian pine (*Pinus nigra* J.F.Arnold), covering an area of 6.2% and 3.2% respectively (Komarek 2018). Scots pine (*Pinus sylvestris* L.) is considered to be native, while Austrian pine (*P. nigra* J.F.Arnold) was introduced to the Pannonian basin, with its first plantations established in the 19th century (Borhidi 2003; Tamás 2003; Bölöni *et al.* 2008).

The effects of such forest plantations on local biodiversity depend on the sensibility and ecological specificity of the site (e.g. the surrounding landscape), in which they are found. Thus, planted

forests – both native and exotic – may have positive, as well as negative effects on biodiversity at the population, stand or landscape level (Humphrey *et al.* 2000; Mack *et al.* 2000; Carnus *et al.* 2006; Cseresnyés and Tamás 2014). Due to the wide ecological tolerance of Scots pine, it is hard to describe the effects of its semi-natural stands on local biodiversity. In this regard, indicator species for these semi-natural stands should be recognized on the local level (Kelly and Conolly 2000). In case of Austrian pine, it is widely accepted that its plantations had detrimental effects on the Hungarian sites where they were planted in the past. Many local species disappeared from the native flora and fauna of such plantations (Török and Tóth 1996; Bartha *et al.* 2004; Cseresnyés and Tamás 2014).

On the other hand, vast diversity of fungi can be seen in plantations of exotic conifers (Newton and Haigh 1998; Humphrey *et al.* 2000). These planted stands could serve as potential substrata for native lignicolous fungal assemblages, which could include rare and threatened species (Humphrey *et al.* 2000; Ryvarden and Melo 2014).

The *Onnia* P. Karst. genus (Hymenochetaceae) contains nine white rot polypore species, from which seven are growing on gymnosperms. *Onnia tomentosa* (Fr.) P. Karst. and *O. leporina* (Fr.) H. Jahn grow mainly on *Picea*, but also can be found on *Pinus*, *Abies* and *Larix*, while *O. triquetra* (Pers.) Imazeki, *O. subtriquetra* Vlasák & Y.C. Dai, *O. microspora* Y.C. Dai & L.W. Zhou, *O. kesiya* Zhou & F.Wu and *O. tibetica* Y.C. Dai & S.H. He occurs mainly on *Pinus* (Dai 2010, 2012; Ryvarden and Melo 2014; Ji *et al.* 2017). These fungi are characterized by annual, sessile or stipitate basidiocarps; pileal surface from yellowish brown to dark brown and velutinate to rough; pore surface from yellowish brown to dark brown; duplex context. Main microscopic characteristics are: monomitic hyphal system with generative hypha bearing simple septa; presence of mostly hooked hymenial setae; and hyaline, thin-walled, smooth, nonamyloid, nondextrinoid, and acyanophilous basidiospores (Dai 2010; Ji *et al.* 2017). The genus is widespread; its distribution extends from boreal to subtropical climates, being present throughout the Northern Hemisphere (Dai 2012; Ryvarden and Melo 2014; Lockman and Kearns 2016; Ji *et al.* 2017).

In Europe, *Onnia tomentosa*, *O. leporina* and *O. triquetra* are considered to be native. In contrast to the first two circumboreal

species, *O. triquetra* is geographically restricted to Europe, growing mainly on *Pinus* (Ryvarden and Gilbertson 1993; Dai 2012; Ji *et al.* 2017). The possible occurrence of *O. triquetra* in Hungary was implied first by Zoltán Igmándy in the 1980's (Igmándy 1989), but since then it was never found.

In Igmándy's own collection, among deposited unidentified specimens, we have found a sample collected from a *P. sylvestris* trunk near Szilvagy (Zala County, Western Hungary) from 1959, which we identified as *O. triquetra*. Almost 60 years later, a recently collected specimen from *P. nigra* in the Búbánat-völgy (Komárom-Esztergom County, North Hungary) was identified to be the same species.

MATERIALS AND METHODS

The historical specimen (Z. Igmándy 1131) collected by L. Haracsi is available at the Institute of Silviculture and Forest Protection, University of Sopron (Sopron, Hungary); the new specimen (Borsicki 100916) is deposited at the personal fungarium of the last author (PV). Macromorphological descriptions are based on field notes. The micromorphological data were obtained using a light microscope following the methods by Papp and Dima (2018). Microscopic characters, measurements and drawings were made from slide preparations stained with Melzer's reagent. The sections were studied at 1000× magnification using a Zeiss Axio Imager A2 light microscope (Zeiss, Göttingen, Germany) attached with an AxioCam HRc camera (Zeiss, Göttingen, Germany). Drawings of the micromorphological characteristics of the basidiocarps were made using a drawing tube. Spores were measured from sections cut from the tubes. For the measurements, we used the Axio Vison Release 4.8. software. The following abbreviations were used in the description of the basidiospores: IKI = Melzer's reagent, IKI- = both inamyloid and indextrinoid, L = mean spore length, W = mean spore width, Q = variation in the L/W ratios, n = number of spores measured.

TAXONOMY

Onnia triquetra (Pers.) Imazeki [as 'triqueter'], Mycol. Fl. Japan, Basidiomycetes 2(4): 386 (1955) – (Figure 1 and 2)

Basionym: *Boletus triqueter* Pers., Observ. mycol. (Lipsiae) 1: 86 (1796)

Synonyms: *Polyporus triqueter* (Pers.) Pers., Mycol. eur. (Erlanga) 2: 57 (1825); *Inoderma triquetrum* (Pers.) P. Karst. [as 'triquetrum'], Meddn Soc. Fauna Flora fenn. 5: 39 (1879); *Ochroporus triqueter* (Pers.) J. Schröt., in Cohn, Krypt.-Fl. Schlesien (Breslau) 3.1(25–32): 485 (1888) [1889]; *Mucronoporus tomentosus* var. *triqueter* (Pers.) Domański, Orloś & Skirg., Flora Polska. Grzyby, II: 321 (1967); *Mucronoporus circinatus* var. *triqueter* (Pers.) Domański, Orloś & Skirg., Fungi, Polyporaceae 2, Mucronoporaceae 2, Revised transl. Ed. (Warsaw): 289 (1973)

Specimens examined: Hungary, Zala County, Szilvág, on *Pinus sylvestris* trunk, 22 Jan 1959, leg. L. Haracsi, herb. Z. Igmándy 1131. Komárom-Esztergom County, Esztergom, Búbánat-völgy, on *Pinus nigra* trunk, 10 Sept 2016, leg. I. Borsicki, herb. Borsicki 100916.

Basidiocarps annual, sessile or laterally stipitate/substipitate, soft corky or corky. Pileus dimidiate, with 3–4 cm diameter and 2–3 cm thickness at centre. Pileal surface golden brown to rust-brown, tomentose to velutinate, concentric zones indistinct; margin sharp to blunt. Pore surface yellowish brown to rusty brown; sterile margin pale yellowish; pores angular, 3–4 per mm, later more angular to semi labyrinthine and up to 1–2 mm wide in places; dissepiments thin, slightly lacerate. Context duplex; upper layer rust-brown, spongy; lower layer umber-brown, corky, demarcation zone indistinct between the two layers; entire context up to 1 cm thick. Tubes pale umber-brown, slightly paler than context and pore surface, hard corky, up to 5 mm long. Stipe rust-brown, hirsute to velutinate; 1–2 cm long, 0.5–1 mm in diameter; pores decurrent on stipe.




Figure 1. *Onnia triquetra* basidiocarp in the habitat (Búbánat-völgy, Komárom-Esztergom County, Hungary) **A.** pileal surface **B.** pore surface. Photographs by I. Borsicki.




Figure 2. Micromorphological structures of *Onnia triquetra*, collected from *Pinus sylvestris* (a-e) and *Pinus nigra* (f-j) **a.** section of trama **b.** setae **c.** hyphae from trama **d.** hyphae from context **e.** basidiospore **f.** setae **g.** narrow setae **h.** hyphae from trama **i.** hyphae from context **j.** basidiospores.

Hypal system monomitic; generative hyphae simple septate; tissue darkening but otherwise unchanged in KOH. Tramal hyphae hyaline to pale yellowish, thin- to slightly thick-walled, occasionally branched and septate, agglutinated, 2.5–6 µm in diam. Contextual hyphae yellowish to golden brown, thin- to slightly thick-walled, rarely branched, with frequent simple septa, regularly arranged, agglutinated, 5–8 µm in diameter. Contextual hyphae at upper

spongy layer pale yellowish, thin-walled, unbranched or rarely branched, with frequent simple septa, interwoven, 3–6 μm in diam; hyphae of lower solid layer pale yellowish, septate, rarely branched, 3–5 μm in diameter. Hyphae from stipe similar to those in context. Setae abundant, mostly hooked, sometimes narrow, sharp-pointed, dark brown, thick-walled, deep-rooting, 28–68 \times 8–13 μm . Basidia clavate, with four sterigmata, simple septate at the base, 9–17 \times 4.8–6 μm . Basidiospores cylindric-ellipsoid, hyaline, thin-walled, smooth, mostly glued together in tetrads, IKI–, (5.84–) 6.15–6.61(–7.03) \times (4.12–)4.24–4.68(–5.02) μm , L=6.4 μm , W= 4.5 μm , Q=1.35–1.50, Q_{av} =1.43 (n=50).

DISCUSSION

The concept of whether a planted and/or non-native forest is beneficial or detrimental for local biodiversity has been much debated. It is often assumed, that tree plantations impoverish local flora and fauna; however, proper forest management can improve the balance between sustainable biodiversity and timber production (Hartley 2002). In case of non-indigenous gymnosperm plantations, both negative (Török and Tóth 1996; Mack *et al.* 2000; Magura *et al.* 2000; Brockerhoff *et al.* 2003; Cseresnyés and Tamás 2014) and positive (Fisher and Goldney 1998; Brockerhoff *et al.* 2003) impacts on natural biodiversity have been observed, related to different ecological and management qualities of the stands.

Plantations of native and non-native gymnosperms have considerable impact on fungal biodiversity. Exotic *Pinus radiata* plantations were found to be less rich in lignicolous saprotrophs and ectomycorrhizal fungi, compared to native oak forests in the Basque Country (Sarrionandia *et al.* 2015). Although the study recorded numerous host specific fungi within the *P. radiata* stands, the overall fungal community consisted mainly of generalist species (i.e. non habitat specialists), presumably because the stands did not reach maturity due to local forest management practices. The authors concluded that exotic plantations in their older, mature stages, with patches of native trees are more capable of contributing to a higher macrofungal diversity. Humphrey *et al.* (2000) came to a similar conclusion in terms of exotic *Picea sitchensis* and native *Pinus sylvestris* plantations in Great Britain. They have met unexpectedly high incidences of rare and

threatened fungi species in stands near to the extant native pinewoods (acting as sources of inoculum). They also emphasized the importance of understory diversity and the volume of fallen deadwood, as well as maintaining the over-mature growth stage of the stands through some form of continuous cover management system. Dejene *et al.* (2017) also confirmed the role of older stands of alien gymnosperms in shaping fungal diversity in case of *Pinus patula* in Ethiopia. Johnston (2010) pointed out, that though introduced *Pinus* species in New Zealand could serve as hosts for indigenous non-lichenised fungi, exotic plants are usually associated with exotic fungi. A similar observation was made by Campi *et al.* (2015) in case of *P. taeda*. Functional composition of wood-decaying assemblages may also differ amongst native and non-native conifers: white rotting or brown rotting fungi could be more or less dominant on the non-indigenous substrata, depending on host species and geographical region (Edman and Fällström 2013).

In Europe, the main distribution area of Scots pine is Central and Northern Europe, while more to the south it is usually restricted to higher elevations (Boratyński 1991). Due to its wide tolerance to climatic and edaphic conditions, this species sustains on many different habitat types (Kelly and Conolly 2000; Köbölkuti *et al.* 2017). In Hungary, Scots pine occurs mainly in the westernmost part of the country (Vendvidék, Őrség, Kemeneshát) in acidophilous coniferous forests. Small stands are also present in the xerophilous forest communities in Central- and Western Transdanubia, on calcareous, sandy substrates, like the habitat in Fenyőfő (Kelly and Conolly 2000; Borhidi 2003; Bölöni *et al.* 2008; Köbölkuti 2018). It has been stated that original Scots pine stands in Hungary are of small size, they do not cover large territories, some being considered relict stands. However, Scots pine forests are preserved by land use that has a crucial role in their survival. Populations in the present are mostly considered to be of planted origin (Völgyi 1955; Mason and Alía 2000; Bölöni *et al.* 2008).

Another important species in Hungary from the *Pinus* genera is Austrian pine. Being an Alpine-Mediterranean tree, Austrian pine is alien to the Pannonian Basin. It was first introduced to Hungary in the 19th century for soil preservation and landscape protection purposes. Trees were planted on hillsides, dolomite slopes and on sandy substrates in the lowlands (Tamás 2003). Today, larger

plantations are found mainly in the Great Hungarian Plain, but there are also plantations in the Transdanubian- and in the Northern Hungarian Mountains (Cseresnyés and Tamás 2014). The establishment of these plantations had a harmful effect on the native vegetation of dolomite slopes, warm tolerant oak forests and sandy grasslands. The above mentioned habitats contain numerous endemic species; the monodominant structure of the plantations resulted in closed canopies, impoverished understory and reduced diversity of the local plant and animal communities (Török and Tóth 1996; Bartha *et al.* 2004; Bíró 2008; Cseresnyés and Tamás 2014). Moreover, the accumulating litter under the canopy of Austrian pine plantations contributed to elevated fire risks. In this regard, both needles, cones and fallen branches could serve as focal points for forest fires due to the relatively slow decomposition rate (Cseresnyés *et al.* 2006).

Both Scots pine and Austrian pine are serving as substrata for numerous native polypore fungi in Europe. Many of these species are found in East- and Central Europe, including Hungary (Ryvarden and Gilbertson 1993, 1994). Conifer-dwelling macrofungi in Hungary were already being studied more than a half century ago. Throughout these decades, numerous poroid species were recorded on hosts from the *Pinus* genera (Igmándy 1954; Völgyi 1955; Pagony 1977). Amongst these, species inhabiting both gymnosperm and angiosperm trees are represented in the highest amount (e.g. *Bjerkandera adusta* (Willd.) P. Karst., *Fomitopsis pinicola* (Sw.) P. Karst., *Pappia fissilis* (Berk. & M.A. Curtis) Zmitr., etc.). The angiosperm hosts include species from natural forest communities, e.g. *Fagus*, *Quercus*, *Carpinus*, *Populus* (Szabó 2012; Siller *et al.* 2013). Thus, polypores inhabiting both native and non-native gymnosperms and angiosperms may be originally considered indigenous to Hungary, with a preference for a broad range of hosts.

A bit smaller amount of *Pinus*-inhabiting polypores in Hungary are restricted only to gymnosperms: e.g. *Neoantrodia serialis* (Fr.) Audet, *Climacocystis borealis* (Fr.) Kotl. & Pouzar, *Gloeophyllum sepiarium* (Wulfen) P. Karst., *Gloeoporus taxicola* (Pers.) Gilb. & Ryvarden, *Osmoporus odoratus* (Wulfen) Singer, *Skeletocutis amorpha* (Fr.) Kotl. & Pouzar. Some species are found on *Pinus* exclusively: e.g. *Antrodia ramentacea* (Berk. & Broome) Donk, *Dichomitus squalens* (P. Karst.) D.A. Reid, *Diplomitoporus flavescens*

(Bres.) Domański, *Porodaedalea pini* (Brot.) Murrill. According to Igmándy (1981), a small number of these species associated with Scots pine were also found on Austrian pine (e.g. *Climacocystis borealis*, *Postia fragilis* (Fr.) Jülich), while some other species were only present on the former host (e.g. *Dichomitus squalens*, *Fuscoporia viticola* (Schwein.) Murrill).

Taken these into consideration, it seems that naturalized and non-native *Pinus* plantations have an important role in conserving indigenous polypore diversity. Igmándy's collection of *Onnia triquetra* from 1959 shows, that even a wider spectrum of poroid species were most probably present since a long time ago in Hungarian semi-natural Scots pine forests. The ongoing shrinkage of these forests (KSH 2013; Komarek 2018) and the recent occurrence of *O. triquetra* on Austrian pine highlights the importance of planted, non-native stands of conifers in preserving fungal biodiversity. These stands could serve as possible surrogate habitats for indigenous fungi, as it was presented in many other cases (Humphrey *et al.* 2000; Johnston 2010; Edman and Fällström 2013; Sarrionandia *et al.* 2015; Dejene *et al.* 2017).

Acknowledgements – The authors wish to thank Jenő Jakab (Institute of Sylviculture and Forest Protection, University of Sopron) for providing additional information on the specimen collected by Zoltán Igmándy. We also wish to thank Dr. Mária Höhn for her additional suggestions and advice related to the topic.

REFERENCES

- BARTHA, S., CAMPETELLA, G., CANULLO, R., BÓDIS, J. & MUCINA, L. (2004). On the importance of fine-scale spatial complexity in vegetation restoration studies. *International Journal of Ecology and Environmental Sciences* **30**(2): 101–116.
- BIRÓ, M. (2008). A Duna-Tisza köze fásszárú vegetációjának átalakulása a 18. század óta, különös tekintettel a száraz homokterületekre. In: KRÖEL-DULAY, GY., KALAPOS, T. & MOJZES, A. (eds.): *Talaj-vegetáció-klíma kölcsönhatások. Köszöntjük a 70 éves Láng Editet*. MTA Ökológiai és Botanikai Kutatóintézete. Vácrátót, pp. 23–38.
- BÖLÖNI, J., MOLNÁR, Z., BIRÓ, M. & HORVÁTH, F. (2008). Distribution of the (semi-) natural habitats in Hungary II. Woodlands and shrublands. *Acta Botanica Hungarica* **50**: 107–148. <https://doi.org/10.1556/ABot.50.2008.Suppl.6>
- BORATYŃSKI, A. (1991). *Range of natural distribution*. In: GIERTYCH, M. & MATYAS, C. (eds.): *Genetics of Scots pine. Developments in Plant Genetics and Breeding*. Elsevier, Amsterdam, pp. 19–30. <https://doi.org/10.1016/B978-0-444-98724-2.50006-7>
- BORHIDI, A. (2003). *Magyarország növénytársulásai*. Akadémiai Kiadó, Budapest,

610 p.

- BROCKERHOFF, E.G., ECROYD, C.E., LECKIE, A.C. & KIMBERLEY, M.O. (2003). Diversity and succession of adventive and indigenous vascular understorey plants in *Pinus radiata* plantation forests in New Zealand. *Forest Ecology and Management* **185**(3): 307–326. [https://doi.org/10.1016/S0378-1127\(03\)00227-5](https://doi.org/10.1016/S0378-1127(03)00227-5)
- CAMPI, M.G., MAUBET, Y.E. & BRITOS, L. (2015). Mycorrhizal fungi associated with plantations of *Pinus taeda* L. from the National University of Asunción, Paraguay. *Mycosphere* **6**(4): 486–492. <https://doi.org/10.5943/mycosphere/6/4/10>
- CARNUS, J.-M., PARROTTA, J., BROCKERHOFF, E., ARBEZ, M., JACTEL, H., KREMER, A., LAMB, D., O'HARA, K. & WALTERS, B. (2006). Planted forests and biodiversity. *Journal of Forestry* **104**(2): 65–77.
- CSERESNYÉS, I., CSONTOS, P. & BÓZSING, E. (2006). Stand age influence on litter mass of *Pinus nigra* plantations on dolomite hills in Hungary. *Canadian Journal of Botany* **84**(3): 363–370. <https://doi.org/10.1139/b06-003>
- CSERESNYÉS, I. & TAMÁS, J. (2014). Evaluation of Austrian Pine (*Pinus nigra*) Plantations in Hungary With Respect To Nature Conservation – a Review. *Tájékológiai Lapok* **12**(2): 267–284.
- DAI, Y.C. (2010). Hymenochaetaceae (Basidiomycota) in China. *Fungal Diversity* **45**(1): 131–343. <https://doi.org/10.1007/s13225-010-0066-9>
- DAI, Y.C. (2012). Polypore diversity in China with an annotated checklist of Chinese polypores. *Mycoscience* **53**(1): 49–80. <https://doi.org/10.1007/s10267-011-0134-3>
- DEJENE, T., ORIA-DE-RUEDA, J.A. & MARTÍN-PINTO, P. (2017). Fungal diversity and succession following stand development in *Pinus patula* Schiede ex Schltdl. & Cham. plantations in Ethiopia. *Forest Ecology and Management* **395**: 9–18. <https://doi.org/10.1016/j.foreco.2017.03.032>
- EDMAN, M. & FÄLLSTRÖM, I. (2013). An introduced tree species alters the assemblage structure and functional composition of wood-decaying fungi in microcosms. *Forest Ecology and Management* **306**: 9–14. <https://doi.org/10.1016/j.foreco.2013.06.023>
- FISHER, A.M. & GOLDNEY, D.C. (1998). Native forest fragments as critical bird habitat in a softwood forest landscape. *Australian Forestry* **61**(4): 287–295. <https://doi.org/10.1080/00049158.1998.10674753>
- HARTLEY, M.J. (2002). Rationale and methods for conserving biodiversity in plantation forests. *Forest Ecology and Management* **155**(1–3): 81–95. [https://doi.org/10.1016/S0378-1127\(01\)00549-7](https://doi.org/10.1016/S0378-1127(01)00549-7)
- HUMPHREY, J.W., NEWTON, A., PEACE, A.J. & HOLDEN, E. (2000). The importance of conifer plantations in northern Britain as a habitat for native fungi. *Biological Conservation* **96**(2): 241–252. [https://doi.org/10.1016/S0006-3207\(00\)00077-X](https://doi.org/10.1016/S0006-3207(00)00077-X)
- IGMÁNDY, Z. (1954). A fenyőtapló előfordulása és károsítása hazánkban. *Erdészeti és Faipari Tudományos Közlemények* **53**: 5–10.
- IGMÁNDY, Z. (1981). *Hazánk csövestapló (Polyporaceae s.l.) flórája és a fajok növénykórtani jelentősége*. Soproni Egyetem, Sopron, 159 p.
- IGMÁNDY, Z. (1989). Vas megye csövestapló (Polyporaceae s. l.) flórája. *Savaria - A Vas Megyei Múzeumok Értesítője* **17–18**: 27–49.
- JI, X.-H., HE, S.-H., CHEN, J.-J., SI, J., WU, F., ZHOU, L.-W., VLASÁK, J., TIAN, X.-M. & DAI, Y.-C.

- (2017). Global diversity and phylogeny of *Onnia* (Hymenochaetaceae) species on gymnosperms. *Mycologia* **109**(1): 27–34.
<https://doi.org/10.1080/00275514.2016.1274619>
- JOHNSTON, P.R. (2010). Causes and consequences of changes to New Zealand's fungal biota. *New Zealand Journal of Ecology* **34**(1): 175–184.
- KELLY, D.L. & CONOLLY, A. (2000). A review of the plant communities associated with Scots Pine (*Pinus sylvestris* L.) in Europe, and an evaluation of putative indicator/specialist species. *Investigación Agraria. Sistemas y Recursos Forestales* **9**(1): 15–40.
- KÖBÖLKUTI, Z.A. (2018). Adaptive variation of selected Scots pine (*Pinus sylvestris* L.) populations from the Carpathians and the Pannonian basin inferred from morphological, anatomical and genomic data. Szent István University, Budapest, 139 p.
- KÖBÖLKUTI, Z.A., TÓTH, E.G., LADÁNYI, M. & HÖHN, M. (2017). Morphological and anatomical differentiation in peripheral *Pinus sylvestris* L. populations from the Carpathian region. *Dendrobiology* **77**: 105–117.
<http://dx.doi.org/10.12657/denbio.077.009>
- KOMAREK, L. (2018). Hungarian Forest Management Tendencies at the Beginning of the XXI Century. *Russian Journal of Agricultural and Socio-Economic Sciences* **78**(6): 7–18. <https://doi.org/10.18551/rjoas.2018-06.01>
- KSH (2013). Az erdőgazdálkodás jellemzői. *Statisztikai Tükör* **8**(95): 1–6.
- LOCKMAN, I.B. & KEARNS, H.S.J. (2016). *Forest Root Diseases Across the United States*. In: LOCKMAN, I.B. & KEARNS, H.S.J. (eds.): *Gen. Tech. Rep. RMRS-GTR-342*. Ogden, UT, U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, 55 p.
- MACK, N.R., DANIEL, S., LONSDALE, W.M., EVANS, H., CLOUT, M. & BAZZAZ, F. (2000). Biotic Invasions: Causes, Epidemiology, Global Consequences and Control. *Ecological Applications* **10**(3): 689–710.
[https://doi.org/10.1890/1051-0761\(2000\)010\[0689:BICEGC\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0689:BICEGC]2.0.CO;2)
- MAGURA, T., TÓTHMÉRÉSZ, B. & BORDÁN, Z. (2000). Effects of nature management practice on carabid assemblages (Coleoptera: Carabidae) in a non-native plantation. *Biological Conservation* **93**(1): 95–102.
[https://doi.org/10.1016/S0006-3207\(99\)00073-7](https://doi.org/10.1016/S0006-3207(99)00073-7)
- MASON, W.L. & ALÍA, R. (2000). Current and Future Status of Scots pine (*Pinus sylvestris* L.) Forests in Europe. *Forest Systems* **9**: 317–335.
- NEWTON, A.C. & HAIGH, J.M. (1998). Diversity of ectomycorrhizal fungi in Britain: a test of the species-area relationship, and the role of host specificity. *New Phytologist* **138**(4): 619–627.
<https://doi.org/10.1046/j.1469-8137.1998.00143.x>
- PAGONY, H. (1977). A gyökérrontó tapló kártétele a somgyi és az alföldi fenyvesekben. *Az Erdő* **26**: 316–319.
- PAPP, V. & DIMA, B. (2018). New systematic position of *Aurantiporus alborubescens* (Meruliaceae, Basidiomycota), a threatened old-growth forest polypore. *Mycological Progress* **17**(3): 319–332.
<https://doi.org/10.1007/s11557-017-1356-3>
- RYVARDEN, L. & GILBERTSON, R.L. (1993). European polypores. Part 1. *Synopsis Fungorum* **6**: 348–350.
- RYVARDEN, L. & GILBERTSON, R.L. (1994). European polypores. Part 2. *Synopsis*

- Fungorum* 7: 394–743.
- RYVARDEN, L. & MELO, I. (2014). Poroid fungi of Europe. *Synopsis Fungorum* 31: 1–450.
- SARRIONANDIA, E., OLARIAGA, I., PICÓN, R., DUÑABEITIA, M., ROBREDO, A., RODRÍGUEZ, N. & SALCEDO, I. (2015). Implications of exotic *Pinus radiata* plantations for macrofungal diversity in the Urdaibai Biosphere Reserve (northern Spain). *Canadian Journal of Forest Research* 45(6): 667–675.
<https://doi.org/10.1139/cjfr-2014-0317>
- SILLER, I., KUTSZEGI, G., TAKÁCS, K., VARGA, T., MERÉNYI, ZS., TURCSÁNYI, G., ÓDOR, P. & DIMA, B. (2013). Sixty-one macrofungi species new to Hungary in Órség National Park. *Mycosphere* 4(5): 871–924. <https://doi.org/10.5943/mycosphere/4/5/3>
- SZABÓ, I. (2012). Poroid Fungi of Hungary in the Collection of Zoltán Igmándy. *Acta Silvatica et Lignaria Hungarica* 8(1): 113–122.
<https://doi.org/10.2478/v10303-012-0009-0>
- TAMÁS, J. (2003). The history of Austrian pine plantations in Hungary. *Acta Botanica Croatica* 62(2): 147–158.
- TÖRÖK, J. & TÓTH, L. (1996). Composition of bird fauna in Nagy-Szénás Protected Area. *Természetvédelmi Közlemények* 3–4: 65–70.
- VÖLGYI, L. (1955). Az akác és fenyő a somogyi homokon. *Az Erdő* 4: 220–224.

(submitted: 21.08.2019, accepted: 14.11.2019)