

Università degli Studi di Salerno
DIPARTIMENTO DI SCIENZE ECONOMICHE E STATISTICHE

Francesco Giordano – Maria Lucia Parrella

A Locally Adaptive Bandwidth Selector For Kernel Based Regression

WORKING PAPER 3.209

A locally adaptive bandwidth selector for kernel based regression

Francesco Giordano Maria Lucia Parrella

December 10, 2009

Abstract

The selection of the smoothing parameter represents a crucial step in the local polynomial regression, because of the implications on the consistency of the nonparametric regression estimator and because of the difficulties in the implementation of the selection procedure. Moreover, to capture the complexity of the unknown regression curve, a local variable bandwidth is needed, which determines an increase in the efficiency and computational costs of such algorithms. This paper focuses on the problem of the automatic selection of a local bandwidth. We propose a slightly different approach with respect to the traditional ones, which does not require additional computational effort. The empirical performance of the method is shown in the paper through a simulation study.

Keywords: *nonparametric regression; variable bandwidth selection; derivative estimation; neural networks; local polynomials; dependent data.*

1 Context and aims

Nonparametric estimation is particularly useful in the context of model selection. Volatility analysis, trend estimation, forecasts, represent only some of the objectives of statistical analysis which imply the selection of a suitable parametric model for the data generating process. As a consequence, reliable considerations about the risk of misspecification and robustness of the results are generally required. In order to validate the analysis through the empirical evidence, the selection of the parametric model is usually done through the comparison with some nonparametric estimation.

Kernel based estimators represent, maybe, the most used nonparametric tool for the estimation of a regression function. Among them, the Nadaraya-Watson estimator and the local polynomial estimator of order p (linear, quadratic, etc...). The good properties of local polynomial estimators (which include the Nadaraya-Watson estimator as a special case) have been analyzed several times. See, above all, the book of Fan & Gijbels (1996). However, such good properties are often challenged by the misspecification of the tuning parameter, the *bandwidth* of the kernel function, which regulates the smoothness of the estimated

function. The difficulties in specifying such tuning parameter may do vanish at all the advantages in using these nonparametric tools. Therefore, many automatic data-driven bandwidth selection procedures have been proposed so far in the literature. We refer to Fan & Gijbels (1996), Loader (1999) and references therein for a deep review on the topic.

The aim of this work is to propose a new procedure for the automatic selection of the smoothing parameter, based on a slightly different approach with respect to the traditional ones. Just as an example of application, we focus on the problem of estimating the volatility function of dependent data through the local polynomial estimator. However, our procedure can be easily adapted to other contexts, such as the estimation of a generic conditional moment or the estimation of some derivative function, both for dependent and independent data. For this reason, we tried to expose the theory by using a notation the more general and standard as possible, in order to make comparisons easier.

The rationale of our proposal is the following. Consider a local polynomial estimation of a regression function. It is usually suggested to choose the bandwidth by minimizing some estimated measure of the mean squared error (MSE). We start from the *plug-in* point of view, that is to consider the asymptotic MSE and then to estimate the unknown functionals which appear therein. Then we adopt the approach proposed in Giordano & Parrella (2007), that is to estimate such functionals through the neural network technique, in order to avoid the use of pilot bandwidths. Finally, we extend here the result in two directions. From the one hand, we consider the problem of local adaptation of the bandwidth; for this problem we base on the idea of Fan & Gijbels (1995) of splitting the support of the function into subintervals and then estimating global bandwidths on each subinterval. On the other hand, we remove the assumption of homoschedasticity in the model, which is instead assumed by Giordano & Parrella (2007) and Fan & Gijbels (1995). This represents an interesting extension of our previous setup, and it is a must when analyzing economic and financial time series.

In the following two sections we give the assumptions which have to be considered in this paper, together with the notation involved with local polynomial regression. Section 4 gives the definition of the asymptotical optimal bandwidth for the local polynomial estimator. Our procedure is described in detail in section 5. Finally, section 6 reports the results of a simulation study devoted to the assessment of the empirical performance of our procedure and some concluding remarks.

2 Setup

Consider the process $\{Y_t, X_t\}$, where $X_t \in \mathbb{S}_X \subseteq \mathbb{R}$ and $Y_t \in \mathbb{R}$, and define the following nonparametric regression model

$$Y_t = m(X_t) + \sigma(X_t)\varepsilon_t, \quad t = 1, 2, \dots \quad (1)$$

The errors ε_t are real random variables independent from X_t , for which $E(\varepsilon_t) = 0$ and $Var(\varepsilon_t) = 1, \forall t$. Given model (1), we consider the generic problem of

estimating the conditional regression function

$$m_\phi(x) = E \{ \phi(Y_t) | X_t = x \}, \quad (2)$$

which includes several special cases by suitable defining the function $\phi(\cdot)$. Note, for example, that the conditional mean function is given by considering $\phi_1(x) = x$; the conditional second moment function is given by considering $\phi_2(z) = x^2$; the indicator function $\phi_3 = \mathbb{I}(X_t \leq x)$ can be used, instead, in order to define the conditional distribution function, and so on. Finally, model (1) can be extended to the nonparametric ARCH model by putting $X_t = Y_{t-1}$. Given a realization of the process $\{Y_t, X_t; t = 1, \dots, n\}$, the unknown function $m_\phi(\cdot)$ and its derivatives can be estimated nonparametrically using the local polynomial estimators (LPE).

Here we present the assumptions we have to consider for the process.

- (a1) The errors have continuous and positive density function f_ε , and

$$E(\varepsilon_t^2) = 1, \quad E(\varepsilon_t) = E(\varepsilon_t^3) = 0, \quad E|\varepsilon_t|^\delta < \infty,$$

for some $\delta \geq 4$.

- (a2) The function $m(\cdot)$ has continuous second derivative on \mathbb{S}_X . The function $\sigma(\cdot)$ is positive and it has continuous second derivative on \mathbb{S}_X .

- (a3) For dependent data, there exist the constants $M_1 > 0$ e $M_2 > 0$ such that, for $y \in \mathbb{S}_X$

$$|m(y)| \leq M_1(1 + |y|), \quad |\sigma(y)| \leq M_2(1 + |y|), \quad M_1 + M_2 [E|\varepsilon_t|^\delta]^{1/\delta} < 1.$$

- (a4) The density function $f_X(\cdot)$ of the probability measure of the process μ exists, and it is bounded, continuous and positive on \mathbb{S}_X .

Remark 1: Under the assumptions (a1)-(a4), it can be shown that the process is geometrically ergodic (see Härdle & Tsybakov, 1997).

Remark 2: The value of δ must be fixed also considering the particular function $\phi(\cdot)$ in the (2), in order to guarantee that $E|\phi(Y_t)|^2 < \infty$.

3 The fundamentals of local polynomial regression.

Let $K(\cdot)$ be a *kernel function*, supposed to be a density function defined on the interval $[-1, 1]$. Assuming that the derivative of order $(p + 1)$ of the function $m_\phi(X_t)$ exists at the point x , we can use the Taylor expansion to approximate $m_\phi(X_t)$ around the point x through the following polynomial of order p

$$m_\phi(X_t) \approx \sum_{j=0}^p \frac{1}{j!} m_\phi^{(j)}(x) (X_t - x)^j \quad (3)$$

where $m_\phi^{(v)}(x)$ is the derivative of order v of the function m_ϕ evaluated at the point x , for $v = 1, \dots, p$. Define $\beta_j(x) = \frac{1}{j!} m_\phi^{(j)}(x)$, so that $m_\phi^{(j)}(x) = j! \beta_j(x)$, for $j = 0, \dots, p$. Now consider the vector $\boldsymbol{\beta}(x) = \{\beta_0(x), \beta_1(x), \dots, \beta_p(x)\}^T$ of length $p + 1$. Using the (3) and the least squares method, the local polynomial estimator derives an estimate of the vector $\boldsymbol{\beta}(x)$ by solving the following weighted least squares problem:

$$\min_{\boldsymbol{\beta}} \sum_{t=1}^n \left\{ \phi(Y_t) - \sum_{j=0}^p \beta_j(x) (X_t - x)^j \right\}^2 K\left(\frac{X_t - x}{h_n}\right).$$

The positive real number h_n represents the *bandwidth* of the kernel estimator. It regulates, through the kernel function, the amount of local averaging and so the smoothness of the estimated function. It also affects the consistency of the estimator, so its selection seems to be very crucial. For simplicity of notation, later on we will omit in the notation the dependence of β on x and h and the dependence of h on n . Moreover, generally we will indicate explicitly if the bandwidth h depends also on the point x , although it will be clear from the context. Using the matrix notation, it is easy to show that

$$\hat{\boldsymbol{\beta}}(x) = (\mathbf{X}^T \mathbf{W} \mathbf{X})^{-1} \mathbf{X}^T \mathbf{W} \mathbf{Y} \quad (4)$$

$$\hat{m}_\phi^{(v)}(x) = v! \mathbf{u}_v^T \hat{\boldsymbol{\beta}}(x), \quad v = 0, \dots, p, \quad (5)$$

where $\mathbf{Y} = (\phi(Y_1), \dots, \phi(Y_n))$, \mathbf{u}_v is a column vector of length $p + 1$ with all elements equal to zero except for a one in position v , and

$$\mathbf{X} = \begin{pmatrix} 1 & (X_1 - x) & \dots & (X_1 - x)^p \\ \vdots & \vdots & \ddots & \vdots \\ 1 & (X_n - x) & \dots & (X_n - x)^p \end{pmatrix}, \quad \mathbf{W} = \text{diag} \left(K\left(\frac{X_i - x}{h_n}\right) \right)_{n \times n}.$$

The (4) and the (5) represent the most general configuration of the non-parametric local polynomial estimator of a regression function, and it includes several different setups by considering different values for p , v and $\phi(\cdot)$. For example, for $p = 0$ we have the Nadaraya-Watson estimator, which is usually considered as the classic kernel regression estimator. For $p = 1$ we have the local linear estimator and for $p = 2$ the local quadratic estimator. By fixing $v = 0$ we estimate the regression function $m_\phi(x)$. The derivatives of the function $m_\phi(x)$ are estimated by considering $v > 0$, but in this case we must fix $p \geq v$. Moreover, as shown before, by considering a particular function $\phi(\cdot)$ we can obtain estimators of several conditional moments of the process.

As concerning the order of the polynomial fit p , it is known that there is a high correlation between p and the size of the smoothing parameter h . It is difficult (and also useless) to optimize the local polynomial estimation with respect to both this tuning parameters. Generally, one is fixed and the optimization is performed with respect to the other. We choose to fix the

parameter p at a value such that $p - v$ is odd. This because of the advantages shown in Ruppert & Wand (1994).

The asymptotic properties of the local polynomial estimator reported in (5) have been derived by several researchers. For mixing processes, see Masry & Fan (1997) and Härdle & Tsybakov (1997).

4 The Mean Squared Error and the optimal bandwidth.

Most of the bandwidth selectors given in the literature are based on the minimization of some measure of the estimation error, for example the mean squared error (MSE). The optimal bandwidth is then the bandwidth which minimizes the MSE of the estimator

$$h_{MSE}^{opt}(x) = \arg \min_h MSE\{\hat{m}_\phi^{(v)}(x; h)\}. \quad (6)$$

The (6) is considered the theoretical optimal bandwidth, that is the one toward which each bandwidth estimator must converge to. The main difficulty is how to estimate the mean squared error $MSE\{\hat{m}_\phi^{(v)}(x; h)\}$. The usual approach uses the asymptotic expansion of the MSE, as a polynomial in the variable h , in order to evaluate the main asymptotic components and to derive the asymptotic mean squared error (*AMSE*). As usual, the *AMSE* of the local polynomial estimator can be decomposed into the sum of the asymptotic variance and squared bias

$$\begin{aligned} AMSE\{\hat{m}_\phi^{(v)}(x; h)\} &= bias_A^2\{\hat{m}_\phi^{(v)}(x; h)\} + Var_A\{\hat{m}_\phi^{(v)}(x; h)\} \\ &= \mathbb{B}^2(x)h^{2(p+1-v)} + \mathbb{V}(x)\frac{1}{nh^{(2v+1)}} \end{aligned} \quad (7)$$

where

$$\mathbb{B}^2(x) = \left\{C_1 m_\phi^{(p+1)}(x)\right\}^2 \quad \mathbb{V}(x) = \frac{C_2 \sigma_\phi^2(x)}{f_X(x)} \quad (8)$$

Note that the only unknown components in the (8) are the variance function $\sigma_\phi^2(x) = Var\{\phi(Y_t)|X_t = x\}$, the derivative function $m_\phi^{(p+1)}(x)$ and the design density $f_X(x)$. The constant values C_1 and C_2 are known and they depends, obviously, on the kernel function, on the degree of the polynomial p and on the order of the derivative v . In particular, $C_1 = v!B_v/(p+1)!$ and $C_2 = v!^2V_v$, where the values V_v and B_v are derived as follows. Denote with

$$\mu_j = \int_{-\infty}^{+\infty} u^j K(u)du, \quad \nu_j = \int_{-\infty}^{+\infty} u^j K^2(u)du, \quad j = 1, \dots, 2p+1$$

the equivalent moments of the Kernel. Arrange these moments into the $(p+1) \times (p+1)$ matrices \mathbf{S} and $\tilde{\mathbf{S}}$, whose (i, j) -th elements are μ_{i+j-2} and ν_{i+j-2} respectively. Consider also the vector $\mathbf{r} = (\mu_{p+1}, \mu_{p+2}, \dots, \mu_{2p+1})^T$. The numbers V_v and B_v are, respectively, equal to the $(v+1)$ th diagonal element of the matrix $\mathbf{S}^{-1}\tilde{\mathbf{S}}\mathbf{S}^{-1}$ and the $(v+1)$ th element of the vector $\mathbf{S}^{-1}\mathbf{r}$.

The asymptotically optimal bandwidth is the bandwidth which minimizes the right-hand part of the (7). Denote such bandwidth with h_{AMSE}^{opt} . It is given by

$$h_{AMSE}^{opt}(x) = \left\{ \frac{(2v+1)\mathbb{V}(x)}{2n(p-v+1)\mathbb{B}^2(x)} \right\}^{1/(2p+3)}. \quad (9)$$

The *plug-in* method derives an estimation of the optimal bandwidth by estimating the unknown functionals $\mathbb{V}(x)$ and $\mathbb{B}^2(x)$, and plugging them into equation (9). Note that these functionals and the optimal bandwidth $h_{AMSE}^{opt}(x)$ depend on the degree of the estimated derivative v by simple and known constants, so the optimal bandwidth for the estimation of the function $m_\phi(x)$ or for the estimation of some derivative $m_\phi^{(v)}(x)$ can be computed in the same way. Note also that $h_{AMSE}^{opt}(x)$ represents an optimal local bandwidth, given its dependence on the point x , so that we have a local bandwidth function. One way to simplify the problem of estimating the optimal bandwidth is to consider the optimal global bandwidth, that is the bandwidth which minimizes a global measure of the estimation error. In this way one has to estimate a single value instead of a bandwidth function. For example, consider the integrated value of the asymptotical mean squared error, for a given weight function $w(x)$,

$$\begin{aligned} AMISE\{\hat{m}_\phi^{(v)}\} &= \int_{\mathbb{S}_X} \left[Bias_A^2\{\hat{m}_\phi^{(v)}(x;h)\} + Var_A\{\hat{m}_\phi^{(v)}(x;h)\} \right] \omega(x) dF_X \\ &= \mathbb{B}_\omega^2 h^{2(p+1-v)} + \mathbb{V}_\omega \frac{1}{nh^{(2v+1)}} \end{aligned} \quad (10)$$

where

$$\mathbb{B}_\omega^2 = C_1^2 R_f\left(m_\phi^{(p+1)}\right), \quad \mathbb{V}_\omega = C_2 R(\sigma_\phi) \quad (11)$$

$$R_f\left(m_\phi^{(p+1)}\right) = \int_{\mathbb{S}_X} [m_\phi^{(p+1)}(x)]^2 \omega(x) dF_X \quad R(\sigma_\phi) = \int_{\mathbb{S}_X} \sigma_\phi^2(x) \omega(x) dx. \quad (12)$$

Note that the two integrals $R_f\left(m_\phi^{(p+1)}\right)$ and $R(\sigma_\phi)$ involves different measures, because in the second the density $f_X(x)$ cancels with the denominator in the (8). The use of the weight function $\omega(x)$ is useful in order to generalize the selection problem to a wide range of cases, for example to select a bandwidth variable on the support, or to control border effects. It can be shown that in this case the optimal global bandwidth is given by

$$h_{AMISE}^{opt} = \left\{ \frac{(2v+1)\mathbb{V}_\omega}{2n(p-v+1)\mathbb{B}_\omega^2} \right\}^{1/(2p+3)}. \quad (13)$$

Here, as expected, the functionals \mathbb{V}_ω , \mathbb{B}_ω^2 and h_{AMISE}^{opt} , do not depend on the point of estimation x , because they are derived by integration on the support \mathbb{S}_X with respect to the weight function $\omega(x)$. Again, an estimate of the functionals \mathbb{V}_ω , \mathbb{B}_ω^2 must be plugged into the (13) in order to derive the plug-in estimate of the optimal global bandwidth.




Figure 1: The two steps of the procedure proposed for the estimation of the *locally global bandwidth*. The plots report the function $\gamma(x) = \mathbb{V}(x)/\mathbb{B}^2(x)$.

As said before, the estimation of the (9) is more difficult than the estimation of the (13), because the first is a function while the second is a number (=mean value). Anyway, at least asymptotically, it is always true that

$$AMSE\{\hat{m}_\phi^{(v)}(x; h_{AMSE}^{opt}(x))\} \leq AMSE\{\hat{m}_\phi^{(v)}(x; h_{AMISE}^{opt})\}, \forall x \in \mathbb{S}_X,$$

so one could ask in what measure it is convenient to consider the local bandwidth $h_{AMSE}^{opt}(x)$ instead of the global bandwidth h_{AMISE}^{opt} , since the kernel estimators are particularly used for local estimations. Moreover, the estimation of the function h_{AMSE}^{opt} would raise some other questions. For example,

- how to smooth the estimated $\hat{h}_{AMSE}^{opt}(x)$ function?
- what is the effective gain in using the (estimated) local bandwidth?
- is it always convenient to afford a more complicated bandwidth estimation problem?
- how to deal with pilot bandwidths?

These will remain as open questions, although some considerations will be made in the last section of this paper.

5 The Neural Network bandwidth selector for the estimation of the volatility function.

We point to deriving a procedure based on two steps. In the first step we should evaluate the effective gain in using a local bandwidth for the estimation of the

function $m_\phi(x)$ instead of a global bandwidth. Then we should split in some way the support \mathbb{S}_X of the function $m_\phi(x)$ into a number of compact subsets $I_X \in \mathbb{S}_X$. In the second stage we estimate the “*locally global bandwidths*”, that are global bandwidths (=constant) which are estimated on each one of the homogeneous subsets by means of a global optimization. Figure 1 shows the rationale of the method. For the moment, we concentrate on the second step of the procedure.

Suppose we want to estimate the function $m_\phi(x)$ on a given compact subset $I_X \subseteq \mathbb{S}_X$. We are interested in the estimation of a global bandwidth which could be considered optimal on that subset. As seen before, we have to estimate the two functionals in (12) conditionally on the subset I_X . Specific problems follow concerning the integration, since we have to consider a conditioned probability measure. In particular, denoting with $f_{X|I_X}$ the density of the process conditioned on the event $X \in I_X$, we have

$$\begin{aligned} AMISE\{\hat{m}_\phi\} &= \int_{I_X} AMSE\{\hat{m}_\phi(x; h)\} f_{X|I_X}(x) dx \\ &= C_1^2 h^{2(p+1)} \int_{I_X} [m_\phi^{(p+1)}(x)]^2 \frac{f_X(x)}{\mu(I_X)} dx + \frac{C_2}{nh} \int_{I_X} \sigma_\phi^2(x) \frac{dx}{\mu(I_X)} \\ &= C_1^2 h^{2(p+1)} \int_{I_X} [m_\phi^{(p+1)}(x)]^2 f_X(x) d\omega_I + \frac{C_2}{nh} \int_{I_X} \sigma_\phi^2(x) d\omega_I \\ &= \mathbb{B}_{\omega_I} h^{2(p+1)} + \mathbb{V}_{\omega_I} \frac{1}{nh} \end{aligned}$$

where $d\omega_I = \frac{dx}{\mu(I_X)}$ and

$$\mathbb{B}_{\omega_I} = C_1^2 R_f^I(m_\phi^{(p+1)}), \quad \mathbb{V}_{\omega_I} = C_2 R^I(\sigma_\phi) \quad (14)$$

$$R_f^I(m_\phi^{(p+1)}) = \int_{I_X} [m_\phi^{(p+1)}(x)]^2 f_X(x) d\omega_I \quad R^I(\sigma_\phi) = \int_{I_X} \sigma_\phi^2(x) d\omega_I. \quad (15)$$

As usual, the optimal bandwidth can be derived by estimating the functionals in (14) and (15) and plugging them into the (13). In order to do that, the problem is now how to estimate the functions $\sigma_\phi^2(x)$ and $m_\phi^{(p+1)}(x)$ on the subset I_X . We propose to use a global estimator based on the neural network technique. For simplicity, we consider here one particular example of application but the procedure can be easily extended to other cases.

Suppose we want to estimate the volatility function $\sigma^2(x)$ for model (1) and suppose that $m(x) = 0$. This is a very typical set up when analyzing financial data. It results that we have to consider $\phi(Y_t) = Y_t^2$ in (2), (4), (5) and following equations. Let the notation $m_r(x)$ denote the conditional moment function $E(Y^r | X_t = x)$. Note that $\sigma^2(x) \equiv m_2(x)$ and

$$\sigma_\phi^2(x) = Var\{Y_t^2 | X_t = x\} = m_4(x) - m_2^2(x). \quad (16)$$

Generally, the nonparametric estimation of the (16) implies two simultaneous (but different) nonparametric estimations of the functions $m_4(x)$ and $m_2(x)$,

as, for example, in Yao & Tong (1994), Härdle & Tsybakov (1997), Fan & Yao (1998) and Franke & Diagne (2006). In particular, denoting with $\hat{\boldsymbol{\eta}}_i$ a generic nonparametric estimator, we should have

$$\hat{\boldsymbol{\eta}}_i = \arg \min_{\boldsymbol{\eta}} \sum_{t=1}^n [g_i(Y_t) - q(X_t; \boldsymbol{\eta})]^2, \quad i = 1, 2 \quad (17)$$

where

- $g_1(z) = z^4$ for the estimation of $m_4(x)$;
- $g_2(z) = z^2$ for the estimation of $m_2(x)$;
- $q(X_t; \boldsymbol{\eta})$ is some approximation function (neural network function, local polynomial function, etc...).

This procedure is somehow inefficient. For example, when the estimation is (again) based on the use of a kernel estimator, it is necessary to select two different *pilot bandwidths*. To avoid the problem of the pilot bandwidths, we could use the neural network technique, as in Franke & Diagne (2006), but still we would have to consider the two estimations in (17), which is particularly inefficient with neural networks. For this reason, we propose an alternative approach based on the following riparametrization:

$$\sigma_{\phi}^2(x) = m_4(x) - m_2^2(x) = m_2^2(x) [m_{4\varepsilon} - 1] \quad (18)$$

where $m_{4\varepsilon} = E(\varepsilon_t^4)$. Then we use a *Feedforward Neural Network* (FNN), with one input layer and one hidden layer, as approximation function in the (17). It is defined as

$$q(X_t; \boldsymbol{\eta}) = \sum_{k=1}^d c_k \Gamma(a_k X_t + b_k) + c_0, \quad (19)$$

where

- $\boldsymbol{\eta} = (c_0, c_1, \dots, c_d, a_1, \dots, a_d, b_1, \dots, b_d)$ is the vector of parameters of the FNN, to be estimated;
- d is the number of nodes in the hidden layer, such that $d = O(\sqrt{n/\log n})$;
- $\Gamma(\cdot)$ is the *logistic activation function*.

Now using (17), (18) and (19), we define the following estimator of the function $\sigma_{\phi}^2(x)$ based on the neural network approach:

$$\hat{\sigma}_{\phi}^2(x) = \hat{m}_2^2(x) [\hat{m}_{4\varepsilon} - 1] \quad \hat{m}_2(x) = q(x, \hat{\boldsymbol{\eta}}_2) \quad \hat{m}_{4\varepsilon} = \frac{\sum_{t=1}^n X_t^4}{\sum_{t=1}^n [\hat{m}_2(X_t)]^2}. \quad (20)$$

The appeal of the estimator proposed in (20) is that we use only one neural network estimator $\hat{\boldsymbol{\eta}}_2$ in (17). This is, in our opinion, a novel approach for the estimation of the function $\sigma_{\phi}(x)$ which has some interests in its own.

Secondly, we need to estimate the derivative function $m_\phi^{(p+1)}(x)$. Considering that

$$m_\phi^{(p+1)}(x) \equiv m_2^{(p+1)}(x),$$

we can use the same previous NN estimate in order to get the estimate of the derivative function:

$$\hat{m}_2^{(p+1)}(x) = q^{(p+1)}(x; \hat{\boldsymbol{\eta}}_2). \quad (21)$$

Finally, given the (20), the (21) and the ergodicity of the process, we propose the following two estimators for the functionals in (15):

$$\hat{R}_f^I(m_\phi^{(p+1)}) = \frac{\sum_{t=1}^n [\hat{m}_2^{(p+1)}(X_t)]^2}{\sum_{t=1}^n \mathbb{I}(X_t \in I_X)}, \quad (22)$$

$$\hat{R}^I(\sigma_\phi) = \frac{\sum_{i=1}^{n^*} \hat{\sigma}_\phi^2(x_i)}{n^*}, \quad (23)$$

In the (23), the points $\{x_1, x_2, \dots, x_{n^*}\}$ are uniformly spaced values from the interval I_X , with $n^* < n$, such as $n^* = \lfloor n/2 \rfloor$ ($\lfloor x \rfloor$ is the integer part of x). Once again, note that we use only one neural network to estimate both the functionals $R_f^I(m_\phi^{(p+1)})$ and $R^I(\sigma_\phi)$.

6 Numerical performance of the NNB bandwidth selector.

Consider an application of LPE of degree $p = 1$ for the estimation of the volatility function of the following time series models (here $\psi(\cdot)$ denotes the density function of the normal $N(0, 1)$, and $\mathbb{I}(A)$ is the indicator function, which is equal to one if condition A is satisfied and zero elsewhere)

<i>Models</i>	<i>errors</i>
1: $Y_t = 0.7\varepsilon_t$	$\varepsilon_t \sim N(0, 1)$
2: $Y_t = [\psi(Y_{t-1} + 1.2) + 1.5\psi(Y_{t-1} - 1.2)] \varepsilon_t$	$\varepsilon_t \sim N(0, 1)$
3: $Y_t = \sqrt{0.1 + 0.3Y_{t-1}^2} \varepsilon_t$	$\varepsilon_t \sim N(0, 1)$
4: $Y_t = \sqrt{0.1 + 0.15Y_{t-1}^2} \varepsilon_t$	$\varepsilon_t \sim t_{(10)}$
5: $Y_t = \sqrt{0.01 + 0.1Y_{t-1}^2 + 0.2Y_{t-1}^2 \mathbb{I}(Y_{t-1} < 0)} \varepsilon_t$	$\varepsilon_t \sim N(0, 1)$

Model 1 is a simple white noise with variance $\sigma_\varepsilon^2 = 0.49$, $Y_t \sim WN(0, 0.47)$. Models 2-4 are autoregressive and heteroschedastic models, $Y_t \sim ARCH(1)$, where the number 1 in the brackets denotes the lags in the model. The last model is a threshold autoregressive model, $Y_t \sim TAR(1)$. As you can note, all of these models suppose that the conditional mean function is equal to zero. Moreover, only model 1 is homoschedastic. Some of these models are modified versions of models considered in other papers. In particular, Härdle & Tsybakov




Figure 2: Boxplots of the neural network estimates of the error moment $m_{4\epsilon} = E(\epsilon)^4$, for models 1-5.

(1997) for model 2; Franke & Diagne (2006) and Franke, Neumann & Stockis (2004) for model 5.

We performed a Monte Carlo simulation study with 100 replications for each model, with three different lengths: $n = (500, 1000, 2000)$. We considered the estimation of the volatility function $\sigma^2(x)$ onto two different subsets (\equiv intervals): the first is denoted with ${}_1I_X = (x_{11}, x_{21})$ and it was selected in such a way that it includes for each model about the 90% of the central points; the second is indicated with ${}_2I_X = (x_{12}, x_{22})$, and it includes about the 50% of the central points. It follows that ${}_2I_X \subset {}_1I_X$.

We implemented the procedure described in section 5 on the simulated data. As concerning the neural network estimations, the number of nodes in the hidden layer was selected by following an automatic procedure (BIC). See Giordano & Parrella (2007) for further remarks on this topic.

Figure 2 shows empirically the consistency of the method. The plots report the neural network estimations of the error moment $m_{4\epsilon} = E(\epsilon)^4$, for all the models. In our opinion, they show a clear and strong good result for our procedure, if we consider the (20) and if we remember that all the estimations are based on the same neural network estimated approximation function. In particular, note how the estimations improve when $n = 2000$, even for those models which cause some trouble when $n = 500$ (models 4 and 5). More specifically, note that model 4 is equivalent to model 3, but it presents t-student innovations instead of normal innovations. This entails asymmetry and fat tails in the box-plots of the estimations. For model 5, note that the assumption (a2)

$n =$	<i>MISE</i>			<i>MEDISE</i>			<i>SDISE</i>		
	500	1000	2000	500	1000	2000	500	1000	2000
<i>Model 1</i>									
${}_1I_X$	0.03	-0.52	-0.56	0.02	-0.46	-0.51	0.07	-0.74	-0.71
${}_2I_X$	0.01	-0.58	-0.54	0.01	-0.64	-0.49	0.01	-0.60	-0.58
<i>Model 2</i>									
${}_1I_X$	0.05	-0.50	-0.51	0.03	-0.56	-0.29	0.11	-0.50	-0.81
${}_2I_X$	0.01	-0.47	-0.44	0.01	-0.46	-0.44	0.01	-0.46	-0.54
<i>Model 3</i>									
${}_1I_X$	0.30	-0.85	-0.54	0.05	-0.46	-0.47	2.77	-0.95	-0.80
${}_2I_X$	0.04	-0.63	-0.67	0.02	-0.69	-0.58	0.04	-0.46	-0.78
<i>Model 4</i>									
${}_1I_X$	0.06	-0.32	-0.47	0.04	-0.44	-0.35	0.09	-0.05	-0.55
${}_2I_X$	0.02	-0.35	-0.42	0.01	-0.22	-0.44	0.04	-0.49	-0.42
<i>Model 5</i>									
${}_1I_X$	0.08	-0.27	-0.65	0.08	-0.12	-0.86	0.04	-0.15	-0.19
${}_2I_X$	0.04	-0.37	-0.74	0.03	-0.37	-0.84	0.03	-0.31	-0.60

Table 1: Results from the simulation study for the integrated squared errors (ISE) of models 1-5. *MISE* is the mean of ISE; *MEDISE* is the median of ISE; *SDISE* is the standard deviation of ISE. ${}_1I_X$ is the interval including about the 90% of the central points; ${}_2I_X$ is the subset including about the 50% of the central points. Numbers in bold refer to the observed values; the other numbers show the relative variation with respect to the previous value of n .

is not completely satisfied, since the second derivative of the volatility function does not exist for $x = 0$. Anyway, the box-plot in figure 2 shows that there is some robustness of the procedure against the violation of the assumption (a2), at least when the set of points interested has zero probability measure.

In table 1 we report some numerical results concerning the integrated squared errors (*ISE*) of models 1-5, which empirically confirm the consistency of the procedure. In particular, the mean value of the integrated squared error (*MISE*) is reported in the first column, for the three different time series lengths; the median value of the integrated squared error (*MEDISE*) is reported in the second column, while the standard deviation value of the integrated squared error (*SDISE*) is reported in the last column. On the rows, for each model, we report the results concerning the two intervals ${}_1I_X$ (about the 90% of the central points) and ${}_2I_X$ (about the 50% of the central points). The numbers written in bold style represent the observed values. The other numbers show the relative variation with respect to the previous value of n , so they indicate the relative variation observed when the length of time series increases. Note that all these numbers are negative, which means that all the values (mean, median and standard deviation of ISE) tend to zero when n increases, so that the procedure is consistent. As expected, models 4 and 5 show the lowest rate of convergence. Also note that the second row of each model (concerning the interval

${}_2I_X$) generally reports the best results.

In order to better clarify things, we present some plots of the true functions. In figure 3 we report the true optimal bandwidth function $h^{opt}(x)$ and the true asymptotic mean squared error function for the estimation of the volatility function of models 2 and 3. In particular, the first and third rows of plots are devoted to the optimal bandwidth function $h^{opt}(x)$, while the second and fourth rows show the mean squared error. On the columns there are the three different time series lengths, $n = 500$, $n = 1000$ and $n = 2000$. In all the plots of the bandwidth (first and third rows), the black line refers to the optimal local bandwidth (optimum), the red line denotes the global bandwidth (usually used) and the dashed blue line indicates the locally global bandwidth (our proposal). In all the plots of the mean squared error (second and fourth rows), the black line refers to the true MSE function evaluated for the true optimal bandwidth (optimum); the red line denotes the function evaluated for the global bandwidth (usually used); the dashed blue line indicates the function evaluated for the locally global bandwidth (our proposal). Figure 6 is organized in the same way, but they refer to models 4 and 5. From these plots we can note that the locally global bandwidth performs always better than the global bandwidth and always quite as well as the local bandwidth. So generally there is a convenience in using the locally global bandwidth. Anyway, for some model we can see a very little gain, like for models 2 and 4. It seems, therefore, that even the use of the local bandwidth sometimes does not produce significantly better results than the use of a simple global bandwidth. If we consider, moreover, that the (unknown) local bandwidth function must be estimated from the realized time series, we can imagine how furthermore this gain vanish.

Finally, figures 6, 6 and 6, report the boxplots of the local polynomial estimations of the volatility function for models 1-5, when $n = 500$. The red line is the true function. The estimations are made for 20 points in the interval ${}_1I_X$. The bandwidth used is our locally global bandwidth, derived on the interval ${}_1I_X$. The blue box in the middle shows the position of the (included) interval ${}_2I_X$. The plot in the second line shows the same as in the first plot, but this time the estimations are based on the interval ${}_2I_X$. The plots in the third and fourth lines are organized as in the first two lines, but they refer to another model. For the problematic model 5 in figure 7, we report also the plots for the case of $n = 2000$, to show how the estimations improves for longer time series.

References

- Fan J., Gijbels I. (1995) Data-driven bandwidth selection in local polynomial fitting: variable bandwidth and spatial adaptation, *Journal of the Royal Statistical Society, series B*, **57**, 371–394.
- Fan J., Gijbels I. (1996) *Local polynomial modelling and its applications*, Chapman and Hall, London.

- Fan J. and Yao Q. (1998), Efficient estimation of conditional variance functions in stochastic regression, *Biometrika*, **85**, 645–660.
- Franke J. and Diagne M. (2006), Estimating market risk with neural networks, *Statistics & Decision*, **24**, 233–253.
- Franke J., Neumann M. H. and Stockis J. P. (2004), Bootstrapping nonparametric estimators of the volatility function, *Journal of Econometrics*, **118**, 189–218.
- Giordano F. and Parrella M.L. (2007), Neural Networks for bandwidth selection in local linear regression of time series, *Computational Statistics & Data Analysis*, **52**, 2435–2450.
- Härdle W. and Tsybakov A. (1997), Local polynomial estimation of the volatility function in nonparametric autoregression, *Econometrica*, **81**, 223–242.
- Loader C.R. (1999) Bandwidth selection: classical or plug-in?, *Annals of Statistics*, **27**, 2, 415–438.
- Masry E., Fan J. (1997) Local polynomial estimation of regression functions for mixing processes, *Scandinavian Journal of Statistics*, **24**, 165–179.
- Ruppert D., Wand P. (1994) Multivariate Locally Weighted Least Squares Regression, *Annals of Statistics*, **22**, 1346–1370.
- Yao Q., Tong H. (1994) Quantifying the influence of initial values on nonlinear prediction, *Journal of Royal Statistical Society*, **56**, 701–725.




Figure 3: First row of plots: the optimal bandwidth function $h^{opt}(x)$ for the estimation of the volatility function of model 2, respectively for $n = 500$, $n = 1000$ and $n = 2000$ (on the columns). The black line refers to the true function; the red line denotes the global bandwidth; the dashed blue line indicates the locally global bandwidth (our proposal). Second row: the asymptotic Mean Squared Error function, for the three different lengths. The black line refers to the true function evaluated for the true optimal bandwidth (optimum); the red line denotes the function evaluated for the global bandwidth; the dashed blue line indicates the function evaluated for the locally global bandwidth (our proposal). Third and fourth rows: the same as the first two rows, but for model 3.




Figure 4: First row of plots: the optimal bandwidth function $h^{opt}(x)$ for the estimation of the volatility function of model 4, respectively for $n = 500$, $n = 1000$ and $n = 2000$ (on the columns). The black line refers to the true function; the red line denotes the global bandwidth; the dashed blue line indicates the locally global bandwidth (our proposal). Second row: the asymptotic Mean Squared Error function, for the three different lengths. The black line refers to the true function evaluated for the true optimal bandwidth (optimum); the red line denotes the function evaluated for the global bandwidth; the dashed blue line indicates the function evaluated for the locally global bandwidth (our proposal). Third and fourth rows: the same as the first two rows, but for model 5.




Figure 5: First plot on the top: box-plots of the local polynomial estimations of the volatility function for model 1 ($n = 500$). The red line is the true function. The estimations are made for 20 points in the interval ${}_1I_X$. The bandwidth used is our locally global bandwidth, derived on the interval ${}_1I_X$. The blue box in the middle shows the position of the (included) interval ${}_2I_X$. Plot in the second row: the same as above, but this time the estimations are based on the interval ${}_2I_X$. The plots in the third and fourth rows are organized as in the first two rows, but they refer to model 2.




Figure 6: First plot on the top: box-plots of the local polynomial estimations of the volatility function for model 3 ($n = 500$). The red line is the true function. The estimations are made for 20 points in the interval ${}_1I_X$. The bandwidth used is our locally global bandwidth, derived on the interval ${}_1I_X$. The blue box in the middle shows the position of the (included) interval ${}_2I_X$. Plot in the second row: the same as above, but this time the estimations are based on the interval ${}_2I_X$. The plots in the third and fourth rows are organized as in the first two rows, but they refer to model 4.




Figure 7: First plot on the top: box-plots of the local polynomial estimations of the volatility function for model 5 ($n = 500$). The red line is the true function. The estimations are made for 20 points in the interval ${}_1I_X$. The bandwidth used is our locally global bandwidth, derived on the interval ${}_1I_X$. The blue box in the middle shows the position of the (included) interval ${}_2I_X$. Plot in the second row: the same as above, but this time the estimations are based on the interval ${}_2I_X$. The plots in the third and fourth rows are organized as in the first two rows, but for $n = 2000$.

WORKING PAPERS DEL DIPARTIMENTO

- 1988, 3.1 Guido CELLA
Linkages e moltiplicatori input-output.
- 1989, 3.2 Marco MUSELLA
La moneta nei modelli di inflazione da conflitto.
- 1989, 3.3 Floro E. CAROLEO
Le cause economiche nei differenziali regionali del tasso di disoccupazione.
- 1989, 3.4 Luigi ACCARINO
Attualità delle illusioni finanziarie nella moderna società.
- 1989, 3.5 Sergio CESARATTO
La misurazione delle risorse e dei risultati delle attività innovative: una valutazione dei risultati dell'indagine CNR- ISTAT sull'innovazione tecnologica.
- 1990, 3.6 Luigi ESPOSITO - Pasquale PERSICO
Sviluppo tecnologico ed occupazionale: il caso Italia negli anni '80.
- 1990, 3.7 Guido CELLA
Matrici di contabilità sociale ed analisi ambientale.
- 1990, 3.8 Guido CELLA
Linkages e input-output: una nota su alcune recenti critiche.
- 1990, 3.9 Concetto Paolo VINCI
I modelli econometrici sul mercato del lavoro in Italia.
- 1990, 3.10 Concetto Paolo VINCI
Il dibattito sul tasso di partecipazione in Italia: una rivisitazione a 20 anni di distanza.
- 1990, 3.11 Giuseppina AUTIERO
Limiti della coerenza interna ai modelli con la R.E.H..
- 1990, 3.12 Gaetano Fausto ESPOSITO
Evoluzione nei distretti industriali e domanda di istituzione.
- 1990, 3.13 Guido CELLA
Measuring spatial linkages: input-output and shadow prices.
- 1990, 3.14 Emanuele SALSANO
Seminari di economia.

- 1990, 3.15 Emanuele SALSANO
Investimenti, valore aggiunto e occupazione in Italia in contesto biregionale: una prima analisi dei dati 1970/1982.
- 1990, 3.16 Alessandro PETRETTO- Giuseppe PISAURO
Uniformità vs selettività nella teoria della ottima tassazione e dei sistemi tributari ottimali.
- 1990, 3.17 Adalgiso AMENDOLA
Inflazione, disoccupazione e aspettative. Aspetti teorici dell'introduzione di aspettative endogene nel dibattito sulla curva di Phillips.
- 1990, 3.18 Pasquale PERSICO
Il Mezzogiorno e le politiche di sviluppo industriale.
- 1990, 3.19 Pasquale PERSICO
Priorità delle politiche strutturali e strategie di intervento.
- 1990, 3.20 Adriana BARONE - Concetto Paolo VINCI
La produttività nella curva di Phillips.
- 1990, 3.21 Emiddio GALLO
Varianze ed invarianze socio-spaziali nella transizione demografica dell'Italia post-industriale.
- 1991, 3.22 Alfonso GAMBARDELLA
I gruppi etnici in Nicaragua. Autonomia politica ed economica.
- 1991, 3.23 Maria SCATTAGLIA
La stima empirica dell'offerta di lavoro in Italia: una rassegna.
- 1991, 3.24 Giuseppe CELI
La teoria delle aree valutarie: una rassegna.
- 1991, 3.25 Paola ADINOLFI
Relazioni industriali e gestione delle risorse umane nelle imprese italiane.
- 1991, 3.26 Antonio e Bruno PELOSI
Sviluppo locale ed occupazione giovanile: nuovi bisogni formativi.
- 1991, 3.27 Giuseppe MARIGLIANO
La formazione del prezzo nel settore dell'intermediazione commerciale.
- 1991, 3.28 Maria PROTO
Risorse naturali, merci e ambiente: il caso dello zolfo.
- 1991, 3.29 Salvatore GIORDANO
Ricerca sullo stato dei servizi nelle industrie del salernitano.

- 1992, 3.30 Antonio LOPES
Crisi debitoria e politiche macroeconomiche nei paesi in via di sviluppo negli anni 80.
- 1992, 3.31 Antonio VASSILLO
Circuiti economici semplici, complessi, ed integrati.
- 1992, 3.32 Gaetano Fausto ESPOSITO
Imprese ed istituzioni nel Mezzogiorno: spunti analitici e modalità di relazione.
- 1992, 3.33 Paolo COCCORESE
Un modello per l'analisi del sistema pensionistico.
- 1994, 3.34 Aurelio IORI
Il comparto dei succhi di agrumi: un caso di analisi interorganizzativa.
- 1994, 3.35 Nicola POSTIGLIONE
Analisi multicriterio e scelte pubbliche.
- 1994, 3.36 Adriana BARONE
Cooperazione nel dilemma del prigioniero ripetuto e disoccupazione involontaria.
- 1994, 3.37 Adriana BARONE
Le istituzioni come regolarità di comportamento.
- 1994, 3.38 Maria Giuseppina LUCIA
Lo sfruttamento degli idrocarburi offshore tra sviluppo economico e tutela dell'ambiente.
- 1994, 3.39 Giuseppina AUTIERO
Un'analisi di alcuni dei limiti strutturali alle politiche di stabilizzazione nei LCDs.
- 1994, 3.40 Bruna BRUNO
Modelli di contrattazione salariale e ruolo del sindacato.
- 1994, 3.41 Giuseppe CELI
Cambi reali e commercio estero: una riflessione sulle recenti interpretazioni teoriche.
- 1995, 3.42 Alessandra AMENDOLA, M. Simona ANDREANO
The TAR models: an application on italian financial time series.
- 1995, 3.43 Leopoldo VARRIALE
Ambiente e turismo: Parco dell'Iguazù - Argentina.

- 1995, 3.44 A. PELOSI, R. LOMBARDI
Fondi pensione: equilibrio economico-finanziario delle imprese.
- 1995, 3.45 Emanuele SALSANO, Domenico IANNONE
Economia e struttura produttiva nel salernitano dal secondo dopoguerra ad oggi.
- 1995, 3.46 Michele LA ROCCA
Empirical likelihood and linear combinations of functions of order statistics.
- 1995, 3.47 Michele LA ROCCA
L'uso del bootstrap nella verosimiglianza empirica.
- 1996, 3.48 Domenico RANESI
Le politiche CEE per lo sviluppo dei sistemi locali: esame delle diverse tipologie di intervento e tentativo di specificazione tassonomica.
- 1996, 3.49 Michele LA ROCCA
L'uso della verosimiglianza empirica per il confronto di due parametri di posizione.
- 1996, 3.50 Massimo SPAGNOLO
La domanda dei prodotti della pesca in Italia.
- 1996, 3.51 Cesare IMBRIANI, Filippo REGANATI
Macroeconomic stability and economic integration. The case of Italy.
- 1996, 3.52 Annarita GERMANI
Gli effetti della mobilitazione della riserva obbligatoria. Analisi sull'efficienza del suo utilizzo.
- 1996, 3.53 Massimo SPAGNOLO
A model of fish price formation in the north sea and the Mediterranean.
- 1996, 3.54 Fernanda MAZZOTTA
RTFL: problemi e soluzioni per i dati Panel.
- 1996, 3.55 Angela SPAGNUOLO
Concentrazione industriale e dimensione del mercato: il ruolo della spesa per pubblicità e R&D.
- 1996, 3.56 Giuseppina AUTIERO
The economic case for social norms.
- 1996, 3.57 Francesco GIORDANO
Sulla convergenza degli stimatori Kernel.
- 1996, 3.58 Tullio JAPPELLI, Marco PAGANO
The determinants of saving: lessons from Italy.

- 1997, 3.59 Tullio JAPPELLI
The age-wealth profile and the life-cycle hypothesis: a cohort analysis with a time series of cross sections of Italian households.
- 1997, 3.60 Marco Antonio MONACO
La gestione dei servizi di pubblico interesse.
- 1997, 3.61 Marcella ANZOLIN
L'albero della qualità dei servizi pubblici locali in Italia: metodologie e risultati conseguiti.
- 1997, 3.62 Cesare IMBRIANI, Antonio LOPES
Intermediazione finanziaria e sistema produttivo in un'area dualistica. Uno studio di caso.
- 1997, 3.63 Tullio JAPPELLI
Risparmio e liberalizzazione finanziaria nell'Unione europea.
- 1997, 3.64 Alessandra AMENDOLA
Analisi dei dati di sopravvivenza.
- 1997, 3.65 Francesco GIORDANO, Cira PERNA
Gli stimatori Kernel per la stima non parametrica della funzione di regressione.
- 1997, 3.66 Biagio DI SALVIA
Le relazioni marittimo-commerciali nell'imperiale regio litorale austriaco nella prima metà dell'800.
I. Una riclassificazione delle Tafeln zur Statistik der Österreichischen Monarchie.
- 1997, 3.67 Alessandra AMENDOLA
Modelli non lineari di seconda e terza generazione: aspetti teorici ed evidenze empiriche.
- 1998, 3.68 Vania SENA
L'analisi econometrica dell'efficienza tecnica. Un'applicazione agli ospedali italiani di zona.
- 1998, 3.69 Domenico CERBONE
Investimenti irreversibili.
- 1998, 3.70 Antonio GAROFALO
La riduzione dell'orario di lavoro è una soluzione al problema disoccupazione: un tentativo di analisi empirica.
- 1998, 3.71 Jacqueline MORGAN, Roberto RAUCCI
New convergence results for Nash equilibria.

- 1998, 3.72 Rosa FERRENTINO
Niels Henrik Abel e le equazioni algebriche.
- 1998, 3.73 Marco MICOCCI, Rosa FERRENTINO
Un approccio markoviano al problema della valutazione delle opzioni.
- 1998, 3.74 Rosa FERRENTINO, Ciro CALABRESE
Rango di una matrice di dimensione K .
- 1999, 3.75 Patrizia RIGANTI
L'uso della valutazione contingente per la gestione del patrimonio culturale: limiti e potenzialità.
- 1999, 3.76 Annamaria NESE
Il problema dell'inefficienza nel settore dei musei: tecniche di valutazione.
- 1999, 3.77 Gianluigi COPPOLA
Disoccupazione e mercato del lavoro: un'analisi su dati provinciali.
- 1999, 3.78 Alessandra AMENDOLA
Un modello soglia con eteroschedasticità condizionata per tassi di cambio.
- 1999, 3.79 Rosa FERRENTINO
Su un'applicazione della trasformata di Laplace al calcolo della funzione asintotica di non rovina.
- 1999, 3.80 Rosa FERRENTINO
Un'applicazione della trasformata di Laplace nel caso di una distribuzione di Erlang.
- 1999, 3.81 Angela SPAGNUOLO
Efficienza e struttura degli incentivi nell'azienda pubblica: il caso dell'industria sanitaria.
- 1999, 3.82 Antonio GAROFALO, Cesare IMBRIANI, Concetto Paolo VINCI
Youth unemployment: an insider-outsider dynamic approach.
- 1999, 3.83 Rosa FERRENTINO
Un modello per la determinazione del tasso di riequilibrio in un progetto di fusione tra banche.
- 1999, 3.84 DE STEFANIS, PORZIO
Assessing models in frontier analysis through dynamic graphics.
- 1999, 3.85 Annunziato GESUALDI
Inflazione e analisi delle politiche fiscali nell'U.E..
- 1999, 3.86 R. RAUCCI, L. TADDEO
Dalle equazioni differenziali alle funzioni e^x , $\log x$, a^x , $\log_a x$, x^x .

- 1999, 3.87 Rosa FERRENTINO
Sulla determinazione di numeri aleatori generati da equazioni algebriche.
- 1999, 3.88 C. PALMISANI, R. RAUCCI
Sulle funzioni circolari: una presentazione non classica.
- 2000, 3.89 Giuseppe STORTI, Pierluigi FURCOLO, Paolo VILLANI
A dynamic generalized linear model for precipitation forecasting.
- 2000, 3.90 Rosa FERRENTINO
Un procedimento risolutivo per l'equazione di Dickson.
- 2000, 3.91 Rosa FERRENTINO
Un'applicazione della mistura di esponenziali alla teoria del rischio.
- 2000, 3.92 Francesco GIORDANO, Michele LA ROCCA, Cira PERNA
Bootstrap variance estimates for neural networks regression models.
- 2000, 3.93 Alessandra AMENDOLA, Giuseppe STORTI
A non-linear time series approach to modelling asymmetry in stock market indexes.
- 2000, 3.94 Rosa FERRENTINO
Sopra un'osservazione di De Vylder.
- 2000, 3.95 Massimo SALZANO
Reti neurali ed efficacia dell'intervento pubblico: previsioni dell'inquinamento da traffico nell'area di Villa S. Giovanni.
- 2000, 3.96 Angela SPAGNUOLO
Concorrenza e deregolamentazione nel mercato del trasporto aereo in Italia.
- 2000, 3.97 Roberto RAUCCI, Luigi TADDEO
Teoremi ingannevoli.
- 2000, 3.98 Francesco GIORDANO
Una procedura per l'inizializzazione dei pesi delle reti neurali per l'analisi del trend.
- 2001, 3.99 Angela D'ELIA
Some methodological issues on multivariate modelling of rank data.
- 2001, 3.100 Roberto RAUCCI, Luigi TADDEO
Nuove classi di funzioni scalari quasiconcave generalizzate: caratterizzazioni ed applicazioni a problemi di ottimizzazione.
- 2001, 3.101 Adriana BARONE, Annamaria NESE
Some insights into night work in Italy.
- 2001, 3.102 Alessandra AMENDOLA, Marcella NIGLIO

Predictive distributions of nonlinear time series models.

- 2001, 3.103 Roberto RAUCCI
Sul concetto di certo equivalente nella teoria HSSB.
- 2001, 3.104 Roberto RAUCCI, Luigi TADDEO
On stackelberg games: a result of unicity.
- 2001, 3.105 Roberto RAUCCI
Una definizione generale e flessibile di insieme limitato superiormente in \mathfrak{R}^n
- 2001, 3.106 Roberto RAUCCI
Stretta quasiconcavità nelle forme funzionali flessibili.
- 2001, 3.107 Roberto RAUCCI
Sugli insiemi limitati in \mathfrak{R}^m rispetto ai coni.
- 2001, 3.108 Roberto RAUCCI
Monotonie, isotonie e indecomponibilità deboli per funzioni a valori vettoriali con applicazioni.
- 2001, 3.109 Roberto RAUCCI
Generalizzazioni del concetto di debole Kuhn-Tucker punto-sella.
- 2001, 3.110 Antonia Rosa GURRIERI, Marilene LORIZIO
Le determinanti dell'efficienza nel settore sanitario. Uno studio applicato.
- 2001, 3.111 Gianluigi COPPOLA
Studio di una provincia meridionale attraverso un'analisi dei sistemi locali del lavoro. Il caso di Salerno.
- 2001, 3.112 Francesco GIORDANO
Reti neurali per l'analisi del trend: un approccio per identificare la topologia della rete.
- 2001, 3.113 Marcella NIGLIO
Nonlinear time series models with switching structure: a comparison of their forecast performances.
- 2001, 3.114 Damiano FIORILLO
Capitale sociale e crescita economica. Review dei concetti e dell'evidenza empirica.
- 2001, 3.115 Roberto RAUCCI, Luigi TADDEO
Generalizzazione del concetto di continuità e di derivabilità.
- 2001, 3.116 Marcella NIGLIO
Ricostruzione dei dati mancanti in serie storiche climatiche.

- 2001, 3.117 Vincenzo VECCHIONE
Mutamenti del sistema creditizio in un'area periferica.
- 2002, 3.118 Francesco GIORDANO, Michele LA ROCCA, Cira PERNA
Bootstrap variable selection in neural network regression models.
- 2002, 3.119 Roberto RAUCCI, Luigi TADDEO
Insiemi debolmente convessi e concavità in senso generale.
- 2002, 3.120 Vincenzo VECCHIONE
Know how locali e percorsi di sviluppo in aree e settori marginali.
- 2002, 3.121 Michele LA ROCCA, Cira PERNA
Neural networks with dependent data.
- 2002, 3.122 Pietro SENESI
Economic dynamics: theory and policy. A stability analysis approach.
- 2002, 3.123 Gianluigi COPPOLA
Stima di un indicatore di pressione ambientale: un'applicazione ai comuni della Campania.
- 2002, 3.124 Roberto RAUCCI
Sull'esistenza di autovalori e autovettori positivi anche nel caso non lineare.
- 2002, 3.125 Maria Carmela MICCOLI
Identikit di giovani lucani.
- 2002, 3.126 Sergio DESTEFANIS, Giuseppe STORTI
Convexity, productivity change and the economic performance of countries.
- 2002, 3.127 Giovanni C. PORZIO, Maria Prosperina VITALE
Esplorare la non linearità nei modelli Path.
- 2002, 3.128 Rosa FERRENTINO
Sulla funzione di Seal.
- 2003, 3.129 Michele LA ROCCA, Cira PERNA
Identificazione del livello intermedio nelle reti neurali di tipo feedforward.
- 2003, 3.130 Alessandra AMENDOLA, Marcella NIGLIO, Cosimo VITALE
The exact multi-step ahead predictor of SETARMA models.
- 2003, 3.131 Mariangela BONASIA
La dimensione ottimale di un sistema pensionistico: means tested vs programma universale.
- 2003, 3.132 Annamaria NESE
Abitazione e famiglie a basso reddito.

- 2003, 3.133 Maria Lucia PARRELLA
Le proprietà asintotiche del Local Polynomial Bootstrap.
- 2003, 3.134 Silvio GIOVE, Maurizio NORDIO, Stefano SILVONI
Stima della prevalenza dell'insufficienza renale cronica con reti bayesiane: analisi costo efficacia delle strategie di prevenzione secondaria.
- 2003, 3.135 Massimo SALZANO
Globalization, complexity and the holism of the italian school of public finance.
- 2003, 3.136 Giuseppina AUTIERO
Labour market institutional systems and unemployment performance in some Oecd countries.
- 2003, 3.137 Marisa FAGGINI
Recurrence analysis for detecting non-stationarity and chaos in economic times series.
- 2003, 3.138 Marisa FAGGINI, Massimo SALZANO
The reverse engineering of economic systems. Tools and methodology.
- 2003, 3.139 Rosa FERRENTINO
In corso di pubblicazione.
- 2003, 3.140 Rosa FERRENTINO, Roberto RAUCCI
Sui problemi di ottimizzazione in giochi di Stackelberg ed applicazioni in modelli economici.
- 2003, 3.141 Carmine SICA
In corso di pubblicazione.
- 2004, 3.142 Sergio DESTEFANIS, Antonella TADDEO, Maurizio TORNATORE
The stock of human capital in the Italian regions.
- 2004, 3.143 Elena Laureana DEL MERCATO
Edgeworth equilibria with private provision of public good.
- 2004, 3.144 Elena Laureana DEL MERCATO
Externalities on consumption sets in general equilibrium.
- 2004, 3.145 Rosa FERRENTINO, Roberto RAUCCI
Su alcuni criteri delle serie a termini non negativi.
- 2004, 3.146 Rosa FERRENTINO, Roberto RAUCCI
Legame tra le soluzioni di Minty e di Stempacenhia nelle disequazioni variazionali.

- 2004, 3.147 Gianluigi COPPOLA
In corso di pubblicazione.
- 2004, 3.148 Massimo Spagnolo
The Importance of Economic Incentives in Fisheries Management
- 2004, 3.149 F. Salsano
La politica monetaria in presenza di non perfetta osservabilità degli obiettivi del banchiere centrale.
- 2004, 3.150 A. Vita
La dinamica del cambiamento nella rappresentazione del territorio. Una mappa per i luoghi della Valle dell'Irno.
- 2004, 3.151 Celi
Empirical Explanation of vertical and horizontal intra-industry trade in the UK: a comment.
- 2004, 3.152 Amendola – P. Vitale
Self-Assessment and Career Choices: An On-line resource for the University of Salerno.
- 2004, 3.153 A. Amendola – R. Troisi
Introduzione all'economia politica dell'organizzazione: nozioni ed applicazioni.
- 2004, 3.154 A. Amendola – R. Troisi
Strumenti d'incentivo e modelli di gestione del personale volontario nelle organizzazioni non profit.
- 2004, 3.155 Lavinia Parisi
La gestione del personale nelle imprese manifatturiere della provincia di Salerno.
- 2004, 3.156 Angela Spagnuolo – Silvia Keller
La rete di accesso all'ultimo miglio: una valutazione sulle tecnologie alternative.
- 2005, 3.157 Davide Cantarelli
Elasticities of Complementarity and Substitution in Some Functional Forms. A Comparative Review.
- 2005, 3.158 Pietro Coretto – Giuseppe Storti
Subjective Expectations in Economics: a Statistical overview of the main findings.
- 2005, 3.159 Pietro Coretto – Giuseppe Storti
Moments based inference in small samples.

- 2005, 3.160 Massimo Salzano
Una simulazione neo-keynesiana ad agenti eterogeni.
- 2005, 3.161 Rosa Ferrentino
Su alcuni paradossi della teoria degli insiemi.
- 2005, 3.162 Damiano Fiorillo
Capitale sociale: uno o molti? Pochi.
- 2005, 3.163 Damiano Fiorillo
Il capitale sociale conta per outcomes (macro) economici?.
- 2005, 3.164 Damiano Fiorillo – Guadalupi Luigi
*Attività economiche nel distretto industriale di Nocera inferiore – Gragnano.
Un'analisi su Dati Tagliacarne.*
- 2005, 3.165 Rosa Ferrentino
Pointwise well-posedness in vector optimization and variational inequalities.
- 2005, 3.166 Roberto Iorio
La ricerca universitaria verso il mercato per il trasferimento tecnologico e rischi per l'Open Science": posizioni teoriche e filoni di indagine empirica.
- 2005, 3.167 Marisa Faggini
The chaotic system and new perspectives for economics methodology. A note.
- 2005, 3.168 Francesco Giordano
Weak consistent moving block bootstrap estimator of sampling distribution of CLS estimators in a class of bilinear models
- 2005, 3.169 Edgardo Sica
Tourism as determinant of economic growth: the case of south-east asian countries.
- 2005, 3.170 Rosa Ferrentino
On Minty variational inequalities and increasing along rays functions.
- 2005, 3.171 Rosa Ferrentino
On the Minty and Stampacchia scalar variational inequalities
- 2005, 3.172 Destefanis - Storti
A procedure for detecting outliers in frontier estimation
- 2005, 3.173 Destefanis - Storti
Evaluating business incentives through DEA. An analysis on capitalia firm data

- 2005, 3.174 Nese – O'Higgins
In and out of the capitalia sample: evaluating attrition bias.
- 2005, 3.175 Maria Patrizia Vittoria
Il Processo di terziarizzazione in Campania. Analisi degli indicatori principali nel periodo 1981-2001
- 2005, 3.176 Sergio Destefanis – Giuseppe Mastromatteo
Inequality and labour-market performance. A survey beyond an elusive trade-off.
- 2007, 3.177 Giuseppe Storti
Modelling asymmetric volatility dynamics by multivariate BL-GARCH models
- 2007, 3.178 Lucio Valerio Spagnolo – Mario Cerrato
No euro please, We're British!
- 2007, 3.179 Maria Carmela Miccoli
Invecchiamento e seconda transizione demografica
- 2007, 3.180 Maria Carmela Miccoli – Antonio Cortese
Le scuole italiane all'estero: una realtà poco nota
- 2007, 3.181 Rosa Ferrentino
Variational inequalities and optimization problems
- 2007, 3.182 Lavinia Parisi
Estimating capability as a latent variable: A Multiple Indicators and Multiple Causes Approach. The example of health
- 2007, 3.183 Rosa Ferrentino
Well-posedness, a short survey
- 2007, 3.184 Roberto Iorio – Sandrine Labory – Daniele Paci
Relazioni tra imprese e università nel biotech-salute dell'Emilia Romagna. Una valutazione sulla base della co-authorship delle pubblicazioni scientifiche
- 2007, 3.185 Lavinia Parisi
Youth Poverty after leaving parental home: does parental income matter?
- 2007, 3.186 Pietro Coretto – Christian Hennig
Identifiability for mixtures of distributions from a location-scale family with uniform
- 2007, 3.187 Anna Parziale
Il fitness landscape: un nuovo approccio per l'analisi del federalismo fiscale
- 2007, 3.188 Christian Di Pietro – Elena L. del Mercato
Seminal contributions to the theory of Knowledge and technological change

- 2007, 3.189 Valeria D'Amato
Pricing di Opzioni esotiche: Rassegna Teorica e Strumenti Informatici per il Prezzamento
- 2007, 3.190 Roberto Iorio – Sandrine Labory – Daniele Paci
The Determinants of Research Quality in Italy: Empirical Evidence using Bibliometric Data in the Biotech Sector
- 2008, 3.191 Luca Romaniello – Roberto Iorio
Soddisfazione ed insoddisfazione nel lavoro. Determinanti individuali dell'insoddisfazione lavorativa ed analisi dei fattori di disagio. Un'analisi del caso del Triveneto
- 2008, 3.192 Antonio Cortese – Maria Carmela Miccoli
L'immigrazione nei paesi dell'Europa mediterranea: il caso del Portogallo
- 2008, 3.193 Marialuisa Restaino
Dropping out of University of Salerno: a Survival Approach
- 2008, 3.194 Mari Carmela Miccoli
Stranieri sempre più numerosi, con figli sempre più istruiti. Le seconde generazioni nel nostro sistema scolastico
- 2008, 3.195 Carlo Capuano – Giuseppe De Feo
Privatization in oligopoly: the Impact of the shadow cost of public funds
- 2008, 3.196 Giuseppe De Feo
Efficiency gains and mergers
- 2008, 3.197 Maria Olivella Rizza
Gunnar Myrdal's Critiques of Utility Theory. Some implications
- 2008, 3.198 Sergio De Stefanis – Giuseppe Mastromatteo
Winds of change and policies. The nequality-Employment trade-off in the OECD
- 2008, 3.199 Giuseppe Giordano – Michele La Rocca – Maria Prosperina Vitale
Strumenti di analisi per esplorare reti di collaborazione scientifica
- 2008, 3.200 Domenico De Stefano – Giancarlo Ragozzini - Maria Prosperina Vitale
Un approccio di rete all'analisi delle relazioni amicali dei disoccupati nella città di Napoli
- 2008, 3.201 Francesco Giordano
Weak consistent moving block bootstrap estimator for the variance of cls estimators in a class of bilinear models

- 2008, 3.202 Antonio Guariglia
L'evoluzione del regime degli scambi nel commercio internazionale agroalimentare: dal GATT alla WTO
- 2008, 3.203 Giovanni Camillo Porzio – Maria Prosperina Vitale
Assessing Linearity in Structural Equation Models through Graphics
- 2009, 3.204 Antonio Cortese
La rilevazione statistica dei senza tetto e delle alter persone non occupanti un'abitazione
- 2009, 3.205 Roberto Iorio – Daniele Paci
La ricerca in collaborazione con l'industria dei docenti universitari: aggiornamento sugli esiti di un questionario
- 2009, 3.206 Rosamaria D'Amore - Roberto Iorio
Internal and external sources of innovation in the Italian biotech sector
- 2009, 3.207 Maria Carmela Miccoli – Giovanni Ancona – Antonella Biscione
Dinamica demografica, crescita economica e povertà in Albania
- 2009, 3.208 Giuseppina Albano – Francesco Giordano – Cira Perna
Parameter Estimation In Continuous Stochastic Volatility Models

Stampa a cura della C.U.S.L. Cooperativa Universitaria Studio e
Lavoro, Via Ponte Don Melillo, Fisciano
Finito di stampare il 14 dicembre 2009