

Lifestyle approaches for people with intellectual disabilities: A systematic multiple case analysis

Steenbergen H.A^{1,5}, Van der Schans C.P.^{1,2,5}, Van Wijck R.⁴, De Jong J.³, Waninge A.¹

1. Hanze University of Applied Sciences Groningen, Groningen, the Netherlands

2. Department of Rehabilitation Medicine, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

3. Hanze University of Applied Sciences Groningen, School of Sport Studies, Groningen, the Netherlands.

4. Center of Human Movement Sciences, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

5. Department of Health Psychology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

Aim

To analyze the intervention components, levels of influence, explicit use of theory, and conditions for sustainability of currently used lifestyle interventions within lifestyle approaches aiming at physical activity and nutrition in healthcare organizations supporting people with Intellectual Disabilities (ID).

Conclusion

Comprehensive, integrated, and theory-driven approaches at multiple levels should be promoted in healthcare organizations for people with ID.

Background

Healthcare organizations carry out a range of interventions to support and improve a healthy lifestyle.

However, it is difficult to implement and maintain active and healthy lifestyle into daily support.

It is unknown to what extent factors for

Methods

Design: descriptive multiple case study

Research units: 9 healthcare organizations

Data collection: newly developed online inventory form (qualitative data).

Data procedure: analysis at four essential factors for implementation in practice:

Results

Nine healthcare organizations, 59 interventions

- Aims (improve); 31% physical activity, 10% nutrition, 59% a combination of both.
- Aimed at; 49% educational component, 19% daily activities, 16% generic activities, 16% evaluation component
- Target groups; 38% individuals with ID, 37% professionals, 25% social levels
- Structurally embedded; 52%
- Theory-driven: 17%
- Lifestyle policy; 44% clear vision,

implementation are taken into consideration by healthcare organizations.

- Multiple intervention components^{1,2}
- Multiple levels of influence^{3,1,2,4}
- Explicit use of theory^{5,2,4}
- Comprehensive written lifestyle policy^{3,6}
- 67% policy on lifestyle, 67% employees, 56% individual lifestyle-plans
- Funding; between 33% and 100% funded by own organizations,

Table 1. The lifestyle policy, the currently used components targeting the individual, professional, and social level, horizontal, within each healthcare organization, vertical. (i = individuals ID, p = professional, s = social). The gray colored cells represent the presence of a vision and policy on lifestyle, the presence of employees responsible for the theme lifestyle, and the presence of individual lifestyle plans, monitoring of adherence,, as well as the presence of the components per level within the healthcare organization (to read by row). The organization with the most components is listed at the top; we listed the organization with the fewest components at the bottom. The second and third row depict the number of interventions, combined strategies, and separate parts of interventions per healthcare organization.

org	Lifestyle	phy	physical activity										nutrition										both physical activity and nutrition																	
	Vision	vision Policy emplo plans			education			daily living			additional activities			inventory / evaluation			education		daily living			additional activities			inventory /			education			daily living			additional activities			inventory / evaluation			
			yees							activities			evaluation								activities			evaluation								activities		conduction						
					i	р	S	i	р	S	i	р	S	i	р	s	i	р	S	i	р	S	i	р	s	i	р	S	i	р	S	i	р	S	i	р	s	i	р	s
I	Yes	yes	yes																																					
П	Yes	yes		yes																																				
III	yes	yes	yes	yes																																				
IV			yes	yes																																				
V			yes																																					
VI																																								
VII	yes	yes	yes	yes																																				
VIII		yes	yes																																					
IX		yes	yes	yes																																				

References

1.Sallis, J. F., Cervero, R. B., Ascher, W., Henderson, K. A., Kraft, M. K., & Kerr, J. (2006). An ecological approach to creating active living communities. Annual Review of Public Health, 27, 297-322. doi:10.1146/annurev.publhealth.27.021405.102100. 2.Bartholomew L. K., Parcel G. S., Kok G., Gottlieb N. H. & Fernández M. E. (2011) Planning Health Promotion Programs; An Intervention Mapping Approach, Trans. Jossey-Bass, San Francisco, California.. 3. Glasgow RE, Vogt TM, Boles SM, (1999), Evaluating the Public Health Impact of Health Promotion Interventions: The RE-AIM Framework. American Journal of Public Health. September 1999, Vol. 89, No. 9. 4. Naaldenberg, J., Kuijken, N., van Dooren, K., & de Valk, H. V. S. L. (2013). Topics, methods and challenges in health promotion for people with intellectual disabilities: A structured review of literature. Research in Developmental Disabilities, 34(12), 4534–4545.. 5. Gardner B, Whittington C, McAteer J, Eccles MP, Michie S: Using theory to synthesise evidence from behaviour change interventions: the example of audit and feedback. Soc Sci Med 2010, 70:1618–1625.. 6. Goodman RM, Steckler A (1989): A model for the institutionalization of health promotion programs. Fam Community Health 11:63-78.



Contact Details Rianne Steenbergen h.a.steenbergen@pl.hanze.nl

share your talent. move the world.



