

August 1, 1995

Descent equations of Yang–Mills anomalies in noncommutative geometry

Edwin Langmann

Theoretical Physics, Royal Institute of Technology, S-10044 Sweden

Abstract

Consistent Yang–Mills anomalies $\int \omega_{2n-k}^{k-1}$ ($n \in \mathbb{N}$, $k = 1, 2, \dots, 2n$) as described collectively by Zumino’s descent equations $\delta\omega_{2n-k}^{k-1} + d\omega_{2n-k-1}^k = 0$ starting with the Chern character $Ch_{2n} = d\omega_{2n-1}^0$ of a principal $SU(N)$ bundle over a $2n$ dimensional manifold are considered (i.e. $\int \omega_{2n-k}^{k-1}$ are the Chern–Simons terms ($k = 1$), axial anomalies ($k = 2$), Schwinger terms ($k = 3$) etc. in $(2n - k)$ dimensions). A generalization in the spirit of Connes’ noncommutative geometry using a minimum of data is found. For an arbitrary graded differential algebra $\Omega = \bigoplus_{k=0}^{\infty} \Omega^{(k)}$ with exterior differentiation d , form valued functions $Ch_{2n} : \Omega^{(1)} \rightarrow \Omega^{(2n)}$ and $\omega_{2n-k}^{k-1} : \underbrace{\Omega^{(0)} \times \dots \times \Omega^{(0)}}_{(k-1) \text{ times}} \times \Omega^{(1)} \rightarrow$

$\Omega^{(2n-k)}$ are constructed which are connected by generalized descent equations $\delta\omega_{2n-k}^{k-1} + d\omega_{2n-k-1}^k = (\dots)$. Here $Ch_{2n} = (F_A)^n$ where $F_A = d(A) + A^2$ for $A \in \Omega^{(1)}$, and (\dots) is not zero but a sum of graded commutators which vanish under integrations (traces). The problem of constructing Yang–Mills anomalies on a given graded differential algebra is thereby reduced to finding an interesting integration \int on it. Examples for graded differential algebras with such integrations are given and thereby noncommutative generalizations of all Yang–Mills anomalies are found.

1 Introduction

Connes' noncommutative geometry (NCG) is a very influential subject in mathematics unifying notions and ideas from geometry and functional analysis [C1, C2] (for review see [C3]). Since recent it is also receiving broad attention from the physics community. This probably was triggered by a convincing new geometric interpretation of the structure of the standard model including Higgs sector on the classical level based on NCG [CL] (see also [C3] and references therein). Even more interesting, however, is the possibility that NCG could provide a natural mathematical framework for *quantum* field theory, considering that a deeper mathematical understanding of renormalization of 3+1 dimensional quantum gauge theories like QCD has been a main challenge in theoretical physics up to this day. Progress in this direction should not only lead to a better understanding of the nonperturbative structure of these fundamental theories of nature but also provide new calculation tools for analyzing them.

One hint that NCG could be useful in this context is that one of the most widely used regularization schemes for quantum gauge theories — dimensional regularization [tHV] — is done by formally extending the models to $(4 - \epsilon)$ -dimensional spacetime, and that NCG provides the means to naturally extend the setting of Yang Mills gauge theories — connections on a principal $SU(N)$ -bundle over a manifold M — to more general situations without underlying manifold. In fact, all that is required is a graded differential algebra (GDA) (Ω, d) which is a differential complex, $\Omega = \bigoplus_{k=0}^{\infty} \Omega^{(k)}$ with a differentiation $d : \Omega \rightarrow \Omega$ mapping $\Omega^{(k)}$ to $\Omega^{(k+1)}$ such that $d^2 = 0$, carrying a compatible algebra structure (for precise definitions see next Section). Then elements $A \in \Omega^{(1)}$ can be interpreted as generalized Yang–Mills connections, and $X \in \Omega^{(0)}$ as infinitesimal gauge transformations acting on $\Omega^{(1)}$ by $A \rightarrow d(X) + [A, X]$. To formulate also the Yang–Mills action one needs in addition an integration \int and a Hodge- $*$ operation on Ω . In case of ordinary Yang–Mills theory, this GDA is, of course, given by the de Rham forms on the spacetime (or space) manifold M , but there are interesting examples for GDA (Ω, d) based on algebras of Hilbert space operators and which naturally generalize the de Rham forms [C1, C2] (see also next Section).

One important aspect of quantized gauge theories is the occurrence of anomalies and other topological terms like axial anomalies, Schwinger terms, or Chern characters (for review see e.g. [J]). An essential requirement to any regularization procedure used in this context is that it correctly reproduces the anomalies. This is nontrivial — we recall the well-known difficulties to derive, or even formulate, anomalies

(topological terms) in lattice gauge theories (see e.g. [Lü]). The motivation for the present work was an example where a generalized Yang–Mills setting (Ω, d) , as discussed above, has been used successfully for regularizing certain fermion Yang–Mills systems and giving a mathematically rigorous quantum field theory derivation of Schwinger terms [MR, M, L1, LM1] (for a more leisurely discussion of the following see [L3]). One basic idea here is to consider not only the gauge group \mathcal{G} and the set of Yang–Mills fields \mathcal{A} , but certain algebras of Hilbert space operators in which \mathcal{G} and \mathcal{A} can be naturally embedded. The defining relations of these algebras are Schatten ideal conditions which can be regarded as abstract characterizations of the degree of ultraviolet divergences occurring in the spacetime dimension under consideration. Interestingly, this leads exactly to the GDA (Ω, d) playing a fundamental role in NCG [C1]. One is thereby naturally lead to a universal Yang–Mills setting based on such GDA of Hilbert space operators (for another application of this in quantum field theory see [R, FR]). The advantage of using this extended setting for regularization is that it forces one to concentrate on essentials. This leads to very transparent, despite more general, constructions. One remarkable result is an universal Schwinger term in (3+1) dimensions [MR] generalizing the Schwinger term occurring in the Gauss’ law commutators in chiral QCD(3+1) [LM1]. This former Schwinger terms has a natural interpretation as generalization of the latter to NCG if one regards the (conditional) Hilbert space trace as generalization of integration of de Rham forms [LM1, L2]. Having this one example where NCG appears in a quantum field theory derivation of anomalies, it is natural to ask whether also other anomalies have similar generalizations to the universal Yang Mills setting, or, more generally, to other GDA. The structure of these generalized anomalies can be expected to directly hint to how they arise from regularizing a quantum gauge theory. For example, the regularization procedure used in [L1] to derive the anomalous Gauss’ law commutators in chiral QCD trivially generalizes from (3+1) to higher dimensions once the higher dimensional universal Schwinger terms are known.

In this paper we give a very general answer to this question treating all kinds of anomalies and arbitrary GDA at once. We show that certain ‘raw data’ for all anomalies and topological terms can be constructed in any GDA (Ω, d) , and these give rise to anomalies (nontrivial cocycles) if and only if there is a nontrivial integration on (Ω, d) . Since the GDA associated with the universal Yang–Mills setting have such nontrivial integrations (the conditional Hilbert space trace), we obtain the universal generalizations of all anomalies as a special case (including, of course, all higher dimensional universal Schwinger terms).

Since many (perhaps all) interesting gauge invariant regularization schemes for

Yang–Mills systems are based on some GDA, our result should be of fundamental interest for quantum gauge theories. It also provides an explanation how anomalies with their rich differential geometric structure can arise from explicit field theory calculations. The latter are done by calculating Feynman diagrams (see [J] and references therein) which can be interpreted as (regularized) traces of certain Hilbert space operators [LM2]; our result makes explicit that the differential geometric structure of anomalies can be present already on the level of Hilbert space operators. We feel that this adds strong support to the expectation that NCG is relevant for quantum gauge theories.

To be more specific, we recall Zumino’s descent equations $\delta\bar{\omega}_{2n-k}^{k-1} + d\bar{\omega}_{2n-k-1}^k = 0$ [Z] providing a concise summary and intriguing relations between all Yang–Mills anomalies (the reason why we put the bar over ω will become clear immediately). These relations imply that the integrals $\int_M \bar{\omega}_{2n-k}^{k-1}$ over $(2n-k)$ dimensional manifolds M without boundary are $(k-1)$ -cocycles, and these are exactly the Chern Simons terms ($k=1$), axial anomalies ($k=2$), Schwinger terms ($k=3$) etc., occurring in $(2n-k)$ dimensional gauge theories. Our result is a generalization of Zumino’s descent equations to all GDA (Ω, d) . For all $X_i \in \Omega^{(0)}$ and $A \in \Omega^{(1)}$ we construct forms $\omega_{2n-k}^{k-1}(X_1, \dots, X_{k-1}; A) \in \Omega^{(2n-k)}$ which are multilinear and antisymmetric in the X_i and obey the generalized descent equations

$$\delta\omega_{2n-k}^{k-1} + d\omega_{2n-k-1}^k = (\dots) \quad k = 1, 2, \dots, 2n, \quad (n \in \mathbb{N}) \quad (1)$$

(for definition of δ see (23), for full equations (29a–d)) where (\dots) is not zero but a sum of graded commutators which vanish under integrations \int . It is interesting to note that these descent equations generalize Zumino’s even for the case of de Rham forms: we recall that the forms mentioned above are traces of matrix valued forms, $\bar{\omega}_{2n-k}^{k-1} = \text{tr}_N(\omega_{2n-k}^{k-1})$, and our equations are without the matrix trace tr_N (they generalize Zumino’s since the terms (\dots) vanish under tr_N). This our generalization is natural since tr_N , which we regard as partial integration, is not available for general GDA.

As mentioned, these forms ω_{2n-k}^{k-1} are raw data from which the anomalies are obtained by integration \int ; the descent equations and basic properties of integration guarantee that they obey the appropriate consistency conditions $\delta\int\omega_{2n-k}^{k-1} = 0$ (= cocycle relations). Thus our result reduces the question of existence of anomalies for general GDA (Ω, d) to existence of a nontrivial integration \int on it. We stress that this latter question is still very nontrivial. In fact, our result makes explicit that all topological information is contained in such an integration: the raw data for

topological terms ω_{2n-k}^{k-1} *always* exist and give rise to $(k-1)$ -cocycles by integration \int ; whether these are nontrivial solely depends on the integration \int and has to be checked for each case individually.

We note that there are other generalizations of Zumino's descent equations in the literature [CM, KS]. These generalize the equation for $\bar{\omega}_{2n-k}^{k-1}$ and therefore also need the notation of (partial) integration. Our result is more general since it does not require an integration. In fact, we believe that it is the most general result in this direction and very much in the spirit of NCG since it uses a minimum amount of input data and strips the descent equations from a differential geometric setting to the bones of simple algebraic relations. (Note also that we do not have a graded double complex since $\delta d + d\delta$ is not zero on forms ω_{2n-k}^{k-1} but only on $\bar{\omega}_{2n-k}^{k-1}$!) Also, our proof is very simple and direct — we give an explicit formula for the forms ω_{2n-k}^{k-1} and prove by explicit calculation that our generalized descent equations are obeyed. We did not try to find a geometric interpretation of these equations (1), but since the nontrivial r.h.s. (\cdots) have a very simple, suggestive form (cf. (29a-d)), we believe that such an interpretation is possible and should be interesting.

There are two important examples for GDA of bounded Hilbert space operators which generalize the de Rham forms to NCG. In one differentiation is given by the graded commutator with the Dirac operator and integration by Dixmier trace [C2], in the other differentiation is graded commutation with the *sign* of the Dirac operator and integration is the Hilbert space trace [C1] (see also [C3]). In this paper we use only the latter as example since it is this GDA which naturally appears in quantum field theory. Also, the noncommutative generalization of anomalies to the former is discussed in [CM].

To avoid confusion, we already point out here that we use Grassmann numbers θ_i for writing down the formulas for the forms ω_{2n-1}^{k-1} , but this is only a matter of convenience and one could do without at the cost of having longer expressions and proofs. However, these Grassmann numbers are a convenient means to keep track of the combinatorics and order of terms — they are a substitute for the symmetrized trace used by Zumino [Z] but which is not available in our general setting.

The plan of this paper is as follows. To make the paper self-contained and have nontrivial examples to which our result can be applied, we summarize the basis definitions and examples of GDA and integrations in Section 2. This Section is also meant as an introduction to basic notions from NCG [C3] relevant for quantum field theory and should be useful for physicists. Of mathematical interest might be our definition of partial integration on GDA (generalizing integration of de Rham forms over manifolds with boundary) which seems somewhat different from the one in [C3].

Our main result and its proof is in Section 3. In the final Section we discuss some interesting special cases including the universal anomalies mentioned.

Notation: We denote as \mathbb{N}_0 the non-negative integers, \mathbb{C} the complex numbers, \mathfrak{gl}_N the complex algebra of $N \times N$ matrices with product the usual matrix multiplication, $[a, b]_{\pm} = ab \pm ba$, $[a, b] = ab - ba$ and $\{a, b\} = ab + ba$.

2 Prerequisites

2.1 Graded differential algebra (GDA)

A GDA (Ω, d) is a \mathbb{N}_0 -graded associative algebra over the \mathbb{C} , $\Omega = \bigoplus_{k=0}^{\infty} \Omega^{(k)}$, with a linear map $d : \Omega \rightarrow \Omega$ such that $d^2 = 0$, and for all $\omega_{k,\ell} \in \Omega^{(k,\ell)}$, $k, \ell \in \mathbb{N}_0$,

$$\omega_k \omega_\ell \in \Omega^{(k+\ell)}, \quad d\omega_k \in \Omega^{(k+1)}, \quad d(\omega_k \omega_\ell) = d(\omega_k) \omega_\ell + (-)^k \omega_k d(\omega_\ell). \quad (2)$$

(we write the algebra product in Ω as $(\omega, \omega') \rightarrow \omega \omega'$).

We denote elements $\omega_k \in \Omega^{(k)}$ as k -forms and d as exterior differentiation. Note that by definition, all linear combinations of elements $u_0 du_1 \cdots du_k$ with $u_i \in \Omega^{(0)}$ are k -forms.

Example 1: The simplest example for a GDA is the de Rham GDA $(\Omega_{\text{dR},d}, d)$ of de Rham forms on \mathbb{R}^d which we define as follows: $\Omega_{\text{dR},d}^{(0)} = C_0^\infty(\mathbb{R}^d; \mathfrak{gl}_N)$ is the algebra of \mathfrak{gl}_N -valued C_0^∞ functions on \mathbb{R}^d , and for $k \geq 1$, $\Omega_{\text{dR},d}^{(k)}$ is the vector space generated by elements $u_0 du_1 \cdots du_n$, $u_i \in \Omega_{\text{dR},d}^{(0)}$, where d is the usual exterior differentiation of de Rham forms.

Example 2: Let \mathcal{H} be a separable Hilbert space. We denote as \mathcal{B} and \mathcal{B}_1 the bounded and trace class operators on \mathcal{H} , respectively, and $\mathcal{B}_p = \{a \in \mathcal{B} \mid (a^* a)^{p/2} \in \mathcal{B}_1\}$ for $p > 0$ are the Schatten classes. We recall that $a \in \mathcal{B}_{\frac{p}{2}}$, $b \in \mathcal{B}_{\frac{q}{2}}$ and $n, p, q > 0$ implies $ab \in \mathcal{B}_{\frac{n}{p+q}}$ [S]. Let $\varepsilon \in \mathcal{B}$ be a grading operator, i.e. $\varepsilon = \varepsilon^* = \varepsilon^{-1}$. We will use the notation

$$a = a_+ + a_-, \quad a_{\pm} = \frac{1}{2}(a \pm \varepsilon a \varepsilon) \quad \forall a \in \mathcal{B} \quad (3)$$

so that $(ab)_+ = a_+ b_+ + a_- b_-$, $(ab)_- = a_+ b_- + a_- b_+$.

Let $d \in \mathbb{N}$. We define

$$p = \frac{1}{2}(d + 1) \quad (4a)$$

and

$$\hat{\Omega}_d^{(0)} = \{a \in \mathcal{B} \mid a_- \in \mathcal{B}_{2p}\}$$

$$\begin{aligned}
\hat{\Omega}_d^{(2k-1)} &= \left\{ a \in \mathcal{B} \mid a_+ \in \mathcal{B}_{\frac{2p}{2k}}, a_- \in \mathcal{B}_{\frac{2p}{2k-1}} \right\} \\
\hat{\Omega}_d^{(2k)} &= \left\{ a \in \mathcal{B} \mid a_+ \in \mathcal{B}_{\frac{2p}{2k}}, a_- \in \mathcal{B}_{\frac{2p}{2k+1}} \right\} \\
& \quad k = 1, 2, \dots
\end{aligned} \tag{4b}$$

Then

$$\hat{d}\omega_k \equiv i \left(\varepsilon \omega_k - (-)^k \omega_k \varepsilon \right) \quad \forall \omega_k \in \hat{\Omega}_d^{(k)} \tag{4c}$$

defines an exterior differentiation \hat{d} on $\hat{\Omega}_d = \bigoplus_{k=0}^{\infty} \hat{\Omega}_d^{(k)}$ which makes $(\hat{\Omega}_d, \hat{d})$ into a GDA.

Note that the multiplication in this GDA is equal to the operator product in \mathcal{B} (the latter follows from $\varepsilon^2 = 1$ which implies $\varepsilon[\varepsilon, a]\varepsilon = -[\varepsilon, a]$ for all $a \in \mathcal{B}$).

For $d = 2n$ even, we introduce another grading operator Γ on \mathcal{H} which anticommutes with ε , $\varepsilon\Gamma = -\Gamma\varepsilon$. For the introduction of integrations below we need to restrict $(\hat{\Omega}_d, \hat{d})$ to a subalgebra $(\hat{\Omega}_{\text{dR},d}, \hat{d})$ defined as $\hat{\Omega}_{\text{dR},d} = \bigoplus_{k=0}^{\infty} \hat{\Omega}_{\text{dR},d}^{(k)}$,

$$\hat{\Omega}_{\text{dR},d}^{(k)} = \left\{ \omega_k \in \hat{\Omega}_d^{(k)} \mid \omega_k \Gamma - (-)^k \Gamma \omega_k = 0 \right\} \quad (d = 2n) \tag{5}$$

(it is easy to see that this indeed defines a GDA). To simplify notation we write $\Gamma^{d-1} = \Gamma$ and 1 (identity) for d even and odd, respectively, and for d odd, $\hat{\Omega}_{\text{dR},d} = \hat{\Omega}_d$.

The above mentioned embedding of the de Rham GDA $(\Omega_{\text{dR},d}, d)$ in $(\hat{\Omega}_{\text{dR},d}, \hat{d})$ is as follows. Consider the Hilbert space

$$\mathcal{H} = L^2(\mathbb{R}^d) \otimes \mathbb{C}_{spin}^{\nu} \otimes \mathbb{C}^N \tag{6}$$

where $\nu = 2^{\lfloor d/2 \rfloor}$ can be interpreted as the number of spin indices ($\lfloor d/2 \rfloor = (d-1)/2$ for d odd and $\lfloor d/2 \rfloor = d/2$ for d even). With γ_i the usual selfadjoint γ -matrices acting on \mathbb{C}_{spin}^{ν} and obeying $\gamma_i \gamma_j + \gamma_j \gamma_i = 2\delta_{ij}$ (for explicit formulas see e.g. [L2]), the free Dirac operator $\mathcal{D}_0 = \sum_{i=1}^d \gamma_i (-i) \frac{\partial}{\partial x_i}$ on \mathbb{R}^d defines a selfadjoint operator on \mathcal{H} which we also denote as \mathcal{D}_0 . Note that for $d = 2n$ even we have an additional γ -matrix γ_{d+1} which defines a grading operator Γ on \mathcal{H} .

We recall that every $X \in C_0^{\infty}(\mathbb{R}^d, \mathfrak{gl}_N)$ can be naturally identified with a bounded operator \hat{X} on \mathcal{H} , $(\hat{X}f)(x) = X(x)f(x)$ for all $f \in \mathcal{H}$ ($x \in \mathbb{R}^d$).

We define $\varepsilon = \text{sign}(\mathcal{D}_0)$ where $\text{sign}(x) = +1(-1)$ for $x \geq 0$ ($x < 0$) (using the spectral theorem for self-adjoint operators [RS]). For $d = 2n$ even, $\Gamma = \gamma_{d+1}$ anticommutes with ε , and it obviously commutes with all \hat{X} for $X \in C_0^{\infty}(\mathbb{R}^d, \mathfrak{gl}_N)$.

One can prove

$$X \in C_0^\infty(\mathbb{R}^d, \mathfrak{gl}_N) \Rightarrow \hat{X} \in \hat{\Omega}_{\text{dR},d}^{(0)} \quad (7a)$$

(see e.g. [MR]) implying that

$$X_0 dX_1 \cdots X_k \mapsto (\mathbf{i})^k \hat{X}_0[\varepsilon, \hat{X}_1] \cdots [\varepsilon, \hat{X}_k] \quad \forall X_i \in \Omega_{\text{dR},d}^{(k)} \quad (7b)$$

provides an embedding $\omega \mapsto \hat{\omega}$ of $(\Omega_{\text{dR},d}, d)$ in $(\hat{\Omega}_{\text{dR},d}, \hat{d})$

We shall refer to $(\hat{\Omega}_{\text{dR},d}, \hat{d})$ as *universal de Rham GDA*.

2.2 Integrations

A GDA with integration is a triple (Ω, d, f) such that (Ω, d) is a GDA, and there is a linear map $f : \Omega \rightarrow \mathbb{C}$ obeying

$$\int d\omega_k = 0, \quad \int \omega_k \omega_\ell = (-)^{k\ell} \int \omega_\ell \omega_k \quad \forall \omega_{k,\ell} \in \Omega^{(k,\ell)}. \quad (8)$$

(The first of these relations is Stokes' theorem and the second graded commutativity of forms under the integration.) If f is nonzero only on $\Omega^{(d)}$ for some $d \in \mathbb{N}_0$ we say that Ω has dimension d and indicate this by writing $\Omega = \Omega_d$.

Remark: A GDA with integration is called *cycle* in [C3].

Example 1: The de Rham GDA $(\Omega_{\text{dR},d}, d)$ has the natural integration defined as $\int \omega_k = 0$ for $\omega_k \in \Omega_{\text{dR},d}^{(k \neq d)}$ and

$$\int \omega_d = \int_{\mathbb{R}^d} \text{tr}_N(\omega_d) \quad \forall \omega_d \in \Omega_{\text{dR},d}^{(d)} \quad (9)$$

(integration of de Rham forms as usual; $\text{tr}_N(\cdot)$ is the trace of $N \times N$ matrices).

Example 2: We define the conditional trace as

$$\text{Tr}_C(a) \equiv \text{Tr}(a_+) \quad \text{for all } a \in \mathcal{B} \text{ with } a_+ \in \mathcal{B}_1 \quad (10)$$

where $\text{Tr}_C(\cdot)$ is the usual Hilbert space trace on \mathcal{H} . Then $\hat{\int} \omega_k = 0$ for all $\omega_k \in \hat{\Omega}_{\text{dR},d}^{(k \neq d)}$ and

$$\hat{\int} \omega_d = \frac{1}{c_d} \text{Tr}_C(\Gamma^{d-1} \omega_d) \quad \forall \omega_d \in \hat{\Omega}_{\text{dR},d}^{(d)}, \quad (11a)$$

where

$$c_d = (2\mathbf{i})^{[d/2]} \frac{1}{d(2\pi)^d} \frac{2\pi^{d/2}}{\Gamma(d/2)}. \quad (11b)$$

is a convenient normalization constant (explained below), defines an integration on $(\hat{\Omega}_{\text{dR},d}, \text{d})$. Indeed, for $\omega_d \in \hat{\Omega}_{\text{dR},d}^{(d)}$,

$$(\Gamma^{d-1} \omega_d)_+ = \begin{cases} (\omega_d)_+ & \text{for } d \text{ odd} \\ \Gamma(\omega_d)_- & \text{for } d \text{ even} \end{cases}$$

which is \mathcal{B}_1 by definition. The relations (8) can be easily checked. (The first relation of (8) (Stokes' theorem) is trivial since

$$(\Gamma^{d-1} \hat{\text{d}}\omega_{d-1})_+ = -\frac{i}{2} \Gamma^{d-1} \varepsilon \hat{\text{d}}^2(\omega_{d-1}) = 0$$

The second follows from

$$\left(\Gamma^{d-1} (\omega_k \omega_{d-k} - (-)^{k(d-k)} \omega_{d-k} \omega_k) \right)_+ = [(\Gamma^{d-1} \omega_k)_+, (\omega_{d-k})_+] + [(\Gamma^{d-1} \omega_k)_-, (\omega_{d-k})_-]$$

and cyclicity of trace (recall that $\text{Tr}([a, b]) = 0$ if at least one of $a, b \in \mathcal{B}$ is compact and ab and ba are \mathcal{B}_1). Note that in case d is even, we used that $\Gamma \omega_k = (-)^k \omega_k \Gamma$; this is why we had to restrict ourselves from $\hat{\Omega}_d$ to $\hat{\Omega}_{\text{dR},d}$.)

The embedding $\omega \mapsto \hat{\omega}$ of de Rham GDA $(\Omega_{\text{dR},d}, \text{d})$ in $(\hat{\Omega}_{\text{dR},d}, \hat{\text{d}})$ as discussed above naturally extends to these integrations. One has

$$\begin{aligned} i^d \text{Tr}_C \left(\Gamma^{d-1} \hat{X}_0[\varepsilon, \hat{X}_1] \cdots [\varepsilon, \hat{X}_d] \right) &= c_d \int \text{tr}_N (X_0 \text{d} X_1 \cdots \text{d} X_d) \\ &\quad \forall X_i \in C_0^\infty(\mathbb{R}^d; \mathfrak{gl}_N) \end{aligned} \quad (12)$$

(for an elementary proof see [L2]) which shows that $\hat{f}\hat{\omega} = f\omega$ for all $\omega \in \Omega_{\text{dR},d}$.

2.3 Partial Integrations

A GDA with partial integration structure (GDAPI) is a quintuple $(\Omega, \text{d}, f, {}^*\Omega, \partial)$ such that, (i) (Ω, d) is a GDA, (ii) ${}^*\Omega = \bigoplus_{k=0}^\infty {}^*\Omega^{(-k)}$ is graded cycle, i.e. a \mathbb{N}_0 -graded vector space with a linear map $\partial : {}^*\Omega \rightarrow {}^*\Omega$ such that $\partial Q^{(-k)} \in {}^*\Omega^{(-k-1)}$ for all $Q^{(-k)} \in {}^*\Omega^{(-k)}$ and $\partial^2 = 0$, (iii) there is a bilinear map $f : {}^*\Omega \times \Omega \rightarrow \mathbb{C}$, $(Q, \omega) \mapsto \int_Q \omega$ such that

$$\int_Q \text{d}\omega_k = \int_{\partial Q} \omega_k, \quad \int_Q \omega_k \omega_\ell = (-)^{k\ell} \int_Q \omega_\ell \omega_k \quad (13)$$

for all $\omega_{k,\ell} \in \Omega^{(k,\ell)}$ and $Q \in {}^*\Omega$. (The first of these is the generalization of Stokes' theorem for manifolds with boundaries.) A GDAPI is of dimension d if f is nonzero only on $\bigoplus_{k=0}^d {}^*\Omega^{(-k)} \times \Omega^{(d-k)}$, and we write in this case $\Omega = \Omega_d$ and ${}^*\Omega = {}^*\Omega_d$.

In our examples below, GDAPI $(\mathcal{S}, d, f, {}^*\mathcal{S}, \partial)$ arise from a given GDA (Ω, d, f) with integration as follows. (i) (\mathcal{S}, d, f) is a subcomplex of, i.e. a GDA contained in, (Ω, d, f) , (ii) ${}^*\mathcal{S} = \bigoplus_{k=0}^{\infty} {}^*\mathcal{S}^{(-k)}$ is a graded complex such that for each $Q \in {}^*\mathcal{S}$ and $0 < \epsilon < 1$ there is a $\eta_\epsilon(Q) \in \Omega$ such that

$$\lim_{\epsilon \searrow 0} \int \eta_\epsilon(Q) \omega \quad (14a)$$

exist for all $\omega \in \mathcal{S}$ and, (iii) for all $\omega_{k,\ell} \in \mathcal{S}^{(k,\ell)}$,

$$\lim_{\epsilon \searrow 0} \int \eta_\epsilon(Q) (\omega_k \omega_\ell - (-)^{k\ell} \omega_\ell \omega_k) = 0. \quad (14b)$$

We then set

$$\int_Q \omega \equiv \lim_{\epsilon \searrow 0} \int \eta_\epsilon(Q) \omega, \quad (15)$$

and ∂Q is defined by

$$\eta_\epsilon(\partial Q)_{-k} \equiv (-)^{-k-1} d\eta_\epsilon(Q)_{-k} \quad (16)$$

(we use the notation $\Omega \ni \eta = \sum_k \eta_{-k}$ with $\eta_{-k} \in \Omega^{(-k)}$). Stokes' theorem in (13) then trivially follows from $\int d(\eta_\epsilon(Q)\omega) = 0$. Thus it is (14b) which is the nontrivial condition determining whether \mathcal{S} and ${}^*\mathcal{S}$ are compatible so as to form a GDAPI.

Obviously $(\mathcal{S}, d, f, {}^*\mathcal{S}, \partial)$ is of dimension d if (Ω, d, f) is.

Remark: One can identify elements in ${}^*\mathcal{S}$ with equivalence classes $Q = \eta / \sim$ of maps $\eta : (0, 1) \rightarrow \Omega, \epsilon \mapsto \eta_\epsilon$ such that $\lim_{\epsilon \searrow 0} \int \eta_\epsilon \omega$ exist for all $\omega \in \mathcal{S}$, with the equivalence relation

$$\eta \sim \eta' \Leftrightarrow \lim_{\epsilon \searrow 0} \int (\eta_\epsilon - \eta'_\epsilon) \omega = 0 \quad \forall \omega \in \mathcal{S}.$$

Example 1: Let $\mathcal{D}_d^{(0)}$ be the set of d -dimensional compact submanifolds D of \mathbb{R}^d with boundary ∂D which is a $(d-1)$ -dimensional compact C^∞ manifold, and $\mathcal{D}_d^{(-k)}$ be the set of all $(d-k)$ -dimensional compact C^∞ manifolds of the form $D^{(-k)} = D_0 \cap \partial D_1 \cap \cdots \cap \partial D_k$ with $D_i \in \mathcal{D}_d^{(0)}$. Then $(\Omega_{d\mathbb{R},d}, d, f, \mathcal{D}, \partial)$ is a GDAPI with ∂ the usual boundary operation and $\int_{D^{(-k)}} \omega_{d-k} \equiv \int_{D^{(-k)}} \text{tr}_N(\omega_{d-k})$ the integration of an $(d-k)$ -form over an k -dimensional submanifold $D^{(-k)}$ of \mathbb{R}^d as usual. With $\chi_D(x)$ the characteristic function of $D \in \mathcal{D}_d^{(0)}$ ($\chi_D(x) = 1$ for $x \in D$ and $= 0$ otherwise) we can formally write

$$\int_{D_0 \cap \partial D_1 \cap \cdots \cap \partial D_k} \text{tr}_N(\omega_{d-k}) = \int_{\partial D_1 \cap \cdots \cap \partial D_k} \text{tr}_N(\chi_{D_0} \omega_{d-k}) =$$

$$\begin{aligned}
& \int_{D_1 \cap \partial D_2 \cap \dots \cap \partial D_k} \mathrm{tr}_N \left(d \left(\chi_{D_0} \omega_{d-k} \right) \right) = \dots \\
& = \int_{\mathbb{R}^d} \mathrm{tr}_N \left(\chi_{D_k} d \chi_{D_{k-1}} \dots d \chi_{D_1} d \left(\chi_{D_0} \omega_{d-k} \right) \right) \\
& = (-)^k \int_{\mathbb{R}^d} \mathrm{tr}_N \left(d \chi_{D_k} \dots d \chi_{D_1} \chi_{D_0} \omega_{d-k} \right)
\end{aligned}$$

where we used repeatedly $\partial^2 = d^2 = 0$ and Stokes' theorem for integration of de Rham forms over manifolds with boundaries. Introducing approximate δ -functions $\delta^{(\epsilon)} \in C_0^\infty(\mathbb{R}^d; \mathbb{R})$ for $0 < \epsilon < 1$ obeying $\delta^{(\epsilon)}(x) \geq 0$, for all $x \in \mathbb{R}^d$, $= 0$ for $|x| \geq \epsilon$ and $\int_{\mathbb{R}^d} d^d x \delta^{(\epsilon)}(x) = 1$, we define $\chi_D^{(\epsilon)}(x) = \int_{\mathbb{R}^d} d^d y \delta^{(\epsilon)}(x-y) \chi_D(y)$. With that, our formal calculation above implies

$$\int_{D^{(-k)}} \mathrm{tr}_N (\omega_{d-k}) = \lim_{\epsilon \searrow 0} \int_{\mathbb{R}^d} \mathrm{tr}_N \left(\eta_\epsilon(D^{(-k)}) \omega_{d-k} \right) \quad (17a)$$

with

$$\eta_\epsilon(D_0 \cap \partial D_1 \cap \dots \cap \partial D_k) = (-)^k d \chi_{D_k}^{(\epsilon)} \dots d \chi_{D_1}^{(\epsilon)} \chi_{D_0}^{(\epsilon)} \in \Omega_{\mathrm{dR},d}^{(k)} \quad \forall \epsilon > 0, \quad (17b)$$

and we see that we have an example of a GDAPI as discussed above.

Example 2: For all $D \in \mathcal{D}^{(0)}$ and $0 < \epsilon < 1$, $\chi_D^{(\epsilon)}$ define operators $Q_D^{(\epsilon)}$ on \mathcal{H} (6), $(Q_D^{(\epsilon)} f)(x) = \chi_D^{(\epsilon)}(x) f(x)$ for all $f \in \mathcal{H}$, and these operators are in $\hat{\Omega}_{\mathrm{dR},d}^{(0)}$. Thus

$$\eta_\epsilon(Q^{(-k)}) = (-i)^k [\varepsilon, Q_{D_k}^{(\epsilon)}] \dots [\varepsilon, Q_{D_1}^{(\epsilon)}] Q_{D_0}^{(\epsilon)} \quad (18)$$

are in $\hat{\Omega}_{\mathrm{dR},d}^{(k)}$ and define a GDAPI $(\hat{S}_{\mathrm{dR},d}, d, f, {}^* \hat{S}_{\mathrm{dR},d}, \partial)$ as discussed above. We do not attempt here to specify $\hat{S}_{\mathrm{dR},d}$ and just note that it at least contains $\Omega_{\mathrm{dR},d}$ as follows from our discussion above. Especially eq. (12) implies that this GDAPI is a natural noncommutative generalization of the GDAPI $(\Omega_{\mathrm{dR},d}, d, f, \mathcal{D}, \partial)$ from Example 1. Other examples are obtained by generating ${}^* \hat{S}_{\mathrm{dR},d}$ with other families of bounded, selfadjoint operators Q_i with $0 \leq Q_i \leq 1$ and determining $\hat{S}_{\mathrm{dR},d}$ such that (14b) holds.

3 Noncommutative Descent Equations

3.1 Result

Definitions: Given a GDA (Ω, d) , we can interpret $\Omega^{(1)}$ as generalized Yang–Mills connections and $\Omega^{(0)}$ as Lie algebra of the gauge group acting on $\Omega^{(1)}$ as

$$\Omega^{(1)} \times \Omega^{(0)} \rightarrow \Omega^{(1)}, (A, X) \mapsto d(X) + [A, X],$$

and on all polynomial functions $f : \Omega^{(1)} \rightarrow \Omega$, $A \rightarrow f(A)$ by Lie derivative

$$\mathcal{L}_X f(A) \equiv \left. \frac{d}{dt} f(A + t(d(X) + [A, X])) \right|_{t=0} \quad \forall X \in \Omega^{(0)}. \quad (19)$$

We also define a modified Lie derivative

$$\hat{\mathcal{L}}_X f(A) \equiv \mathcal{L}_X f(A) - [f(A), X] \quad (20)$$

which obviously also obeys the Leibnitz rule $\hat{\mathcal{L}}_X(ff') = \hat{\mathcal{L}}_X(f)f' + f\hat{\mathcal{L}}_X(f')$.

In the following we consider k -chains which we define as polynomial functions

$$f^k : \underbrace{\Omega^{(0)} \times \cdots \times \Omega^{(0)}}_{k \text{ times}} \times \Omega^{(1)} \rightarrow \Omega, \quad (X_1, \cdots, X_k, A) \mapsto f^k(X_1, \cdots, X_k; A) \quad (21)$$

which are multilinear in the first k arguments, and antisymmetric, i.e. for all permutations π of $(1, \cdots, k)$,

$$f^k(X_{\pi(1)}, \cdots, X_{\pi(k)}; A) = (-)^{\deg(\pi)} f^k(X_1, \cdots, X_k; A) \quad (22)$$

where $\deg(\pi)$ is 0 and 1 for even and odd permutations π , respectively. On such chains we define the operator

$$\begin{aligned} \delta f^{k-1}(X_1, \cdots, X_k; A) &= \sum_{j=1}^k (-)^{j+1} \mathcal{L}_{X_j} f^{k-1}(X_1, \cdots, \cancel{X_j}, \cdots, X_k; A) \\ &+ \sum_{\substack{i,j=1 \\ i < j}}^k (-)^{i+j} f^k([X_i, X_j], X_1, \cdots, \cancel{X_i}, \cdots, \cancel{X_j}, \cdots, X_k; A) \end{aligned} \quad (23)$$

where $\cancel{X_j}$ means that X_j is omitted. δ is the usual operation acting on chains and satisfying $\delta^2 = 0$. We also introduce the operator

$$\mathcal{J} f^{k-1}(X_1, \cdots, X_k; A) = \sum_{j=1}^k (-)^{j+1} [X_j, f^{k-1}(X_1, \cdots, \cancel{X_j}, \cdots, X_k; A)] \quad (24)$$

also satisfying $\mathcal{J}^2 = 0$.

For positive, even integers $2n$, we define the $2n$ -forms

$$Ch_{2n}(A) \equiv (F_A)^n, \quad F_A \equiv d(A) + A^2 \quad \forall A \in \Omega^{(1)} \quad (25)$$

where F_A is the curvature of the Yang–Mills connection A .

To write our formulas in compact form we introduce Grassmann variables θ_i and integration $\int d\theta_i$ over them satisfying usual relations

$$\begin{aligned} \{\theta_i, \theta_j\} &= \{\theta_i, d\theta_j\} = \{d\theta_i, d\theta_j\} = [\theta_i, \omega] = [d\theta_i, \omega] = 0 \\ \int d\theta_i \theta_j &= \delta_{ij}, \quad \int d\theta_i = 0 \quad \forall i, j = 0, 1, 2, \dots, 2n, \omega \in \Omega. \end{aligned} \quad (26)$$

For $A \in \Omega^{(1)}$, $X_i \in \Omega^{(0)}$, $0 \leq t \leq 1$ we set

$$\nu_{2n-k-1}^k(X_1, \dots, X_k; A, t) \equiv \int d\theta_k \cdots d\theta_0 \left(F_{tA} + \theta_0 A + \sum_{i=1}^k (t-1)\theta_i d(X_i) \right)^n \quad k = 0, 1, \dots, n-1 \quad (27a)$$

where $F_{tA} = td(A) + t^2 A^2$, and

$$\begin{aligned} \nu_{2n-k-1}^k(X_1, \dots, X_k; A, t) &\equiv (-)^n \int d\theta_k \cdots d\theta_0 \left(\sum_{i=0}^k t\theta_i d(X_i) + \sum_{\substack{i,j=1 \\ i < j}}^k (t^2 - t)\theta_j [X_i, X_j] \right. \\ &\quad \left. + \sum_{i=1}^k \theta_0 \theta_i X_i \right)^n, \quad k = n, n+1, \dots, 2n-1 \end{aligned} \quad (27b)$$

(the ν_{2n-k-1}^k for $k > n$ are actually independent of A). Here and in the following we write $\int d\theta_k \cdots d\theta_0$ short for $\int d\theta_k \int d\theta_{k-1} \cdots \int d\theta_0$.

Using this we then define k -chains

$$\omega_{2n-k-1}^k(X_1, \dots, X_k; A) \equiv \int_0^1 dt \nu_{2n-k-1}^k(X_1, \dots, X_k; A, t), \quad (28a)$$

$$\begin{aligned} \tilde{\omega}_{2n-k-1}^k(X_1, \dots, X_k; A) &\equiv \int_0^1 dt t \nu_{2n-k-1}^k(X_1, \dots, X_k; A, t) \\ &k = 0, 1, \dots, 2n-1 \end{aligned} \quad (28b)$$

which are easily seen to be forms of the degrees indicated (this follows from (26)).

Theorem: For arbitrary graded differential algebra (GDA) (Ω, d) and $n = 1, 2, \dots$, the forms defined above obey the following generalized descent equations,

$$Ch_{2n} = d(\omega_{2n-1}^0) + \{A, \tilde{\omega}_{2n-1}^0\} \quad (29a)$$

$$\begin{aligned}\delta\omega_{2n-k}^{k-1} + d(\omega_{2n-k-1}^k) &= \mathcal{J}\omega_{2n-k}^{k-1} - [A, \tilde{\omega}_{2n-k-1}^k]_\sigma \\ \sigma &= (-)^k, \quad k = 1, 2, \dots, n-1\end{aligned}\tag{29b}$$

$$\begin{aligned}\delta\omega_{2n-k}^{k-1} + d(\omega_{2n-k-1}^k) &= \mathcal{J}\tilde{\omega}_{2n-k}^{k-1} \\ k &= n, n+1, \dots, 2n-1\end{aligned}\tag{29c}$$

$$\delta\omega_0^{(2n-1)} = \mathcal{J}\tilde{\omega}_0^{(2n-1)}.\tag{29d}$$

(We recall that $[a, b]_\sigma = ab + \sigma ba$ for $\sigma = \pm$.)

Corollary: For all differential graded algebras with partial integration structure (GDAPI) $(\Omega, d, f, {}^*\Omega, \partial)$ the forms defined above obey the following relations

$$\begin{aligned}\delta \int_Q \omega_{2n-k}^{k-1} &= \int_{\partial Q} \omega_{2n-k-1}^k \\ \forall k &= 1, 2, \dots, 2n, n = 1, 2, \dots, Q \in {}^*\Omega\end{aligned}\tag{30}$$

(we set $\omega_{-1}^{2n} \equiv 0$, and $\delta \int_Q \equiv \int_Q \delta$).

3.2 Proof

We will divide the proof in three parts (a)–(c).

For the following we note that every element $\omega_k \in \Omega^{(n)}$ naturally defines an operator $L_{\omega_k} : \Omega \rightarrow \Omega, \omega \mapsto L_{\omega_k}(\omega) \equiv \omega_k \omega$ (left multiplication), and obviously $dL_{\omega_k} = L_{d\omega_k} + (-)^k L_{\omega_k} d$. Since the distinction of L_{ω_k} from ω_k would clutter notation, we simply write $L_{\omega_k} = \omega_k$. The former relation then is written as $d\omega_k = d(\omega_k) + (-)^k \omega_k d$. It is therefore important in the following to distinguish between $d\omega$ and $d(\omega)$ ($\omega \in \Omega$). Thus for $A \in \Omega^{(1)}$, $d(A) = \{d, A\}$, and since $d^2 = 0$ we can write $F_A = (d + A)^2$. In this notation Bianchi identity is trivial, $[d + A, F_A] = 0$. Similarly for $X \in \Omega^{(0)}$, $d(X) = [d, X]$.

(a) We write ($\partial_t \equiv \partial/\partial t$)

$$Ch_{2n}(A) = \int_0^1 dt \partial_t (F_{tA})^n = \int_0^1 dt \sum_{\nu=0}^{n-1} (F_{tA})^{n-1-\nu} \partial_t (F_{tA}) (F_{tA})^\nu$$

(note that $F_0 = 0$). Since $\partial_t (F_{tA}) = d(A) + 2tA^2 = \{d + tA, A\}$ and $d + tA$ commutes with F_{tA} (Bianchi identity), this is equivalent to

$$Ch_{2n}(A) = \int_0^1 dt \{d + tA, \sum_{\nu=0}^{n-1} (F_{tA})^{n-1-\nu} A (F_{tA})^\nu\}.$$

We note that (cf. (27a))

$$\nu_{2n-1}^0(A; t) = \int d\theta_0 (F_{tA} + \theta_0 A)^n = \sum_{\nu=0}^{n-1} (F_{tA})^{n-1-\nu} A (F_{tA})^\nu, \quad (31)$$

and with our definition (28a) we obtain (29a).

(b) Here $k = 1, 2, \dots, n$.

We shall use

$$\mathcal{L}_X(A) = [d + A, X], \quad \mathcal{L}_X(F_A) = [F_A, X]. \quad (32)$$

This implies

$$\mathcal{L}_X(F_{tA}) = [F_{tA}, X] + (t-1)\{d + tA, [d, X]\}. \quad (33)$$

(To see this, note that $F_{tA} = tF_A + (t^2 - t)A^2$, thus

$$\begin{aligned} \mathcal{L}_X(F_{tA}) &= t[F_A, X] + (t^2 - t)\{A, [d + A, X]\} = \\ &= [tF_A, X] + [(t^2 - t)A^2, X] + (t-1)\{tA, [d, X]\} \end{aligned}$$

where we used $\{A, [A, X]\} = [A^2, X]$. Adding $(t-1)\{d, [d, X]\} = 0$ yields (33).)

With Grassmann numbers θ_j as above we introduce

$$\mathcal{A} \equiv \theta_0 A + \sum_{j=1}^k (t-1)\theta_j [d, X_j], \quad \mathcal{F}_\pm \equiv F_{tA} \pm \mathcal{A}, \quad (34)$$

the operator

$$\Delta = \theta_0 \partial_t + \sum_{j=1}^k \theta_j \hat{\mathcal{L}}_{X_j}, \quad (35)$$

and we calculate $\Delta(\mathcal{F}_+) = (\cdot)_1 + (\cdot)_2$. Using $\partial_t(F_{tA}) = \{d + tA, A\}$,

$$(\cdot)_1 \equiv \theta_0 \partial_t(\mathcal{F}_+) = \{d + tA, \theta_0 A\} + \sum_{j=1}^k \theta_0 \theta_j [d, X_j].$$

The relations above and the definition (20) of $\hat{\mathcal{L}}_X$ imply

$$\hat{\mathcal{L}}_X(A) = \{d, X\}, \quad \hat{\mathcal{L}}_X(F_{tA}) = (t-1)\{d + tA, [d, X]\}$$

and $\hat{\mathcal{L}}_X(Y) = -[Y, X] = [X, Y]$ if Y is independent of A , thus

$$\begin{aligned} (\cdot)_2 &\equiv \sum_{i=1}^k \theta_i \hat{\mathcal{L}}_{X_i}(\mathcal{F}_+) = \sum_{i=1}^k \theta_i ((t-1)\{d + tA, [d, X_i]\} + \theta_0 [d, X_i] + \\ &+ \sum_{j=1}^k (t-1)\theta_j [X_i, [d, X_j]]) = \{d + tA, \sum_{i=1}^k (t-1)\theta_i [d, X_i]\} + \sum_{i=1}^k \theta_i \theta_0 [d, X_i] + \Xi \end{aligned}$$

where we introduced $\Xi = \sum_{i,j=1}^k \theta_i \theta_j (t-1) [X_i, [d, X_j]]$, or equivalently (using $\{\theta_i, \theta_j\} = 0$, and the Jacobi identity and antisymmetry of $[\cdot, \cdot]$.)

$$\Xi = \sum_{\substack{i,j=1 \\ i < j}}^k (t-1) \theta_i \theta_j [d, [X_i, X_j]]. \quad (36)$$

Putting this together we see that the second terms in $(\cdot)_1$ and $(\cdot)_2$ cancel each other, and we obtain

$$\Delta(\mathcal{F}_+) = \{d + tA, \mathcal{A}\} + \Xi. \quad (37)$$

To proceed it is useful to use another Grassmann number $\theta \equiv \theta_{k+1}$ which obviously satisfies $\theta \mathcal{A} = -\mathcal{A} \theta$. This allows to rewrite (37) as $\theta \Delta(\mathcal{F}_+) = [\theta(d + tA), \mathcal{A}] + \theta \Xi$, or equivalently (since $[\theta(d + tA), F_{tA}] = 0$)

$$\theta \Delta(\mathcal{F}_+) = [\theta(d + tA), \mathcal{F}_+] + \theta \Xi.$$

Since $\theta \Delta \mathcal{F}_+ = \theta \Delta(\mathcal{F}_+) + \mathcal{F}_+ \theta \Delta$, this implies

$$\begin{aligned} \theta \Delta((\mathcal{F}_+)^n) &= \sum_{\nu=0}^{n-1} (\mathcal{F}_+)^{n-1-\nu} \theta \Delta(\mathcal{F}_+) (\mathcal{F}_+)^{\nu} \\ &= \sum_{\nu=0}^{n-1} (\mathcal{F}_+)^{n-1-\nu} ([\theta(d + tA), \mathcal{F}_+] + \theta \Xi) (\mathcal{F}_+)^{\nu} \\ &= [\theta(d + tA), (\mathcal{F}_+)^n] + \sum_{\nu=0}^{n-1} (\mathcal{F}_+)^{n-1-\nu} \theta \Xi (\mathcal{F}_+)^{\nu} \end{aligned}$$

as $\sum_{\nu=0}^{n-1} (\mathcal{F}_+)^{n-1-\nu} [\cdot, \mathcal{F}_+] (\mathcal{F}_+)^{\nu} = [\cdot, (\mathcal{F}_+)^n]$. To get rid of θ again we use $\mathcal{F}_+ \theta = \theta \mathcal{F}_-$ (cf. (34)), thus

$$\Delta((\mathcal{F}_+)^n) = (d + tA)(\mathcal{F}_+)^n - (\mathcal{F}_-)^n (d + tA) + \sum_{\nu=0}^{n-1} (\mathcal{F}_-)^{n-1-\nu} \Xi (\mathcal{F}_+)^{\nu}. \quad (38)$$

We now apply integration $\int d\theta_k \cdots d\theta_0$ to this equation. In the following calculation we keep in mind that we can always forget about terms containing higher powers than 1 of at least one of the θ_i 's. Thus,

$$\begin{aligned} \int d\theta_k \cdots d\theta_0 \Delta((\mathcal{F}_+)^n) &= \int d\theta_k \cdots d\theta_0 \theta_0 \partial_t \left(F_{tA} + \sum_{j=1}^k (t-1) \theta_j [d, X_j] \right)^n \\ &\quad + \sum_{i=1}^k \hat{\mathcal{L}}_{X_i} \int d\theta_k \cdots d\theta_0 \theta_i \left(F_{tA} + \theta_0 A + \sum_{\substack{j=1 \\ j \neq i}}^k (t-1) \theta_j [d, X_j] \right)^n. \end{aligned}$$

Using

$$\int d\theta_k \cdots d\theta_0 \theta_j = (-)^j \int d\theta_k \cdots d\theta_j \cdots d\theta_0 \quad j = 0, 1 \cdots k$$

and the definition (27a) of ν_{2n-k}^{k-1} we can write this

$$\begin{aligned} \int d\theta_k \cdots d\theta_0 \Delta((\mathcal{F}_+)^n) &= \partial_t \nu_{2n-k}^{k-1}(X_2, \dots, X_k; (t-1)[d, X_1], t) \\ &+ \sum_{j=1}^k (-)^j \hat{\mathcal{L}}_{X_j} \nu_{2n-k}^{k-1}(X_1, \dots, X_j \cdots, X_k; A, t). \end{aligned}$$

Moreover, since $\int d\theta_k \cdots d\theta_0 (\mathcal{F}_\pm)^n = (\pm)^{k+1} \nu_{2n-k-1}^k(X_1, \dots, X_k; A, t)$ (cf. (27a) and (34)), we have

$$\begin{aligned} &\int d\theta_k \cdots d\theta_0 ((d+tA)(\mathcal{F}_+)^n - (\mathcal{F}_-)^n(d+tA)) \\ &= [d+tA, \nu_{2n-k-1}^k(X_1, \dots, X_k; A, t)]_\sigma, \quad \sigma = (-)^k. \end{aligned}$$

Finally, (we use $\mathcal{F}_- \theta_i = \theta_i \mathcal{F}_+$; cf. (36))

$$\begin{aligned} &\int d\theta_k \cdots d\theta_0 \sum_{\nu=0}^{n-1} (\mathcal{F}_-)^{n-1-\nu} \Xi(\mathcal{F}_+)^{\nu} \\ &= \sum_{\substack{i,j \\ i < j}}^{n-1} \sum_{\nu=0}^{n-1} \int d\theta_k \cdots d\theta_0 \theta_i (\mathcal{F}_+)^{n-1-\nu} (t-1) \theta_j [d, [X_i, X_j]] (\mathcal{F}_+)^{\nu} \\ &= \int d\theta_k \cdots d\theta_0 \theta_i \sum_{\substack{i,j \\ i < j}} \left(F_{tA} + \theta_0 A + (t-1) \theta_j [d, [X_i, X_j]] + \sum_{\substack{\ell=1 \\ \ell \neq i,j}}^k (t-1) \theta_\ell [d, X_\ell] \right)^n \\ &= \sum_{\substack{i,j \\ i < j}} (-)^{i+j} \nu_{2n-k}^{k-1}([X_i, X_j], X_1, \dots, X_i \cdots, X_j \cdots, X_k; A, t) \end{aligned}$$

where we used

$$\int d\theta_k \cdots d\theta_0 \theta_i = (-)^{i+j} \int d\theta_k \cdots d\theta_i \cdots d\theta_j \cdots d\theta_0.$$

Using the definitions (23) and (24) we can therefore write $\int d\theta_k \cdots d\theta_0$ (38) as

$$\begin{aligned} &\hat{\delta} \nu_{2n-k}^{k-1}(X_1, \dots, X_k; A, t) + [d+tA, \nu_{2n-k-1}^k(X_1, \dots, X_k; A, t)]_\sigma \\ &= \partial_i \nu_{2n-k-1}^k(X_2, \dots, X_k; (t-1)[d, X_1]; A, t), \quad \sigma = (-)^k \end{aligned} \quad (39)$$

where $\hat{\delta} = \delta - \mathcal{J}$. Our descent equations for $k < n$ are obtained by applying integration $\int_0^1 dt$ to (39). We first note that for $k < n$, $\nu_{2n-k-1}^k(X_2, \dots, X_k; (t-1)[d, X_1]; A, t)$ is zero for $t = 1$ (trivially) and $t = 0$ (since $F_0 = 0$), thus $\int_0^1 dt \partial_t \nu_{2n-k-1}^k = 0$. Thus with the definitions (28a),

$$\delta \omega_{2n-k}^{k-1} - \mathcal{J} \omega_{2n-k}^{k-1} + [d, \omega_{2n-k-1}^k]_\sigma + [A, \tilde{\omega}_{2n-k-1}^k]_\sigma = 0 \quad k = 1, 2, \dots, n-1 \quad (40)$$

equivalent to (29b). For $k = n$ we observe that ν_{n-1}^n as defined in (27a) is zero, thus

$$\delta \omega_n^{n-1} - \mathcal{J} \omega_n^{n-1} = N_n^n \quad (41)$$

where $N_n^n = \int_0^1 dt \partial_t \nu_{n-1}^n$ is nonzero,

$$N_n^n(X_1, \dots, X_n) = (-)^{n+1} \int d\theta_n \cdots d\theta_1 \left(\sum_{i=1}^n \theta_i d(X_i) \right)^n. \quad (42)$$

We will show below that N_n^n equal to $-d\omega_n^{n-1}$.

(c) Here $k = n, n+1, \dots, 2n-1$.

We introduce

$$\mathcal{V} = \sum_{i=1}^k \theta_i X_i, \quad \tilde{\mathcal{F}}_\pm = t d(\mathcal{V}) + (t^2 - t) \mathcal{V}^2 \pm \theta_0 \mathcal{V} \quad (43)$$

and the operator

$$\tilde{\Delta} = \theta_0 \partial_t + t \hat{I}, \quad \hat{I} = \sum_{i=1}^k \theta_i [X_i, \cdot]. \quad (44)$$

We observe

$$\mathcal{V}^2 = \frac{1}{2} \sum_{i,j=1}^k \theta_i \theta_j [X_i, X_j] = \sum_{\substack{i,j=1 \\ i < j}}^k \theta_i \theta_j [X_i, X_j]$$

implying $\hat{I}(\mathcal{V}) = 2\mathcal{V}^2$. Similarly,

$$\hat{I}(d(\mathcal{V})) = \frac{1}{2} \sum_{i,j=1}^k \theta_i \theta_j ([X_i, d(X_j)] - [X_j, d(X_i)]) = d(\mathcal{V}^2),$$

and

$$\begin{aligned} \hat{I}(\mathcal{V}^2) &= \sum_{\ell=1}^k \sum_{\substack{i,j=1 \\ i < j}}^k \theta_\ell \theta_i \theta_j [X_\ell, [X_i, X_j]] \\ &= \frac{1}{6} \sum_{i,j,\ell=1}^k \theta_\ell \theta_i \theta_j ([X_\ell, [X_i, X_j]] + [X_i, [X_j, X_\ell]] + [X_j, [X_\ell, X_i]]) = 0 \end{aligned}$$

where we used $\theta_\ell\theta_i\theta_j = \theta_i\theta_j\theta_\ell$ etc. and the Jacobi identity for $[\cdot, \cdot]$. Thus

$$t\hat{I}(\tilde{\mathcal{F}}_+) = t^2d(\mathcal{V}^2) - 2t\theta_0\mathcal{V}^2$$

(note that $\hat{I}\theta_0 = -\theta_0\hat{I}$). Obviously $\theta\partial_t(\tilde{\mathcal{F}}_+) = \theta_0d(\mathcal{V}) + (2t-1)\theta_0\mathcal{V}^2$. Combining the last two equations we obtain

$$\tilde{\Delta}(\tilde{\mathcal{F}}_+) = d(\tilde{\mathcal{F}}_+) + \tilde{\Xi}, \quad \tilde{\Xi} = td(\mathcal{V}^2) - (t^2-t)\hat{I}(\mathcal{V}^2) - \theta_0\mathcal{V}^2 \quad (45)$$

where we added terms $d(td(\mathcal{V})) = -2(t^2-t)\hat{I}(\mathcal{V}^2) = 0$. Similarly as in (b) above this implies

$$\tilde{\Delta}((\tilde{\mathcal{F}}_+)^n) = d((\tilde{\mathcal{F}}_+)^n) + \sum_{\nu=0}^{n-1} (\tilde{\mathcal{F}}_+)^{n-1-\nu} \tilde{\Xi}(\tilde{\mathcal{F}}_+)^{\nu} \quad (46)$$

(we used $d\tilde{\mathcal{F}}_+ = d(\tilde{\mathcal{F}}_+) + \tilde{\mathcal{F}}_+d$). Noting that our definition (27b) can be written as $\nu_{2n-k-1}^k = (-)^n \int d\theta_k \cdots d\theta_0 (\tilde{\mathcal{F}}_+)^n$, we now apply $(-)^n \int d\theta_k \cdots d\theta_0$ to this equation. The following calculation is very similar to the one above in (b),

$$\begin{aligned} (-)^n \int d\theta_k \cdots d\theta_0 \tilde{\Delta}((\mathcal{F}_+)^n) &= \partial_t \tilde{\nu}_{2n-k-1}^k(X_1, \dots, X_k; t) \\ &+ \sum_{i=1}^k (-)^i t [X_i, \nu_{2n-k}^{k-1}(X_1, \dots, X_j, \dots, X_k; t)] \end{aligned}$$

where we introduced

$$\tilde{\nu}_{2n-k-1}^k(X_1, \dots, X_k; t) = (-)^n \int d\theta_k \cdots d\theta_1 (td(\mathcal{V}) + (t^2-t)\mathcal{V}^2)^n \quad (47)$$

and used (27b). Similarly,

$$(-)^n \int d\theta_k \cdots d\theta_0 d((\mathcal{F}_+)^n) = d(\nu_{2n-k-1}^k(X_1, \dots, X_k; t)).$$

To calculate the last term we write

$$\tilde{\Xi} = \sum_{\substack{i,j=1 \\ i < j}}^k \theta_i \left(t\theta_j d([X_i, X_j]) + \sum_{\substack{\ell=1 \\ \ell \neq i,j}}^k \theta_j \theta_\ell [[X_i, X_j], X_\ell] + \theta_0 \theta_j [X_i, X_j] \right)$$

and similarly as in (b),

$$\begin{aligned} &(-)^n \int d\theta_k \cdots d\theta_0 \sum_{\nu=0}^{n-1} (\tilde{\mathcal{F}}_+)^{n-1-\nu} \tilde{\Xi}(\tilde{\mathcal{F}}_+)^{\nu} \\ &= \sum_{\substack{i,j \\ i < j}} (-)^{i+j} \nu_{2n-k}^{k-1}([X_i, X_j], X_1, \dots, X_i, \dots, X_j, \dots, X_k; t). \end{aligned}$$

Recalling the definitions (23) and (24) we can therefore write $(-)^n \int d\theta_k \cdots d\theta_0$ (46) as

$$\delta\nu_{2n-k-1}^{k-1} + d\nu_{2n-k-1}^k = \partial_t \tilde{\nu}_{2n-k-1}^k - t \mathcal{J} \nu_{2n-k-1}^k \quad (48)$$

where we suppressed the common argument $(X_1, \dots, X_k; t)$. Again the descent equations are obtained by integrating $\int_0^1 dt$ this equation. Since for $k > n$, $\tilde{\nu}_{2n-k-1}^k$ is zero for $t = 0$ and $t = 1$ (cf. (27b)) we get

$$\delta\omega_{2n-k-1}^{k-1} + d\omega_{2n-k}^k = -\mathcal{J} \tilde{\omega}_{2n-k-1}^{k-1} \quad (49)$$

proving (29c) for $k = n + 1, \dots, 2n - 1$. Obviously this equation remains also true for $k = 2n$ if we set $d\omega_{(-1)}^{2n} = 0$. This implies (29d).

For $k = n$, ν_n^{n-1} as defined in (27b) is zero. Moreover, it is easy to see that $\int_0^1 dt \partial_t \tilde{\nu}_{n-1}^n = -N_n^n$ as defined in (42) (cf. (47)), thus

$$d\omega_{n-1}^n = -N_n^n. \quad (50)$$

Combining this with equation (41) this gives (29c) for $k = n$ which completes the proof of the theorem.

The corollary then follows from our definitions and the defining relations (13) of integrations on GDAPI.

4 Final Remarks

Given a GDA (Ω, d) one can consider a \mathbb{N}_0 -graded vector space $C = \bigoplus_{k=0}^{\infty} C^{(k)}$ of polynomial maps

$$c^k : \underbrace{\Omega^{(0)} \times \cdots \times \Omega^{(0)}}_{k \text{ times}} \times \Omega^{(1)} \rightarrow \mathbb{C}, \quad (X_1, \dots, X_k, A) \mapsto c^k(X_1, \dots, X_k; A)$$

which are multilinear and antisymmetric in the first k arguments, with the operator $\delta : C^{(k)} \rightarrow C^{(k+1)}$ defined as in (23). One easily checks that $\delta^2 = 0$, thus (C, δ) is a graded differential complex. A map $c^k \in C^{(k)}$ is called a k -cocycle if $\delta c^k = 0$, and a k -cocycle c^k is nontrivial if it is not a k -boundary i.e. not of the form δb^{k-1} with $b^{k-1} \in C^{(k-1)}$. Anomalies in quantum gauge theories are nontrivial cocycles, e.g. the axial anomalies and Schwinger terms are nontrivial 1- and 2-cocycles, respectively [J].

The result of this paper in Section 3 provides an explicit construction of k -cocycles

$$c_{2n-k-1}^k = \int \omega_{2n-k-1}^k \quad (51)$$

for arbitrary GDA (Ω, d) with integration \int . The cocycle property is a trivial consequence of our generalized descent equations and basis properties of integration. Whether or not c_{2n-k-1}^k is nontrivial depends on the integration \int and the precise definition of $C^{(k-1)}$ — in most cases it will be trivial or even zero. Well-known examples for (51) being nontrivial is for the de Rham GDA $(\Omega_{\text{dR}, 2n-k-1}, d)$ with integration $\int_{\mathbb{R}^{2n-k-1}} \text{tr}_N$, or more generally, the de Rham GDA $(\Omega_{\text{dR}, d}, d)$, d big enough, with integration $\int_M \text{tr}_N$ over a $(2n - k - 1)$ dimensional submanifold M of \mathbb{R}^d . According to our discussion in Section 2, these examples immediately generalize to universal Rham GDA (= Example 2; the hats in the following are to indicate that we are considering universal de Rham forms). We conjecture that all the universal cocycles \hat{c}_{2n-k-1}^k for $(\hat{\Omega}_{\text{dR}, 2n-k-1}, \hat{d}, \hat{f})$ are nontrivial since this is known to be true for $k = 2$ and $n = 2$ and $n = 3$ (work on proving this in general is in progress). Indeed, \hat{c}_1^2 is proportional to the Lundberg cocycle [Lb] known to be nontrivial, and \hat{c}_3^2 is proportional to the Mickelsson–Rajeev cocycle which is also nontrivial [MR] (see also [L3]).

To have one explicit example and also demonstrate how formulas can be written without using Grassmann numbers, we finally give a more explicit expression for the universal Schwinger terms generalizing the Mickelsson–Rajeev cocycle to higher dimensions,

$$\begin{aligned} \hat{c}_{2n-3}^2(\hat{X}_1, \hat{X}_2; \hat{A}) &= \frac{3}{c_{2n-3}} \sum_{\substack{i_\nu \geq 0 \\ \sum_\nu i_\nu = n-3}} (i)^2 \int_0^1 dt (t-1)^2 \\ &\times \text{Tr}_C \left((\hat{F}_{tA})^{i_1} \hat{A} (\hat{F}_{tA})^{i_2} [\varepsilon, \hat{X}_1] (\hat{F}_{tA})^{i_3} [\varepsilon, \hat{X}_2] (\hat{F}_{tA})^{i_4} - (\hat{X}_1 \leftrightarrow \hat{X}_2) \right) \end{aligned} \quad (52)$$

where $\hat{F}_{tA} = i\{\varepsilon, \hat{A}\}t + \hat{A}^2 t^2$ used cyclicity under conditional Hilbert space trace; for notation see Section 2); the \hat{X}_i and \hat{A} here are bounded operators on a Hilbert space \mathcal{H} such that $[\varepsilon, \hat{X}_i]$ and $[\varepsilon, \hat{A}]$ are in the Schatten class \mathcal{B}_{2n-2} and $\{\varepsilon, \hat{A}\} \in \mathcal{B}_{n-1}$. Restricting to operators $\hat{A} = (F - \varepsilon)$ where $F = F^{-1} = F^*$ is a grading operator (as usually done in this context [L1]), we have $\hat{F}_{tA} = (t^2 - t)(F - \varepsilon)^2$, and (52) should be proportional to the universal Schwinger terms given in [FT].

Acknowledgments

I am grateful to D. Burghelea, A. Carey, H. Grosse, G. Landi, and especially G. Kelnhofer and J. Mickelsson for useful discussions. This paper was written at the the Erwin Schrödinger International Institute in Vienna. I would like to thank the ESI for hospitality and P. Marmo and P. Michor for inviting me there.

References

- [C1] Connes A., ‘Noncommutative differential geometry,’ *I.H.E.S. Publ. Math.* **62**, 257 (1985)
- [C2] Connes A., ‘The action functional in non-commutative geometry,’ *Commun. Math. Phys.* **117**, 117 (1988)
- [C3] Connes A., *Noncommutative Geometry*, Academic Press (1994)
- [CL] Connes A. and Lott J., ‘Particle models and noncommutative geometry,’ *Nucl.Phys. B* **18** (1990) *suppl.*, 29 (1991)
- [CM] Connes A. and Moscovici H., ‘Transgression du caractère de Chern et cohomologie cyclique,’ *C.R. Acad. Sc. Paris Sér. I Math.* **303** 913 (1986)
- [F] Faddeev L. D., ‘Operator anomaly for the Gauss law,’ *Phys. Lett.* **145B**, 81 (1984)
- [FR] Ferretti G. and Rajeev S. G., ‘Universal Dirac–Yang–Mills theory,’ *Phys. Lett.* **244B**, 265 (1990)
- [FT] Fujii K. and Tanaka M., ‘Universal Schwinger cocycle and current algebras in $(D + 1)$ - dimensions: geometry and physics,’ *Commun. Math. Phys.* **129**, 267 (1990)
- [J] Jackiw R., ‘Topological investigations of quantized gauge theories.’ in: *Relativity, Groups and Topology II*, Les Houghes 1983, DeWitt B. S. and Stora R. (eds.), North Holland, Amsterdam (1984)
- [KS] Kastler D., and Stora R., ‘A differential geometric setting for BRS transformations and anomalies I,’ *J. Geom. and Phys.* **3**, 438 (1986)
- [L1] Langmann E., ‘Fermion current algebras and Schwinger terms in $(3+1)$ -Dimensions,’ *Commun. Math. Phys.* **162**, 1 (1994)

- [L2] Langmann E., ‘Noncommutative integration calculus,’ *J. Math. Phys.* **36**, 3822 (1995)
- [L3] Langmann E., ‘On anomalies and noncommutative geometry,’ Proc. of “34. Internationale Universitätswochen für Kern- und Teilchenphysik Schladming”, Schladming March 1995, Springer Verlag (to appear); [hep-th/9507088](#)
- [Lb] Lundberg L.-E. ‘Quasi-free “second quantization”,’ *Commun. Math. Phys.* **50**, 103 (1976)
- [Lü] Lüscher M., ‘Topology of lattice gauge fields,’ *Commun. Math. Phys.* **85**, 39 (1982)
- [LM1] Langmann E., and Mickelsson J., ‘(3+1)-dimensional Schwinger terms and non-commutative geometry,’ *Phys. Lett. B* **338**, 241 (1994)
- [LM2] Langmann E., and Mickelsson J., ‘Elementary derivation of the chiral anomaly,’ *Lett. Math. Phys.* (in press), [hep-th/9412129](#)
- [M] Mickelsson J., ‘Current algebra representations for 3+1 dimensional Dirac-Yang-Mills theory,’ *Commun. Math. Phys.* **117**, 261 (1988); ‘Commutator anomalies and the Fock bundle,’ *Commun. Math. Phys.* **127**, 285 (1990)
- [MR] Mickelsson J., and Rajeev S. G., ‘Current algebras in $d + 1$ -dimensions and determinant bundles over infinite dimensional Grassmannians,’ *Commun. Math. Phys.* **116**, 400 (1988)
- [S] Simon B., *Trace Ideals and Their Applications*. Cambridge University Press, Cambridge, UK (1979)
- [R] Rajeev S. G., ‘Universal gauge theory,’ *Phys. Rev. D* **42**, 2779 (1990); ‘Embedding Yang–Mills theory into universal Yang–Mills theory,’ *Phys. Rev. D* **44**, 1836 (1991)
- [RS] Reed R., and Simon B., *Methods of Modern Mathematical Physics I. Functional Analysis*, Academic Press, New York (1968)
- [tHV] t’Hooft G. and Veltman M., ‘Regularization and renormalization of gauge fields,’ *Nucl. Phys. B* **44**, 189 (1972)

- [Z] Zumino B., 'Cohomology of gauge groups: cocycles and Schwinger terms,'
Nucl.Phys. B **253**, 477 (1985)