

are removed from the quiescent cold disc. In this case the heating front would arrive at its usual speed to the inner disc edge but then the ‘missing’ UV emitting regions would be rebuilt in a viscous time, comparable to the observed UV delay.

A weak magnetic field ($\lesssim 10^5$ G) would disrupt the inner disc at a distance of few to ten white dwarf radii. Livio & Pringle (1992) showed that including a white dwarf magnetic field $\sim 10^4$ G into the standard DIM may account for the observed UV delay.

Meyer & Meyer-Hofmeister (1994) studied the structure of the corona above a quiescent dwarf nova disc. They found that the frictionally heated corona evaporates because of the inefficient radiative cooling of the optically thin gas. As a result a hole forms in the inner disc. A similar idea has been applied to quiescent Soft X-ray Transients (SXT) by Narayan et al. (1995). It is more difficult than in magnetic field case to find a general formula for the inner radius of the evaporated disc but calculations suggest that its value will be also from few to ten white dwarf radii.

2.2. The quiescent UV and X-ray flux problem

A ‘hole’ in the inner disc would also help to remove another difficulty of the DIM: in some quiescent DN systems one observes an UV and hard X-ray emission that corresponds to accretion at rates higher than those allowed by the model. The DIM requires the surface density of the quiescent disc Σ to be less than the critical surface density Σ_{\max} everywhere in the disc (see Fig. 1). This implies that an inequality has to be satisfied by the accretion rate in a quiescent disc (Lasota 1995a):

$$\dot{M}(r) \approx 4.1 \times 10^{14} t_6^{-1} r_9^{3.11} M_1^{-0.37} \left(\frac{\alpha}{0.01} \right)^{-0.79} \text{ g s}^{-1} \quad (1)$$

where ν is the kinematic viscosity coefficient, α the viscosity parameter, M_1 the white dwarf mass in solar units, t_6 is the recurrence time in units of 10^6 s and r_9 is the radius in 10^9 cm. In several systems however the observed UV and hard X-ray flux imply accretion rates at the inner disc edge of at least one order of magnitude higher than the one given by Eq. (1) for $r_{\text{in}} = r_9 = 1$ (see Meyer & Meyer-Hofmeister 1994). As the maximum allowed accretion rate in a quiescent disc varies as $r^{3.11}$, the removal of the inner disc regions by magnetic stress or evaporation could easily solve the problem of DIM constraints on the quiescent accretion rate without assuming an unusually small viscosity parameter α .

Both the UV delay problem and the too high quiescent UV and X-ray flux problem can be overcome by only one effect: the removal of the inner disc in quiescence.

The superhumps observed during SU UMa superoutbursts are due to a tidal distortion of the outer disc resulting from the presence of the 3:1 resonance inside the disc (see e.g. King 1994). It requires mass ratios $q = M_2/M_1 < q_{\text{crit}} \approx 0.25 - 0.33$; It is remarkable that U Gem showed a very long (45 day) outburst which looks like a superoutbursts (see Mason et al. 1988) but was not classified as such because a superhump was absent. The mass ratio for this system is ~ 0.46 .

According to Osaki (1989) superoutbursts are pure disc phenomena. His tidal-thermal instability model makes several predictions that do not seem to be confirmed by observations (Smak 1991, Whitehurst 1994). In addition the thermal-tidal instability model cannot explain the ‘superoutburst’ of U Gem and such very long and rare outbursts in systems with $q > q_{\text{crit}}$ would require a different explanation.

A (moderate) mass-transfer enhancement model for superoutbursts was proposed by Whitehurst (1994). Such enhancements are observed in several systems (Smak 1995). EMT would explain the length of the superoutburst. The exact cause of EMT is not known but there is growing evidence that mass transfer rate in CVs is subject to substantial fluctuations on timescales from days to months. In the pure EMT model, however, it is difficult to generate the disc excentricity which is supposed to give rise to the superhump (Lubow 1994). It seems that a ‘hybrid’ model might be necessary to explain superoutbursts (Smak 1995).

WZ Sge has the particularity of showing only superoutbursts with a recurrence time of ~ 33 years and no normal outbursts have been observed in this system. A few other systems show similar but less extreme behaviour. In order to account for such long recurrence time, DIM requires WZ Sge to have an extremely low quiescent viscosity: $\alpha_{\text{cold}} < 5 \times 10^{-5}$ (Smak 1993) as compared to the usual value of ~ 0.01 . The physical reason for such a small value would require an explanation. A similar difficulty is encountered when DIM is applied to SXTs (Lasota 1995a).

3. Dwarf novae at very low mass transfer rates

At a given radius r , thermal equilibria of accretion discs in cataclysmic variables form, on the $\Sigma-\dot{M}$ (or $\Sigma-T_{\text{eff}}$) plane, a S-shaped curve, where the middle branch with the negative slope is thermally (and viscously) unstable. Hot stable equilibria exist therefore only for $\Sigma > \Sigma_{\min}$ and cold ones only for $\Sigma < \Sigma_{\max}$ (see e.g. Cannizzo 1993). For the disc to be globally unstable, i.e. unstable at some radius $r_{\text{in}} \leq r \leq r_{\text{out}}$ one requires the mass transfer rate \dot{M}_T to satisfy the inequality

$$\dot{M}(\Sigma_{\max}(r_{\text{in}})) < \dot{M}_T < \dot{M}(\Sigma_{\min}(r_{\text{out}})) \quad (2)$$

The range of unstable mass transfer rates therefore depends on the size of the disc. Fig. 1 shows surface density

M_1 (Hameury et al. 1995). The accretion rate \dot{M} is constant for each curve.




Fig. 1. Density profiles of stationary accretion discs around white dwarfs with $M_1 = 0.4$ (upper panel), $M_1 = 0.7$ (middle panel) and $M_1 = 1$ (lower panel), for $\dot{M} = 10^{13}, 10^{14}, 10^{15}, 10^{16}$ and 10^{17} g s^{-1} (except for $M_1 = 0.4$ for which the 10^{13} profile is not plotted). The viscosity parameter α is equal to 0.01. The stable parts of cold equilibria are represented by a thick line, the unstable part by a dotted line and the hot stable equilibria by a continuous thin line. For $M_1 = 0.4$ the 10^{15} g s^{-1} profile is represented by a dashed-dotted line when it coincides with Σ_{\max} i.e. when the disc is marginally stable.

The part of the curve which is above the Σ_{\max} line cannot represent a cold thermal equilibrium so that the disc cannot be globally in equilibrium ($\dot{M} = \text{const}$) for the corresponding mass transfer rate.

From Fig. 1 one can see that there exist two types of equilibria represented by the $\Sigma(r)$ curves: for $\dot{M} \lesssim \dot{M}'$ (where \dot{M}' is some value of accretion rate depending on M_1 and α ; in the Fig. 1 $\dot{M}' \sim 10^{13} - 10^{14} \text{ g s}^{-1}$) the surface density is always less than Σ_{\max} . For such low \dot{M} the disc is globally in cold stable equilibrium, from r_{out} down

part of the disc can be in a cold stable equilibrium. Starting at some r_{out} from $\Sigma < \Sigma_{\max}$, the $\Sigma(r)$ curve reaches, with decreasing r , the value Σ_{\max} at some radius r_{crit} . The disc can be in cold equilibrium only for $r_{\text{crit}} < r < r_{\text{out}}$. The next segment of the curve between Σ_{\max} and Σ_{\min} is unstable while the curve to the left of Σ_{\min} represent hot equilibria. For $\dot{M} > \dot{M}'$ the disc is therefore globally unstable if it extends down to the surface of the white dwarf, and a cold steady state cannot exist. There exists however a radius $r_{\text{in}} = r_{\text{crit}}$ for which is disc is globally stable at a $\dot{M} = \text{const}$. As mentioned above both the magnetically disrupted and the evaporated accretion discs have inner radii $\approx 3 \times 10^9 \text{ cm}$. Fig. 2 shows $\dot{M}(\Sigma_{\max}(r)) \equiv \dot{M}_{\text{crit}}(r)$. One can see that for $r_{\text{in}} \sim 2.5 - 5 \times 10^9$ (depending on M_1) the critical mass transfer rate below which the disc is globally stable is $\dot{M}_{\text{crit}} \approx 10^{15} \text{ g s}^{-1}$.

If models assuming a ‘hole’ in the inner disc were to apply to CVs transferring mass at low rates, those systems in quiescence would be globally stable and outbursts could only be triggered by an *increase of mass transfer rate* from the companion. One should stress here that the resulting outburst would be due to the ‘usual’ thermal instability, as the only effect of the enhanced mass transfer is to bring the disc into a globally unstable state (and to add some mass into the disc). The proposed model scheme is therefore a ‘hybrid’ (EMT-DIM) model.




Fig. 2. $\dot{M}(\Sigma_{\max}(r)) \equiv \dot{M}_{\text{crit}}(r)$ for three values of $M_1 = 0.4, 0.7$ and 1. At a given r' , mass transfer rates $\dot{M} < \dot{M}_{\text{crit}}(r')$ correspond to global, stable disc equilibria for $r > r'$.

There are no reasons for the disrupted or evaporated disc models not to apply to systems at low mass transfer rates. On the contrary, both types of models were devised to explain the behaviour of discs at low accretion rates.

One of such systems is WZ Sge for which Smak (1993) estimated the mass transfer rate to be $\sim 10^{15} \text{ g s}^{-1}$ and the white dwarf mass $M_1 = 0.4$. Fig. 2 shows that at this

ally stable even a very weak magnetic field would make it stable. This system should not show dwarf nova outbursts. Indeed, WZ Sge shows only superoutbursts with a very long recurrence time. Those superoutbursts would be due to a thermal instability triggered by EMT from the companion. It is interesting to note that Patterson (1994) suggested that the white dwarf in WZ Sge is weakly magnetized.

Our model allows a unified scheme for SU UMa's and WZ Sge type systems. If SU UMa's superoutbursts are triggered by enhanced mass transfer, the difference between those systems and WZ Sge and similar binaries would be due only to the value of the mass transfer from the companion. In SU UMa's mass transfer rates are close to 10^{16} g s^{-1} and \dot{M}_T can be even larger than \dot{M}_{crit} ($r_{\text{crit}} \gtrsim r_{\text{out}}$) so that, contrary to WZ Sge, their discs are globally unstable for any value of the inner radius and they undergo 'normal' outbursts and superoutbursts are triggered by EMT. There is no need in the proposed scheme to assume, for unknown reasons, extremely low values of α in WZ Sge. True, our scheme involves mass transfer modulations whose reasons are also unknown but at least such modulations are observed in many systems. One possible reason for those modulations could be appearance of magnetic spots at the inner Lagrange point (Livio & Pringle 1994).

Of special interest is the system HT Cas which show rare outbursts (Wenzel 1987) of which at least one was a superoutburst (Zhang et al. 1986). According to Wood et al. (1992) the mass transfer rate in this system is $\sim 2 \times 10^{15} \text{ g s}^{-1}$. Eclipse mapping shows a very flat slope of the temperature in the inner disc. This could be due to magnetic disruption and/or to nonradiative energy loss (see e.g. Rutten et al. 1992). HT Cas was also observed in a 'low state' with accretion (and mass transfer) rate of $\sim 6 \times 10^{13} \text{ g s}^{-1}$ (Wood et al. 1995). We suggest that the strange 'erratic' behaviour of HT Cas is due to the fact that its mass transfer rate is very close to \dot{M}_{crit} and quite often the disc is globally stable. In any case the mass transfer rate in this system *is observed* to vary on short time scales.

As shown by Narayan et al. (1995) quiescent spectra of soft X-ray transients can be explained by advection dominated flows near the central black hole. A scheme similar to the one proposed here may be used to explain SXTs outbursts (Yi et al. 1995, Lasota 1995b).

4. Conclusions

We have shown that the current ideas on the inner accretion disc structure of quiescent dwarf novae imply that CVs with very low mass transfer rates from the companion are globally stable with respect to the thermal instability. Thus, outbursts can occur only during episodes of enhanced mass transfer.

no (or only rare) 'normal' outbursts. The viscosity in the discs of those systems would be comparable to the one in binaries with slightly higher mass transfer rate. The proposed scheme naturally fits into the models in which superoutbursts are triggered by EMT.

References

- Cannizzo J.K., 1993, The Limit Cycle Instability in Dwarf Nova Accretion Disks. In: Craig Wheeler J. (ed) *Accretion Disks in Compact Stellar Systems*. World Scientific Publishing, Singapore, p.6
- Frank J., King A.R., Raine D., 1992. *Accretion Power in Astrophysics*. Cambridge University Press, Cambridge
- Hameury J.M., Huré J.M., Lasota J.P., 1995 *A&A*, in preparation
- King, A.R. 1994, Tidal Insabilities in Accretion Disks. In: Holt S.S., Day C. (eds) *The Evolution of X-ray Binaries*. AIP Conference Proceedings 308, New York, p. 515
- Lasota J.P., 1995a, Mechanisms for Dwarf Nova Outbursts and Soft X-ray Transients. In: van den Heuvel E.P.J, van Paradijs J.(eds) *Compact Stars in Binaries*. Proceedings of the IAU Symposium 165, Kluwer, in press
- Lasota J.P., 1995b, Models of Soft X-ray Transients and Dwarf Novae. In Wood J.H. et al. (eds) *Cataclysmic Variables and Related Objects*. Proceedings of IAU Coll. 158, in press
- Livio M., Pringle J.E., 1992 *MNRS* 259, 23P
- Livio M., Pringle J.E., 1994 *ApJ* 427, 956
- Lubow S.H., 1994 *ApJ* 432, 224
- Mason K.O., Cordova F.A., Watson M.G., King, A.R., 1988 *MNRS* 232, 779
- Meyer F., Meyer-Hofmeister E., 1989 *A&A* 221, 36
- Meyer F., Meyer-Hofmeister E., 1994 *A&A* 288, 175
- Narayan R., Yi I., 1995 *ApJ*, in press
- Narayan R., McClintock J.E., Yi I., 1995 *ApJ*, submitted
- Osaki Y., 1985 *ã144369*
- Osaki Y., 1989 *PASJ* 41, 1005
- Patterson, J. 1994 *PASP* 106, 209
- Pringle, J.E., Verbunt, F. & Wade, R.A. 1986 *MNRS* 221, 169
- Rutten R.G.M., van Paradijs J., Tinbergen J., 1992 *A&A* 260, 213
- Smak J.I., 1991 *Acta. Astron.* 41, 269
- Smak J.I., 1993 *Acta. Astron.* 43, 101
- Smak J.I., 1995 *Acta. Astron.* 45, 355
- Smak J.I., 1995, Dwarf Nova Outbursts and Superoutbursts. In Wood J.H. et al. (eds) *Cataclysmic Variables and Related Objects*. Proceedings of IAU Coll. 158, in press
- Warner B., 1995, *Cataclysmic Variable Stars*. Cambridge University Press, Cambridge
- Wenzel W., 1987 *Astron. Nachr.* 308, 75
- Whitehurst R., 1994 *MNRS* 266, 35
- Whitehurst R., King A.R., 1991 *MNRS* 249, 25
- Wood J.H., Horne K., Vennes S., 1992 *ApJ* 385, 294
- Wood J.H, Naylor T., Hassall B.J.M., Ramseyer T.F., 1995 *MNRS* 273, 772
- Zhang E.-H., Robinson E.I., Nather E.R., 1986 *ApJ* 305, 740