

**Study of vector mesons in dimuon production in a large kinematic region in
p-W and S-W interactions at 200 GeV/c/nucleon**

A. L. S. Angelis^{2,a)}, J. Antos³⁾, M. Beaulieu⁴⁾, H. Beker²⁾, G. Briskin^{2,b)}, J. Bystricky⁸⁾,
M.-G. Catanesi¹⁾, P. Cerello⁹⁾, S. Dagan^{2,c)}, G. Dellacasa⁹⁾, M. Diehl⁸⁾, S. Di Liberto⁷⁾,
B. Dolgoshein⁵⁾, M. Esten^{2,d)}, C. W. Fabjan²⁾, A. Gaidot⁸⁾, M. Gallio¹⁾, P. Giubellino⁹⁾,
U. Goerlach²⁾, C. Guerra⁸⁾, L.-A. Hamel⁴⁾, S. Konovalov⁶⁾, I. Kralik³⁾, G. London⁸⁾, F.
Martelli⁹⁾, J.-P. Martin⁴⁾, M. Masera⁹⁾, M.-A. Mazzoni⁷⁾, F. Meddi⁷⁾, M.-T. Muciaccia¹⁾,
S. Muraviev⁶⁾, A. Nomerotsky²⁾, Y. Oren^{2,e)}, J.-P. Pansart⁸⁾, G. Poulard²⁾, L. Ramello⁹⁾,
L. Riccati⁹⁾, G. Rosa⁷⁾, L. Sandor³⁾, M. Sarris²⁾, E. Scomparin⁹⁾, A. Shmeleva⁶⁾, S.
Simone¹⁾, S. Smirnov⁵⁾, P. Taras⁴⁾, J. Urban³⁾, E. Vercellin⁹⁾

The HELIOS/3 collaboration

(To be submitted to Zeit. fur Physik)

Abstract

Results are presented on $\rho + \omega$, ϕ and J/Ψ production in p-W and ^{32}S -W interactions at 200 GeV/c/nucleon measured via the dimuon decay in a large kinematic region. The data are normalized to the charged particle multiplicity in the same rapidity interval. They have been collected using the HELIOS/3 muon spectrometer at the CERN SPS. The ratio $\phi/(\rho + \omega)$ increases between proton and sulphur projectiles, and is further enhanced going from peripheral to central nucleus-nucleus interactions. This results from an increase in the number of produced ϕ 's per charged particle. The increase in the ratio $\phi/(\rho + \omega)$ is not clearly dependent on p_T , but appears larger at higher rapidities, due to kinematic effects in conjunction with varying acceptance. J/Ψ production, likewise normalized to charged multiplicity, is significantly lower in S-W compared to p-W interactions.

¹⁾ University of Bari and INFN, Bari, Italy

²⁾ CERN, CH-1211 Geneva, Switzerland

³⁾ Slovak Academy of Sciences, Kosice, Slovak Republic

⁴⁾ University of Montreal, Montreal, Canada

⁵⁾ Lebedev Institute of Physics, Moscow, Russia

⁶⁾ Institute of Physics and Engineering, Moscow, Russia

⁷⁾ University of Rome "La Sapienza" and INFN Rome Italy

⁸⁾ DAPNIA, CE-Saclay, Gif/Yvette, France

⁹⁾ University of Turin and INFN Turin Italy

^{a)} also McGill University, Montreal, Canada

^{b)} presently at Tel Aviv University, Tel Aviv, Israel

^{c)} also Saclay; on leave of absence from Tel Aviv University, Tel Aviv, Israel

^{d)} also University College, London, Great Britain

^{e)} on leave of absence from Tel Aviv University, Tel Aviv, Israel

INTRODUCTION

The study of vector meson production in the environment of large hadron densities achieved in interactions of ultrarelativistic ions can be used to probe phenomena such as chiral symmetry restoration, strangeness enhancement and J/Ψ suppression in a quark gluon plasma (QGP). In the limit of chiral symmetry restoration, the quark masses tend to zero and consequently vector meson masses and widths should change [1].

Rafelski and Muller [2] have suggested that a quark-gluon plasma created in nuclear interactions would result in an enhanced production of strange quarks. In this picture, the enhancement is due to the large number of gluons which are responsible for a fast rate of flavor equilibration. The measurement of the ratio $\phi/(\rho + \omega)$ is a way to study an enhancement of strangeness production [3]. The ϕ and ω are identical in their quantum numbers and are comparable in mass; they differ primarily in their quark content. The ϕ/ω ratio in p-p interactions is $\approx 8\%$. In S-W interactions, it could be as large as 1 [3]. Another estimation including hadronic absorption but no QGP [4] predicts a smaller enhancement of this ratio, but still 6-10 times its p-p value. The latter analysis links the observation of ϕ enhancement and J/Ψ suppression. Finally, another hadronic model [5] with no QGP but with final state absorption predicts a factor of two increase of the ratios ρ/π and ϕ/π and a 40% decrease of ω/π , going from p-p to central S-W interactions.

The HELIOS/3 experiment at the CERN SPS was designed to explore the especially interesting spectrum of low mass and low transverse momentum dimuons as well as the relative production of vector mesons ϕ , ρ , and ω in a large kinematic region.

In this article, we present our measurements of the ϕ , $\rho + \omega$ and J/Ψ vector mesons, normalized to the charged particles produced in approximately the same pseudorapidity acceptance. The data are presented as a function of transverse momentum, rapidity and centrality, the latter estimated by the accompanying charged multiplicity [6].

EXPERIMENTAL SET-UP AND TRIGGER

The experimental configuration, which was installed in the H2-beam line of the CERN SPS, is shown in Figure 1 [6]. The target region with beam counters, silicon ring counters and a petal hodoscope is detailed in the inset.

The beam signal was defined by a anti-coincidence of the veto counters, LV and SV, with the beam scintillators LB and SB, or alternatively with the quartz counter Q and scintillator LB for protons and sulphur projectiles, respectively (insert in Figure 1). A gas threshold Čerenkov counter in the beam line was used to veto pions during the proton running. The six scintillators of the petal hodoscope covering the pseudorapidity region $1.8 \leq \eta \leq 3.9$ provided the interaction trigger referred to as "INT" in the following. At least 3 counters of the hodoscope were required, each being hit by 1 or by 5 to 6 particles in proton or in sulphur runs, respectively.

The use of silicon counters SiR₁ and SiR₂ was twofold. They provided a fast trigger on the charged multiplicity and detailed off-line information on the charged particle multiplicity of the event. Within the rigorously reduced space allowed for π and K decays in front of the absorber, the lay-out of the silicon pad detectors was optimized to measure the flux of particles strongly increasing in the forward rapidity region. The first counter SiR₁ consisted of 384 pads in 12 sectors covering the pseudorapidity region of $1.6 \leq \eta \leq 3.7$; the second one SiR₂, with 288 pads in 24 sectors, covered the region of $2.6 \leq \eta \leq 5.5$ (Figure 2).

The multiplicity signal in the forward part of SiR₂ ($3.7 \leq \eta \leq 5.2$, approximately the muon spectrometer acceptance) was used as a trigger condition to select the centrality

of the events. Three discrimination levels on the digital Si multiplicity were used in the sulphur runs to enhance the number of central event triggers; they correspond to the beginning of the plateau, the knee and the tail of the multiplicity distribution. As seen previously [7], the correlation of hits in the two silicon counters (Figure 3) demonstrates the proportionality of the forward and central multiplicities for low multiplicity values and a saturation of the forward region for very central events. In addition, events triggered by the interaction trigger signal (“INT”) alone were recorded (\sim minimum bias trigger).

A tungsten rod target of 1 mm diameter and of 5 or 10 mm length for sulphur or proton runs, respectively, was placed 25 cm before the hadron absorber. The two target lengths correspond to about 13% of an interaction length using the S–W inelastic cross section and about 12% of an interaction length using the p–W inelastic cross section. The absorber was optimized for low transverse mass dimuon measurements and consisted of an Al₂O₃ cone 180 cm long (6 interaction lengths) followed by 100 cm of iron (more than 5 interaction lengths). An 8 mrad conic hole centered on the beam axis, followed by a 60 cm long tungsten rod starting at 146 cm from the target, insured that most ion fragments interacted far from the target.

The large acceptance muon spectrometer consisted of a dipole magnet with a vertical field of 4.1 Tm and 7 multiwire proportional chambers for a total of 32 X,Y,U,V planes. A scintillator hodoscope H1 was placed at the exit of the magnet; after the last chamber, an 80 cm iron hadron filter was sandwiched between two trigger hodoscopes H3 and H2.

Each hodoscope, H3 and H2, was made up of two layers of scintillators, Front and Back. Each layer consisted of several rows in the (vertical) Y direction, and each row contained two (Left and Right) horizontal scintillator slabs. H3 and H2 were vertically offset by 1/2 slab height.

H3 and H2 provided the first-level dimuon trigger. Trigger rows were defined in each hodoscope by an ‘OR’ of the 4 slabs: Front/Left, Front/Right, Back/Left, Back/Right, except for the two rows in H3 immediately above and below the beam axis where the Left and Right slabs were not ‘ORed’. The muon first-level trigger required at least two coincidences between H3 and H2, a hit in an H3 row to be in coincidence with a hit in either of the H2 rows which overlap with it. The dimuon trigger was completed by requiring the reconstruction of at least two X-Z track projections in chambers PC3, PC5 and PC6. To eliminate events which are difficult to reconstruct, dimuon triggers were eliminated for most of the sulphur runs when more than six of the 11 slabs of H1 fired.

To summarize, our dimuon trigger was defined by

$$(H2 \cdot H3 \geq 2) \cdot (PC3/5/6 \geq 2) \quad or \quad (H2 \cdot H3 \geq 2) \cdot (PC3/5/6 \geq 2) \cdot (\overline{H1 > 6})$$

for the proton together with the first part of the sulphur running and for the second larger part of the sulphur running, respectively. It was used in coincidence with a multiplicity requirement, either using the interaction signal or a signal based on the second silicon ring counter, with $3.7 \leq \eta \leq 5.2$. We define a 2μ event trigger as the dimuon trigger together with its multiplicity requirement. In parallel, we accumulated events defined by the multiplicity trigger only, labeled 0μ , appropriately downscaled.

DATA REDUCTION AND ANALYSIS

During nineteen days of data taking in 1990 with the 200 GeV/c sulphur beam and the tungsten target we collected 9×10^6 dimuon triggers. This number was reduced to 3.2×10^5 events by requiring a clean sulphur beam signal, compatibility of the two silicon

ring detector information and exactly two reconstructed tracks in the spectrometer. The target selection by vertex cuts, and a restricted kinematic region (see section below on acceptance) decreased the number to 2.7×10^5 dimuon events used in the analysis, of which 68.4% were $\mu^+\mu^-$, 14.4% $\mu^+\mu^+$ and 17.2% $\mu^-\mu^-$. Similar selection criteria resulted in 44200 dimuon events for the proton beam run with 95.6% $\mu^+\mu^-$, 3.3% $\mu^+\mu^+$ and 1.1% $\mu^-\mu^-$ pairs.

Beam Selection

The information from the various beam counters was used off-line to reject events from ions other than sulphur as well as pile-up events. A multihit TDC recorded the arrival time of beam particles neighboring the one which caused the trigger. Events were kept when no other particle was found closer than 40 ns before or 300 ns after the trigger. The fraction of events rejected varies from 7 to 15% depending on the beam intensity. The quartz counter and the LB scintillator information were combined in order to define the incoming beam charge. First a correction for the pulse-height dependence on the beam intensity was applied to both counters. The pulse-heights of the two counters were then normalized and averaged, with a resolution ranging from 5 to 7%. The sulphur beam was defined by a 3 sigma cut around the peak of the weighted mean; about 5% of the events, mainly residual pile-up, were rejected. We could not separate S^{16+} from P^{15+} , however the latter contamination appears negligible in the analyzed sample.

Interaction Clean-up

Information from the 2 silicon ring counters was used to reduce the contamination due to interactions occurring out of the target. A very good correlation in the hit multiplicity between the two counters is expected in the events originating from the target. In our data, a small fraction of the events ($\simeq 1\%$) showed a high multiplicity in the second counter and a very small multiplicity in the first. These events were eliminated since they were probably due to interactions in the first ring counter.

Events from upstream interactions were rejected by a cut on the pattern correlation between the two counters. The cut was defined by looking at data taken with and without the target and then verified with a simple geometrical Monte Carlo program. This procedure removed $\simeq 3\%$ of the events.

Track Reconstruction

Track reconstruction begins behind the magnet (chambers PC3-6) by finding two straight lines in each of the X-Z and Y-Z projections. Those projections for which the χ^2 fit probability exceeds 0.1% were then associated by using the inclined U and V planes. Hits were not shared among different projections, but the same projection can be used more than once to find tracks in space. These space tracks were projected upstream, through the magnet, and associated to hits in PC0-PC2. Hits in these chambers can be shared by different tracks. The complete tracks were then fitted and the best fits retained. An overall $P(\chi^2) > 1\%$ cut was used. A muon candidate is a reconstructed track which, in addition, has associated H3 and H2 slabs, taking into account multiple scattering in the iron filter.

Determination of kinematic quantities

The momentum reconstructed in the spectrometer was corrected for the average energy loss in the absorber determined by a detailed simulation of the muon trajectories

in the absorber using GEANT 3.14; for details see [8]. The reconstructed angles for each muon in the spectrometer just behind the absorber are estimators of the true angles, but are greatly influenced by the multiple scattering in the absorber which produces two effects, a change in the true angle and an apparent track displacement. For tracks originating at the target (see next section), another independent estimator for each angle can be determined [9] by extrapolating the reconstructed track to a Z -plane at which the two effects are uncorrelated; the angles were then determined from the target position and the intersection of the track with this plane. The two independent estimators were averaged.

The mass scale was calibrated for the ϕ and J/Ψ , the first being more sensitive to energy loss and the second to the central value of the magnetic field.

Target Reconstruction

The z -coordinate of the point of closest approach to the beam axis relative to the target, Z_μ , was calculated for each muon track together with its estimated error σ_{Z_μ} whose average value increases with higher Z_μ .

A problem specific to low transverse mass dimuon measurements is an apparent forward shift of the production vertex. This shift is due to the multiple scattering of muons in the absorber material: a small opening angle between the muons is more affected than a bigger one. Thus the apparent vertex shift is particularly large for dimuons of small transverse masses, see Figure 4a. This shift is largely but not completely compensated when normalized to the error, see Figure 4b. The phenomenon is well understood and reproduced in simulation programs[9].

We kept only muons satisfying the requirement, $Z_\mu/\sigma_{Z_\mu} < 3$ and $Z_\mu < 250$ cm. These criteria were effective in rejecting dimuons which did not come from the target but they were mass-dependent retaining more dimuons at higher transverse mass; the latter effect has been taken into account in the acceptance evaluation (see below).

Acceptance and Resolution

The acceptance and resolution of the apparatus were extensively studied by Monte Carlo methods [10]. A 19x17x13 acceptance grid of m, p_T, y was determined, assuming an isotropic decay angle distribution of the dimuon. A pair was accepted if both muons were successfully reconstructed in the spectrometer. Intermediate points were calculated using an interpolation scheme based on the reconstructed kinematical variables of the muon pair. The difference between the generated and reconstructed values determined the resolution function. For example, the mass resolution at the ϕ and J/Ψ mesons was 82 and 90 MeV/ c^2 , respectively.

The acceptance for the $\rho \rightarrow \mu\mu$ is shown in Figure 5 as contour lines in the m_T vs. y plane. The acceptance is mainly determined by the momentum cutoff of the absorber due to energy losses and by the strength of the dipole magnet of the spectrometer. This leads to an acceptance which is mainly determined by the transverse mass m_T and the rapidity. In order to remove events with very low acceptance, we restricted ourselves to the kinematic region

$$m_T \geq 4(7 - 2y) \quad \text{and} \quad m_T \geq \sqrt{(2m_\mu)^2 + \left(\frac{2P_{min}}{\cosh y}\right)^2}$$

motivated by symmetric dimuon decays, where $P_{min} = 7.5$ GeV/ c is well above the minimum accepted momentum determined by the energy loss in the absorber and the deflection

of the magnetic field. These cuts reduce the kinematic acceptance for the analysis, as indicated in the figure. The remaining events are weighted by their acceptance which varies by about a factor 6.

Multiplicity Analysis

The multiplicity of charged particles measured in the pseudorapidity region $3.7 < \eta < 5.2$ was used to group our data according to their centrality. The measurement was performed starting from the number of pads having a pulse-height exceeding a threshold defined as 60% of the minimum ionizing particle signal in $300\mu\text{m}$ of silicon[11]. The most important factors that limited our resolution were the high occupancy ($\sim 70\%$ of the pads for central events) and the contamination due to secondary processes, mainly electron pair production in the tungsten target. The fluctuations due to the second effect play a very important role at low multiplicities.

The determination of the relationship of the raw multiplicity to the real multiplicity, event-by-event, relied on Monte Carlo methods. A good reproduction of the raw multiplicity data was obtained using VENUS 3.11 [12] as generator and GEANT 3.14 for a full detector simulation. We divided the data in classes of overall occupancy as measured by the total number of firing pads in the detector. For each class we tabulated the relationship between the number of firing pads belonging to a given ring, hence all in the same rapidity region, and the mean number of charged particles produced in the primary collision in the same region. These look-up tables were then used to correct the raw multiplicity. The achieved resolution, defined as the standard deviation of the ratio $(Mult_{corr.} - Mult_{true})/Mult_{true}$ ranged from less than 15% for very central interactions to about 50% for peripheral interactions. For protons, this value was 54%; the mean multiplicity in the p-W data was 4.6 with an RMS = 2.5. The statistical error on the mean was determined from the standard deviation and the number of events, while the systematic error on the mean was determined to be about 10% for classes 2–5 and 15% for classes 1 and 6.

Table 1: Charged multiplicity class vs. number of projectile participants.

class	multiplicity $3.7 < \eta < 5.2$	$\langle MULT \rangle$	Nr. of proj. part. mean (r.m.s.)
1	$MULT \leq 25$	14.0	3.7 (3.0)
2	$25 < MULT \leq 70$	43.5	14.5 (5.6)
3	$70 < MULT \leq 100$	80.8	22.7 (5.5)
4	$100 < MULT \leq 130$	115.8	26.6 (4.8)
5	$130 < MULT \leq 160$	147.0	29.4 (2.6)
6	$160 < MULT$	165.1	30.6 (1.7)
central coll.	$100 < MULT$	134.6	28.2 (3.5)

We grouped our data into six multiplicity classes, requiring that the triggers contributing to each class gave comparable average multiplicities. For each class, the average number of projectile participants was estimated by means of VENUS 3.11 (and checked with FRITIOF[13]). A central collision class was constructed from classes 4–6 by weighting each class with its fraction as determined by the minimum bias trigger. In Table 1 one can find the average number of projectile participants corresponding to each class. In this paper, class 1 was not used because of lack of statistics.

Combinatorial Background

An important background is due to uncorrelated muon pairs originating from π and K meson decays. Under the hypothesis that all like-sign muon pairs, N^{++} and N^{--} , are of this kind, the combinatorial background in the unlike-sign sample, N_c^{+-} , is proportional to the geometric mean of the like-sign samples with the proportionality constant defined as:

$$R \equiv N_c^{+-} / 2\sqrt{N^{++}N^{--}} \quad (1)$$

The constant R depends to zeroth order on the number of positive and negative particles, tending towards unity in the limit of very large multiplicities. The precise value of R was studied extensively by a full Monte Carlo simulation, using the VENUS 3.11 generator. The result is $R = 1.14 \pm 0.02$ and 1.57 ± 0.10 for the sulphur and proton runs, respectively. The results presented in this article are not sensitive to a change of 10% in R .

To obtain the precise shape of the combinatorial background, reducing the bin-to-bin statistical errors, we used μ pairs formed from muons in different like-sign events within the same multiplicity class. The same muon was used more than once depending on the proportion of $\mu^+\mu^+$ and $\mu^-\mu^-$ events. For example, in the proton sample, the same μ^+ was used 35 times and the same μ^- 100 times, while in the sulphur sample, the numbers are 26 times and 22 times, respectively. The overall normalization for the unlike-sign combinatorial background constructed by this procedure was given by real like-sign events via equation (1). More details and references concerning the combinatorial background can be found in [14].

Absorber Background

A different background of true muon pairs, contributing mainly to the low mass region, is the dilepton production in the hadron absorber by two sources: interacting hadrons and projectile fragments from the primary interaction in the target [15]. The cuts described above on the z-coordinate of the point of closest approach removed a large part of this background.

The spectral shape and the absolute rate of the remaining absorber background was estimated using a generator describing the hadron production in the target and subsequent dimuon production in the absorber (more details can be found in [16]). The primary hadron production simulation was based on the $dN^{charged}/d\eta$ measured by the HELIOS/2 collaboration [17] and a rapidity-dependent gaussian p_T distribution [15]. The hadron was tracked until its interaction in the absorber. The generator parameters were tuned to describe the experimental data taken in special runs with π beams at 20, 50 and 100 GeV/c impinging on the front face of the absorber.

The second source contributing to this type of background comes from non-interacting projectile nucleons (“fragments”). The number of fragment nucleons depends on the multiplicity class (centrality). The estimation of this background is based on the fits to distributions measured in a special run of 200 GeV/c protons hitting the tungsten plug, scaled by $A^{2/3}$ [18], like the inelastic hadronic cross section.

Since the reconstruction algorithm for muon pairs assumes the origin of the pair at the target, the reconstructed polar angles of pairs produced by the above processes are smaller than the true ones. Therefore, the effective dimuon mass and p_T of the absorber background are shifted towards lower values, and the rapidity to higher values. In the

reconstructed dimuon mass spectrum this background appears therefore only below the ρ resonance and plays a small role in the analysis presented in this paper.

Normalization to Charged Particle Production

In each multiplicity class, the sample of 0μ triggers was used to normalize the number of dimuons to the number of charged particles in the pseudorapidity interval $3.2 < \eta < 5.2$. The $\mu\mu/\text{charged}$ ratios were calculated at the trigger level, so that the multiplicity trigger acceptances cancelled. The yield of muon pairs normalized to the charged particle multiplicity is defined as:

$$\left(\frac{\mu\mu_{\text{reconstructed}}}{\text{charged}_{(3.2 < \eta < 5.2)}} \right)_{\text{trig,mult}} = \frac{1}{\langle M_{\text{trig,mult}} \rangle} \frac{F_{2\mu,\text{trig}}}{F_{0\mu,\text{trig}}} \frac{W_{\text{trig,mult}}^{\text{corr}}}{N_{\text{trig,mult}}^{\text{corr}}}$$

where

trig = trigger type

mult = multiplicity class

$\langle M_{\text{trig,mult}} \rangle$ = average charged multiplicity ($3.2 < \eta < 5.2$)

$F_{2\mu,\text{trig}}, F_{0\mu,\text{trig}}$ = 2μ and 0μ dead time correction factors

$N_{\text{trig,mult}}^{\text{corr}}$ = 0μ triggers corrected for downscaling per beam

$W_{\text{trig,mult}}^{\text{corr}}$ = sum of 2μ weights due to acceptance corrected for downscaling per beam

The events with a cut on the multiplicity in the H1-hodoscope in the 2μ trigger were corrected class-by-class using the sample with no H1 cut.

The results of the various triggers contributing to each multiplicity class were combined using weights corresponding to the overall errors of each trigger, including the errors of $\langle M_{\text{trig,mult}} \rangle$, $F_{2\mu,\text{trig}}$, $F_{0\mu,\text{trig}}$, $N_{\text{trig,mult}}^{\text{corr}}$, $W_{\text{trig,mult}}^{\text{corr}}$ [19]. The point-to-point errors of individual mass bins were calculated from corresponding parts of $W_{\text{trig,mult}}^{\text{corr}}$. The error on the overall normalization is determined by the errors of the quantities $\langle M_{\text{trig,mult}} \rangle$, $F_{2\mu,\text{trig}}$, $F_{0\mu,\text{trig}}$, $N_{\text{trig,mult}}^{\text{corr}}$, H_1 cut corrections and the background estimation, taking into account the correlations.

The Fits to the Resonances

The dimuon mass spectra were fitted between 0.3 and 3.6 GeV/c² by the function

$$F(m) = P_1[0.6BW_\rho(m) + G_\omega(m) + P_2G_\phi(m)] + P_3PSI(m) + C(m)$$

where G_i are gaussians describing the ω and ϕ meson peaks, BW_ρ a p-wave Breit-Wigner function [20, 21], $PSI(m)$ a GEANT-determined J/Ψ line shape and $C(m)$ an empirical function describing the continuum. The constant 0.6, representing $(\sigma B_{\mu\mu})_\rho / (\sigma B_{\mu\mu})_\omega$, was determined using $\sigma(\rho) = \sigma(\omega)$, as measured in p-p interactions [22], and the dimuon branching ratios, $B_{\mu\mu}$, from the Particle Data Group [23]. The results of this paper are not sensitive within errors to this assumption.

The Breit-Wigner function $BW_\rho(m)$ has a mass-dependent width. Relative to $\pi\pi$, the $\mu\mu$ partial width has an additional m^{-4} factor from the photon propagator suppressing the high mass shoulder of the Breit-Wigner function. The ρ meson shape in the dimuon mass spectrum was taken as:

$$BW_\rho(m) = \sigma(m) \frac{\pi^{-1} m^2 \Gamma_{\rho \rightarrow \mu\mu}(m)}{(m^2 - m_\rho^2)^2 + m^2 \Gamma_{\rho \rightarrow \pi\pi}^2(m)} \quad (2)$$

where

$$\Gamma_{\rho \rightarrow \mu\mu}(m) = \frac{k}{k_0} \frac{(2m_\mu^2 + m^2)}{(2m_\mu^2 + m_\rho^2)} \frac{m_\rho^6}{m^6} \Gamma_{\rho \rightarrow \mu\mu}^0 \quad \text{and} \quad \Gamma_{\rho \rightarrow \pi\pi}(m) = \left(\frac{k}{k_0}\right)^3 \left(\frac{m_\rho}{m}\right)^2 \Gamma_{\rho \rightarrow \pi\pi}^0$$

$$k = \sqrt{0.25m^2 - m_\mu^2} \quad \text{and} \quad k_0 = \sqrt{0.25m_\rho^2 - m_\mu^2}$$

and $\sigma(m)$ is the cross section for the production of a stable particle with mass m (summed over all spin states).

The functional form of this cross section was found in analogy with the approach used for ρ meson searches in the $\pi\pi$ channel, where it is assumed to behave like the nonresonant background under the resonance. To avoid double counting the final state phase space (see e.g. [20]), we used

$$\sigma(m) = PS(m) \frac{m}{k_\pi}. \quad (3)$$

where $PS(m)$ is the uncorrelated $\pi\pi$ spectrum. The QGSM model [24] for p-p interactions at 200 GeV/c was used to obtain the form of $PS(m)$ by parametrizing it as

$$PS(m) = \left(\frac{k_\pi}{m}\right)^\beta \exp(ak_\pi + bk_\pi^2)$$

$$k_\pi = \sqrt{0.25m^2 - m_\pi^2},$$

$\beta = 0.5725$, $a = -4.5729$ and $b = 0.3130$. In order that $PS(m)$ could be defined also below the $\pi\pi$ threshold mass, the pion mass was replaced with muon mass. Within errors, the results of this paper are also not sensitive to these assumptions.

Equation (2) was convoluted with the experimental resolution.

The mean masses of the ω and ϕ resonances were fixed at their nominal values, while the gaussian variances were equal to the experimental resolution. The results of the fits indicate that the experimental resolution is well understood.

The function $PSI(m)$ was determined between 2.5 and 3.5 GeV/c² by a full GEANT simulation.

Equation (4) represents the continuum function chosen for its behavior at threshold and high mass (i.e. Drell-Yan-like).

$$C(m) = (P_4 e^{P_5 m} + \frac{P_6}{m^3} e^{P_7 m}) (1 - \frac{2m_\mu}{m})^{P_8} \quad (4)$$

As examples of the data and the fits, Figure 6 displays the minimum bias p-W and central S-W dimuon mass spectra with the fits and the individual contributions superimposed: continuum, J/Ψ , ρ , ω and ϕ .

RESULTS

In the following, we present the dependence on centrality, transverse momentum p_T and rapidity y of vector meson production as well as the ratio $\phi/(\rho + \omega)$. Furthermore, we present some results on J/Ψ production.

The measured projectile participant and multiplicity dependence of the $\phi/(\rho + \omega)$ ratio is given in Table 2 and Figure 7, respectively. The data in Figure 7 have been normalized to the proton value. The horizontal error bars reflect the binning of the charged

Table 2: Centrality dependence of vector mesons

projectile participants	$\phi/(\rho + \omega)$	$(\phi/charged)$ $\times 10^6$	$(\rho + \omega/charged)$ $\times 10^6$
1	0.138 ± 0.009	1.00 ± 0.06	6.83 ± 0.17
14.5	0.224 ± 0.082	1.88 ± 0.65	8.32 ± 2.34
22.7	0.253 ± 0.086	1.63 ± 0.50	6.44 ± 1.30
26.6	0.217 ± 0.032	1.72 ± 0.23	7.92 ± 0.63
29.4	0.318 ± 0.033	2.15 ± 0.18	6.77 ± 0.47
30.6	0.317 ± 0.045	2.15 ± 0.24	6.75 ± 0.61

multiplicity. The ratio $\phi/(\rho + \omega)$ clearly increases with centrality in S–W interactions and going from p–W to central S–W interactions.

For proton and central sulphur interactions, three p_T and three y bins were considered. Note that these bins refer only to the dimuons, not to the charged particles used for the normalization. The ratios evaluated in different intervals of transverse momentum are presented in Table 3 and show an enhancement of the ratio $\phi/(\rho + \omega)$ from proton to central sulphur data, with no clear dependence on p_T , while the rapidity results (Table 4) show that this enhancement increases strongly with rapidity. We will discuss the influence of kinematics on this result in the next section.

Table 3: Transverse momentum dependence of $\phi/(\rho + \omega)$.

	central S–W	p–W	(S–W)/(p–W)
$p_T \leq 0.35$	0.266 ± 0.041	0.137 ± 0.017	1.94 ± 0.38
$0.35 < p_T \leq 0.6$	0.204 ± 0.042	0.145 ± 0.015	1.41 ± 0.32
$0.6 < p_T$	0.315 ± 0.032	0.136 ± 0.013	2.32 ± 0.32

Table 4: Rapidity dependence of $\phi/(\rho + \omega)$.

	central S–W	p–W	(S–W)/(p–W)
$y \leq 3.9$	0.314 ± 0.031	0.232 ± 0.017	1.35 ± 0.17
$3.9 < y \leq 4.4$	0.289 ± 0.040	0.078 ± 0.009	3.71 ± 0.67
$4.4 < y$	0.202 ± 0.021	0.038 ± 0.008	5.32 ± 1.25

NA38 measurements [25] of the ratio $\phi/(\rho + \omega)$ performed at relatively high transverse momentum, complementary to our experiment, have shown an increase relative to centrality which is consistent with our result; their result corresponds approximately to our first rapidity bin combined with our last transverse momentum bin.

To study separately the ϕ and $(\rho + \omega)$ behavior, we have determined the $\phi/charged$ and $(\rho + \omega)/charged$ ratios as a function of centrality (Table 2 and Figure 8). The values in the figures have been normalized to the proton point. We observe that there is an increase in the $\phi/charged$ ratio, while the $(\rho + \omega)/charged$ ratio remains constant. We have also determined the rapidity variation of $\phi/charged$ (Table 5) and $(\rho + \omega)/charged$

(Table 6). In all cases, there is a strong decrease at higher rapidity but this decrease is quite different for proton and central sulfur interactions, depending on the resonance. In particular, the ϕ falls about half as much in S–W than in p–W interactions, while the $\rho + \omega$ falls twice as fast for S–W than for p–W interactions.

The behavior of increased ϕ production has also to be seen in the context of increased production in heavy ion interactions of other strange particles, like kaons and strange baryons. However, for a quantitative interpretation of the observed enhancement, our dimuon acceptance and kinematic effects between p–W and S–W collisions have to be taken into account. We have attempted this for two specific models and will present our results in the next section of this paper.

Table 5: Rapidity dependence of $\phi/charged$

rapidity region	central S–W $\times 10^6$	p–W $\times 10^6$	(S–W)/(p–W)
$y \leq 3.9$	1.58 ± 0.13	0.83 ± 0.05	1.90 ± 0.19
$3.9 < y \leq 4.4$	0.50 ± 0.15	0.17 ± 0.02	2.94 ± 0.94
$4.4 < y$	0.20 ± 0.02	0.052 ± 0.011	3.85 ± 0.90

Table 6: Rapidity dependence of $(\rho + \omega)/charged$

rapidity region	central S–W $\times 10^6$	p–W $\times 10^6$	(S–W)/(p–W)
$y \leq 3.9$	5.04 ± 0.37	3.47 ± 0.14	1.45 ± 0.12
$3.9 < y \leq 4.4$	1.56 ± 0.49	2.15 ± 0.08	0.72 ± 0.23
$4.4 < y$	1.00 ± 0.08	1.37 ± 0.05	0.73 ± 0.06

The $J/\Psi/charged$ ratio in our kinematic region is $(1.20 \pm 0.10)10^{-7}$ in p–W interactions. Table 7 gives the dependence on centrality relative to p–W interactions and qualitatively confirms the suppression seen by [26]. Note however that our normalization to charged particles (produced by soft processes) differs from their normalization to the continuum (produced by hard processes).

For completeness, Table 8 shows the rapidity dependence of J/Ψ production in minimum bias p–W and central S–W interactions.

Table 7: Ratio of $J/\Psi/charged$ production for central S–W interactions relative to minimum bias p–W interactions as a function of centrality.

Nr. of proj. part. mean	S–W/p–W $J/\Psi/charged$
26.6	0.66 ± 0.14
29.4	0.65 ± 0.10
30.6	0.71 ± 0.15

Table 8: Rapidity dependence of $J/\Psi/charged \times 10^7$.

rapidity region	central S-W	p-W
$y \leq 3.9$	0.57 ± 0.07	1.03 ± 0.09
$3.9 < y \leq 4.4$	0.095 ± 0.020	0.11 ± 0.02
$4.4 < y$	0.0092 ± 0.0053	0.0098 ± 0.0062

We emphasize that our vector meson results are presented as ratios in a fixed rapidity window, sometimes with respect to each other and sometimes with respect to charged particles in a similar but not identical pseudorapidity window. The A^α -dependence of particle production usually depends on Feynman's x_F with a specific $\alpha(x_F)$ for each particle; for example, see [27, 28]. Therefore, the interpretation of our results must necessarily entail a comparison with models.

COMPARISON WITH MODELS AND DISCUSSION

Quantitative comparisons were made with two physics generators, DTUNUC 1.02 [29] and VENUS 4.12 [12], each based on the dual parton model (DPM) [30] with the addition of an intranuclear cascade model. Version 4.12 of VENUS was used (as opposed to version 3.11) since it keeps both the parent vector meson as well as its decay products which are necessary for the normalization to charged particles. The naive treatment of intranuclear cascade processes overestimates particle yields in particular in the region of high hadron density where independent binary interactions are implausible. One way to modulate the treatment is by the introduction of a formation zone which suppresses the cascading of high-energy secondaries. This is introduced via a simple formation time concept [29]. Another way is to introduce cluster formation and decay [12]. In either case, the treatment is ad-hoc.

Within the DPM, the high-energy projectile undergoes a multiple scattering process. Particles are produced by the fragmentation of two colorless parton-parton chains constructed from valence quarks and sea- $q\bar{q}$ pairs of the multiply interacting particles. In this model, the number of chains with sea-quarks at their ends grows with the complexity of the target and projectile nuclei. For example, in p-p interactions at our energies, there are no chains with sea quarks to first order, while they exist in p-A collisions, increasing with A. In particular, since the masses of these chains are in the GeV energy range, there is no reason to suppose strange quark suppression. Therefore an SU(3) flavor symmetry is possible for these sea-quarks. This mechanism leads to a rise in the relative yield of strange particles as the final state multiplicity increases.

The predictions of the two models, normalized to the proton value, are shown on Figure 7 and Figure 8. Relative to the data, VENUS predicts a more rapid rise of $\phi/(\rho+\omega)$ as a function of centrality. The discrepancy is due to the larger rise in $\phi/chgd$ while the $(\rho+\omega)/chgd$ ratio is well reproduced. The DTUNUC predictions are more consistent with the data.

For VENUS, the data are about 1.5-2 times the absolute predictions (not shown) of $(\rho+\omega)/chgd$ and $\phi/chgd$ ratios while the $\phi/(\rho+\omega)$ ratios are quite consistent with the data, except that the proton data point is about 1.5 times the prediction. The latter seems due to a larger underestimation of the $\phi/chgd$ compared to the data than $(\rho+\omega)/chgd$ in p-W interactions. It should be pointed out that VENUS overestimates the charged particle production as compared to data [31]. For DTUNUC, the absolute predictions of $(\rho+\omega)/chgd$ and $\phi/chgd$ ratios are similarly low but somewhat independent of projectile.

Other models predict that the ratio $\phi/(\rho + \omega)$ should rise either by about 6-10 [4] or 1.8 [5] over its p-p value, quite higher or somewhat lower than our data, respectively. The prediction [32] that this rise appears only at high p_T is invalidated by the results of this paper.

CONCLUSIONS

In this paper we have presented a measurement of $\rho + \omega$, ϕ and J/Ψ vector meson production in p-W and S-W interactions at 200 GeV/c/nucleon for rapidities larger than 3.2, the rapidity threshold depending on the transverse mass. Within this kinematic range, the ϕ resonance is enhanced in heavy ion collisions by a factor of about two relative to the other low mass resonances. As a function of the charged particle multiplicities in our acceptance, the $\rho + \omega$ combination is unchanged, the ϕ is enhanced and J/Ψ is suppressed.

From the comparisons to the DTUNUC and VENUS models, we conclude that there is no compelling evidence in our data on vector meson production for physics processes not already included in these models, such as the quark-gluon plasma. To make this a firmer conclusion, these models should reproduce the unusual multiplicity dependence of ϕ production measured in pBe interactions at 450 GeV/c [33] as well as the measurements of other particles containing strange quarks, such as hyperons.

Acknowledgements

The HELIOS/3 collaboration wishes to thank the staff of the PS-SPS accelerator complex for their outstanding performance. The valuable contributions of the technical staff of CERN and the collaborating institutes are gratefully acknowledged. We are grateful for support by the Direction des Sciences de la Matière (CEA, France) and the Istituto Nazionale di Fisica Nucleare of Italy.

References

- [1] R. Pisarsky, Phys. Lett. **B110** (1982) 155
- [2] J. Rafelsky and B. Muller, Phys. Lett. **B48** (1982) 1066 and J. Rafelsky, Nucl. Phys. **A418** (1984) 215
- [3] A. Shor, Phys. Rev. Lett. **54** (1985) 1122
- [4] P. Koch, U. Heinz, J. Pisut, Phys. Lett. **B243** (1990) 149; Z. Phys. **C47** (1990) 477
- [5] M. Berenguer, H. Sorge, and W. Greiner, UFTP preprint **356** (1994) and references therein, and H. Sorge et al., Z. Phys. **C59** (1993) 85
- [6] more details are to be found in F. Martelli, Produzione di mesoni vettoriali in interazioni S-W e p-W a 200 GeV/nucleone, Tesi del Dottorato di Ricerca in fisica, V Ciclo, (1993)
- [7] P. Giubellino et al., Nucl. Instr. Meth. **A275** (1989) 89
- [8] M. Sarris, HELIOS/3 note 26
- [9] G. London, HELIOS note 319, and references therein
- [10] S. Dagan, HELIOS/3 note 33; J. Bystricky and G. London, HELIOS/3 note 34
- [11] T. Åkesson et al., Nucl. Phys. **B333** (1990) 48
- [12] K. Werner, Phys. Rep. **232** (1993) 88 and references therein; K. Werner, University of Heidelberg preprint **HD-TVP-93-1** (1993).
- [13] B. Andersson, G. Gustafsson and B. Nilsson-Almqvist, Nucl. Phys. **B281** (1987) 289
- [14] J. Bystricky et al., to be submitted for publication to Nucl. Instr. Methods
- [15] G. London, HELIOS/3 note 83

- [16] “Excess of continuum dimuon production at masses between threshold and the J/Ψ in S–W interactions at 200 GeV/c/nucleon”, A.L.S.Angelis et al., to be submitted for publication to Zeit. fur Phys.
- [17] M. Masera, *24th Rencontres de Moriond*, New results in hadronic interactions, J. Tran Thanh Van editor, Editions Frontieres (1989) 477
- [18] G. Baroni et al., Nucl. Phys **A531** 1991 691
- [19] J. Bystricky, HELIOS/3 note 82
- [20] J.D.Jackson, Il Nuovo Cimento **34** (1964) 1644.
- [21] Z.Y.Fang, G.Lopez Castro and J.Pestieau, Use and Misuse of the Breit-Wigner Formula. Preprint **UCL-IPT 87-07**. May 1987.
- [22] T. Åkesson et al., First measurement of branching ratio $\omega^0 \rightarrow \mu^+\mu^-$, to be published
- [23] Review of Particle Properties, Phys. Rev. **D50** (1994) 1173
- [24] N.S.Amelin, K.K.Gudima and V.D.Toneev, NATO ASI series B: Physics **2116** (1989) 473, and preprint **GSI 89-52**
- [25] C. Baglin et al., Phys. Lett. **B272** (1991) 449
- [26] C. Baglin et al., Phys. Lett. **B270** (1991) 105
- [27] W. Geist, Nucl. Phys. **A525** (1991) 149c
- [28] R. Salmeron, Nucl. Phys. **A566** (1994) 199
- [29] J.Ranft, SSC-report **150** (1987),
H.-J.Möhring, J.Ranft, Z. Phys. **C52** (1991) 643,
I.Kawrakow, H.-J.Möhring, J.Ranft, Z. Phys. **C56** (1992) 115.
- [30] A. Capella et al., Nucl. Phys. **B241** (1984) 75
- [31] T. Åkesson et al., Nucl. Phys. **B333** (1989) 1
- [32] F. Grassi, H. Heiselberg, Phys. Lett. **B267** (1991) 1
- [33] T. Åkesson et al., Z. Phys. **C68** (1995) 47

HELIOS/3 set-up






Figure 1: Overall experimental setup with target region inset.

Ring counter 1



Ring counter 2



1 cm

Figure 2: Silicon ring layout




Figure 3: Central multiplicity vs. forward multiplicity. Notice that the forward multiplicity begins to saturate at high central multiplicity.




Figure 4: Dimuon transverse mass vs. a) reconstructed origin along beam and b) reconstructed origin along beam normalized by its error.




Figure 5: Acceptance contours on the m_T vs. y plot for the central mass value of the ρ meson. The region at high rapidity and transverse mass is outside the kinematic limit.




Figure 6: Dimuon mass spectra for (a) minimum bias p-W and (b) central S-W interactions with fits and individual contributions superimposed. The J/Ψ and ρ contributions have been added to the continuum, followed by the ω contribution and, finally, the ϕ contribution. The log mass scale was chosen to emphasize the ρ, ω, ϕ region




Figure 7: The $\phi/(\rho + \omega)$ ratio, normalized to the proton point, vs. the charged multiplicity. A comparison with the predictions of VENUS 4.12 and DTUNUC 1.02 is made.





Figure 8: The $\phi/\text{charged}$ and $(\rho+\omega)/\text{charged}$ ratio, normalized to the proton point, vs. the charged multiplicity. A comparison with the predictions of VENU S 4.12 and DTUNUC 1.02 is made.