

CERN-TH/98-129  
RU98-3-B  
LAPTH683/98  
LP THE ORSAY 98/31

## UNIVERSALITY OF LOW-ENERGY SCATTERING IN (2+1) DIMENSIONS

**Khosrow Chadan**

Laboratoire de Physique Théorique et Hautes Energies<sup>1</sup>  
Université Paris-Sud, F-91405 Orsay, France

**N. N. Khuri**

Department of Physics  
The Rockefeller University, New York, New York 10021

**André Martin**

TH Division, CERN, Geneva, Switzerland,  
Laboratoire de Physique Théorique ENSLAPP  
F-74941 Annecy-le-Vieux, France, and  
Department of Physics, The Rockefeller University  
New York, New York 10021

**Tai Tsun Wu**

Gordon McKay Laboratory, Harvard University  
Cambridge, Massachusetts 02138-2901, and  
TH Division, CERN, CH - 1211 Geneva 23

### ABSTRACT

We prove that, in (2+1) dimensions, the  $S$ -wave phase shift,  $\delta_0(k)$ ,  $k$  being the c.m. momentum, vanishes as either

$$\delta_0 \rightarrow \frac{c}{\ln(k/m)} \quad \text{or} \quad \delta_0 \rightarrow O(k^2)$$

as  $k \rightarrow 0$ . The constant  $c$  is universal and  $c = \pi/2$ . This result is established first in the framework of the Schrödinger equation for a large class of potentials, second for a massive field theory from proved analyticity and unitarity, and, finally, we look at perturbation theory in  $\phi_3^4$  and study its relation to our non-perturbative result. The remarkable fact here is that in  $n$ -th order the perturbative amplitude diverges like  $(\ln k)^n$  as  $k \rightarrow 0$ , while the full amplitude vanishes as  $(\ln k)^{-1}$ . We show how these two facts can be reconciled.

CERN-TH/98-129  
RU98-3-B  
LAPTH683/98  
LP THE ORSAY 98/31  
April 1998

---

<sup>1</sup>Laboratoire associé au CNRS URA 63.

# 1. INTRODUCTION

Quantum field theories in 2+1 dimensions provide us with a useful field of investigation not only for theoretical and mathematical issues, but also in some cases for actual physical problems. Scattering in 2+1 dimensions has the advantage of resembling scattering in 3+1 dimensions much more than 1+1 dimensions in the sense that the scattering amplitude depends on two variables and that, also, the scattering matrix is non-trivial only if production processes exist. As we shall see in Section 3, the analyticity-unitarity programme follows the same lines as in 3+1 dimensions. On the other hand, a non-trivial  $\lambda\phi^4$  massive field theory can be constructed in 2+1 dimensions [1]. Another merit of two space dimensions is that it covers situations occurring in condensed matter physics.

A remarkable property, which is completely different from what happens in 3+1 dimensions is the threshold behaviour of the scattering amplitude. If  $\delta_0$  is the  $S$  wave phase shift, then, as  $k \rightarrow 0$  either

$$\delta_0 \sim \frac{\pi}{2 \ln k} \quad (1.1)$$

or

$$\delta_0 = 0(k^2) \quad (1.2)$$

In this paper we prove this result in three situations:

- i) in the framework of the Schrödinger equation for a very large class of potentials under conditions which will be specified in Section 2 and which are presumably very difficult to weaken; in this case, it appears that the generic case is given by Eq. (1.1), Eq. (1.2) representing an exceptional case; this is done in Section 2;
- ii) in massive axiomatic field theory, in combination with unitarity and a standard assumption of smooth behaviour of the scattering amplitude; this is carried in Section 3;
- iii) starting from the perturbation expansion of  $\lambda\phi^4$  and resumming the leading logs, while individual terms diverge like  $(\ln k)^n$  near  $k = 0$ . This is done in Section 4.

To us and to many of our friends and colleagues, property (1.1)-(1.2) was a great surprise even though a related property appeared already in the theory of antennas [2].

However, it turned out that we had predecessors. Concerning i), the potential case, Bollé and Gestezy [3] found property (1.1) for a class of potentials exponentially decreasing at infinity and Averbush [4] treated the case of a potential with a strictly finite range. However, these authors missed the exceptional case (1.2).

Concerning ii) and iii), Bros and Iagolnitzer [5], in the framework of general  $S$  matrix theory, studying primarily the Riemann sheet structure of scattering amplitudes near threshold, obtained for the two-body  $\rightarrow$  two-body case an equation which implies the alternative between (1.1) and (1.2). If one combines their paper with the early work of Bros, Epstein and Glaser on analyticity of the scattering amplitude near physical points [6] one can consider this as

an axiomatic proof. They also carry the resummation of a subclass of perturbation graphs, without, however, showing that they are the ones associated to the leading logs.

In spite of this, we believe that we have the duty to present our own results with our own methods, which we believe sometimes more accessible, covering in a synthetic way the potential case and the field theoretical case and, in some instances, improving or correcting previous work.

## 2. TWO-DIMENSIONAL POTENTIAL SCATTERING

In two dimensions the partial wave expansion of the scattering amplitude  $T(k, \theta)$  is given by

$$T(k, \theta) = \frac{1}{\sqrt{k}} \sum_{n=0}^{\infty} \epsilon_n (e^{i\delta_n} \sin \delta_n) \cos n\theta, \quad (2.1)$$

where  $\epsilon_0 = 1$ ,  $\epsilon_n = 2$  for  $n \geq 1$ . The phase shifts  $\delta_n(k)$  are obtained in the standard way from the solutions of the Schrödinger equation. In this paper we are interested mainly in the term  $n = 0$ .

The  $n = 0$  solutions,  $u(k, r)$ , satisfy

$$\left[ \frac{d^2}{dr^2} + \frac{1}{4r^2} + k^2 - gV(r) \right] u(k, r) = 0. \quad (2.2)$$

Without loss of generality,  $g$  is taken to be non-negative. Equation (2.2), under conditions on  $V(r)$  to be specified below, has two independent solutions: behaving like  $\sqrt{r}$  and  $\sqrt{r} \ln r$  as  $r \rightarrow 0$ . We take as a regular solution

$$u(k, 0) = 0, \quad u(k, r) \sim \sqrt{r}, \quad (2.3)$$

corresponding to a finite wave function at the origin. For a discussion of this choice, see Appendix A.

The phase shift,  $\delta_0(k)$ , is defined by

$$u(k, r) \xrightarrow{r \rightarrow \infty} c\sqrt{r} [\cos \delta_0 J_0(kr) - \sin \delta_0 Y_0(kr)]. \quad (2.4)$$

The sign of the second term is chosen to correspond to the definition of  $\delta_0$  in the three-dimensional case, i.e.,  $u \rightarrow c\sqrt{\pi/2} \cos(kr - \pi/4 + \delta_0)$  as  $r \rightarrow \infty$ . By rearranging terms in Eq. (2.4), we get

$$u(k, r) \xrightarrow{r \rightarrow \infty} c\sqrt{r} e^{-i\delta_0} [H_0^{(2)}(kr) + e^{2i\delta_0} H_0^{(1)}(kr)]. \quad (2.5)$$

We can always choose  $u(k, r)$  such that

$$u(k, r) \xrightarrow{r \rightarrow \infty} -\frac{1}{2i\sqrt{k}} [e^{-i(kr-\pi/4)} + S(k)e^{+i(kr-\pi/4)}], \quad (2.6)$$

where we have used the asymptotic formulas for  $H_0^{(1),(2)}(z)$  for large  $|z|$ , and

$$S(k) \equiv e^{2i\delta_0(k)}. \quad (2.7)$$

The Jost functions in this case are solutions of (2.2) finite at  $r = 0$ , which we denote as  $f_{\pm}(k, r)$  with the asymptotic behaviour

$$f_{\pm}(k, r) \xrightarrow{r \rightarrow \infty} e^{\mp i(kr - \pi/4)}. \quad (2.8)$$

We can thus write

$$u(k, r) = -\frac{1}{2i\sqrt{k}} [f_+(k, r) + S(k)f_-(k, r)]. \quad (2.9)$$

It is convenient to follow a method of treating singular potentials [7]. We shall see below how this simplifies the task of taking the limit  $k \rightarrow 0$ . Following ref. [7], we define  $g(k, r)$  as

$$g(k, r) \equiv \frac{1}{2i\sqrt{k}} [f_+(k, r) + f_-(k, r)]. \quad (2.10)$$

The sign here is different from that in the three-dimensional case. From Eq. (2.9) we now have

$$u(k, r) = -[g(k, r) + A(k)f_-(k, r)], \quad (2.11)$$

where  $A(k)$  is the  $n = 0$  scattering amplitude,

$$A(k) \equiv \frac{1}{2i\sqrt{k}} [S(k) - 1] \equiv \frac{1}{\sqrt{k}} e^{i\delta_0} \sin \delta_0. \quad (2.12)$$

The condition  $u(k, r) \rightarrow 0$  as  $r \rightarrow 0$  gives us

$$A(k) = -\lim_{r \rightarrow 0} [g(k, r)/f_-(k, r)]. \quad (2.13)$$

Notice that this limit is always finite. This is because  $f_-$ , being a combination of  $\text{Re } f_-$  and  $\text{Im } f_-$ , i.e., of two linearly independent solutions of (2.2), has to behave as  $f_- \sim \sqrt{r} \ln r$  as  $r \rightarrow 0$ .

The asymptotic behaviour of  $u(k, r)$  can be written as

$$u(k, r) \xrightarrow{r \rightarrow \infty} \frac{i \cos(kr - \pi/4)}{\sqrt{k}} - A(k)e^{i(kr - \pi/4)}. \quad (2.14)$$

This follows from Eqs. (2.8) and (2.9).

Following ref. [7], we introduce a Green's function  $G(r, r')$  for  $r, r' > 0$ , defined by

$$\left[ \frac{d^2}{dr^2} + \frac{1}{4r^2} + k^2 \right] G(r, r') \equiv \delta(r - r'). \quad (2.15)$$

This  $G$  is given explicitly by

$$G(r, r') = \frac{\pi}{2} \sqrt{rr'} [J_0(kr)Y_0(kr') - J_0(kr')Y_0(kr)]\theta(r' - r), \quad (2.16)$$

where  $J_0$  and  $Y_0$  are the standard Bessel functions of the first and second kind.

The next step is to introduce a  $u_0(k, r)$  which is a solution of the free,  $V = 0$ , Schrödinger equation. We set

$$u_0(k, r) \equiv u(k, r) - g \int_0^\infty dr' G(r, r')V(r')u(k, r'). \quad (2.17)$$

From Eq. (2.15) it is now obvious that

$$\left[ \frac{d^2}{dr^2} + \frac{1}{4r^2} + k^2 \right] u_0(k, r) = 0. \quad (2.18)$$

As  $r \rightarrow \infty$ ,  $u_0 \rightarrow u$ , and from Eq. (2.14) it is clear that  $u_0$  is given by

$$u_0(k, r) = \sqrt{\frac{\pi}{2}} i\sqrt{r} J_0(kr) - \sqrt{\frac{\pi}{2}} A(k)\sqrt{kr} H_0^{(1)}(kr). \quad (2.19)$$

The integral equation for  $u$  can now be written as

$$u(k, r) = u_0(k, r) + g \int_r^\infty dr' \tilde{G}(k; r, r')V(r')u(k, r'), \quad (2.20)$$

with

$$\tilde{G}(k; r, r') = \frac{\pi}{2} \sqrt{rr'} [J_0(kr)Y_0(kr') - J_0(kr')Y_0(kr)]. \quad (2.21)$$

Using Eqs. (2.11) and (2.19), we can get from (2.20) two separate integral equations for  $g(k, r)$  and  $f_-(k, r)$ . These are

$$g(k, r) = -i\sqrt{\frac{\pi}{2}} \sqrt{r} J_0(kr) + g \int_r^\infty dr' \tilde{G}(k; r, r')V(r')g(k, r') \quad (2.22)$$

and

$$f_-(k, r) = \sqrt{\frac{\pi}{2}} \sqrt{kr} H_0^{(1)}(kr) + g \int_r^\infty dr' \tilde{G}(k; r, r')V(r')f_-(k, r'). \quad (2.23)$$

These last two equations are the same except for the inhomogeneous term. We are interested in studying them in the limit of small  $k$ . Before we can do that, it is convenient to remove a  $\sqrt{k}$  factor from  $f_-$  and define  $\tilde{f}_-(k, r)$  as

$$\tilde{f}_-(k, r) \equiv \frac{1}{\sqrt{k}} f_-(k, r). \quad (2.24)$$

With this definition, Eq. (2.13) becomes

$$e^{i\delta_0(k)} \sin \delta_0(k) = -\lim_{r \rightarrow 0} [g(k, r)/\tilde{f}_-(k, r)]. \quad (2.25)$$

We now take the  $k \rightarrow 0$  limit of Eq. (2.22) and the equation corresponding to (2.23) for  $\tilde{f}_-$ . Using

$$\frac{\pi}{2} [J_0(kr)Y_0(kr') - J_0(kr')Y_0(kr)] = \ln \frac{r'}{r} + O(k^2) \quad (2.26)$$

for small  $k$ , we get

$$g(k, r) = -i\sqrt{\frac{\pi}{2}} \sqrt{r} + g \int_r^\infty dr' \sqrt{rr'} \left( \ln \frac{r'}{r} \right) V(r') g(k, r') + O(k^2) \quad (2.27)$$

and

$$\begin{aligned} \tilde{f}_-(k, r) &= i\sqrt{\frac{2}{\pi}} \left( \ln k + \ln r - \ln 2 + \gamma - i\frac{\pi}{2} \right) \sqrt{r} \\ &\quad + g \int_r^\infty dr' \sqrt{rr'} \left( \ln \frac{r'}{r} \right) V(r') \tilde{f}_-(k, r') + O(k^2), \end{aligned} \quad (2.28)$$

where  $\gamma$  is Euler's constant. For  $r > 0$ , taking the  $k \rightarrow 0$  limit under the integral sign is allowed if we assume

$$\int_a^\infty r' dr' (1 + |\ln r'|^2) |V(r')| < \infty, \quad a > 0. \quad (2.29)$$

We shall discuss this condition in more detail later.

At this stage, we introduce two functions,  $A(r)$  and  $B(r)$ , defined by the following integral equations:

$$A(r) = 1 + g \int_r^\infty r' dr' \left( \ln \frac{r'}{r} \right) V(r') A(r') \quad (2.30)$$

and

$$B(r) = \ln r + g \int_r^\infty r' dr' \left( \ln \frac{r'}{r} \right) V(r') B(r'). \quad (2.31)$$

It is clear from inspecting Eqs. (2.27) and (2.28) that

$$A(r) \equiv \lim_{k \rightarrow 0} \left[ \frac{ig(k, r)}{\sqrt{\pi/2} \sqrt{r}} \right] \quad (2.32)$$

and

$$\left[ \frac{-i\tilde{f}_-(k, r)}{\sqrt{2/\pi} \sqrt{r}} \right] \equiv \left[ A(r) \left( \ln k - \ln 2 + \gamma - i\frac{\pi}{2} \right) + B(r) \right] + O(k^2). \quad (2.33)$$

Thus, for small  $k$  we have

$$- \left[ \frac{g(k, r)}{\tilde{f}_-(k, r)} \right] = \frac{(\pi/2)A(r)}{A(r) \left( \ln k - \ln 2 + \gamma - i\frac{\pi}{2} \right) + B(r)} + O(k^2). \quad (2.34)$$

Our task is now to study the existence of solutions  $A(r)$  and  $B(r)$  of the two integral equations (2.30) and (2.31), and more specifically, to study the behaviour of  $A$  and  $B$  for small  $r$ .

In Appendix B, we shall prove that for the general class of potentials,  $V(r)$ , satisfying

$$\text{A)} \quad \int_0^\infty r' dr' |V(r')| (|\ln r'| + 1) < \infty \quad (2.35)$$

and

$$\text{B)} \quad \int_a^\infty r' dr' |V(r')| (\ln r')^2 < \infty, \quad a > 1, \quad (2.36)$$

the solutions  $A(r)$  and  $B(r)$  exist for all  $r > 0$ , and furthermore, near  $r = 0$  one has the behaviour

$$A(r) = [-gC_a(g) + o(1)] \ln r \quad (2.37)$$

and

$$B(r) = [1 - gC_b(g) + o(1)] \ln r. \quad (2.38)$$

Here,

$$C_a(g) = \int_0^\infty r dr V(r) A(r) \quad (2.39)$$

and

$$C_b(g) = \int_0^\infty r dr V(r) B(r). \quad (2.40)$$

Both integrals for  $C_a$  and  $C_b$  are absolutely convergent since one can easily show that, as  $r \rightarrow \infty$ ,  $A$  and  $B$  have the bounds

$$|A(r)| < \text{Const.}, \quad |B(r)| < \text{Const.} |\ln r|, \quad (2.41)$$

for  $r > r_0 > 1$ . The convergence of Eqs. (2.39) and (2.40) at  $r = 0$  is guaranteed by Eqs. (2.35), (2.37), and (2.38).

Going back to Eq. (2.34), we write for the neighborhood of  $r \approx 0$ :

$$-\frac{g(k, r)}{\tilde{f}_-(k, r)} = \frac{(\pi/2)gC_a(g) \ln r + O(1)}{gC_a(g) \ln r \left( \ln k - \ln 2 + \gamma - i \frac{\pi}{2} \right) + [gC_b(g) - 1] \ln r + O(1)} + O(k^2). \quad (2.42)$$

This result leads to

$$e^{i\delta_0(k)} \sin \delta_0(k) = \frac{\pi}{2} \left[ \frac{gC_a(g)}{gC_a(g) \left( \ln k - \ln 2 + \gamma - i \frac{\pi}{2} \right) + [gC_b(g) - 1]} \right] + O(k^2). \quad (2.43)$$

There are now two cases to consider,  $C_a(g) \neq 0$  and  $C_a(g) = 0$ . For  $C_a(g) \neq 0$ , we have the universal result as  $k \rightarrow 0$ ,

$$\delta_0(k) = \frac{\pi}{2 \ln k} + O\left(\frac{1}{(\ln k)^2}\right). \quad (2.44)$$

One should note that  $C_b(g)$  is finite. A somewhat stronger form of (2.44) is that, as  $k \rightarrow 0$ ,

$$e^{i\delta_0(k)} \sin \delta_0(k) = \frac{\pi}{2 \ln k - i\pi} + O\left(\frac{1}{(\ln k)^{2,3}}\right), \quad (2.45)$$

meaning that the real part of the first term is accurate to order  $(\ln k)^{-2}$  while the imaginary part is accurate to  $(\ln k)^{-3}$ .

The second case,  $C_a(g) = 0$ , is clearly exceptional. If  $C_a(g) = 0$  for any interval  $g_1 < g < g_2$ , then  $V \equiv 0$ . For  $V \neq 0$ ,  $C_a(g)$  can only vanish for discrete values of  $g$ . In this case, because of (2.38) and (2.39),  $(1 - gC_b)$  cannot vanish. Hence, it follows from (2.42) that, as  $k \rightarrow 0$ ,

$$\delta_0(k) = O(k^2). \quad (2.46)$$

Equation (2.43) also implies the uniform formula

$$\delta_0(k) = \frac{\xi}{\xi + 1} \frac{\pi}{2 \ln k - i\pi} + O\left(\frac{1}{(\ln k)^{2,3}}\right) \quad (2.47)$$

in the same sense as (2.45), where

$$\xi = \frac{gC_a(g) \left(\ln k - i \frac{\pi}{2}\right)}{gC_b(g) - 1}. \quad (2.48)$$

### 3. THRESHOLD BEHAVIOUR IN (2+1) DIMENSIONS: THE FIELD THEORETICAL CASE

We take as our starting point axiomatic local field theory with a minimum non-zero mass. There is then very little difference between 2 + 1 and 3 + 1 dimensions. In both cases, the on-shell scattering amplitude depends on two variables. The analyticity domain of the scattering amplitude is obtained, in both cases, in two steps: i) analytic continuation of the off-shell amplitude [6],[8] and ii) use of the positivity of the absorptive part to enlarge the analyticity domain [9]. The partial wave expansion in the (2 + 1)-dimensional case is given in terms of Chebyshev polynomials and not Legendre polynomials. Indeed, for the (2 + 1)-dimensional case, we have

$$T(s, \cos \theta) = 16 \sum_{n=0}^{\infty} \epsilon_n f_n(s) \cos n\theta. \quad (3.1)$$

Here,  $s$  is the square of the center-of-mass energy, and  $\theta$  is the scattering angle. In the elastic region,  $f_n(s)$  is related to the phase shifts by

$$f_n(s) = \sqrt{s} e^{i\delta_n} \sin \delta_n. \quad (3.2)$$

This and the factor of 8 in (3.1) are chosen to give  $T(s, \cos \theta) = -g + O(g^2)$  in  $\phi_3^4$  perturbative field theory with a  $(g/4!)\phi^4$  interaction.

The absorptive part of  $T$  is

$$A_s(s, \cos \theta) = 16 \sum_{n=0}^{\infty} \epsilon_n \cdot \text{Im} f_n(s) \cos n\theta, \quad (3.3)$$



with  $\text{Im } f_n(s) \geq 0$ , from the unitarity condition. From Eq. (3.3), it is easy to obtain

$$\left| \left( \frac{d}{d \cos \theta} \right)^n A_s(s, \cos \theta) \right| \leq \left( \frac{d}{d \cos \theta} \right)^n A_s(s, \cos \theta) \Big|_{\cos \theta=1}; \quad s \geq 4m^2, \quad (3.4)$$

for all  $\theta$  such that  $-1 \leq \cos \theta \leq +1$ . This last inequality is precisely what made the enlargement of the analyticity domain in the  $3+1$  case possible [9]. Therefore, one gets the same enlargement in  $2+1$  dimensions.

For simplicity, we consider a case with the kinematics and symmetry of pion-pion scattering although our results are much more general. We use the Mandelstam variables

$$\begin{aligned} s &= 4(k^2 + m^2), \\ t &= 2k^2(\cos \theta - 1), \\ u &= 4m^2 - s - t. \end{aligned} \quad (3.5)$$

For any fixed  $t$ ,  $|t| < 4m^2$ ,  $T(s, t)$  is analytic in the doubly cut  $s$ -plane with cuts along

$$\begin{aligned} s &= 4m^2 + \lambda, \\ u &= 4m^2 + \lambda; \quad \lambda > 0. \end{aligned} \quad (3.6)$$

For fixed  $s$ , the absorptive part,  $A_s(s, \cos \theta)$ , is analytic inside an ellipse in the  $\cos \theta$ -plane, which is an enlargement of the Lehmann ellipse [10]. The foci are at  $\cos \theta = \pm 1$  and the right extremity is at  $\cos \theta = 1 + 4m^2/2k^2$ .

The partial wave amplitudes,  $f_n(s)$ , are defined as

$$f_n(s) = \frac{1}{16\pi} \int_{-1}^{+1} T(s, \cos \theta) \cos n\theta \frac{d(\cos \theta)}{\sin \theta}. \quad (3.7)$$

The  $f_n$ 's are analytic in a region that contains

$$|s - 4m^2| < 4m^2, \quad (3.8)$$

excluding a cut along  $4m^2 \leq s \leq 8m^2$ . A major difference with the  $(3+1)$ -dimensional case is the kinematical factor  $\sqrt{s}$  which comes from unitarity as explicitly shown in Eq. (3.2), a point clarified with the help of R. Stora [11].

Thus, the unitarity condition in  $2+1$  dimensions is

$$\text{Im } f_n(s) \geq \frac{1}{\sqrt{s}} |f_n(s)|^2, \quad \forall s > 4m^2. \quad (3.9)$$

In the elastic region,  $4m^2 \leq s < 16m^2$ ,

$$\text{Im } f_n(s) = \frac{1}{\sqrt{s}} |f_n(s)|^2. \quad (3.10)$$

This slightly changed form of the unitarity condition given in Eq. (3.9), gives a different Froissart bound [9],[12] in the  $(2+1)$  case. The number of partial waves effectively contributing to the scattering amplitude is still bounded by

$$L = C\sqrt{s} \ln s, \quad (3.11)$$

for large  $s$ . However, the Froissart bound in  $2+1$  dimensions is

$$|F(s, \cos \theta)| < Cs \ln s, \quad -1 \leq \cos \theta \leq +1. \quad (3.12)$$

This is instead of the  $s \ln^2 s$  in the  $3+1$  case. The number of subtractions in the dispersion relations, for  $|t| < 4m^2$ , is still at most 2, as in the  $3+1$  case [13].

The general properties outlined so far are sufficient to determine the singularity of  $f_n(s)$  at  $k = 0$ . For simplicity, we restrict ourselves to the  $S$ -wave case, although our method applies to the higher waves. It is convenient to change variables and define

$$f_0(s) = F_0(k). \quad (3.13)$$

We also set the mass  $m = 1$ . In the variable  $k$ , the analyticity domain of  $F_0(k)$  contains the half circle  $\Gamma$ ,

$$\Gamma : \quad \{|k| < 1, \quad \text{and} \quad \text{Im } k > 0\}. \quad (3.14)$$

A very important property of  $T(s, t)$  is the reality property:  $T$  is real for  $s < 4$ ,  $t < 4$ ,  $u < 4$ . From this property, it follows that  $f_0(s)$  is real for  $0 < s < 4$ , and hence  $F_0(k)$  is real for  $k = i\kappa$ ,  $0 < \kappa < 1$ . By Schwarz's reflection principle, for  $k \in \Gamma$ , we have

$$F_0(k) = F_0^*(-k^*). \quad (3.15)$$

The unitarity condition, Eq. (3.10), can be written in a form suitable for analytic continuation. With initially  $k = k^*$ , we write

$$F_0(k) - F_0^*(k^*) = \frac{2i}{\sqrt{s}} F_0(k) F_0^*(k^*). \quad (3.16)$$

This gives

$$F_0(k) = \frac{F_0^*(k^*)}{1 - \frac{2i}{\sqrt{s}} F_0^*(k^*)}, \quad (3.17)$$

and defines a function analytic in the second sheet. This function will be the continuation to the semicircle,  $|k| < 1$ ,  $\text{Im } k < 0$ , through the line  $0 < k < 1$ . The only thing to prevent that would be an accumulation of zeros of  $[1 - (2i/\sqrt{s})F_0^*(k^*)]$  along this line, giving a natural boundary. There is nothing in the general axioms to prevent that [14]. However, it is sufficient to assume that  $F_0(k)$  is continuous on  $0 < k < 1$  in order to avoid this catastrophe. We thus get the continuation of  $F_0(k)$  to the second sheet [15], which, using the reality condition (3.15), can be written as

$$F_0(k) = \frac{F_0(-k)}{1 - \frac{2i}{\sqrt{s}} F_0(-k)}. \quad (3.18)$$

Hence,  $F_0(k)$  is meromorphic for  $|k| < 1$ , outside the origin.

Let us introduce  $G_0(k)$  as

$$G_0(k) = \frac{1}{F_0(k)}. \quad (3.19)$$

We get

$$G_0(k) = G_0(-k) - \frac{2i}{\sqrt{s}}. \quad (3.20)$$

Next, we define  $H_0(k)$  as

$$H_0(k) \equiv G_0(k) - \frac{2}{\pi\sqrt{s}} \left( \ln k - i \frac{\pi}{2} \right). \quad (3.21)$$

$H_0(k)$  is again real for  $k = i\kappa$ ,  $0 < \kappa < 1$ . Using Eq. (3.21), we get

$$H_0(k) = H_0(-k). \quad (3.22)$$

$H_0$  is therefore an even function of  $k$ , i.e.,

$$H_0(k) \equiv K_0(k^2). \quad (3.23)$$

$K_0(k^2)$  is a meromorphic function of  $k^2$ , and the  $S$ -wave amplitude can be written as

$$F_0(k) = \frac{1}{K_0(k^2) + \frac{2}{\pi\sqrt{s}} \left( \ln k - i \frac{\pi}{2} \right)}. \quad (3.24)$$

If  $K_0(k^2)$  has no pole at the origin, the  $\ln k$  dominates the denominator as  $k \rightarrow 0$ , and we get

$$F_0(k) \simeq \frac{\pi}{2} \sqrt{s} \left( \frac{1}{\ln k} \right). \quad (3.25)$$

The phase shift then behaves as

$$\delta_0(k) \simeq \frac{\pi}{2 \ln k}, \quad (3.26)$$

which is precisely the behaviour obtained in the potential case. As in the potential case, the existence of a pole of  $K_0(k^2)$  at  $k^2 = 0$  cannot be excluded.

The derivation we presented above also applies to higher waves, but it can be proved that what is hopefully an exception for  $n = 0$  turns out to be the rule for  $n \geq 1$ .  $K_n(k^2)$  has a pole, and we shall show in a future publication that  $\delta_n \sim k^{2n}$  for  $n \geq 1$ .

For the restricted class of potentials such that

$$\int_0^\infty r dr \left| 1 + |\ln r| \right| |V(r)| \exp \mu r < \infty,$$

the derivation of the dispersion relations for  $|t| < \mu^2$  obtained first by one of us [16] in the (3+1) case also holds in 2 + 1 dimensions. It implies that the partial wave amplitude is analytic in

$|k| < \mu/2$ ,  $\text{Im } k > 0$ , and therefore the derivation presented in this section applies also to this potential case.

Equation (3.17) was also obtained by Bros and Iagolnitzer in ref. [5], in a more general but less elementary approach based on a postulated analyticity of the  $S$ -matrix which, however, becomes axiomatic by using, as we said in the introduction, ref. [6]. These authors emphasize the Riemann sheet structure at the threshold rather than the actual behaviour of the physical scattering amplitude.

## 4. PERTURBATION THEORY FOR $\phi_3^4$

It is of importance to compare our result with perturbation theory. We are fortunate that in  $(2+1)$  dimensions we have a rigorously defined super-renormalizable theory [1] with a mass gap, namely,  $\phi_3^4$ .

Taking

$$\mathcal{L}_{\text{int}}(\phi) = \frac{-g}{4!} : \phi^4(x) :$$

we obtain up to order  $g^2$  for  $T(p_1, p_2; -p_3, -p_4)$

$$T(s, t) = -g + g^2[f(s) + f(t) + f(u)] + O(g^3), \quad (4.1)$$

where  $f(s)$ ,  $s = (p_1 + p_2)^2$ , is given by the Feynman diagram shown in Figure 1,

$$f(s) = \left(\frac{-i}{2}\right) \int \frac{d^3k}{(2\pi)^3} \frac{1}{(k^2 - \mu^2 + i\epsilon)((p_1 + p_2 - k)^2 - \mu^2 + i\epsilon)}. \quad (4.2)$$

The factor  $(\frac{1}{2})$  is for identical outgoing particles, and the  $(-i)$  follows from  $S = 1 + iT$ ,  $S$  being the  $S$ -matrix.

This last integral can be easily evaluated in the Euclidean region,  $s < 4\mu^2$ , by carrying out a Wick rotation, and the result is

$$f(s) = -\frac{1}{16\pi\sqrt{s}} \ln\left(\frac{2\mu - \sqrt{s}}{2\mu + \sqrt{s}}\right), \quad 0 < s < 4\mu^2. \quad (4.3)$$

The normalization of  $T$  is chosen such that elastic unitarity is given by

$$\frac{1}{2i}(T - T^*) = \frac{1}{16\sqrt{s}} \int_0^{2\pi} \frac{d\theta}{2\pi} |T(s, \theta)|^2, \quad 4\mu^2 \leq s < 16\mu^2. \quad (4.4)$$

The partial wave expansion is then

$$T(s, \theta) = 16\sqrt{s} \sum_n \epsilon_n \cos n\theta e^{i\delta_n} \sin \delta_n. \quad (4.5)$$

As  $s \rightarrow 4\mu^2$ ,  $k \rightarrow 0$ , then for physical  $\theta$ ,  $t \rightarrow 0$ ,  $u \rightarrow 0$ , and the leading log term comes from (4.3), since  $f(0)$  is finite.

We get for  $k \rightarrow 0$

$$T = -g - \frac{g^2}{32\pi\mu} \ln \frac{k^2}{\mu^2} + O(1)g^2 + O(g^3). \quad (4.6)$$

The first thing to notice is that at order  $g^2$ ,  $T$  diverges as  $k \rightarrow 0$ . This is just the opposite of the full result we obtained in the previous section where  $T \rightarrow 0$  as  $k \rightarrow 0$ .

In third order, the leading  $\ln k$  behaviour comes from the two-bubble diagram shown in Figure 2. The triangle diagram in Figure 3 is only of order  $(\ln k)$ . We conjecture that this continues in higher orders, and the leading  $(\ln k)$  approximation is given by

$$T \cong -g \sum_{n=0}^{\infty} \left( \frac{g \ln(k/\mu)}{16\pi\mu} \right)^n, \quad k \rightarrow 0. \quad (4.7)$$

This sum is divergent for  $k < \mu \exp(-16\pi\mu/g)$ . Thus the present perturbation calculation does not give a meaningful result. If we ignore this divergence and sum the geometric series formally, the result is

$$T \cong -g \left\{ \frac{1}{1 - \left( \frac{g \ln(k/\mu)}{16\pi\mu} \right)} \right\}. \quad (4.8)$$

However, as  $k \rightarrow 0$ ,  $s \rightarrow 4\mu^2$ ,

$$\frac{T}{16\sqrt{s}} \cong e^{i\delta_0} \sin \delta_0. \quad (4.9)$$

We thus recover the potential scattering result as  $k \rightarrow 0$ ,

$$e^{i\delta_0(k)} \sin \delta_0(k) \sim \frac{-g}{32\mu} \left( \frac{1}{-\frac{g \ln(k/\mu)}{16\pi\mu}} \right) \quad (4.10)$$

and

$$\delta_0(k) \sim \frac{\pi}{2} \frac{1}{\ln(k/\mu)}, \quad k \rightarrow 0. \quad (4.11)$$

$\phi_3^4$  is a well-defined theory, both perturbatively and non-perturbatively, and it is clear from our results that as  $k \rightarrow 0$  perturbation theory gives the wrong answer. It is perhaps interesting to note that  $\phi_3^4$  is asymptotically free. If our conjecture on the higher-order  $(\ln k)$  behaviour is correct, then this would be the first completely rigorous demonstration of how perturbation theory order by order could be extremely misleading. Such a resummation is also present in ref. [5].

## 5. REMARKS AND CONCLUSIONS

We close this paper with three significant remarks.

i) The power of elastic unitarity together with analyticity is clearly demonstrated by the following remark stressed to us by Porrati [17]. Once we are given a phase-shift behaviour such that

$$a_0(k) = e^{i\delta_0(k)} \sin \delta_0(k) = \frac{c}{\ln k - i\frac{\pi}{2}} + O\left(\frac{1}{(\ln k)^{1+\epsilon}}\right), \quad k \rightarrow 0, \quad (5.1)$$

then unitarity alone fixes  $c$  to be  $c = \pi/2$ , since

$$a_0^*(k) = a_0(-k) = \frac{c}{\ln |k| - i\frac{\pi}{2}} + O\left(\frac{1}{(\ln -k)^{1+\epsilon}}\right). \quad (5.2)$$

The factors  $(i\pi/2)$  are necessary to keep  $a_0(k)$  real for  $k$  purely imaginary and  $\text{Im } k > 0$ . Hence we get

$$\text{Im } a_0(k) = \frac{\pi}{2} \frac{c}{(\ln k)^2} + O\left(\frac{1}{(\ln k)^{2+\epsilon}}\right). \quad (5.3)$$

From  $\text{Im } a_0 = |a_0|^2$ , we obtain, when  $c \neq 0$ ,

$$c = \pi/2. \quad (5.4)$$

It should be pointed out, however, that this argument requires analyticity in  $k$  in a semicircle in  $\text{Im } k > 0$ , and hence only applies to exponentially decreasing potentials.

ii) In one dimension, the simplest potential is the  $\delta$ -function potential. In two or three dimensions, the corresponding simplest potential is the so-called point interaction, which is the same as the Fermi pseudopotential. There is a large literature on the Fermi pseudopotential.

Recently, Jackiw [18] obtained the phase shift  $\delta_0(k)$  for the point interaction in two dimensions. Although this potential does not belong to the class considered in Section 2, his result for  $k \rightarrow 0$  agrees with that of ref. [1] and ours; see Eq. (3.26) in his paper. It should be stressed, however, that our relativistic result holds for any  $2+1$  field theory with the standard analyticity and without zero-mass particles; we are not restricted to  $\phi_3^4$ .

iii) In a  $\phi^4$ -type field theory, the renormalized coupling constant is defined by the value of the  $2 \rightarrow 2$  scattering amplitude,  $T(s, t, u)$ , evaluated at some Euclidean point  $(s, t, u) < 4\mu^2$ , often for convenience taken to be the symmetric point  $s = t = u = 4\mu^2/3$ . In  $(3+1)$  dimensions, given the well-established analyticity and unitarity properties of  $T$ , it has been shown in many papers [19] that the coupling constant is bounded. Some of these bounds are surprisingly strong. In  $\phi_3^4$ , Glimm and Jaffe [1] obtained bounds directly from constructive field theory, but their results are weaker than what can be obtained from analyticity and unitarity.

The general methods used in the papers cited in ref. [19] for the  $(3+1)$  case can be easily modified to apply to  $(2+1)$  dimensions. Only the kinematic factor outside the partial wave

expansion is different. The results of this paper thus present us with a new and significant challenge. We have now a new piece of information on the scattering amplitude which is exact. Namely, we know that

$$T(s, t, u) \ln \frac{\sqrt{s - 4\mu^2}}{2\mu} \rightarrow 16\pi\mu \quad \text{as } s \rightarrow 4\mu^2, \quad t \rightarrow 0, \quad u \rightarrow 0,$$

i.e., at certain points on the Mandelstam triangle. Given the power of unitarity and analyticity, we are quite confident that this new input will improve the bounds on the coupling constant. Only the magnitude of the improvement is in question. Work on this problem is in progress.

## ACKNOWLEDGEMENTS

We thank our colleagues H. Grosse, H. Lehmann, M. Porrati, and H. C. Ren for their useful comments, and R. Stora for his crucial help. We also thank J. Bros for pointing out to us his previous work. Two of us, N.N.K. and T.T.W., are grateful to the CERN Theory Division for its kind hospitality. This work was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-91ER40651, Task B, and under Grant No. DE-FG02-84ER40158.

## Appendix A.

In this appendix, we study briefly the equation (2.2), together with the Dirichlet boundary condition (2.3). We start with the free equation,

$$\left(\frac{d^2}{dr^2} + \frac{1}{4r^2} + k^2\right)u(k, r) = 0, \quad (\text{A. 1})$$

with  $u(k, 0) = 0$ . Because of the presence of the attractive singular potential,  $-1/(4r^2)$ , one must be careful in the extension of the differential operator,  $-(d^2/dr^2) - (1/4r^2)$ , to a self-adjoint operator on  $L^2(0, \infty)$ . This has been thoroughly studied in the literature [20],[21]. We quote the result here. The two independent fundamental solutions of (A. 1) are  $\sqrt{r} J_0(kr)$  and  $\sqrt{r} Y_0(kr)$ . Both vanish at the origin. Every other solution, being a linear combination of these two, also vanishes at  $r = 0$ . Therefore, we are in the limit-circle case for the differential operator with a Dirichlet boundary condition at  $r = 0$ . There exist an infinite number of self-adjoint extensions of the symmetric differential operator, depending on one (real) parameter. Each self-adjoint extension is defined by the amount of mixing of the two fundamental solutions. Among all these extensions, there exists a “distinguished” one, which corresponds to taking the pure Bessel solution  $\sqrt{r} J_0(kr)$ . These generalized eigenfunctions are less singular, behaving like  $\sqrt{r}$  at the origin, as compared to the eigenfunctions of all other extensions, which behave like  $\sqrt{r} \ln r$  as  $r \rightarrow 0$ . Moreover, it can be shown that the “distinguished” extension corresponds to the Friedrichs extension [21],[22]. But, for the physicist, the more important fact is this: in all the other self-adjoint extensions, there exists, besides the continuum, a negative energy eigenvalue. In other words, there exists always a real bound state with negative energy,  $E_0 = k_0^2 < 0$  [20],[21].

The extension  $H_\lambda$  is defined by taking the behaviour, as  $r \rightarrow 0$ ,

$$u(r) \rightarrow \sqrt{r} + \lambda\sqrt{r} \ln r; \quad \lambda \text{ real.} \quad (\text{A. 2})$$

It is then easy to check that if we define a solution such that

$$\sqrt{r} [J_0(kr) + Y_0(kr)] \xrightarrow{r \rightarrow 0} \sqrt{r} + \lambda\sqrt{r} \ln r, \quad (\text{A. 3})$$

then an elementary calculation shows that, by setting  $k = +i\kappa_0$ , we get

$$\ln \kappa_0 = \frac{1 - \lambda(\gamma - \ln 2)}{\lambda}, \quad (\text{A. 4})$$

where  $\gamma$  is the Euler constant. Thus, for any real  $\lambda$ ,  $\lambda > 0$ , we have a bound state at  $E = -\kappa_0^2(\lambda)$ .

There is no such bound state in the “distinguished” extension. However, in this case we are just at the threshold of having a bound state. More precisely, in the “distinguished” extension, if we add to the free Hamiltonian a purely attractive (negative) potential, no matter how weak it happens to be, there appears a true bound state. This fact is well established in the literature using a variational argument.



As an aside here we give the upper bound of Setô [23] on the number  $N$  of bound states for dimension  $d = 2$ , and  $l = 0$ . This is the 2-dimensional version of the old Bargmann inequality for  $d = 3$ . The Setô bound is

$$N_2^0 \leq 1 + \frac{\frac{1}{2} \int_0^\infty dr \int_0^\infty dr' \left| \ln \frac{r}{r'} \right| V(r) V(r')}{-\int_0^\infty r V(r) dr}, \quad (\text{A. 5})$$

where, given our assumptions on  $V(r)$ , all the integrals are finite. The fact that there is always a bound state, regardless of how weak an attractive potential  $V$  may be, is somehow reflected by the presence of 1 in the right-hand side of (A. 5). This cannot be improved.

In any case, this last property of the “distinguished” extension of the free differential operator to a self-adjoint operator *without a bound state* is the most important criterion by which we must choose this extension, and discard all others. As physicists, we do not have the freedom to start with a “free Hamiltonian” that binds a free particle. Mathematicians have this luxury.

We finally come to the equation (2.2) itself. Starting from the “distinguished” extension of the free Hamiltonian, and adding to it a potential  $V$ , does not alter the self-adjointness, provided  $V$  is “weak” in the sense of Kato and others [22],[24]. The condition defining this “weak” class is expressed precisely in the following integrability condition on the potential:

$$\int_0^\infty r dr (1 + |\ln r|) |V(r)| < \infty. \quad (\text{A. 6})$$

This ensures the semi-boundedness of the total Hamiltonian, and the finiteness of the number of bound states. Note that (A. 6) is precisely the condition (2.35) which we had to use in section II. We shall need it in Appendix B to establish the existence and study the properties of the solutions of the two integral equations (2.30) and (2.31).

To conclude this appendix, let us point out that an extension different from the “distinguished” one can be used to simulate a renormalized delta-function interaction, as was done by Jackiw [18].

## Appendix B.

In this appendix we study the integral equations (2.30) and (2.31). For the class of potentials satisfying Eqs. (2.35) and (2.36), we first prove that the solutions  $A(r)$  and  $B(r)$  exist and are bounded, as  $r \rightarrow \infty$ , as in Eq. (2.41). Next, we prove that the behaviour of  $A(r)$  and  $B(r)$  as  $r \rightarrow 0$  is given by Eqs. (2.37) and (2.38), respectively. We will only give the details for Eq. (2.31). The procedure for Eq. (2.30) is easier and very similar.

Our starting point is the integral equation

$$B(r) = \ln r + g \int_r^\infty r' dr' \left( \ln \frac{r'}{r} \right) V(r') B(r'). \quad (\text{B. 1})$$

We can first consider the case  $r' \geq r \geq 1$ , where we have the inequality

$$0 \leq \ln \frac{r'}{r} \leq \ln r'. \quad (\text{B. 2})$$

Therefore, an upper bound  $\bar{B}$  is obtained for  $B$  by replacing the integral equation (B. 1) by

$$\bar{B}(r) = \ln r + g \int_r^\infty r' dr' |V(r')| \ln r' \bar{B}(r'), \quad r > 1. \quad (\text{B. 3})$$

The solution of (B. 3) can be obtained by standard methods and is given by

$$\bar{B}(r) = \left[ \int_1^r \frac{dt}{t} \exp\left(-g \int_t^\infty u |V(u)| \ln u du\right) + C \right] \exp\left(g \int_r^\infty t |V(t)| \ln t dt\right). \quad (\text{B. 4})$$

The constant  $C$  is given by

$$C = \int_1^\infty \frac{1}{r} \left[ 1 - \exp\left(-g \int_r^\infty t |V(t)| \ln t dt\right) \right] dr, \quad (\text{B. 5})$$

which is finite given (2.35). Using this result in (B. 4), we find that

$$\bar{B}(r) = [1 + o(1)] \ln r, \quad \text{as } r \rightarrow \infty. \quad (\text{B. 6})$$

This establishes the bound on  $B(r)$  for  $r \geq 1$ ,

$$|B(r)| \leq C_1 \ln r + D_1, \quad (\text{B. 7})$$

where  $C_1$  and  $D_1$  are positive constants depending on  $g$ .

By the same technique, we arrive at similar conclusions for  $A(r)$ . This time, the bounding condition for  $\bar{A}(r)$  is  $\bar{A}(\infty) = 1$ . We obtain

$$\bar{A}(r) = 1 + o(1), \quad \text{as } r \rightarrow \infty \quad (\text{B. 8})$$

and

$$|A(r)| \leq \bar{A}(r) \leq D_2, \quad r \geq 1, \quad (\text{B. 9})$$

where  $D_2$  is a positive constant.

From these bounds one can easily get, as  $r \rightarrow \infty$ ,

$$A(r) = 1 + o(1); \quad B(r) = [1 + o(1)] \ln r. \quad (\text{B. 10})$$

It is important to note that for the first estimate we need only the condition (2.35), whereas for the second we need (2.36).

Finally, we consider the region  $r < 1$  for both  $A(r)$  and  $B(r)$ . The case for  $B(r)$  is more delicate (singular), and we treat it first.

We can write (B. 1) as

$$B(r) = \ln r + g \int_r^1 r' \left( \ln \frac{r'}{r} \right) V(r') B(r') dr' + g \int_1^\infty r' \left( \ln \frac{r'}{r} \right) V(r') B(r') dr'. \quad (\text{B. 11})$$

In the second integral, since  $r' \geq 1$  and  $r < 1$ , we can use the bound (B. 7) and get, using condition (2.29),

$$\left| \int_1^\infty r \ln \frac{r'}{r} V(r') B(r') dr' \right| < C + D \left( \ln \frac{1}{r} \right), \quad (\text{B. 12})$$

where  $C$  and  $D$  are positive constants. In the first integral, we have

$$\left| \ln \frac{r'}{r} \right| \leq |\ln r|, \quad r < r' \leq 1. \quad (\text{B. 13})$$

An upper bound,  $\bar{\bar{B}}(r)$ , for  $B(r)$  in  $r \leq 1$  is now obtained by substituting (B. 12) and (B. 13) in (B. 11). We obtain the integral equation

$$\bar{\bar{B}}(r) = C_2 + D_2 |\ln r| + g |\ln r| \left[ \int_r^1 r' |V(r')| \bar{\bar{B}}(r') dr' \right], \quad (\text{B. 14})$$

with some positive constants  $C_2$  and  $D_2$ .

The solution of (B. 14) can be obtained by elementary methods. It is

$$\begin{aligned} \bar{\bar{B}}(r) = & Z(r) g |\ln r| \left[ C_3 + \int_r^1 r' |V(r')| [C_2 + D_2 |\ln r'|] (Z^{-1}(r')) dr' \right. \\ & \left. + C_2 + D_2 |\ln r| \right], \end{aligned} \quad (\text{B. 15})$$

where

$$Z(r) = \exp \left[ \int_r^1 dr' g r' |\ln r'| |V(r')| \right]. \quad (\text{B. 16})$$

Noting that  $Z(r)$  is bounded for  $0 \leq r \leq 1$ , from the condition (2.35), we get

$$|B(r)| \leq \bar{\bar{B}}(r) < \lambda + \mu |\ln r|. \quad (\text{B. 17})$$

In the same way, we can analyze the integral equation (2.30) for  $A(r)$ . We again find that, for  $r \rightarrow 0$ ,

$$|A(r)| \leq \lambda_1 |\ln r| + \mu_1. \quad (\text{B. 18})$$

Using these two bounds, we can now prove the asymptotic estimates (2.37) and (2.38). From Eq. (2.30), we get, as  $r \rightarrow 0$ ,

$$A(r) = -g C_a \ln r + g \int_r^\infty r' \ln r' V(r') A(r') dr' + 1. \quad (\text{B. 19})$$

This can be written as

$$A(r) = -g C_a \ln r + g \int_r^1 r' \ln r' V(r') A(r') dr' + O(1). \quad (\text{B. 20})$$

The integral in (B. 20) could diverge as  $r \rightarrow 0$ . However, setting

$$I(r) = g \int_r^1 r' \ln r' V(r') A(r') dr', \quad (\text{B. 21})$$

and using (B. 18), we get

$$\begin{aligned} |I(r)| &< g\lambda_1 \int_r^1 r' |\ln r'|^2 |V(r')| dr' + g\mu_1 \int_r^1 r' |\ln r'| |V(r')| dr' \\ &< g\lambda_1 \int_r^1 r' |\ln r'|^2 |V(r')| dr' + O(1). \end{aligned} \quad (\text{B. 22})$$

Next we define

$$F(r) \equiv r^2 |\ln r|^2 |V(r)|. \quad (\text{B. 23})$$

From the condition (2.35), we have

$$\int_0^1 dr' r' |\ln r'| V(r) = \int_0^1 \frac{dr'}{r' |\ln r'|} \cdot F(r') < \text{const.} \quad (\text{B. 24})$$

This implies that  $F(r) \rightarrow 0$  as  $r \rightarrow 0$ . From Eqs. (B. 22) and (B. 23), we get

$$|I(r)| \leq g\lambda_1 \int_r^1 \frac{dr'}{r'} F(r') + O(1), \quad (\text{B. 25})$$

and, hence, since  $F(r')$  vanishes as  $r' \rightarrow 0$ ,

$$|I(r)| = |\ln r| o(1). \quad (\text{B. 26})$$

This establishes Eq. (2.37). For Eq. (2.38), the derivation is similar.

It is important to notice that, if  $A(r)/\ln r \rightarrow 0$  as  $r \rightarrow 0$ , then  $B(r)/\ln r$  cannot approach zero as  $r \rightarrow 0$ . This is because  $A$  and  $B$  are solutions of the *same* differential equation,

$$\frac{d}{dr} \left( r \frac{dX}{dr} \right) = -grV(r)X(r),$$

and are thus linearly independent.

## References

- [1] J. Glimm and A. Jaffe, *Quantum Physics—A Functional Integral Point of View*, 2nd ed. (Springer-Verlag, New York, 1987).
- [2] R.W.P. King, *Theory of Linear Antennas* (Harvard University Press, Cambridge, MA, 1956).
- [3] D. Bollé and F. Geztesy, *Phys. Rev. Lett.* **52**, 1469 (1984).
- [4] P.G. Averbuch, *J. Phys.* **A19**, 2325 (1986).
- [5] J. Bros and D. Iagolnitzer, *Comm. Math. Phys.* **85**, 197 (1982);  
J. Bros and D. Iagolnitzer, *Phys. Rev.* **D27**, 811 (1983);  
See also Saclay Preprint (1988).

- [6] J. Bros, H. Epstein, and V. Glaser, *Nuovo Cimento* **31**, 1265 (1964).
- [7] A. Pais and T.T. Wu, *Phys. Rev.* **134**, B1303 (1964).
- [8] J. Bros, H. Epstein, and V. Glaser, *Comm. Math. Phys.* **1**, 240 (1965);  
N.N. Bogoliubov, B.V. Medvedev, and M. K. Polivanov, *Voprosy teorii dispersionnykh sootnoshenii* (Gos. izd-vo Fiziko-matematicheskoi lit-ry, Moscow, 1958);  
K. Symanzik, *Phys. Rev.* **105**, 743 (1957).
- [9] A. Martin, *Nuovo Cimento* **42**, 930 (1965);  
A. Martin, *Nuovo Cimento* **44**, 1219 (1966).
- [10] H. Lehmann, *Nuovo Cimento* **10**, 579 (1958).
- [11] R. Stora, private communication.
- [12] M. Froissart, *Phys. Rev.* **123**, 1053 (1961).
- [13] Y.S. Jin and A. Martin, *Phys. Rev.* **B135**, 1375 (1964).
- [14] A. Martin, in *Problems in Theoretical Physics*, dedicated to N.N. Bogoliubov,  
D.I. Blokintseff et al., eds. (NAUKA, Moscow, 1969), p. 113;  
A.S. Wightman, in *Proceedings of the 14th International Conference on High Energy Physics, Vienna, 1968*, J. Prentki and J. Steinberger, eds. (CERN, Geneva, 1968), p. 434.
- [15] W. Zimmermann, *Nuovo Cimento* **21**, 249 (1961).
- [16] N.N. Khuri, *Phys. Rev.* **107**, 1148 (1957).
- [17] M. Porrati, private communication.
- [18] R. Jackiw, in *M.A.B. Bég Memorial Volume*, A. Ali and P. Hoodbhoy, eds. (World Scientific, Singapore, 1991), pp. 35–53.
- [19] A. Martin, in *High Energy and Elementary Particles, ICTP, Trieste* (I.A.E.A., Vienna, 1965), p. 155;  
L. Lukaszuk and A. Martin, *Nuovo Cimento* **A47**, 265 (1967);  
J.B. Healy, *Phys. Rev.* **D8**, 1904 (1973);  
G. Auberson, L. Epele, G. Mahoux, and F. R.A. Simão, *Nucl. Phys.* **B94**, 344 (1975);  
B. Bonnier, C. Lopez, and G. Mennessier, *Phys. Lett.* **60B**, 63 (1975);  
C. Lopez and G. Mennessier, *Nucl. Phys.* **B118**, 426 (1977).
- [20] E.C. Titchmarsh, *Eigenfunction Expansions, I*, 2nd ed. (Oxford University Press, 1962).
- [21] H. Narnhofer, *Acta Physica Austriaca* **40**, 306–322 (1974), and references quoted there.
- [22] G. Hellwig, *Differential Operators of Mathematical Physics* (Addison-Wesley, Reading, MA, 1967).

- [23] N. Setô, *Publ. of RIMS* **9**, 429–461 (1974). See also,  
R.G. Newton, *Scattering Theory of Waves and Particles*, 2nd ed. (Springer-Verlag, New York, 1982).
- [24] T. Kato, *Perturbation Theory of Linear Operators* (Springer-Verlag, Heidelberg, 1976).




FIG. 1. Second-order diagram.




FIG. 2. Third-order diagram behaving as  $(\ln k)^2$  as  $k \rightarrow 0$ .




FIG. 3. Third-order diagram behaving as  $\ln k$  as  $k \rightarrow 0$ .