

Variational calculations with improved energy functionals in gauge theories

C. Heinemann¹, E. Iancu², C. Martin³ and D. Vautherin¹

¹*LPNHE, LPTPE, Universités Paris VI/VII, case 127, F-75252, Paris Cedex 05, France*

²*Theory Division, CERN, CH-1211 Geneva 23, Switzerland*

³*Institut de Physique Nucléaire, F-91406, Orsay Cedex, France*

(November 30, 1999)

Abstract

For a $SU(N)$ Yang-Mills theory, we present variational calculations using gaussian wave functionals combined with an approximate projection on gauge invariant states. The projection amounts to correcting the energy of the gaussian states by subtracting the spurious energy associated with gauge rotations. Based on this improved energy functional, we perform variational calculations of the interaction energy in the presence of external electric and magnetic fields. We verify that the ultraviolet behaviour of our approximation scheme is consistent, as it should, with that expected from perturbation theory. In particular, we recover in this variational framework the standard one-loop beta function, with a transparent interpretation of the screening and anti-screening contributions.

PACS numbers: 11.15.Tk, 12.38.Lg, 14.70.Pj - LPNHE 99-07, CERN 99-360

Typeset using REVTeX

]

I. INTRODUCTION

The functional Schrödinger picture has proven to be a privileged tool in exploring a rich variety of aspects of gauge theories which are beyond the scope of perturbation theory [1]. It is a useful starting point for developing non perturbative calculations based on the variational approach. In the case of scalar field theories, static as well as dynamical variational calculations have been performed by using trial wave functionals of the gaussian type [2–8]. In the case of gauge theories, some early investigations along these lines can be found in [9–12]. However the application of variational methods to gauge theories is generally plagued by the difficulty to implement in a calculable way the requirement of gauge invariance of physical states [12–14].

Gaussian wave functionals allow for analytic calculations, but are not gauge invariant except in the Abelian case. In principle, one can construct gauge invariant states by averaging gaussian wave functionals over all gauge rotations. This results in an effective non-linear sigma model where the fields are the group elements of the gauge transformations [14]. However, to make progress with this theory, further approximations are necessary both in the choice of the kernel of the gaussian, and in the evaluation of the functional integrals over the gauge group [14–18]. Such approximations, which go beyond the variational principle, are not always under control. In particular, in the perturbative regime, they fail to completely reproduce the one-loop beta function [15] (see also [16,18]).

In this paper we shall propose a different strategy which is inspired by techniques developed in 1962 by Thouless and Valatin [19] to deal with the restoration of rotational invariance when deformed solutions are obtained in nuclear Hartree-Fock calculations. Rather than using gauge invariant variational states, we shall limit ourselves to gaussian wave functionals, but we shall correct the associated energy functional by a non-local term, which approximately corresponds to the energy gain when projecting on gauge invariant states. In the

Abelian case, this amounts to removing the contribution of the longitudinal part of the gaussian kernel to the energy. In the Yang-Mills theory, the corrective energy term is itself determined by the variational principle, and the ensuing variational calculation is a priori non-perturbative.

Our main purpose here is to provide the framework for such a calculation and verify that, in the perturbative regime (i.e., when supplemented with an expansion in powers of g), it leads to results consistent with ordinary perturbation theory. As a first check, we shall show that our variational calculation reproduces the standard one-loop beta function of Yang-Mills theory, with a transparent interpretation of the various contributions in terms of screening and anti-screening phenomena. As a further test, we shall compute the vacuum energy in the presence of an external magnetic field B , and find that it exhibits a minimum at a non vanishing value of B , in agreement with the perturbative calculation in [20].

The paper is organized as follows. In Sec. II we briefly review the functional Schrödinger picture and the variational principle in field theory. In Sec. III, we present the Thouless-Valatin formalism and consider, as a simple illustration, its application to quantum electrodynamics (QED). In Sec. IV we present a variational calculation of the one-loop beta function which is based on the construction of the interaction energy between external electrostatic charges. In Sec. V we consider the energy of the QCD vacuum in the presence of a constant magnetic field. This provides an alternative computation of the beta function, and also of the gluon condensate which is found to satisfy the trace anomaly relation. A summary of our results and a discussion of further possible extensions and applications is presented in Sec. VI.

II. VARIATIONAL CALCULATIONS FOR GAUGE FIELDS

We consider the functional Schrödinger description of the $SU(N)$ Yang-Mills theory. In the temporal gauge $A_a^0 = 0$, the canonical coordinates are the vector potentials $A_a^i(\mathbf{x})$ and the electric fields $E_a^i(\mathbf{x})$, which we shall often write as color matrices in the adjoint

representation: e.g., $A^i \equiv A_b^i T^b$ (the color indices a, b, \dots run from 1 to $N^2 - 1$). The generators T^a of the color group are taken to be Hermitian and traceless; they satisfy

$$[T^a, T^b] = if^{abc}T^c, \quad \text{Tr}(T^a T^b) = N\delta^{ab}, \quad (T^a)_{bc} = -if^{abc}.$$

The Hamiltonian density reads (g denotes the coupling constant) :

$$\mathcal{H}(\mathbf{x}) = \frac{1}{2} \left\{ g^2 E_a^i E_a^i(\mathbf{x}) + \frac{1}{g^2} B_a^i B_a^i(\mathbf{x}) \right\}, \quad (\text{II.1})$$

with the color magnetic field:

$$\mathbf{B}_a = \nabla \times \mathbf{A}_a + \frac{1}{2} f_{abc} \mathbf{A}_b \times \mathbf{A}_c. \quad (\text{II.2})$$

Note that our conventions are such that the QCD coupling constant is absorbed in the normalization of the vector potentials. With these conventions, the covariant derivative reads $D_i = \partial_i - iA_i$, and the electric fields E_a^i are canonically conjugate to the vector potentials A_a^i : $[E_a^i(\mathbf{x}), A_b^j(\mathbf{y})] = i\delta^{ij}\delta_{ab}\delta^{(3)}(\mathbf{x} - \mathbf{y})$.

In the Schrödinger representation, the states are represented by functionals of $A_a^i(\mathbf{x})$, $\Psi[\mathbf{A}]$, and the electric field is acting on such states by functional differentiation:

$$E_a^i(\mathbf{x})\Psi[\mathbf{A}] = i \frac{\delta}{\delta A_a^i(\mathbf{x})} \Psi[\mathbf{A}]. \quad (\text{II.3})$$

The Hamiltonian H commutes with the generator \mathcal{G} of time-independent gauge transformations,

$$\mathcal{G}(\mathbf{x}) \equiv \nabla \cdot \mathbf{E}(\mathbf{x}) + i[A^i, E^i], \quad (\text{II.4})$$

so it is possible to diagonalize H and \mathcal{G} simultaneously.

The physical states are constrained by Gauss' law:

$$\mathcal{G}(\mathbf{x})\Psi[\mathbf{A}] = 0, \quad (\text{II.5})$$

which is the requirement of gauge invariance. [More precisely, eq. (II.5) shows that physical states must be invariant under “small” (i.e., topologically trivial [1]) gauge transformations.

We shall not be concerned with the topological aspects of the gauge symmetry in what follows.] More generally, in the presence of matter fields represented by an external color source with density ρ^a , Gauss' law gets modified as follows:

$$\mathcal{G}_a(\mathbf{x})\Psi[\mathbf{A}] = \rho_a(\mathbf{x})\Psi[\mathbf{A}]. \quad (\text{II.6})$$

The ground state of QCD is the eigenstate Ψ_{vac} of H of minimal energy which satisfies Gauss' constraint (II.5). It can be constructed, at least in principle, by using the Ritz variational principle, which states that:

$$\langle H \rangle \equiv \frac{\langle \Psi | H | \Psi \rangle}{\langle \Psi | \Psi \rangle} \geq E_{vac}, \quad (\text{II.7})$$

with the minimum achieved for $\Psi = \Psi_{vac}$. Here, $\Psi[\mathbf{A}]$ is any wave functional from the physical Hilbert space (i.e., which satisfies eq. (II.5)), and E_{vac} is the energy of the ground state Ψ_{vac} , assumed to be non-degenerate. In practice, however, one has to restrict oneself to *gaussian* wave functionals, the only ones which allow an analytical computation of $\langle H \rangle$.

These have the form

$$\begin{aligned} \Psi_0[\mathbf{A}] = & \mathcal{N}^{-1} \exp \left\{ -\frac{1}{4g^2} \int d^3x d^3y \left[A_a^i(\mathbf{x}) - \bar{A}_a^i(\mathbf{x}) \right] \right. \\ & \left. \times (G^{-1})_{ij}^{ab}(\mathbf{x}, \mathbf{y}) \left[A_b^j(\mathbf{y}) - \bar{A}_b^j(\mathbf{y}) \right] \right\}, \end{aligned} \quad (\text{II.8})$$

where the background field $\bar{A}_a^i(\mathbf{x})$ and the kernel G^{-1} (with matrix elements $(G^{-1})_{ij}^{ab}(\mathbf{x}, \mathbf{y})$) are the variational parameters. We expect that $\bar{A} = 0$ in the vacuum state, and this is the case that we shall consider mostly in this paper. Still, non-vanishing values of \bar{A} will be also considered, in Sec. V below, in a study of the vacuum stability in the presence of a background color magnetic field (in the same spirit as, e.g., in Ref. [20]).

The expectation value of the Hamiltonian density in the gaussian state Ψ_0 is [12]

$$\begin{aligned}
\langle \Psi_0 | \mathcal{H}(\mathbf{x}) | \Psi_0 \rangle &= \frac{1}{2g^2} \bar{\mathbf{B}} \cdot \bar{\mathbf{B}}(\mathbf{x}) + \frac{1}{8} \text{Tr} \langle \mathbf{x} | G^{-1} | \mathbf{x} \rangle \\
&+ \frac{1}{2} \text{Tr} \langle \mathbf{x} | K G | \mathbf{x} \rangle \\
&+ \frac{g^2}{8} (\text{Tr} \{ S_i T^a \langle \mathbf{x} | G | \mathbf{x} \rangle \})^2 \\
&+ \frac{g^2}{4} \text{Tr} \{ S^i T^a \langle \mathbf{x} | G | \mathbf{x} \rangle S^i T^a \langle \mathbf{x} | G | \mathbf{x} \rangle \}.
\end{aligned} \tag{II.9}$$

In this equation $\bar{\mathbf{B}}$ is the magnetic field associated to the center $\bar{\mathbf{A}}$ and S^i is the spin one matrix whose elements (j, k) are given by $i\varepsilon_{ijk}$. The notation Tr in equation (II.9) implies a summation over the discrete (color and spatial) indices. For instance, $\text{Tr} \langle \mathbf{x} | G^{-1} | \mathbf{x} \rangle = \sum_{i,a} (G^{-1})_{ii}^{aa}(\mathbf{x}, \mathbf{x})$. Finally, the operator K is the second derivative of the classical energy with respect to the center \bar{A}_i^a . It reads, in matrix notations,

$$K = (-i\mathbf{S} \cdot \mathcal{D})^2 - \mathbf{S} \cdot \bar{\mathbf{B}}, \tag{II.10}$$

where

$$\mathcal{D}^i \equiv \partial^i - i\bar{A}^i \tag{II.11}$$

denotes the covariant derivative defined by the background field $\bar{A}^i \equiv \bar{A}_a^i T^a$. In particular,

$$K_{ij}(p) = p^2 \delta_{ij} - p_i p_j \quad \text{for } \bar{A} = 0. \tag{II.12}$$

In the case of non-Abelian gauge theories, however, gaussian functionals like eq. (II.8) suffer from a major drawback: they do not satisfy the requirement of gauge invariance (II.5). It is in principle possible to construct gauge invariant states by *projection*, i.e. by averaging a gaussian functional over all its gauge transformations. This is achieved by means of the formula

$$\Psi[\mathbf{A}] = \frac{1}{\mathcal{N}} \int \mathcal{D}[U(\mathbf{x})] \Psi_U[\mathbf{A}], \tag{II.13}$$

where the functional integration is performed over the unitary $N \times N$ matrix field $U(\mathbf{x})$, with the adequate group invariant measure, and \mathcal{N} is a normalization factor. The integrand in eq. (II.13) is the gauge-transform of the gaussian state $\Psi_0[\mathbf{A}]$:

$$\Psi_U[\mathbf{A}] = \Psi_0[U\mathbf{A}U^+ + iU\nabla U^+]. \quad (\text{II.14})$$

The expectation value E_P of the energy in the projected state is given by the following formula

$$E_P = \frac{\int \mathcal{D}[U(\mathbf{x})] \langle \Psi_0 | H | \Psi_U \rangle}{\int \mathcal{D}[U(\mathbf{x})] \langle \Psi_0 | \Psi_U \rangle}, \quad (\text{II.15})$$

which should replace eq. (II.9) in practical calculations. Unfortunately, eq. (II.15) cannot be evaluated in closed form because the functional integral over the group is not gaussian. Various approximations to eq. (II.15) have been considered in Ref. [14–18]. In what follows, we shall propose a different approximation method which is inspired from techniques used in nuclear physics to calculate the zero point rotational energy of deformed nuclei [19,21–23].

III. APPROXIMATE PROJECTION

The fact that our variational ground state, namely the gaussian Ψ_0 in eq. (II.8), is not gauge invariant introduces a spurious degeneracy in the problem: Ψ_0 is degenerate with all its gauge-transforms defined in eq. (II.14). This leads to the existence of spurious excitations of zero energy which corresponds to rotations of Ψ_0 in the gauge space; these are, of course, the Goldstone bosons associated to the spontaneous breaking of the gauge symmetry by Ψ_0 . Accordingly, the expectation value (II.9) of the Hamiltonian in the *deformed* state Ψ_0 includes unphysical contributions expressing the kinetic energy of the gauge rotations. The Thouless-Valatin formalism [19] provides us with a method to estimate, and thus subtract away, such unphysical contributions.

A. The case of rotations

This formalism is best explained on the example of the collective rotations of a deformed nucleus. There, the equivalent of our present variational calculation with gaussian wave functionals is the so-called *Hartree-Fock approximation* where the nuclear wave function is represented by a Slater determinant formed with A single-particle wave functions $\varphi_k(x)$ (for A nucleons) [23]. The latter are determined by solving the Hartree-Fock equations, i.e., the variational equations obtained by minimizing the expectation value of the Hamiltonian in the subspace of Slater determinants. Although the Hamiltonian H is rotationally invariant, the Hartree-Fock equations may lead to non-invariant solutions describing nuclear deformations. If this is the case, then the spectrum develops a ground state rotational band,

$$E_J = E_0 + \frac{J(J+1)}{2I}, \quad (\text{III.1})$$

where $J(J+1)$ is the eigenvalue of the angular momentum operator \mathbf{J}^2 and I is the moment of inertia. One may then conclude that the operator:

$$\tilde{H} \equiv H - \frac{\mathbf{J}^2}{2I} \quad (\text{III.2})$$

is the Hamiltonian of the *intrinsic* (that is, non-rotational) motion: to the ground state rotational band of H corresponds now a single eigenvalue E_0 of \tilde{H} . Then, eq. (III.2) provides an approximate separation of the dynamics into intrinsic and rotational motion, which is reminiscent of the familiar separation of the center-of-mass motion by the formula:

$$\tilde{H} \equiv H - \frac{\mathbf{P}^2}{2M}, \quad (\text{III.3})$$

where

$$\mathbf{P} = \mathbf{p}_1 + \mathbf{p}_2 + \dots + \mathbf{p}_A, \quad (\text{III.4})$$

is the total momentum operator, and $M = Am$ is the nuclear mass. Thus, a mean field description of the intrinsic motion can be given by performing variational (or Hartree-Fock)

calculations for the subtracted Hamiltonian \tilde{H} . The only question is, what is the value of the moment-of-inertia parameter I ?

One can answer this question by studying rotations of the deformed mass distribution. Assume that the nucleus has axial symmetry with respect to the z -axis, and consider an uniform rotation with angular velocity ω about the x -axis. If $\Psi_\omega(t)$ is the *exact* (time-dependent) state describing such a rotation, then

$$\Psi_\omega(t) = e^{-i\omega t J_x} e^{-iE_\omega t} \Psi_\omega(0), \quad (\text{III.5})$$

which satisfies the time-dependent Schrödinger equation,

$$i \frac{\partial \Psi_\omega}{\partial t} = H \Psi_\omega, \quad (\text{III.6})$$

provided $\Psi_\omega(0)$ is a solution to the following time-independent problem:

$$(H - \omega J_x) \Psi_\omega(0) = E_\omega \Psi_\omega(0). \quad (\text{III.7})$$

This is, of course, just the familiar transformation to the rotating frame of reference, which leads us to consider the variational problem for the following Hamiltonian:

$$H_\omega = H - \omega J_x. \quad (\text{III.8})$$

This is equivalent to a *constrained* variational problem with the subsidiary condition that $\langle J_x \rangle$ has a given value; ω plays then the role of the Lagrange multiplier. The constrained Hartree-Fock calculation determines the optimal independent nucleon wave function $\Psi_\omega \equiv \Psi_\omega(0)$ in the rotating frame. (As $\omega \rightarrow 0$, $\Psi_\omega \rightarrow \Psi_0$ which is the deformed Hartree-Fock ground state.) Once the optimal wave function is found it is possible to obtain an estimate of the moment of inertia I of the nucleus by considering the limit

$$I = \lim_{\omega \rightarrow 0} \frac{\langle \Psi_\omega | J_x | \Psi_\omega \rangle}{\omega}. \quad (\text{III.9})$$

This finally allows us to estimate the zero-point rotational energy ΔE_{TV} in the deformed state Ψ_0 :

$$\Delta E_{TV} = \frac{\langle \Psi_0 | \mathbf{J}^2 | \Psi_0 \rangle}{2I} \quad (\text{III.10})$$

Eq. (III.10) is the expected gain in energy when projecting the deformed Hartree-Fock ground state Ψ_0 onto a rotationally invariant state. That is, the corrected average energy after projection, which is the energy of the intrinsic motion (cf. eq. (III.2)), reads

$$\tilde{E} = E_0 - \Delta E_{TV}, \quad (\text{III.11})$$

where $E_0 \equiv \langle \Psi_0 | H | \Psi_0 \rangle$.

The Thouless-Valatin method is an approximation which is expected to be valid for large deformations, or more precisely when the deformation produces a large expectation value of the square of the angular momentum in the Hartree-Fock ground state [21]. Indeed in this case it can be shown that the projection onto invariant states can be accurately performed because the overlap between two states differing in their orientation by an angle θ is sharply peaked near $\theta=0$ thus allowing for an expansion in the vicinity of this point [22].

B. Application to QED

When going to gauge theories in the variational method, the Hartree-Fock ground state Ψ_0 is replaced by the gaussian trial wave functional (cf. eq. (II.8)), and the angular momentum operator is replaced by the generator of the gauge transformations, $\mathcal{G}^a(\mathbf{x})$ (cf. eq. (II.4)).

As a first illustration let us consider a variational calculation in QED, with the gaussian variational Ansatz

$$\Psi_0[\mathbf{A}] = \mathcal{N}^{-1} \exp\left\{-\langle A | \frac{1}{4G} | A \rangle\right\}, \quad (\text{III.12})$$

where $\mathcal{N} = (\det G)^{1/4}$ and the expression in the exponent is a condensed notation for the convolution

$$\langle A | \frac{1}{4G} | A \rangle \equiv \frac{1}{4} \int d^3x d^3y A^i(\mathbf{x}) G_{ij}^{-1}(\mathbf{x}, \mathbf{y}) A^j(\mathbf{y}). \quad (\text{III.13})$$

This wave functional is gauge invariant provided its kernel G^{-1} is transverse: $\partial_i G_{ij}^{-1} = 0$. Let us assume, however, that this is not the case, and see what the Thouless-Valatin correction

would predict in this case. The operator playing the role of the angular momentum is the charge density operator

$$\mathcal{G}(\mathbf{x}) = \nabla \cdot \mathbf{E}(\mathbf{x}), \quad (\text{III.14})$$

and the generalization of the Thouless-Valatin formula (III.10) for the energy correction reads

$$\Delta E_{TV} = \int d^3x d^3y \langle \Psi_0 | \mathcal{G}(\mathbf{x}) \mathcal{G}(\mathbf{y}) | \Psi_0 \rangle \langle \mathbf{x} | \frac{1}{2\mathcal{I}} | \mathbf{y} \rangle, \quad (\text{III.15})$$

where the "moment of inertia" \mathcal{I} is now a matrix in coordinate space, with matrix elements $\langle \mathbf{x} | \mathcal{I} | \mathbf{y} \rangle \equiv \mathcal{I}(\mathbf{x}, \mathbf{y})$. This is obtained via a constrained variational calculation with Hamiltonian $H_\omega = H - H_{ext}$, where

$$H = \frac{1}{2} \int d^3x \{ \mathbf{E}^2(\mathbf{x}) + (\nabla \times \mathbf{A})^2 \}, \quad (\text{III.16})$$

and the external constraint

$$H_{ext} = \int d^3x \omega(\mathbf{x}) \nabla \cdot \mathbf{E}(\mathbf{x}). \quad (\text{III.17})$$

In the present context, the Lagrange multiplier $\omega(\mathbf{x})$ plays the role of the temporal component $A^0(\mathbf{x})$ of the gauge vector potential. The solution to this constrained variational problem is of the form¹

$$\Psi_\omega[\mathbf{A}] = \exp\{-i \langle \mathbf{F} | \mathbf{A} \rangle\} \Psi_0[\mathbf{A}], \quad (\text{III.18})$$

where the vector field $\mathbf{F}(\mathbf{x})$ is a new variational parameter, which expresses the expectation value of the electric field in the state (III.18), $F^i = \langle \Psi_\omega | E^i | \Psi_\omega \rangle$, and is determined by minimizing

¹Eq. (III.18) can be simply understood by recalling that, in the presence of a constraint of the form $H_{ext} = \alpha p$, the ground state wavefunction of an harmonic oscillator is modified by a factor $\exp(ip_0x)$, but its width remains unchanged.

$$\begin{aligned}
E_\omega &\equiv \langle \Psi_\omega | H - H_{ext} | \Psi_\omega \rangle \\
&= E_0 + \frac{1}{2} \int d^3x (\mathbf{F} + \nabla\omega)^2 - \frac{1}{2} \int d^3x (\nabla\omega)^2.
\end{aligned} \tag{III.19}$$

We have denoted here (V is the total volume of the space)

$$\begin{aligned}
E_0 &\equiv \langle \Psi_0 | H | \Psi_0 \rangle \\
&= \frac{V}{2} \int \frac{d^3p}{(2\pi)^3} \left\{ \frac{1}{4} G_{ii}^{-1}(\mathbf{p}) + (p^2\delta_{ij} - p_i p_j) G_{ij}(\mathbf{p}) \right\}.
\end{aligned} \tag{III.20}$$

Note that the magnetic piece of the energy (III.20) (the second term between parentheses) involves only the transverse components of G_{ij} , while the electric piece involves also its longitudinal component.

The functional $E_\omega[\mathbf{F}]$ in eq. (III.19) attains its minimum for $\mathbf{F} = -\nabla\omega$, in which case

$$\langle \Psi_\omega | \nabla \cdot \mathbf{E} | \Psi_\omega \rangle = \nabla \cdot \mathbf{F} = -\Delta\omega. \tag{III.21}$$

According to (III.9), the moment of inertia is obtained as (with $\langle \nabla \cdot \mathbf{E} \rangle_\omega \equiv \langle \Psi_\omega | \nabla \cdot \mathbf{E} | \Psi_\omega \rangle$):

$$\begin{aligned}
\langle \mathbf{x} | \mathcal{I} | \mathbf{y} \rangle &= \frac{\delta \langle \nabla \cdot \mathbf{E}(\mathbf{x}) \rangle_\omega}{\delta \omega(\mathbf{y})} \\
&= \langle \mathbf{x} | -\Delta | \mathbf{y} \rangle,
\end{aligned} \tag{III.22}$$

whose inverse is simply the Coulomb propagator:

$$\langle \mathbf{x} | \frac{1}{I} | \mathbf{y} \rangle = \frac{1}{4\pi|\mathbf{x} - \mathbf{y}|}. \tag{III.23}$$

We thus obtain the following expression for the Thouless-Valatin correction (III.15) in QED:

$$\begin{aligned}
\Delta E_{TV} &= \frac{1}{2} \int d^3x d^3y \\
&\langle \Psi_0 | \nabla \cdot \mathbf{E}(\mathbf{x}) \nabla \cdot \mathbf{E}(\mathbf{y}) | \Psi_0 \rangle \frac{1}{4\pi|\mathbf{x} - \mathbf{y}|},
\end{aligned} \tag{III.24}$$

which is recognized as the electrostatic energy in the state Ψ_0 . For the gaussian state (III.12), this gives

$$\Delta E_{TV} = \frac{V}{8} \int \frac{d^3p}{(2\pi)^3} \frac{p_i p_j}{p^2} G_{ij}^{-1}(\mathbf{p}), \tag{III.25}$$

which simply subtracts the longitudinal piece of the electric energy in eq. (III.20).

The corrected energy $\tilde{E} \equiv E_0 - \Delta E_{TV}$ reads therefore (cf. eqs. (III.20) and (III.25))

$$\tilde{E} = V \int \frac{d^3p}{(2\pi)^3} \left\{ \frac{1}{4} G_T^{-1}(p) + p^2 G_T(p) \right\}, \quad (\text{III.26})$$

and involves only the transverse piece $G_T \equiv \frac{1}{2}(\delta_{ij} - \hat{p}_i \hat{p}_j) G_{ij}$ of the kernel G_{ij} (we have written here $p \equiv |\mathbf{p}|$ and $\hat{p}_i \equiv p_i/p$). Then, the variational equation $\delta\tilde{E}/\delta G_T = 0$ gives G_T in the expected form

$$G_T(p) = \frac{1}{2p}. \quad (\text{III.27})$$

Together, eqs. (III.26) and (III.27) yield an energy density

$$\tilde{E} = V \int \frac{d^3p}{(2\pi)^3} |\mathbf{p}|, \quad (\text{III.28})$$

which is indeed the correct result for the QED ground state [1]. Thus, in the case of QED, the approximate projection method of Thouless and Valatin correctly subtracts the contribution of the unphysical, gauge, degrees of freedom from the average energy. Actually, since QED without fermions is a free theory, the variational solution above coincides with the exact solution [1]: the exact ground state is a gaussian wave functional like eq. (III.12) with a transverse kernel determined by eq. (III.27):

$$\begin{aligned} \Psi_{vac}[\mathbf{A}] &= \exp \left\{ - \int \frac{d^3p}{(2\pi)^3} \frac{p}{2} A^i(\mathbf{p}) (\delta_{ij} - \hat{p}_i \hat{p}_j) A^j(-\mathbf{p}) \right\} \\ &= \exp \left\{ - \int d^3x d^3y \frac{\mathbf{B}(\mathbf{x}) \cdot \mathbf{B}(\mathbf{y})}{4\pi^2 |\mathbf{x} - \mathbf{y}|^2} \right\}. \end{aligned} \quad (\text{III.29})$$

Moreover, wave functionals of the type shown in eq. (III.18) — i.e., gaussian states with a transverse kernel and a non-trivial phase factor — correspond to physical *charged* states, i.e., states of the quantum Maxwell theory in the presence of static, classical, external sources. Indeed, any such a state (which we denote here as Ψ_c) satisfies:

$$\nabla \cdot \mathbf{E}(\mathbf{x}) \Psi_c[\mathbf{A}] = \rho(\mathbf{x}) \Psi_c[\mathbf{A}], \quad (\text{III.30})$$

with the charge density $\rho(\mathbf{x}) = \nabla \cdot \mathbf{F}(\mathbf{x})$. The corresponding energy includes the Coulomb energy, as expected:

$$\begin{aligned}
\langle \Psi_c | H | \Psi_c \rangle &= E_0 + \frac{1}{2} \int d^3x \mathbf{F}^2(\mathbf{x}) \\
&= E_0 + \frac{1}{2} \int d^3x d^3y \frac{\rho(\mathbf{x})\rho(\mathbf{y})}{4\pi|\mathbf{x}-\mathbf{y}|}.
\end{aligned} \tag{III.31}$$

Non-Abelian charged states will be considered in Sec. IV.B below.

For other applications and a more complete study of the Thouless-Valatin method in the context of quantum field theory, see [24].

C. Approximate projection in QCD

Let us now consider the case of non-Abelian gauge theories. The corresponding "moment of inertia" is now a color matrix defined as the polarization tensor (cf. eq. (II.4))

$$\mathcal{I}^{ab}(\mathbf{x}, \mathbf{y}) = \frac{\delta \langle \mathcal{G}^a(\mathbf{x}) \rangle_\omega}{\delta \omega^b(\mathbf{y})} \Big|_{\omega=0}, \tag{III.32}$$

in the presence of an external constraint

$$\begin{aligned}
H_\omega &= H - H_{ext} \\
H_{ext} &= \int d^3x \omega^a(\mathbf{x}) \mathcal{G}^a(\mathbf{x}).
\end{aligned} \tag{III.33}$$

The analog of eq. (III.10), i.e. the gain in energy when projecting a wave functional $\Psi_0[\mathbf{A}]$ onto the subspace of gauge invariant states, reads:

$$\Delta E_{TV} = \int d^3x d^3y \langle \Psi_0 | \mathcal{G}^a(\mathbf{x}) \mathcal{G}^b(\mathbf{y}) | \Psi_0 \rangle \langle a, \mathbf{x} | \frac{1}{2\mathcal{I}} | b, \mathbf{y} \rangle \tag{III.34}$$

The energy functional to be used in variational calculations is therefore

$$\tilde{E} = E_0 - \Delta E_{TV}. \tag{III.35}$$

It is also possible to perform a projection on a subspace with a given color charge distribution $\langle \mathcal{G}^a(\mathbf{x}) \rangle_c \neq 0$ (The subscript c refers to expectation values over charged states). In such a case, however, the gauge generator appearing in the previous correction formula has to be replaced by its deviation $\hat{\mathcal{G}}$ away from the desired value:

$$\hat{\mathcal{G}}^a(\mathbf{x}) = \mathcal{G}^a(\mathbf{x}) - \langle \mathcal{G}^a(\mathbf{x}) \rangle_c. \quad (\text{III.36})$$

This modification guarantees that there is no correction for a state which is an exact eigenstate of the charge operator. The functional to be minimized in the subspace of gaussian functionals is thus

$$\tilde{E} = E_c - E_{\text{ext}} - \Delta \hat{E}_{TV}, \quad (\text{III.37})$$

where $E = \langle H \rangle_c$, $E_{\text{ext}} = \langle H_{\text{ext}} \rangle_c$ is the energy of the external constraint generating the charged state², and

$$\Delta \hat{E}_{TV} = \int d^3x d^3y \langle \hat{\mathcal{G}}^a(\mathbf{x}) \hat{\mathcal{G}}^b(\mathbf{y}) \rangle_c \langle a, \mathbf{x} | \frac{1}{2\mathcal{I}} | b, \mathbf{y} \rangle. \quad (\text{III.38})$$

This procedure is again reminiscent of the elimination of the center-of-mass motion in the mean field description of a composite system of A particles [19]. For a system characterized by a set of single-particle wave functions $\varphi_1, \varphi_2, \dots, \varphi_A$ the optimal state in the *center-of-mass* frame is obtained by minimizing the functional

$$\tilde{E} = \langle H \rangle - \frac{\langle \mathbf{P}^2 \rangle}{2M}, \quad (\text{III.39})$$

The Thouless-Valatin prescription for the total mass M in eq. (III.39) is to use the relation $\langle \mathbf{P} \rangle \equiv M\mathbf{v}$, where $\langle \mathbf{P} \rangle$ is the expectation value of the total momentum (III.4) in the presence of the external constraint $H_{\text{ext}} = \mathbf{v} \cdot \mathbf{P}$. This prescription gives the desired result $M = mA$ where m is the mass of the individual constituents. In a *moving* frame with velocity \mathbf{v} the single particle wave functions become

$$\varphi_i(\mathbf{x}) \rightarrow e^{i\chi(\mathbf{x})} \varphi_i(\mathbf{x}), \quad (\text{III.40})$$

with $\chi(\mathbf{x}) = m\mathbf{v} \cdot \mathbf{x}$. Individual momentum operators in the moving frame are obtained by the gauge transformation

²That is, an exact charged state Ψ_c is defined as an eigenstate of $H - H_{\text{ext}}$.

$$\nabla \rightarrow \nabla + i(\nabla\chi), \quad (\text{III.41})$$

i.e. $\mathbf{p}_i \rightarrow \mathbf{p}_i - \langle \mathbf{p}_i \rangle$. The functional providing the adequate state at the minimum is

$$\tilde{E} = \langle H \rangle - \mathbf{v} \cdot \mathbf{P} - \frac{\langle (\mathbf{P} - \langle \mathbf{P} \rangle)^2 \rangle}{2M}, \quad (\text{III.42})$$

in agreement with eq. (III.37). This procedure to implement Gauss's law will be important in Sec. IV when applied to the calculation of the interaction energy of color charges.

Still in the case of charged states, the chromostatic energy E_{chromo} is given, in our approximation scheme, by the classical chromostatic energy corrected by the Thouless-Valatin term :

$$E_{chromo} = \int d^3x d^3y \langle \mathcal{G}^a(\mathbf{x}) \rangle_c \langle \mathcal{G}^b(\mathbf{y}) \rangle_c \langle a, \mathbf{x} | \frac{1}{2\mathcal{I}} | b, \mathbf{y} \rangle + \Delta \hat{E}_{TV} \quad (\text{III.43})$$

so that

$$E_{chromo} = \int d^3x d^3y \langle \mathcal{G}^a(\mathbf{x}) \mathcal{G}^b(\mathbf{y}) \rangle_c \langle a, \mathbf{x} | \frac{1}{2\mathcal{I}} | b, \mathbf{y} \rangle. \quad (\text{III.44})$$

(This is the analog of using $E_{rot} = \langle \mathbf{J}^2 \rangle / 2I$ as an approximation for the rotational energy of a deformed nucleus.) In the case of QED this identification is obvious on equations like (III.24) or (III.31).

To conclude, the central result of this section is the non-local energy functional (III.35) (or (III.37) in the case of charged states) which approximately corrects for the lack of gauge symmetry when working with gaussian states. This energy functional is the starting point of the variational method we propose for gauge theories. Note that the corrective energy term ΔE_{TV} in eq. (III.34) is a priori of a non-perturbative nature. Our aim in what follows is to check the ultraviolet behaviour of this approximation scheme. We shall thus consider the variational calculations in the perturbative regime $g \ll 1$.

IV. ONE-LOOP BETA FUNCTION FROM VARIATIONAL CALCULATIONS

In this section, we shall use eq. (III.43) to estimate the electrostatic energy E_{chromo} of a non-Abelian charged state, up to order g^2 in perturbation theory. This will allow us to recover the standard expression for the QCD beta function in the one-loop approximation.

A. Moment of inertia for color rotations

The first step is the calculation of the moment of inertia for color rotations, $\mathcal{I}_{ab}(\mathbf{x}, \mathbf{y})$, which enters eq. (III.43). Unlike QED, where this quantity has been computed exactly (cf. eq. (III.23)), in QCD we shall give only a perturbative estimate of I , valid to the order of interest (i.e., up to order g^2). To this aim, it is sufficient to perform variational calculations in the vicinity of the perturbative vacuum.

To zeroth order, the vacuum of the Yang-Mills theory is the same as for the Maxwell theory, namely (cf. eq. (III.29)):

$$\Psi_0[\mathbf{A}] = \mathcal{N}^{-1} \exp \left\{ - \left\langle A \left| \frac{G^{-1}}{4g^2} \right| A \right\rangle \right\}, \quad (\text{IV.1})$$

where

$$(G^{-1})_{ij}^{ab}(\mathbf{k}) = \delta_{ab} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \right) G_{\mathbf{k}}^{-1}, \quad (\text{IV.2})$$

and $G_{\mathbf{k}}^{-1} = 2k$. In the calculation of I below, we shall never need to go beyond this leading order approximation for $G_{\mathbf{k}}$.

Note that, even with such a transverse kernel, the functional (IV.1) is still not invariant under *non-Abelian* gauge transformations; that is, this is a deformed state, according to the terminology in Sec. III. In order to compute its moment of inertia under color rotations, one has to study the response of this state to an external constraint of the form (III.33). The trial wave functional in the presence of this constraint reads :

$$\Psi_\omega[\mathbf{A}] = \mathcal{N}^{-1} e^{-i\langle \mathbf{F} | \mathbf{A} \rangle} \exp \left\{ - \left\langle A \left| \frac{G^{-1}}{4g^2} - i\Sigma \right| A \right\rangle \right\}, \quad (\text{IV.3})$$

which involves two additional variational parameters: the vector field $F_a^i(\mathbf{x})$ (which fixes the expectation value of the electric field: $\langle \mathbf{E}_a \rangle = \mathbf{F}_a$), and the matrix $\Sigma_{ab}^{ij}(\mathbf{x}, \mathbf{y})$, which is taken to be transverse in its spatial indices. The emergence of Σ is a hallmark of the non-Abelian behaviour (recall that $\Sigma = 0$ in the corresponding Abelian problem; cf. eq. (III.18)).

We shall shortly see that, as $\omega \rightarrow 0$, F and Σ are linear in ω , while the first correction to G^{-1} is only quadratic³, and therefore does not matter for the calculation of \mathcal{I} (cf. eq. (III.32)). Thus, for the present purposes, we can take G^{-1} as in eq. (IV.2). The variational parameters in (IV.3) are obtained by minimizing the following functional

$$\begin{aligned} E_\omega &\equiv \langle \Psi_\omega | H - H_{ext} | \Psi_\omega \rangle \\ &= \langle H \rangle_\omega - \int d^3x \omega^a \langle \mathcal{G}^a \rangle_\omega, \end{aligned} \quad (IV.4)$$

with respect to variations in F_a^i and Σ_{ab}^{ij} . A straightforward calculation yields:

$$\langle H \rangle_\omega = \frac{g^2}{2} \int d^3x \{ \mathbf{F}_a \cdot \mathbf{F}_a(\mathbf{x}) + 4g^2 \text{Tr} \langle \mathbf{x} | \Sigma G \Sigma | \mathbf{x} \rangle \}, \quad (IV.5)$$

where we have kept only the terms involving the variational parameters. Similarly,

$$\langle \mathcal{G}^a(\mathbf{x}) \rangle_\omega = \nabla \cdot \mathbf{F}^a(\mathbf{x}) - ig^2 \text{Tr} \langle \mathbf{x} | T^a[\Sigma, G] | \mathbf{x} \rangle. \quad (IV.6)$$

After inserting these expressions in eq. (IV.4), and taking variations with respect to F and Σ , we derive the following expressions for the variational parameters (in momentum space) :

$$F_a^i(\mathbf{q}) = -\frac{iq^i}{g^2} \omega_a(\mathbf{q}), \quad (IV.7)$$

and, for the transverse⁴ components of Σ ,

$$\begin{aligned} \langle k' b | \Sigma | k c \rangle &= \frac{(2\pi)^3}{V} \delta(\mathbf{k}' - \mathbf{k} - \mathbf{q}) \\ &\times \frac{1}{2g^2} \omega^a(\mathbf{q}) f_{abc} \left(\frac{G_{\mathbf{k}} - G_{\mathbf{k}'}}{G_{\mathbf{k}} + G_{\mathbf{k}'}} \right), \end{aligned} \quad (IV.8)$$

³Such quadratic corrections occur since, in contrast to QED, the non-Abelian constraint (III.33) generates a coupling between G and F in the variational equations.

⁴By which we mean transversality with respect to both \mathbf{k} and \mathbf{k}' , as requested by the expressions (IV.5) and (IV.6).

where V is the total volume and $G_{\mathbf{k}}$ is defined in (IV.2).

By using eqs. (IV.6), (IV.7) and (IV.8), we can finally express the average color charge $\langle \mathcal{G} \rangle_\omega$ in terms of ω^a . This is conveniently decomposed into an ‘‘Abelian’’ and a ‘‘non-Abelian’’ piece, as corresponding to the two pieces in the r.h.s. of eq. (IV.6): $\langle \mathcal{G}^a \rangle_\omega = \rho_A^a + \rho_{NA}^a$, with:

$$\rho_A^a(\mathbf{q}) = i\mathbf{q} \cdot \mathbf{F}^a(\mathbf{q}) = \frac{1}{g^2} \mathbf{q}^2 \omega^a(\mathbf{q}), \quad (\text{IV.9})$$

and, respectively,

$$\begin{aligned} \rho_{NA}^a(\mathbf{q}) = g^2 f_{abc} \int \frac{d^3k}{(2\pi)^3} & \left(\delta_{ij} - \frac{k'_i k'_j}{k'^2} \right) \left(\delta_{ij} - \frac{k_i k_j}{k^2} \right) \\ & \times (G_{\mathbf{k}} - G_{\mathbf{k}'}) \langle \mathbf{k}'_c | \Sigma | \mathbf{k} \mathbf{b} \rangle, \end{aligned} \quad (\text{IV.10})$$

where $\mathbf{k}' = \mathbf{k} + \mathbf{q}$. From eqs. (IV.8)–(IV.10) we note that the ‘‘non-Abelian’’ charge density in eq. (IV.10) is a correction of order g^2 relative to the ‘‘Abelian’’ one in eq. (IV.9). Thus, as anticipated after eq. (IV.2), it is consistent to evaluate this correction with the free kernel $G_{\mathbf{k}} = 1/2k$.

The integral in eq. (IV.10) is logarithmically ultraviolet divergent, so it must be evaluated with an upper cutoff. It turns out that this divergence is a part of the charge renormalization in QCD (see Sec. IV.B below). To reconstruct the associated beta function, we need, as usual, only the coefficient of the divergent logarithm. The latter is insensitive to the details of the UV regularization, so we shall consider, for simplicity, a sharp momentum cutoff Λ .

Also, in order to isolate the leading logarithm, we can perform kinematical approximations relying on the inequality $k \gg q$ (since the external momentum q is fixed, while the leading contribution to the integral in eq. (IV.10) comes from relatively large momenta). Physically, we are indeed interested in smooth charge distributions. This allows us to replace $\mathbf{k} + \mathbf{q}$ by \mathbf{k} and thus $(\delta_{ij} - \frac{k'_i k'_j}{k'^2})(\delta_{ij} - \frac{k_i k_j}{k^2})$ by 2. By also using eq. (IV.8) for Σ , we then obtain :

$$\rho_{NA}^a(\mathbf{q}) = \frac{C_N}{2} \omega^a(\mathbf{q}) X(\mathbf{q}), \quad (\text{IV.11})$$

where $C_N = \text{Tr}(T^3 T^3) = N$ for $SU(N)$, and

$$X(\mathbf{q}) \equiv \int \frac{d^3 k}{(2\pi)^3} \frac{[\varepsilon(\mathbf{k}') - \varepsilon(\mathbf{k})]^2}{\varepsilon(\mathbf{k})\varepsilon(\mathbf{k}')(\varepsilon(\mathbf{k}) + \varepsilon(\mathbf{k}'))}, \quad (\text{IV.12})$$

with $\varepsilon(\mathbf{k}) \equiv |\mathbf{k}|$. Here again we can replace $\mathbf{k}' \simeq \mathbf{k}$ everywhere except in the numerator which must be expanded to second order in q :

$$[\varepsilon(\mathbf{k} + \mathbf{q}) - \varepsilon(\mathbf{k})]^2 \simeq q^2 \cos^2 \theta. \quad (\text{IV.13})$$

where θ is the angle between the space vectors \mathbf{k} and \mathbf{q} . The angular average yields $\langle \cos^2 \theta \rangle = \int_0^\pi d\theta \sin \theta \cos^2 \theta = 2/3$, so, finally:

$$X(\mathbf{q}) = \frac{q^2}{12\pi^2} \int \frac{dk}{k}, \quad (\text{IV.14})$$

which is logarithmically divergent in the ultraviolet, as expected, but also in the infrared: the infrared divergence is an artifact of the previous manipulations (in a more careful calculation, this would be screened by q), and to the order of interest we can just regulate it with an ad-hoc infrared cutoff μ . This yields $\rho_{NA}^a = \alpha \rho_A^a$, with

$$\alpha \equiv \frac{g^2 C_N}{48\pi^2} \ln \frac{\Lambda^2}{\mu^2}. \quad (\text{IV.15})$$

The resulting value of the moment of inertia $\mathcal{I}_{ab} \equiv \delta\rho^a / \delta\omega^b$ reads finally:

$$\mathcal{I}_{ab}(\mathbf{q}) = \frac{\delta_{ab}}{g^2} q^2 (1 + \alpha). \quad (\text{IV.16})$$

This should be compared to the corresponding Abelian result⁵ $I(q) = q^2$. We see that the quantum fluctuations in QCD produce an increase of the moment of inertia, which corresponds to the *screening* of color charges by quantum fluctuations. The size of the screening effect that we have obtained agrees indeed with the results of other approaches [26].

⁵The overall factor $1/g^2$ in eq. (IV.16) is simply a consequence of our different normalizations for the field strengths in QCD and QED; compare, in this respect, eqs. (III.12) and (IV.1).

B. Interaction energy in the presence of a background electric field

Interesting properties of the vacuum include its response to an external chromo-electric field, which can be generated by an external constraint of the form

$$H_{\text{ext}} = g^2 \int d^3x \mathbf{E}_{\text{ext}}^a(\mathbf{x}) \cdot \mathbf{E}^a(\mathbf{x}). \quad (\text{IV.17})$$

For this constraint, we shall compute the induced electric mean field and charge density, and the associated electrostatic energy. By comparing the latter with the bare Coulomb interaction, we shall then identify the chromo-electric susceptibility, or charge renormalization. As we shall see, the variational formalism provides a transparent picture of the underlying phenomena of screening and anti-screening.

The optimal state Ψ_c in the presence of the constraint (IV.17) is of the form

$$\Psi_c[\mathbf{A}] = \mathcal{N}^{-1} e^{-i\langle \mathbf{F} | \mathbf{A} \rangle} \exp \left\{ - \left\langle A \left| \frac{G^{-1}}{4g^2} \right| A \right\rangle \right\}, \quad (\text{IV.18})$$

where the parameter F_i^a (the electric mean field) will be related to the external field E_{ext}^i by the variational equations (see eq. (IV.23) below). Note that, in contrast to eq. (IV.3), there is no Σ term in eq. (IV.18) above; this is so because the external perturbation here is different (compare eqs. (IV.17) and (III.33)): it contains a term linear in \mathbf{E}^a , but no non-Abelian term like $[A^i, E^i]$. This situation is analogous to the case of an anharmonic oscillator with an external constraint $H_{\text{ext}} = \alpha p$. The only changes are a factor e^{ip_0x} and, in higher orders in g , a modification of the real part of the width.

According to the discussion in Sec. III (see especially eqs. (III.37) and (III.38)), the energy functional to be minimized in this case is $\tilde{E} = E_c - \langle H_{\text{ext}} \rangle_c - \Delta \hat{E}_{TV}$. The terms involving F_a^i in this functional read:

$$\begin{aligned} E_c - E_{\text{ext}} - \Delta \hat{E}_{TV} = & g^2 \int d^3x \left\{ \frac{1}{2} \mathbf{F}^a \cdot \mathbf{F}^a - \mathbf{E}_{\text{ext}}^a \cdot \mathbf{F}^a \right\} \\ & - g^2 f_{acd} f_{bef} \int d^3x d^3y F_i^d(\mathbf{x}) F_j^f(\mathbf{y}) \\ & \times \langle \mathbf{x} | G_{ij}^{ce} | \mathbf{y} \rangle \langle a, \mathbf{x} | \frac{1}{2\mathcal{L}} | b, \mathbf{y} \rangle \end{aligned} \quad (\text{IV.19})$$

The last term in the r.h.s. corresponds to the Thouless-Valatin correction, eq. (III.38)). Note that, because of the subtraction of the average charge in $\hat{\mathcal{G}}^a \equiv \mathcal{G}^a - \langle \mathcal{G}^a \rangle_c$, it is only the non-Abelian part of the Gauss operator (i.e., the term $i[A^i, E^i]$ in eq. (II.4)) which contributes to eq. (IV.19). To evaluate this contribution, we first rewrite it in momentum space:

$$\begin{aligned} \Delta \hat{E}_{TV} &= g^2 f_{acd} f_{bef} \int \frac{d^3 k}{(2\pi)^3} \frac{d^3 q}{(2\pi)^3} F_i^d(\mathbf{q}) F_j^f(-\mathbf{q}) \\ &\quad \times G_{ij}^{ce}(\mathbf{k}) \langle a | \frac{1}{2\mathcal{I}(\mathbf{k} + \mathbf{q})} | b \rangle. \end{aligned} \quad (\text{IV.20})$$

To the order of interest, we can replace the moment of inertia in the equation above by its leading order expression: $\mathcal{I}(\mathbf{k}) \simeq k^2/g^2$. Then, by performing similar manipulations as in the calculation of the X term in eqs. (IV.10)–(IV.14), we finally obtain:

$$\begin{aligned} \Delta \hat{E}_{TV} &\simeq \frac{g^4 C_N}{3} \int \frac{d^3 k}{(2\pi)^3} \frac{d^3 q}{(2\pi)^3} F_i^a(\mathbf{q}) F_i^a(-\mathbf{q}) \frac{1}{2k} \frac{1}{k^2} \\ &\equiv \frac{g^2 \delta}{2} \int d^3 x F_i^a(\mathbf{x}) F_i^a(\mathbf{x}), \end{aligned} \quad (\text{IV.21})$$

where

$$\delta \equiv \frac{g^2 C_N}{6\pi^2} \int \frac{dk}{k} = \frac{g^2 C_N}{12\pi^2} \ln \left(\frac{\Lambda^2}{\mu^2} \right). \quad (\text{IV.22})$$

After inserting (IV.21) in (IV.19) and minimizing with respect to $F_i^a(\mathbf{x})$, we obtain

$$(1 - \delta) F_a^i(\mathbf{x}) = E_{\text{ext } a}^i(\mathbf{x}). \quad (\text{IV.23})$$

At this point it is convenient to introduce the charge distribution associated to the external field E_{ext}^i ,

$$\rho_{\text{ext}}^a(\mathbf{x}) \equiv \nabla \cdot \mathbf{E}_{\text{ext}}^a(\mathbf{x}), \quad (\text{IV.24})$$

to be referred to as the *external charge* in what follows: this would be the charge in the system in the absence of polarization effects. The actual charge is rather

$$\begin{aligned} \rho^a(\mathbf{x}) &\equiv \langle \Psi_c | \mathcal{G}^a(\mathbf{x}) | \Psi_c \rangle = \nabla \cdot \mathbf{F}^a \\ &= \frac{\rho_{\text{ext}}^a}{1 - \delta} \simeq \rho_{\text{ext}}^a (1 + \delta), \end{aligned} \quad (\text{IV.25})$$

where the second line follows from eq. (IV.24). Note that this relation implies an *antiscreening* of the external charge, since ρ^a is bigger than ρ_{ext}^a . The difference $\rho - \rho_{\text{ext}} = \rho_{\text{ext}} \delta$ may be interpreted as an *induced charge* (see also Sec. IV.C in Ref. [25] and Appendix A for an alternative computation of this quantity).

We are finally in position to compute the chromostatic interaction E_{chromo} in the optimal state Ψ_c . This is given by eq. (III.43) which, together with the above expressions (IV.25) for $\langle \mathcal{G}^a(\mathbf{x}) \rangle_c$, and (IV.21) for $\Delta \hat{E}_{TV}$, implies:

$$E_{\text{chromo}} \simeq g^2 \frac{1+3\delta}{1+\alpha} \int \frac{d^3q}{(2\pi)^3} \rho_{\text{ext}}^a(\mathbf{q}) \rho_{\text{ext}}^a(-\mathbf{q}) \frac{1}{2\mathbf{q}^2}, \quad (\text{IV.26})$$

up to corrections of higher order in g . There is here a factor $(1+\delta)^2 \simeq 1+2\delta$ arising from the induced charge (cf. eq. (IV.25)), another one arising from the Thouless-Valatin correction (IV.21), and a factor $(1+\alpha)$ due to the moment of inertia (cf. eq. (IV.16)). The interaction energy (IV.26) is still Coulomb like,

$$E_{\text{chromo}} = g_R^2 \int \frac{d^3q}{(2\pi)^3} \rho_{\text{ext}}^a(\mathbf{q}) \rho_{\text{ext}}^a(-\mathbf{q}) \frac{1}{2\mathbf{q}^2}, \quad (\text{IV.27})$$

but with a modified coupling constant given by

$$g_R^2(\mu) = g^2 \frac{1+3\delta}{1+\alpha} \quad (\text{IV.28})$$

or, to first order in g^2 ,

$$\frac{1}{g_R^2(\mu)} = \frac{1}{g^2} - \frac{11C_N}{48\pi^2} \ln \frac{\Lambda^2}{\mu^2}. \quad (\text{IV.29})$$

This is the correct one-loop value for the renormalized coupling constant [26]. Note that, in the present calculation, this involves three types of contributions: indeed, the factor 11 in the last equation has arised as $11 = 8 + 4 - 1$, where the 8 corresponds to anti-screening by the induced charge (cf. eq. (IV.25)), the 4 is another anti-screening contribution due to the Thouless-Valatin correction (IV.21), and the (-1) is a screening contribution arising via the correction of order g^2 to the moment of inertia.

V. VACUUM ENERGY IN A MAGNETIC BACKGROUND FIELD

In the previous section, we have studied the electric sector of the vacuum of the Yang-Mills theory, by using a combination of variational and perturbative techniques. In what follows, we shall perform a similar analysis of the magnetic sector. To this aim, we consider the Yang-Mills theory in the presence of a (constant) magnetic background field \bar{B}_a^i , and compute the background field energy by using the variational principle. The final result is not new (it coincides with the one-loop result by Savvidy [20]), but it rather serves as a test for our variational method in the magnetic sector, and in the perturbative regime.

The relevant trial wave variational is the gaussian functional Ψ_0 in eq. (II.8) with the “center” field $\bar{A}_a^i(\mathbf{x})$ chosen so as to reproduce the desired magnetic field \bar{B}_a^i (a convenient choice will be given later). This state is not gauge-invariant, so its energy $E_0 \equiv \langle \Psi_0 | H | \Psi_0 \rangle$, eq. (II.9), must be corrected with the Thouless-Valatin energy ΔE_{TV} , to be computed in the next subsection. Then, by applying the variational principle to the corrected energy $\tilde{E} = E_0 - \Delta E_{TV}$, we shall determine the kernel of the Gaussian (in Sec. V.B). Finally, in Sec. V.C, we shall compute the energy of the magnetic field and the associated gluon condensate, and verify that these quantities are related by the trace anomaly relation, as it should. In this calculation, the standard one-loop beta function will emerge once again.

A. The Thouless-Valatin energy in the background field

We start by computing the moment of inertia $\mathcal{I}_{ab}(\mathbf{x}, \mathbf{y})$ in the presence of the background field $\bar{A}_a^i(\mathbf{x})$. As already explained, this requires constructing the variational ground state Ψ_ω for the constrained Hamiltonian H_ω in eq. (III.33). This state is of the form (compare to eq. (IV.3))

$$\begin{aligned} \Psi_\omega = & \mathcal{N}^{-1} e^{-i \langle F | A - \bar{A} \rangle} \\ & \times \exp \left\{ - \left\langle A - \bar{A} \left| \frac{G^{-1}}{4} - i \Sigma \right| A - \bar{A} \right\rangle \right\}, \end{aligned} \quad (\text{V.1})$$

where the parameters F_a^i and Σ_{ij}^{ab} are related to ω^a by the variational principle (recall the discussion in Sec. IV.A). It turns out that the matrix Σ will not play any role in what follows: indeed, below we shall need the moment of inertia only to leading order in g , while Σ counts starting with order g^2 (cf. Sec. IV.A). We then write, as in eq. (IV.4),

$$E_\omega = \langle H \rangle_\omega - \int d^3x \omega^a(\mathbf{x}) \langle \mathcal{G}^a(\mathbf{x}) \rangle_\omega, \quad (\text{V.2})$$

with $\langle H \rangle_\omega$ given by eq. (IV.5), and

$$\langle \mathcal{G}^a(\mathbf{x}) \rangle_\omega = (\mathcal{D}_i F^i)^a(\mathbf{x}) + \mathcal{O}(g^2), \quad (\text{V.3})$$

where $\mathcal{D}^i \equiv \partial^i - i\bar{A}^i$ is the covariant derivative defined by the background field (cf. eq. (II.11)), and the neglected terms, of $\mathcal{O}(g^2)$, would involve Σ (cf. eq. (IV.6)). The variation with respect to F_a^i yields then

$$F_a^i(\mathbf{x}) = -\frac{1}{g}(\mathcal{D}_i \omega)^a(\mathbf{x}), \quad (\text{V.4})$$

which differs from the corresponding expression in eq. (IV.7) only by the replacement of the ordinary derivative ∂_i by the covariant one \mathcal{D}_i . Together with eq. (V.3), this provides the moment of inertia to the order of interest:

$$\begin{aligned} \mathcal{I}^{ab}(\mathbf{x}, \mathbf{y}) &\equiv \frac{\delta \langle \mathcal{G}^a(\mathbf{x}) \rangle_\omega}{\delta \omega^b(\mathbf{y})} = -\frac{1}{g^2} (\mathcal{D}_x^2)^{ab} \delta^{(3)}(\mathbf{x} - \mathbf{y}) + \mathcal{O}(1) \\ &= \frac{1}{g^2} \langle a, \mathbf{x} | \Pi^2 | b, \mathbf{y} \rangle + \mathcal{O}(1). \end{aligned} \quad (\text{V.5})$$

We have introduced here the *kinetic momentum* $\Pi_j \equiv i\mathcal{D}_j = i\delta_j + \bar{A}_j$, and $\Pi^2 \equiv \Pi_j \Pi_j$.

Within the same accuracy, one has also:

$$\begin{aligned} \langle \Psi_0 | \mathcal{G}^a(\mathbf{x}) \mathcal{G}^b(\mathbf{y}) | \Psi_0 \rangle &\approx \frac{1}{4g^2} \mathcal{D}_{i,\mathbf{x}}^{ac} \mathcal{D}_{db}^{j,\mathbf{y}} (G^{-1})_{ij}^{cd}(\mathbf{x}, \mathbf{y}) \\ &= \frac{1}{4g^2} \langle a, \mathbf{x} | \Pi_i G_{ij}^{-1} \Pi_j | b, \mathbf{y} \rangle. \end{aligned} \quad (\text{V.6})$$

We are now in position to compute the Thouless-Valatin energy ΔE_{TV} , cf. eq. (III.34): by combining eqs. (V.5) and (V.6), one obtains:

$$\Delta E_{TV} \approx \frac{1}{8} \int d^3x \langle a, \mathbf{x} | \left(\Pi_i \frac{1}{\Pi^2} \Pi_j \right) G_{ij}^{-1} | a, \mathbf{x} \rangle, \quad (\text{V.7})$$

up to corrections of order g^2 .

B. The variational equation for G

The improved energy functional $\tilde{E} = E_0 - \Delta E_{TV}$ reads therefore (cf. eqs. (II.9) and (V.7))

$$\begin{aligned} \tilde{E} = \int d^3\mathbf{x} \left\{ \frac{1}{2g^2} \bar{B}_i^a(\mathbf{x}) \bar{B}_i^a(\mathbf{x}) \right. \\ \left. + \frac{1}{8} \langle a, \mathbf{x} | (\delta_{ij} - \Pi_i \frac{1}{\Pi^2} \Pi_j) G_{ij}^{-1} | a, \mathbf{x} \rangle \right. \\ \left. + \frac{1}{2} \text{Tr} [K G(\mathbf{x}, \mathbf{x})] + \mathcal{O}(g^2) \right\}. \end{aligned} \quad (\text{V.8})$$

Note that the last two terms in (II.9) do not contribute to this order. As obvious from this equation, the Thouless-Valatin correction makes the kinetical part of the energy *covariantly* transverse. Since the operator K_{ij} is transverse as well (cf. eq. (II.10)),

$$K_{ij} = \Pi^2 \delta_{ij} - \Pi_i \Pi_j + 2[\Pi_i, \Pi_j], \quad (\text{V.9})$$

it follows that the projected energy (V.8) involves only the *transverse* components of the kernel G . Thus, without loss of generality, we can restrict ourselves to a (covariantly) transverse kernel in what follows:

$$\Pi^i G_{ij}^{-1} = 0 = G_{ij}^{-1} \Pi^j. \quad (\text{V.10})$$

To formalize this, it is convenient to introduce transverse and longitudinal projectors as follows:

$$\hat{P}_{ij} \equiv \Pi_i \frac{1}{\Pi^2} \Pi_j, \quad \hat{Q} \equiv 1 - \hat{P}. \quad (\text{V.11})$$

They satisfy :

$$\begin{aligned} \hat{P}_{ij}^2 &= \Pi_i \frac{1}{\Pi^2} \Pi_k \Pi^k \frac{1}{\Pi^2} \Pi_j = \hat{P}_{ij}, \\ \hat{Q}_{ij}^2 &= (1 - \hat{P})_{ij}^2 = (1 - \hat{P})_{ij} = \hat{Q}_{ij}. \end{aligned} \quad (\text{V.12})$$

Then, a transverse kernel is one satisfying $G = \hat{Q}G\hat{Q}$ (and similarly for G^{-1}). For such a kernel, the variational principle (i.e., the minimization of \tilde{E} , eq. (V.8), with respect to G) produces the following gap equation:

$$\frac{1}{4G^2} \simeq K, \quad (\text{V.13})$$

which determines G to the order of interest. In particular, as $\bar{A} \rightarrow 0$, G reduces to the free, or Abelian, expression in eqs. (IV.2) and (III.27). Thus, the only non-trivial effects which are taken here into account are those associated with the background field.

Note that eq. (V.13) can only be valid at sufficiently high energy, or small coupling constant: indeed, the operator K admits negative modes [20,27,28]. We thus assume that an infrared cutoff has been set - this does not affect the ultraviolet behaviour of the theory, which is our main interest here.

C. The energy of the background field

The previous equations provide the optimal gaussian kernel for a given background field \bar{A} and thus the effective potential $V(\bar{A})$ which is the expectation value of the energy in this state. The next step in our variational approach is to find the minimum of the effective potential. Constructing V for an arbitrary background is however a difficult task. For this reason we now consider a restricted variational space defined by the following background field :

$$\bar{A}_x = 0 ; \bar{A}_y = xBT^3 ; \bar{A}_z = 0. \quad (\text{V.14})$$

This corresponds to a constant magnetic field in the z -direction and in the third color.

With this choice of the background field, we are now able to compute the energy (V.8) in the optimal variational state, which is the gaussian state (II.8) with a transverse (in the sense of eq. (V.10)) kernel G^{-1} satisfying eq. (V.13). The latter equation shows that, at the minimum, the following identity holds:

$$\text{Tr}\{K\langle\mathbf{x}|G|\mathbf{x}\rangle\} = \frac{1}{4} \text{Tr}\langle\mathbf{x}|G^{-1}|\mathbf{x}\rangle. \quad (\text{V.15})$$

That is, magnetic and electric fluctuations have equal energies in our variational ground states, which is merely the virial theorem in the present context. Thus,

$$\tilde{E}_{min} \simeq \int d^3\mathbf{x} \left\{ \frac{1}{2g^2} \bar{B}_i^a(\mathbf{x}) \bar{B}_i^a(\mathbf{x}) + \frac{1}{4} \langle a, \mathbf{x} | G^{-1} | a, \mathbf{x} \rangle \right\}. \quad (\text{V.16})$$

This involves the matrix element $\langle \mathbf{x} | G^{-1} | \mathbf{x} \rangle$, which we shall compute in Appendix A by using the Schwinger proper-time representation (cf. eq. (V.13)) :

$$\langle \mathbf{x} | G^{-1} | \mathbf{x} \rangle = \frac{1}{\sqrt{\pi}} \int_0^\infty \frac{dt}{t^{3/2}} \langle \mathbf{x} | (1 - e^{-tK}) | \mathbf{x} \rangle \quad (\text{V.17})$$

This integral develops ultraviolet divergences as $t \rightarrow 0$, which we shall regularize by shifting the lower bounds of the integral from 0 to $1/\Lambda^2$. As in the electric case, we are mainly interested in the ultraviolet renormalization of the energy (V.16); to this aim, it is sufficient to extract the terms which diverge when $\Lambda \rightarrow \infty$ in eq. (V.17). This is described in detail in Appendix A, from which we quote here the final result:

$$\text{Tr} \langle \mathbf{x} | G^{-1} | \mathbf{x} \rangle = (\dots) \Lambda^4 - \frac{11C_N}{48\pi^2} B^2 \ln \frac{\Lambda^2}{B} + \mathcal{O}(g^2), \quad (\text{V.18})$$

(the coefficient in front of Λ^4 is an uninteresting field-independent number that will be omitted in what follows). Remarkably, there is no divergent term in Λ^2 (which, for dimensional reasons, would be necessarily of the form $\Lambda^2 B$): this is so because of rotational and gauge symmetries which require the magnetic field to enter only in the scalar product $\mathbf{S} \cdot \mathbf{B}_a T^a = BS_z T^3$, whose trace is however zero (see Appendix A for more details).

Note also the numerical factor in front of the logarithmic divergence in eq. (V.18): this is the factor leading to the correct one-loop beta function after renormalization (see below). The projection on (covariantly) transverse gaussian states has been crucial in getting this factor right: without this, we would have obtained a factor $\frac{7}{2}$ instead of the correct factor $\frac{11}{3}$ (compare in this respect eqs. (B.19) and (B.20) in Appendix A).

Finally note that the field strength B appears as an infrared cutoff in eq. (V.18). This is expected from equations (B.6) and (B.7) where the proper time variable always appears in the combination tB . However a complete derivation of (V.18) requires a detailed treatment of unstable modes.

To conclude,

$$\tilde{E}/V \simeq \frac{1}{2g^2} B^2 - C_N \frac{11}{48\pi^2} \frac{B^2}{2} \ln \frac{\Lambda^2}{B}, \quad (\text{V.19})$$

showing that the background field energy has no field dependent UV divergences other than the logarithmic one which can be absorbed into the renormalization of the coupling constant.

We then write, as usual (μ is the subtraction scale),

$$\frac{1}{g_R^2(\mu)} = \frac{1}{g^2} - C_N \frac{11}{48\pi^2} \ln \frac{\Lambda^2}{\mu^2} \quad (\text{V.20})$$

which provides the correct one-loop beta function, as anticipated. The renormalized field energy density reads then:

$$\mathcal{H}(B) = \frac{B^2}{2g_R^2(\mu)} + \frac{B^2}{2} \frac{11}{48\pi^2} C_N \ln \frac{B}{\mu^2}, \quad (\text{V.21})$$

which coincides with the result obtained by Savvidy in perturbation theory [20]. An advantage of the present approach, however, is that it can be improved by using a larger variational space, which is expected to cure the difficulties associated with negative modes [27,28].

As discussed in [20], the energy density (V.21) exhibits a minimum for a non-zero value $B = B_{min}$ of the background field, with

$$B_{min} = \frac{\mu^2}{\sqrt{e}} \exp\left(-\frac{16\pi^2}{g_R^2(\mu)} \frac{3}{11C_N}\right) \quad (\text{V.22})$$

The value of the energy density at this minimum is :

$$\langle \mathcal{H} \rangle_{min} = -\frac{1}{64\pi^2} \frac{11}{3} C_N B_{min}^2 \quad (\text{V.23})$$

which is indeed negative. Our variational vacuum state is therefore characterized by a magnetic field condensate (see, however, Refs. [27,28] for potential problems with such a state).

From the previous results, it is now straightforward to evaluate the gluon condensate in our variational vacuum:

$$\begin{aligned} \langle F^{\mu\nu} F_{\mu\nu} \rangle &\equiv 2g^2 \left(\frac{1}{g^2} \langle B_i^a B_i^a \rangle - \langle g^2 E_i^a E_i^a \rangle \right) \\ &= 2B_{min}^2, \end{aligned} \quad (\text{V.24})$$

where the second line follows from the aforementioned “virial theorem” (V.15). Eqs. (V.23) and (V.24) can be combined into:

$$\langle \mathcal{H} \rangle_{min} = -\frac{11C_N}{128\pi^2} \langle F^{\mu\nu} F_{\mu\nu} \rangle_{min} \quad (\text{V.25})$$

which is consistent, as it should, with the trace anomaly relation:

$$\langle \theta_\mu^\mu \rangle = \frac{\beta(g)}{2g^3} \langle F^{\mu\nu} F_{\mu\nu} \rangle. \quad (\text{V.26})$$

Indeed, with $\langle \mathcal{H} \rangle = \frac{1}{4} \langle \theta_\mu^\mu \rangle$, and the one-loop beta function (which here is a consequence of eq. (V.20))

$$\beta(g) = -\frac{11}{48\pi^2} C_N g^3, \quad (\text{V.27})$$

eq. (V.26) becomes identical to eq. (V.25).

An attractive feature of the formula (V.25) is that it involves two quantities which are independently accessible experimentally (at least indirectly). Indeed, the left hand side of this equation is the energy density of the vacuum, which can be identified with the fourth power of the bag constant, $\mathcal{B}^4 = -(240\text{MeV})^4$ [29], whereas the right hand side depends on the gluon condensate which is known from Ref. [30] to be 0.5GeV^4 . These values are compatible with eq. (V.25) within a 20 percent accuracy.

VI. CONCLUSION

In this paper we have proposed an improved energy functional for variational calculations in gauge field theories. This functional contains a non local term which approximately corresponds to the energy gain when projecting on gauge invariant states. This allows one to use gaussian states as trial functionals and thus perform analytic calculations for physical observables such as the chromoelectric and chromomagnetic susceptibilities, energy expectation values and the gluon condensate.

The main purpose of this work was to check the ultraviolet behavior of our approximation scheme. By performing variational calculations near the perturbative vacuum we have shown

that divergences can be eliminated by a renormalization of the coupling constant. This has allowed us to recover the familiar one loop beta function in a way which makes transparent the various screening and antiscreening contributions. In particular the screening term arises naturally in our formalism, which was not the case in earlier variational approaches [15]. We have also tested our variational method in the magnetic sector, checking that it reproduces the one-loop result by Savvidy [20] for the background field energy. This calculation provides us with another derivation of the one-loop beta function.

Thus, our formalism appears to correctly reproduce the expected behavior of non-Abelian theories in the ultraviolet sector. This strongly encourages us to study its predictions in the non perturbative regime. Indeed, as a variational approach, it is not at all restricted to the vicinity of the perturbative ground state, nor to small values of the coupling constant. We would like to also emphasize that the Thouless-Valatin correction is the first step in an approximation scheme which can be constructed systematically. Indeed, it is the first order term [23] in boson expansion methods which have been constructed by Schwinger [31], Dyson, [32] Holstein-Primakoff [33], Blaizot-Marshalek [34].

An attractive feature of the variational picture is that it allows one to treat situations where instabilities occur. This is the case when the trial state is not the lowest one and where saddle points are reached. An example of such a situation is the occurrence of negative modes generally found in the presence of a constant magnetic background field [27,28]. In this case it would be interesting to work out what is the optimal gaussian kernel in our approach. This state should be reached by allowing the variational space to include the subspace spanned by the negative modes.

Another attractive question is the investigation of the infrared behaviour. It may provide some information on the confinement mechanism and the generation of mass scales [18]. Sum rules and the gluon condensate at finite temperature also appear to be a promising field of investigation.

Acknowledgements

We are most grateful to A. K. Kerman for valuable remarks, regarding in particular the

physical significance of the moment of inertia. Stimulating discussions with J. Polonyi and L. McLerran are also gratefully acknowledged.

APPENDIX A: PHYSICAL INTERPRETATION OF SECTION IV.B. IN TERMS OF THE INDUCED CHARGE

Let us show an other interpretation (and computation) of the relation between ρ^a and the external charge. What follows is directly inspired from Gottfried and Weisskopf in section IV.C of [25]. Let us assume that the system is in presence of a given distribution of external charges $\rho_{ext}^a(\mathbf{x}) = \nabla \cdot \mathbf{E}_{ext}^a(\mathbf{x})$ and compute the corresponding induced charge created by the quantum fluctuations of the gauge field ; it is given by the mean value $\langle -f_{abc}A_i^b(\mathbf{x})(E_L^i)^c(\mathbf{x}) \rangle$ where \mathbf{E}_L is the longitudinal part of the chromoelectric field operator \mathbf{E} [25]. This operator is a non-dynamical variable which is fixed by Gauss' law:

$$\nabla \cdot \mathbf{E}_L^a(\mathbf{x}) = \rho_{ext}^a(\mathbf{x}) - f_{abc}A_i^b(\mathbf{x})(E_L^i)^c(\mathbf{x}) \quad (\text{A.1})$$

which can be solved perturbatively, setting $\mathbf{E}_L^a(\mathbf{x}) = \mathbf{E}_L^{a(0)}(\mathbf{x}) + \mathbf{E}_L^{a(1)}(\mathbf{x}) + \mathbf{E}_L^{a(2)}(\mathbf{x}) + \dots$. In fact, only the first two terms will be needed for a development of the total charge in first order of g^2 . They verify the following set of equations :

$$\begin{aligned} \nabla \cdot \mathbf{E}_L^{(0)a}(\mathbf{x}) &= \rho_{ext}^a(\mathbf{x}) \\ \nabla \cdot \mathbf{E}_L^{(1)a}(\mathbf{x}) &= -f_{abc}A_i^b(\mathbf{x})(E_L^i)^{(0)c}(\mathbf{x}). \end{aligned} \quad (\text{A.2})$$

The first equation shows that $\mathbf{E}_L^{(0)}(\mathbf{x}) = \mathbf{E}_{ext}(\mathbf{x})$. Then the second equation is solved by:

$$(E_L^i)^{(1)c}(\mathbf{x}) = - \int \frac{d^3\mathbf{y}}{4\pi} \frac{x_i - y_i}{|\mathbf{x} - \mathbf{y}|^3} f_{cde}A_j^d(\mathbf{y})(E_{ext}^j)^e(\mathbf{y}) \quad (\text{A.3})$$

The total charge reads therefore :

$$\begin{aligned} \rho_{tot}^a(\mathbf{x}) &= \langle \mathcal{G}^a(\mathbf{x}) \rangle \\ &= \rho_{ext}^a(\mathbf{x}) + \langle -f_{abc}A_i^b(\mathbf{x})(E_{ext}^i)^c(\mathbf{x}) \rangle \\ &\quad + \langle -f_{abc}A_i^b(\mathbf{x})(E_L^i)^{(1)c}(\mathbf{x}) \rangle \\ &\equiv \rho_{ext}^a(\mathbf{x}) + \rho_{ind}^a(\mathbf{x}). \end{aligned} \quad (\text{A.4})$$

The first term in ρ_{ind}^a vanishes since linear in A_i , while the second term yields

$$\begin{aligned}\rho_{ind}^a(\mathbf{x}) &\equiv \langle -f_{abc}A_i^b(\mathbf{x})(E_L^i)^{(1)c}(\mathbf{x}) \rangle_c \\ &= f_{abc}f_{cde} \int \frac{d^3y}{4\pi} \frac{x_i - y_i}{|\mathbf{x} - \mathbf{y}|^3} G_{ij}^{bd}(\mathbf{x} - \mathbf{y})(E_{ext})_j^e(\mathbf{y}),\end{aligned}\tag{A.5}$$

or, after a Fourier transform,

$$\rho_{ind}^a(\mathbf{q}) = f_{abc}f_{cde}(E_{ext})_j^e(\mathbf{q}) \int \frac{d^3x}{4\pi} e^{-i\mathbf{q}\cdot\mathbf{x}} \frac{x_i}{|\mathbf{x}|^3} G_{ij}^{bd}(\mathbf{x}).\tag{A.6}$$

By also using

$$\int \frac{d^3x}{4\pi} e^{i\mathbf{k}\cdot\mathbf{x}} \frac{x_i}{|\mathbf{x}|^3} = i \frac{k_i}{\mathbf{k}^2},\tag{A.7}$$

we finally deduce:

$$\rho_{ind}^a(\mathbf{q}) = f_{abc}f_{cde}i(E_{ext})_j^e(\mathbf{q}) \int \frac{d^3k}{(2\pi)^3} \frac{(k - q)_i}{|\mathbf{k} + \mathbf{q}|^2} G_{ij}^{bd}(\mathbf{k}).\tag{A.8}$$

Since G is transverse this expression reduces to

$$\rho_{ind}^a(\mathbf{q}) = -f_{abc}f_{cde}i(E_{ext})_j^e(\mathbf{q}) \int \frac{d^3k}{(2\pi)^3} \frac{q_i}{|\mathbf{k} + \mathbf{q}|^2} v G_{ij}^{bd}(\mathbf{k}).\tag{A.9}$$

Furthermore for a smooth charge distribution we can approximate $\mathbf{k} + \mathbf{q}$ by \mathbf{k} in the above integral. Noting that the result vanishes unless $i=j$ we have

$$\rho_{ind}^a(\mathbf{q}) = \gamma \rho_{ext}^a(\mathbf{q}),\tag{A.10}$$

with

$$\gamma = g^2 C_N \frac{1}{6\pi^2} \int_0^\infty \frac{dk}{\varepsilon(\mathbf{k})} = \delta.\tag{A.11}$$

The total charge is then given by

$$\begin{aligned}\rho_{tot}^a(\mathbf{q}) &= \rho_{ext}^a(\mathbf{q}) + \rho_{ind}^a(\mathbf{q}) \\ &= \rho_{ext}^a(\mathbf{q})(1 + \delta),\end{aligned}\tag{A.12}$$

which is precisely the expression obtained in section IV.B (cf. eq. (IV.25)).

APPENDIX B: PROPER-TIME CALCULATION OF THE ENERGY DENSITY

Let us present here in some detail the calculation of the quantity $\text{Tr}G^{-1}$ which enters the energy of the magnetic field in Sec. V.C. According to the Schwinger proper-time representation (V.17), one needs the matrix element $\langle \mathbf{x} | e^{-tK} | \mathbf{x} \rangle$. Since, moreover, we are mainly interested in the ultraviolet behaviour of the energy, this expression is needed only at small values of t , which allows us to perform expansions in powers of t whenever necessary.

As explained in Sec. V.C, we shall use the background field in eq. (V.14) for which:

$$[\mathcal{D}_i, \mathcal{D}_j]^{ab} = -f^{ab3} \epsilon^{3ij} B = T^3 S_z B, \quad (\text{B.1})$$

and therefore $[\Pi_i, \Pi_j]^{ab} = -[\mathcal{D}_i, \mathcal{D}_j]^{ab} = -(T^3)_{ab} (S_z)_{ij} B$. It is convenient to define the following operator (cf. eq. (V.9)):

$$\tilde{K}_{ij} \equiv K_{ij} + \Pi_i \Pi_j = \Pi^2 \delta_{ij} + 2[\Pi_i, \Pi_j], \quad (\text{B.2})$$

in terms of which K_{ij} can be rewritten as follows :

$$K = \tilde{K} \hat{Q}. \quad (\text{B.3})$$

By also using $[\tilde{K}, \hat{Q}] = 0$, and $\hat{Q} + \hat{P} = 1$, we deduce

$$e^{-tK} = e^{-t\tilde{K}} \hat{Q} + \hat{P}. \quad (\text{B.4})$$

We thus have to compute the matrix element $\langle \mathbf{x} | e^{-t\tilde{K}} \hat{Q} | \mathbf{x} \rangle$, with :

$$\begin{aligned} \tilde{K} &= \Pi^2 - 2T^3 S_z B \\ \langle \mathbf{x} | e^{-t\tilde{K}} | \mathbf{x} \rangle &= \langle \mathbf{x} | e^{-t\Pi^2} e^{2tT^3 S_z B} | \mathbf{x} \rangle, \end{aligned} \quad (\text{B.5})$$

where the second line follows since $[\Pi^2, T^3 S_z] = 0$.

The computation of $\langle \mathbf{x} | e^{-t\Pi^2} | \mathbf{x} \rangle$ is well-known in the literature [35], with the result :

$$\langle \mathbf{x} | e^{-t\Pi^2} | \mathbf{x} \rangle = \left(\frac{1}{4\pi t} \right)^{3/2} \frac{tT^3 B}{\sinh(tT^3 B)} \quad (\text{B.6})$$

We thus obtain:

$$\langle \mathbf{x} | e^{-t\tilde{K}} | \mathbf{x} \rangle = \left(\frac{1}{4\pi t} \right)^{3/2} \frac{tT^3 B}{\sinh(tT^3 B)} e^{2tT^3 S_z B}. \quad (\text{B.7})$$

By also using $\hat{Q} = 1 - \hat{P}$ and $\Pi_i \Pi^2 = \tilde{K} \Pi_i$, one then rewrite $\langle \mathbf{x} | e^{-t\tilde{K}} \hat{Q} | \mathbf{x} \rangle$ as :

$$\langle \mathbf{x} | e^{-t\tilde{K}} | \mathbf{x} \rangle - \langle \mathbf{x} | e^{-t\tilde{K}} \Pi_i \Pi_j \frac{1}{\tilde{K}} | \mathbf{x} \rangle \quad (\text{B.8})$$

The last term of this equation can be obtained from $\langle \mathbf{x} | e^{-t\tilde{K}} \Pi_i \Pi_j | \mathbf{x} \rangle$ by integration over t .

To calculate $\langle \mathbf{x} | e^{-t\tilde{K}} \Pi_i \Pi_j | \mathbf{x} \rangle$, we follow Schwinger's method [36] : We work in Heisenberg's representation with $t = is$ and deduce

$$\begin{aligned} \langle \mathbf{x} | e^{-t\Pi^2} \Pi_i \Pi_j | \mathbf{x} \rangle &= \langle \mathbf{x} | e^{-is\Pi^2} \Pi_i \Pi_j | \mathbf{x} \rangle \\ &= \langle \mathbf{x}(s) | \Pi_i(0) \Pi_j(0) | \mathbf{x}(0) \rangle \end{aligned} \quad (\text{B.9})$$

where :

$$\begin{aligned} x_i(s) &= e^{is\Pi^2} x_i(0) e^{-is\Pi^2}, \\ \Pi(s) &= e^{is\Pi^2} \Pi(0) e^{-is\Pi^2} \end{aligned} \quad (\text{B.10})$$

The operator $U(s) = e^{-is\Pi^2}$ can be interpreted as the evolution operator of a particle governed by the Hamiltonian Π^2 . We have

$$\begin{aligned} \frac{dx_i}{ds} &= i[\Pi^2, x_i](s) \\ &= 2\Pi(s) \end{aligned} \quad (\text{B.11})$$

and

$$\frac{d\Pi}{ds} = i[\Pi^2, \Pi](s) \quad (\text{B.12})$$

Using $[\Pi^2, \Pi](s) = 2[\Pi_k, \Pi_i] \Pi^k = -2i\mathcal{F}_{ik} \Pi^k$, where

$$\mathcal{F}_{ij}^{ab} \equiv i[\mathcal{D}_i, \mathcal{D}_j]^{ab} = iT^3 S_z B \quad (\text{B.13})$$

one obtains $\Pi(s) = (e^{2s\mathcal{F}})_{ik} \Pi^k(0)$, and thus :

$$\begin{aligned} x_i(s) - x_i(0) &= \left(\frac{e^{2s\mathcal{F}} - 1}{\mathcal{F}} \right)_{ij} \Pi^j(0) \\ &\equiv R_{ij}^{-1} \Pi^j(0) \end{aligned} \quad (\text{B.14})$$

The matrix element $\langle \mathbf{x} | e^{-t\Pi^2} \Pi_i \Pi_j | \mathbf{x} \rangle$ can be now computed as (cf. eq. (B.9)):

$$\begin{aligned}
& \langle \mathbf{x}(s) | \Pi_i(0) \Pi_j(0) | \mathbf{x}(0) \rangle \\
&= \langle \mathbf{x}(s) | R_{im} R_{jn} (x_m(s) - x_m(0))(x_n(s) - x_n(0)) | \mathbf{x}(0) \rangle \\
&= R_{im} R_{jn} \langle \mathbf{x}(s) | x_m(s)x_n(s) - x_m(s)x_n(0) \\
&\quad - x_n(s)x_m(0) + x_m(0)x_n(0) + [x_n(s), x_m(0)] | \mathbf{x}(0) \rangle \\
&= R_{im} R_{jn} R_{nk}^{-1} [\Pi_k(0), x_m(0)] \langle \mathbf{x}(s) | \mathbf{x}(0) \rangle \\
&= -i R_{ij} \langle \mathbf{x}(s) | \mathbf{x}(0) \rangle
\end{aligned} \tag{B.15}$$

Returning to the variable $s = -it$, we obtain

$$\langle \mathbf{x} | e^{-t\Pi^2} \Pi_i \Pi_j | \mathbf{x} \rangle = \left(\frac{-i\mathcal{F}}{e^{-2it\mathcal{F}} - 1} \right)_{ij} \langle \mathbf{x} | e^{-t\Pi^2} | \mathbf{x} \rangle, \tag{B.16}$$

and thus, finally,

$$\begin{aligned}
& \langle \mathbf{x} | e^{-t\tilde{K}} \Pi_i \Pi_j | \mathbf{x} \rangle \\
&= \frac{T^3 S_z B}{2 \sinh(tT^3 S_z B)} \left(\frac{1}{4\pi t} \right)^{3/2} \frac{tT^3 B}{\sinh(tT^3 B)} e^{tT^3 S_z B} \\
&= \frac{1}{2t} \left(\frac{1}{4\pi t} \right)^{3/2} \left\{ 1 + T^3 B S_z t \right. \\
&\quad \left. + \left(\frac{(T^3)^2 B^2 S_z^2}{3} - \frac{(T^3)^2 B^2}{6} \right) t^2 + \dots \right\}
\end{aligned} \tag{B.17}$$

where in writing the second line we have developed up to the second order in t .

The last equation can be now integrated over t to obtain the second term in (B.8):

$$\begin{aligned}
\langle \mathbf{x} | e^{-t\tilde{K}} \hat{P} | \mathbf{x} \rangle &= \left(\frac{1}{4\pi t} \right)^{3/2} \left\{ \frac{1}{3} + T^3 B S_z t \right. \\
&\quad \left. + \left(\frac{(T^3)^2 B^2}{6} - \frac{(T^3)^2 B^2 S_z^2}{3} \right) t^2 + \dots \right\}
\end{aligned} \tag{B.18}$$

Developing $\langle \mathbf{x} | e^{-t\tilde{K}} | \mathbf{x} \rangle$ up to the second order in t :

$$\begin{aligned}
\langle \mathbf{x} | e^{-t\tilde{K}} | \mathbf{x} \rangle &= \left(\frac{1}{4\pi t} \right)^{3/2} \left\{ 1 + 2T^3 B S_z t \right. \\
&\quad \left. + (2(T^3)^2 B^2 S_z^2 - \frac{(T^3)^2 B^2}{6}) t^2 + \dots \right\},
\end{aligned} \tag{B.19}$$

we deduce the following expression for $U(t) \equiv \langle \mathbf{x} | e^{-t\hat{K}} \hat{Q} | \mathbf{x} \rangle$:

$$U(t) = \left(\frac{1}{4\pi t}\right)^{3/2} \left\{ \frac{2}{3} + T^3 B S_z t \right. \\ \left. + \left(\frac{7}{3}(T^3)^2 B^2 S_z^2 - \frac{1}{3}(T^3)^2 B^2\right) t^2 + \dots \right\}. \quad (\text{B.20})$$

Thus, finally,

$$\text{Tr} \langle \mathbf{x} | G^{-1} | \mathbf{x} \rangle = \frac{1}{\sqrt{\pi}} \int_{1/\Lambda^2}^{\infty} \frac{dt}{t^{3/2}} (\text{Tr} \langle \mathbf{x} | \hat{Q} | \mathbf{x} \rangle - \text{Tr} U(t)) \\ = (\dots) \Lambda^4 - \frac{C_N}{8\pi^2} \left(\frac{11}{3} B^2\right) \ln \frac{\Lambda^2}{B} + \mathcal{O}(g^2), \quad (\text{B.21})$$

where $\langle \mathbf{x} | \hat{Q} | \mathbf{x} \rangle$ has been obtained by setting $t = 0$ in eq. (B.20). This is the result announced in eq. (V.18).

REFERENCES

- [1] R. Jackiw, *Analysis of infinite-dimensional manifolds - Schrödinger representation for quantized fields*, Field Theory and Particle Physics - World Scientific, Singapore, 1990.
- [2] L.I. Schiff, Phys. Rev. **130** (1963) 458; G. Rosen Phys. Rev. **173** (1968) 1632.
- [3] J. Kuti, in J. Cornwall, R. Jackiw, E. Tomboulis, Phys. Rev. **D10** (1974) 2428.
- [4] T. Barnes, G. I. Ghandour, Phys. Rev. **D22** (1980) 924.
- [5] W. A. Bardeen, M. Moshe, Phys. Rev. **28** (1983) 1372.
- [6] F. Cooper, E. Mottola, Phys. Rev. **36** (1987) 3114.
- [7] S. Y. Pi, M. Samiullah, Phys. Rev. **36** (1987) 3128.
- [8] O. Eboli, R. Jackiw, S. Y. Pi, Phys. Rev. **D37** (1988) 3557.
- [9] J. Greensite, Nucl. Phys **158** (1979) 469.
- [10] J. Cornwall, Phys. Rev. **D38** (1988) 656.
- [11] M. Danos, D. Gogny and D. Iracane, *Models for the non perturbative QCD vacuum*, US Department of Commerce, National Institute of Standards and Technology, NISTIR 4555, April 1991.
- [12] A. K. Kerman and D. Vautherin, Ann. Phys. **192** (1989) 408.
- [13] R.P. Feynman, in *Variational Calculations in Quantum Field Theory*, Proceedings of the Workshop, Wangerooge, West Germany, 1987, edited by L. Polley and D. Pottinger (World Scientific, Singapore, 1988)
- [14] I. I. Kogan and A. Kovner, Phys. Rev. **D52** (1995) 3719.
- [15] W.E. Brown and I. I. Kogan, Int. J. Mod. Phys. **A14** (1999) 799.
- [16] W.E. Brown, Int. J. Mod. Phys. **A13** (1998) 5219.

- [17] K. Zarembo, Phys. Lett. **B421** (1998) 325 ; *ibid.* Mod.Phys.Lett. A13 (1998) 2317.
- [18] D. Diakonov, Lecture at the 4-th St.Petersburg Winter School in Theoretical Physics, Feb. 22-28, 1998, hep-th/9805137.
- [19] D. J. Thouless and J. G. Valatin, Nucl. Phys. **31** (1962) 211.
- [20] G. K. Savvidy, Phys. Lett. **B71** (1977) 133.
- [21] F. M. H. Villars, Proceedings of the International School of Physics “Enrico Fermi”, Varenna, 1966.
- [22] R. E. Peierls and J. Yoccoz, Proc. Phys. Soc. (London) **A70** (1957) 381.
- [23] P. Ring and P. Schuck, *The nuclear many-body problem*, Springer, Berlin, 1981.
- [24] C. Heinemann, Ph.D.Thesis, Université Pierre et Marie Curie, 2000, unpublished.
- [25] K. Gottfried and V. F. Weisskopf, *Concepts of Particle Physics*, Volume II, Oxford University Press, Oxford, 1986.
- [26] T. D. Lee, *Particle Physics and Introduction to Field Theory*, Harwood, Chur, 1981, Sec. 18.5.
- [27] N. K. Nielsen and P. Olesen, Nucl. Phys. **B144** (1978) 376.
- [28] H. B. Nielsen and P. Olesen, Nucl. Phys. **B160** (1979) 380.
- [29] A. Chodos, R. L. Jaffee, K. Johnson, C. B. Thorn and V. F. Weisskopf, Phys. Rev. **D9** (1974) 3471
- [30] M. Shifman, A. Vainshtein and V. I. Zakharov, Nucl. Phys. **B147** (1979) 385
- [31] J. Schwinger, *Quantum Theory of angular momentum*, L. Biedenharn and H. Van Dam, eds., Academic Press, New York, 1965.
- [32] J.F. Dyson, Phys. Rev. **102** (1956) 1217.

- [33] T. Holstein and H. Primakoff, Phys. Rev. **58** (1940) 1098.
- [34] J.P. Blaizot and E.R. Marshalek, Nucl.Phys. **A309** (1978) 422 ; Phys.Lett **79B** (1978) 1.
- [35] E. Brézin and C. Itzykson, Phys. Rev. **D2** (1970) 1191.
- [36] J. Schwinger, Phys. Rev. **82** (1951) 664.