

Supersymmetric Electroweak Corrections to $W^\pm H^\mp$ Associated Production at the CERN Large Hadron Collider

Ya Sheng Yang ^a, Chong Sheng Li ^a, Li Gang Jin ^a, and Shou Hua Zhu ^{b,c}

^a Department of Physics, Peking University, Beijing 100871, China

^b CCAST(World Laboratory), Beijing 100080, China

^c Institute of Theoretical Physics, Academia Sinica, Beijing 100080, China

ABSTRACT

The $O(\alpha_{ew}m_{t(b)}^2/m_W^2)$ and $O(\alpha_{ew}m_{t(b)}^4/m_W^4)$ supersymmetric electroweak corrections to the cross section for $W^\pm H^\mp$ associated production at the LHC are calculated in the minimal supersymmetric standard model. Those corrections arise from the quantum effects which are induced by the Yukawa couplings from the Higgs sector and the chargino-top(bottom)-sbottom(stop) couplings, neutralino-top(bottom)-stop(sbottom) couplings and charged Higgs-stop-sbottom couplings. The numerical results show that the Yukawa corrections arising from the Higgs sector can decrease the total cross sections significantly for low $\tan\beta$ ($= 1.5$ and 2) when $m_{H^\pm} (< 300)\text{GeV}$, which exceed -12% . For high $\tan\beta$ the Yukawa corrections become negligibly small. The genuine supersymmetric electroweak corrections can increase or decrease the total cross sections depending on the supersymmetric parameters, which can exceed -25% for the favorable supersymmetric parameter values. We also show that the genuine supersymmetric electroweak corrections depend strongly on the choice of $\tan\beta$, A_t , $M_{\tilde{Q}}$ and μ . For large values of A_t , or large values of μ and $\tan\beta$, one can get much larger corrections. The corrections can become very small, in contrast, for larger values of $M_{\tilde{Q}}$.

PACS number: 12.60.Jv, 12.15.Lk, 14.80.Cp, 14.70.Fm

1. Introduction

One of the most important objectives of the CERN Large Hadron Collider (LHC) is the search for Higgs boson. In various extensions of the Higgs sector of the standard model(SM), for example, in the two-Higgs-doublet models(THDM)[1], particularly the minimal supersymmetric standard model(MSSM)[2], there are physical charged Higgs bosons, which do not belong to the spectrum of the SM and therefore their discovery would be instant evidence of new physics. In much of the parameter space preferred by the MSSM, namely $m_{H^\pm} > m_W$ and $1 < \tan\beta < m_t/m_b$ [3,4], the LHC will provide the greatest opportunity for the discovery of charged Higgs boson. Previous studies have shown that for a relatively light charged Higgs boson, $m_{H^\pm} < m_t - m_b$, the dominate production processes at the LHC are $gg \rightarrow t\bar{t}$ and $q\bar{q} \rightarrow t\bar{t}$ followed by the decay sequence $t \rightarrow bH^+ \rightarrow b\tau^+\nu_\tau$ [5], and for a heavier charged Higgs boson the dominate production process is $gb \rightarrow tH^-$ [6,7,8]. Besides the processes mentioned above, in Ref.[9] Dicus et al. also studied the production of a charged Higgs boson in association with a W boson via $b\bar{b}$ annihilation at the tree level and gg fusion at one loop at hadron colliders. Since the leptonic decays of W boson would serve as a spectacular trigger for the charged Higgs boson search, these processes seem attractive. But the authors of Ref.[9] only considered the case where the value of $\tan\beta$ to be in the range $0.3 - 2.3$. Recently Barrientos Bendezu and Kniehl[10] further studied these processes and presented theoretical predictions for the $W^\pm H^\mp$ production cross section at the LHC and Tevatron's Run II, where they generalize the analysis of Ref.[9] for arbitrary values of $\tan\beta$ and to update it. They found that the $W^\pm H^\mp$ production would have a sizeable cross section and its signal should have a significant rate at the LHC unless m_{H^\mp} is very large.

As analyzed in Ref.[7,11], the search for heavy charged Higgs bosons with $m_{H^+} > m_t + m_b$ at a hadron collider is seriously complicated by QCD backgrounds. For example, the processes suggested in Ref.[10] suffer from the irreducible background due to top quark pair production, $q\bar{q} \rightarrow t\bar{t}$ and $gg \rightarrow t\bar{t}$ with subsequent decay through the intermediate state $b\bar{b}W^+W^-$, and heavy charged Higgs boson produced

in association with W^\pm gauge bosons cannot be resolved at the LHC, via semileptonic W^+W^- decays, for charged Higgs boson masses in the range between $2m_t$ and 600GeV at neither low nor high $\tan\beta$ [11]. However, recent analyses[12,13] have shown that the decay mode $H^+ \rightarrow \tau^+\nu$, indeed dominant for light charged Higgs bosons below the top threshold for any accessible $\tan\beta$ [14], provides an excellent signature for a heavy charged Higgs boson in searches at the LHC. The discover region for H^\pm is far greater than had been thought for a large range of the $(m_{H^\pm}, \tan\beta)$ parameter space, extending beyond $m_{H^\pm} \sim 1\text{TeV}$ and down to at least $\tan\beta \sim 3$, and potentially to $\tan\beta \sim 1.5$, assuming the latest results for the SM parameters and parton distribution functions as well as using kinematic selection techniques and the tau polarization analysis[13]. Of course, it is just a theoretical analysis and no experimental simulation has been performed to make the statement very reliable so far.

Since the contributions to the $W^\pm H^\mp$ production cross section due to $b\bar{b}$ annihilation at the tree level are greater than ones due to gg fusion which proceeds at one-loop, it is important to calculate the one-loop radiative corrections to the $W^\pm H^\mp$ production via $b\bar{b}$ annihilation for more accurate theoretical predictions for the cross sections. In this paper we present the calculations of the $O(\alpha_{ew}m_{t(b)}^2/m_W^2)$ and $O(\alpha_{ew}m_{t(b)}^4/m_W^4)$ supersymmetric(SUSY) electroweak(EW) corrections to this $W^\pm H^\mp$ associated production process at the LHC in the MSSM. These corrections arise from the quantum effects which are induced by potentially large Yukawa couplings from the Higgs sector and the chargino-top(bottom)-sbottom(stop) couplings, neutralino- top(bottom)-stop(sbottom) couplings and charged Higgs-stop-sbottom couplings which will contribute at the $O(\alpha_{ew}m_{t(b)}^4/m_W^4)$ to the self-energy of the charged Higgs boson. The relevant QCD corrections are expected to be larger, but not yet available.

The arrangement of this paper is as follows. In Sec.II we give the analytic results. In Sec.III we present some numerical examples and discuss the implications of our results. Some notations used in this paper and the lengthy expressions of the form factors are summarized in Appendix A, B.

2. Calculations

The Feynman diagrams for the charged Higgs boson production via $b(p_1)\bar{b}(p_2) \rightarrow W^\pm(k)H^\mp(p_3)$, which include the SUSY EW corrections to the process, are shown in Fig.1 and Fig.2. We carried out the calculation in the t'Hooft-Feynman gauge and used dimensional reduction, which preserves supersymmetry, for regularization of the ultraviolet divergences in the virtual loop corrections using the on-mass-shell renormalization scheme[15], in which the fine-structure constant α_{ew} and physical masses are chosen to be the renormalized parameters, and finite parts of the counterterms are fixed by the renormalization conditions. The coupling constant g is related to the input parameters e , m_W , and m_Z via $g^2 = e^2/s_w^2$ and $s_w^2 = 1 - m_W^2/m_Z^2$. As far as the parameters β and α , for the MSSM we are considering, they have to be renormalized, too. In the MSSM they are not independent. Nevertheless, we follow the approach of Mendez and Pomarol[16] in which they consider them as independent renormalized parameters and fixed the corresponding renormalization constants by a renormalization condition that the on-mass-shell $H^+\bar{l}\nu_l$ and $h\bar{l}l$ couplings keep the forms of Eq.(3) of Ref.[16] to all order of perturbation theory.

We define the Mandelstam variables as

$$\begin{aligned}\hat{s} &= (p_1 + p_2)^2 = (k + p_3)^2, \\ \hat{t} &= (p_1 - k)^2 = (p_2 - p_3)^2, \\ \hat{u} &= (p_1 - p_3)^2 = (p_2 - k)^2.\end{aligned}\tag{1}$$

The relevant renormalization constants are defined as

$$\begin{aligned}m_{W_0}^2 &= m_W^2 + \delta m_W^2, & m_{Z_0}^2 &= m_Z^2 + \delta m_Z^2, \\ \tan \beta_0 &= (1 + \delta Z_\beta) \tan \beta, \\ \sin \alpha_0 &= (1 + \delta Z_\alpha) \sin \alpha, \\ W_0^{\pm\mu} &= (1 + \delta Z_W)^{1/2} W^{\pm\mu} + i Z_{H^\pm W^\pm}^{1/2} \partial^\mu H^\mp, \\ H_0^\pm &= (1 + \delta Z_{H^\pm})^{1/2} H^\pm, \\ Z_0^\mu &= (1 + \delta Z_Z)^{1/2} Z^\mu + i Z_{Z A}^{1/2} \partial^\mu A, \\ A_0 &= (1 + \delta Z_A)^{1/2} A,\end{aligned}$$

$$\begin{aligned}
H_0 &= (1 + \delta Z_H)^{1/2} H + Z_{Hh}^{1/2} h, \\
h_0 &= (1 + \delta Z_h)^{1/2} h + Z_{hH}^{1/2} H.
\end{aligned} \tag{2}$$

Taking into account the $O(\alpha_{ew} m_{t(b)}^2/m_W^2)$ and $O(\alpha_{ew} m_{t(b)}^4/m_W^4)$ SUSY EW corrections, the renormalized amplitude for $b\bar{b} \rightarrow W^- H^+$ can be written as

$$\begin{aligned}
M_{ren} &= M_0^{(s)} + M_0^{(t)} + [\delta \hat{M}^{V_1(s)} + \delta \hat{M}^{S(s)} + \delta \hat{M}^{V_2(s)}](H_i) + [\delta \hat{M}^{V_1(s)} \\
&\quad + \delta \hat{M}^{S(s)} + \delta \hat{M}^{V_2(s)}](A) + \delta \hat{M}^{V_1(t)} + \delta \hat{M}^{S(t)} + \delta \hat{M}^{V_2(t)} + \delta M^{box}, \tag{3}
\end{aligned}$$

where $M_0^{(s)}$ and $M_0^{(t)}$ are the tree-level amplitudes arising from Fig.1(a) and Fig.1(b), respectively, which are given by

$$M_0^{(s)} = -i \sum_i \frac{gh_b \alpha_{2i} \varphi_{11}}{\sqrt{2}(\hat{s} - m_{H_i}^2)} \sum_{j=1}^4 M_j + \frac{igh_b \beta_{12}}{\sqrt{2}(\hat{s} - m_A^2)} (M_1 - M_2 + M_3 - M_4) \tag{4}$$

and

$$M_0^{(t)} = \frac{ig}{\sqrt{2}(\hat{t} - m_t^2)} (2h_b \beta_{12} M_2 - h_b m_b \beta_{12} M_5 + h_t m_t \beta_{11} M_6 - h_b \beta_{12} M_{12}). \tag{5}$$

Here $h_b \equiv gm_b/\sqrt{2}m_W \cos \beta$ and $h_t \equiv gm_t/\sqrt{2}m_W \sin \beta$ are the Yukawa couplings from the bottom and top quarks, p_1 and p_2 denote the momentum of incoming quarks b and \bar{b} , respectively, while k and p_3 are used for the outgoing W^- Boson and H^+ Boson, respectively. The notations α_{ij} , β_{ij} and φ_{ij} used in the above expressions are defined in Appendix A, and H_i stands for Higgs Bosons h with $i = 1$ and H with $i = 2$. M_i are the standard matrix elements, which are defined by

$$\begin{aligned}
M_1 &= \bar{v}(p_2) P_R u(p_1) p_1 \cdot \varepsilon(k), \\
M_2 &= \bar{v}(p_2) P_L u(p_1) p_1 \cdot \varepsilon(k), \\
M_3 &= \bar{v}(p_2) P_R u(p_1) p_2 \cdot \varepsilon(k), \\
M_4 &= \bar{v}(p_2) P_L u(p_1) p_2 \cdot \varepsilon(k), \\
M_5 &= \bar{v}(p_2) \not{\varepsilon}(k) P_R u(p_1), \\
M_6 &= \bar{v}(p_2) \not{\varepsilon}(k) P_L u(p_1), \\
M_7 &= \bar{v}(p_2) \not{\varepsilon}(k) P_R u(p_1) p_1 \cdot \varepsilon(k),
\end{aligned}$$

$$\begin{aligned}
M_8 &= \bar{v}(p_2) \not{\epsilon}(k) P_L u(p_1) p_1 \cdot \epsilon(k), \\
M_9 &= \bar{v}(p_2) \not{\epsilon}(k) P_R u(p_1) p_2 \cdot \epsilon(k), \\
M_{10} &= \bar{v}(p_2) \not{\epsilon}(k) P_L u(p_1) p_2 \cdot \epsilon(k), \\
M_{11} &= \bar{v}(p_2) \not{k} \not{\epsilon}(k) P_R u(p_1), \\
M_{12} &= \bar{v}(p_2) \not{k} \not{\epsilon}(k) P_L u(p_1),
\end{aligned} \tag{6}$$

where $P_{L,R} \equiv (1 \mp \gamma_5)/2$. The vertex and self-energy corrections to the tree-level process are included in $\delta\hat{M}^{V,S}$, which are given by

$$\begin{aligned}
\delta\hat{M}^{V_1(s)}(H_i) &= -\frac{igh_b}{\sqrt{2}} \left\{ \sum_{i=1,2} \frac{\alpha_{2i}\varphi_{i1}}{\hat{s} - m_{H_i}^2} \left[\frac{\delta h_b}{h_b} + \frac{1}{2}\delta Z_L^b + \frac{1}{2}\delta Z_R^b + \frac{1}{2}\delta Z_{H_i} \right] \right. \\
&\quad + \frac{\sin(\beta - \alpha) \sin \alpha}{\hat{s} - m_H^2} (\tan \alpha \delta Z_\alpha + Z_{hH}^{1/2}) - \frac{\cos(\beta - \alpha)}{\hat{s} - m_h^2} (\sin \alpha \delta Z_\alpha \\
&\quad \left. - \cos \alpha Z_{Hh}^{1/2}) \right\} \sum_{j=1}^4 M_j + \delta M^{V_1(s)}(H), \\
\delta\hat{M}^{V_1(s)}(A) &= -\frac{igh_b \sin \beta}{\sqrt{2}(\hat{s} - m_A^2)} \left[\frac{\delta h_b}{h_b} + \cos^2 \beta \delta Z_\beta + \frac{1}{2}\delta Z_L^b + \frac{1}{2}\delta Z_R^b + \frac{1}{2}\delta Z_A \right] \\
&\quad + \frac{im_W}{\tan \beta \cos \theta_W} Z_{hH}^{1/2} (M_1 - M_2 + M_3 - M_4) + \delta M^{V_1(s)}(A), \\
\delta\hat{M}^{S(s)}(H_i) &= \frac{igh_b}{\sqrt{2}} \sum_{i=1,2} \frac{\alpha_{2i}\varphi_{i1}}{(\hat{s} - m_{H_i}^2)^2} [\delta m_{H_i}^2 - (\hat{s} - m_{H_i}^2) \delta Z_{H_i} - (\hat{s} \\
&\quad - m_H^2) Z_{Hh}^{1/2} - (\hat{s} - m_h^2) Z_{hH}^{1/2}] \sum_{j=1}^4 M_j + \delta M^{S(s)}(H), \\
\delta\hat{M}^{S(s)}(A) &= \frac{igh_b \sin \beta}{\sqrt{2}(\hat{s} - m_A^2)} [\delta m_A^2 - (\hat{s} - m_A^2) \delta Z_A] (M_1 - M_2 + M_3 - M_4) \\
&\quad + \delta M^{S(s)}(A), \\
\delta\hat{M}^{V_2(s)}(H_i) &= -\frac{igh_b}{\sqrt{2}} \left\{ \sum_{i=1,2} \frac{\alpha_{2i}\varphi_{i1}}{\hat{s} - m_{H_i}^2} \left(\frac{\delta g}{g} + \frac{1}{2}\delta Z_{W^-} + \frac{1}{2}\delta Z_{H^+} + \frac{1}{2}Z_{H_i} \right) \right. \\
&\quad - \frac{\cos \alpha \cos(\beta - \alpha)}{\hat{s} - m_H^2} (\sin \beta \cos \beta \delta Z_\beta - \tan \alpha \delta Z_\alpha - Z_{hH}^{1/2}) \\
&\quad + m_W Z_{HW}^{1/2} + \frac{\sin \alpha \sin(\beta - \alpha)}{\hat{s} - m_h^2} (\sin \beta \cos \beta \delta Z_\beta \\
&\quad \left. - \tan \alpha \delta Z_\alpha + Z_{Hh}^{1/2} + m_W Z_{HW}^{1/2}) \right\} \sum_{j=1}^4 M_j + \delta M^{V_2(s)}(H), \\
\delta\hat{M}^{V_2(s)}(A) &= -\frac{igh_b \sin \beta}{\sqrt{2}(\hat{s} - m_A^2)} \left[\frac{\delta g}{g} + \frac{1}{2}\delta Z^A + \frac{1}{2}\delta Z_{H^+} \right]
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2} \delta Z_{W^-}] (M_1 - M_2 + M_3 - M_4) + \delta M^{V_2^{(s)}}(A), \\
\delta \hat{M}^{V_1^{(t)}} &= \frac{ig}{\sqrt{2}(\hat{t} - m_t^2)} (2h_b \beta_{12} M_2 - h_b m_b \beta_{12} M_5 + h_t m_t \beta_{11} M_6 \\
& - h_b \beta_{12} M_{12}) \left(\frac{\delta g}{g} + \frac{1}{2} \delta Z_L^t + \frac{1}{2} \delta Z_L^b + \frac{1}{2} \delta Z_{W^-} \right) + \delta M^{V_1^{(t)}}, \\
\delta \hat{M}^{S^{(t)}} &= \frac{ig}{\sqrt{2}(\hat{t} - m_t^2)^2} \left[(2m_t^2 \frac{\delta m_t}{m_t} + m_t^2 \delta Z_L^t - \hat{t} \delta Z_L^t) (2h_b \beta_{12} M_2 \right. \\
& - h_b m_b \beta_{12} M_5 - h_t \beta_{12} M_{12} + \frac{1}{2} h_t m_t \beta_{11} M_6) + \frac{1}{2} (2\hat{t} \frac{\delta m_t}{m_t} \\
& + m_t^2 \delta Z_R^t - \hat{t} \delta Z_R^t) h_t m_t \beta_{11} M_6 \left. \right] + \delta M^{S^{(t)}}, \\
\delta \hat{M}^{V_2^{(t)}} &= \frac{ig^2}{2m_W(\hat{t} - m_t^2)} \left[m_t^2 \cot \beta \left(\frac{\delta h_t}{h_t} - \cos^2 \beta \delta Z_\beta + \frac{1}{2} \delta Z_L^b + \frac{1}{2} \delta Z_R^t \right. \right. \\
& + \frac{1}{2} \delta Z_{H^+} + \frac{m_W}{\cot \beta} Z_{HW}^{1/2} \left. \right) M_6 + m_b \tan \beta \left(\frac{\delta h_b}{h_b} + \sin^2 \beta \delta Z_\beta + \frac{1}{2} \delta Z_L^t \right. \\
& \left. \left. + \frac{1}{2} \delta Z_R^b + \frac{1}{2} \delta Z_{H^+} - \frac{m_W}{\tan \beta} Z_{HW}^{1/2} \right) (2M_2 - M_{12} - m_b M_5) \right] + \delta M^{V_2^{(t)}}, \quad (7)
\end{aligned}$$

with

$$\begin{aligned}
\frac{\delta g}{g} &= \frac{\delta e}{e} + \frac{1}{2} \frac{\delta m_Z^2}{m_Z^2} - \frac{1}{2} \frac{\delta m_Z^2 - \delta m_W^2}{m_Z^2 - m_W^2}, \\
\frac{\delta h_b}{h_b} &= \frac{\delta g}{g} + \frac{\delta m_b}{m_b} - \frac{1}{2} \frac{\delta m_W^2}{m_W^2} + \cos^2 \beta \delta Z_\beta, \\
\frac{\delta h_t}{h_t} &= \frac{\delta g}{g} + \frac{\delta m_t}{m_t} - \frac{1}{2} \frac{\delta m_W^2}{m_W^2} - \sin^2 \beta \delta Z_\beta, \\
\delta Z_\beta &= -\frac{\delta g}{g} + \frac{1}{2} \frac{\delta m_W^2}{m_W^2} - \frac{1}{2} \delta Z_{H^+} - \frac{m_W}{\tan \beta} Z_{HW}^{1/2}, \\
\delta Z_\alpha &= -\frac{\delta g}{g} + \frac{1}{2} \frac{\delta m_W^2}{m_W^2} - \frac{1}{2} \delta Z_h - \cot \alpha Z_{Hh}^{1/2} - \sin^2 \beta \delta Z_\beta. \quad (8)
\end{aligned}$$

The $\delta e/e$ appearing in Eq.(8) does not contain the $O(\alpha_{ew} m_{t(b)}^2/m_W^2)$ corrections and needs not be considered in our calculations. And $\delta M^{V_1^{(s)}}(H_i)$, $\delta M^{V_1^{(s)}}(A)$, $\delta M^{S^{(s)}}(H_i)$, $\delta M^{S^{(s)}}(A)$, $\delta M^{V_2^{(s)}}(H_i)$, $\delta M^{V_2^{(s)}}(A)$, $\delta M^{V_1^{(t)}}$, $\delta M^{S^{(t)}}$, $\delta M^{V_2^{(t)}}$ and δM^{box} represent the irreducible corrections arising, respectively, from the $b\bar{b}H(h)$ vertex diagrams shown in Fig.1(c)–1(d), the $b\bar{b}A$ vertex diagrams shown in Fig.1(c)–1(d), the H and h boson self-energy diagrams in Fig.1(i)–1(k), the A boson self-energy diagrams shown in Fig.1(i)–1(k), the $H(h)W^-H^+$ vertex diagrams shown in Fig.1(f)–1(h), the AW^-H^+ vertex diagrams shown in Fig.1(f)–1(h), the btW^- vertex diagrams

Fig.1(l) – 1(o), the top quark self-energy diagrams Fig.1(r), the $t\bar{b}H^+$ vertex diagrams Fig.1(p) – 1(q), and the box diagrams Fig.1(s) – 1(x). All above $\delta M^{V,S}$ and δM^{box} can be written in the form

$$\delta M^{V,S,box} = i \sum_{i=1}^{12} f_i^{V,S,box} M_i, \quad (9)$$

where the $f_i^{V,S,box}$ are form factors, which are given explicitly in Appendix B.

Calculating the self-energy diagrams in Fig.2, we can get the explicit expressions of all the renormalization constants as following:

$$\begin{aligned} \frac{\delta m_t}{m_t} &= \sum_i \frac{-h_t^2}{32\pi^2} [\alpha_{1i}^2 (-B_0^{ttH_i} + B_1^{ttH_i}) + \beta_{1i}^2 (B_0^{ttA_i} + B_1^{ttA_i})] \\ &\quad - \sum_i \frac{1}{32\pi^2 m_t} [(h_t^2 m_t \beta_{1i}^2 + h_b^2 m_b \beta_{2i}^2) B_1^{tbH_i^+} + 2h_b h_t \beta_{1i} \beta_{2i} B_0^{tbH_i^+}] \\ &\quad + \sum_{i,j} \frac{h_t^2}{32\pi^2 m_t} [m_t |N_{j4}|^2 (B_0^{t\bar{t}_i \bar{\chi}_j^0} + B_1^{t\bar{t}_i \bar{\chi}_j^0}) + m_{\bar{\chi}_j^0} \theta_{i1}^t \theta_{i2}^t (N_{j4}^2 + N_{j4}^{*2}) B_0^{t\bar{t}_i \bar{\chi}_j^0}] \\ &\quad + \sum_{i,j} \frac{1}{32\pi^2 m_t} \{m_t [h_t^2 (\theta_{i1}^b)^2 |V_{j1}|^2 + h_b^2 (\theta_{i2}^b)^2 |U_{j2}|^2] (B_0^{t\bar{t}_i \bar{\chi}_j^+} + B_1^{t\bar{t}_i \bar{\chi}_j^+}) \\ &\quad + h_b h_t m_{\bar{\chi}_j^+} \theta_{i1}^b \theta_{i2}^b (U_{j2} V_{j2} + U_{j2}^* V_{j2}^*) B_0^{t\bar{t}_i \bar{\chi}_j^+} \}, \\ \delta Z_L^t &= \sum_i \frac{h_b^2 \beta_{2i}^2}{16\pi^2} B_1^{tbH_i^+} - \sum_{i,j} \frac{h_t^2 (\theta_{i2}^t)^2}{16\pi^2} |N_{j4}|^2 (B_0^{t\bar{t}_i \bar{\chi}_j^0} + B_1^{t\bar{t}_i \bar{\chi}_j^0}) \\ &\quad - \sum_{i,j} \frac{h_b^2 (\theta_{i2}^b)^2}{16\pi^2} |U_{j2}|^2 (B_0^{t\bar{t}_i \bar{\chi}_j^+} + B_1^{t\bar{t}_i \bar{\chi}_j^+}) + \delta^t, \\ \delta Z_R^t &= \sum_i \frac{h_t^2 \beta_{1i}^2}{16\pi^2} B_1^{tbH_i^+} - \sum_{i,j} \frac{h_t^2 (\theta_{i1}^t)^2}{16\pi^2} |N_{j4}|^2 (B_0^{t\bar{t}_i \bar{\chi}_j^0} + B_1^{t\bar{t}_i \bar{\chi}_j^0}) \\ &\quad - \sum_{i,j} \frac{h_t^2 (\theta_{i1}^b)^2}{16\pi^2} |V_{j2}|^2 (B_0^{t\bar{t}_i \bar{\chi}_j^+} + B_1^{t\bar{t}_i \bar{\chi}_j^+}) + \delta^t, \\ \delta^t &= \sum_i \frac{h_t^2}{32\pi^2} \{ \alpha_{1i}^2 [B_1^{ttH_i} - 2m_t^2 (B_0^{ttH_i} - B_1^{ttH_i})] + \beta_{1i}^2 [B_1^{ttA_i} + 2m_t^2 (B_0^{ttA_i} + B_1^{ttA_i})] \} \\ &\quad + \sum_i \frac{m_t}{16\pi^2} [m_t (h_t^2 \beta_{1i}^2 + h_b^2 \beta_{2i}^2) B_0'^{tbH_i^+} + 2h_b h_t m_b \beta_{1i} \beta_{2i} B_0'^{tbH_i^+}] \\ &\quad - \sum_{i,j} \frac{h_t^2 m_t}{16\pi^2} [m_t |N_{j4}|^2 (B_0'^{t\bar{t}_i \bar{\chi}_j^0} + B_1'^{t\bar{t}_i \bar{\chi}_j^0}) + m_{\bar{\chi}_j^0} \theta_{i1}^t \theta_{i2}^t (N_{j4}^2 + N_{j4}^{*2}) B_0'^{t\bar{t}_i \bar{\chi}_j^0}] \\ &\quad - \sum_{i,j} \frac{m_t}{16\pi^2} \{m_t [h_t^2 (\theta_{i1}^b)^2 |V_{j1}|^2 + h_b^2 (\theta_{i2}^b)^2 |U_{j2}|^2] (B_0'^{t\bar{t}_i \bar{\chi}_j^+} + B_1'^{t\bar{t}_i \bar{\chi}_j^+}) \\ &\quad + h_b h_t m_{\bar{\chi}_j^+} \theta_{i1}^b \theta_{i2}^b (U_{j2} V_{j2} + U_{j2}^* V_{j2}^*) B_0'^{t\bar{t}_i \bar{\chi}_j^+} \}, \end{aligned}$$

$$\begin{aligned}
\delta m_W^2 &= \frac{g^2}{16\pi^2} \left\{ (m_b^2 - m_t^2) \left(1 + \frac{m_b^2 - m_t^2 - 2m_W^2}{2m_W^2} B_0^{0bt} \right) - 2m_t^2 B_0^{0tt} \right. \\
&\quad \left. - \frac{1}{2m_W^2} [(m_b^2 - m_t^2)^2 + (m_b^2 + m_t^2)m_W^2] B_0^{Wbt} \right\}, \\
\delta Z_W &= \frac{g^2}{32\pi^2 m_W^2} \left\{ \frac{(m_b^2 - m_t^2)^2}{m_W^2} (B_0^{0bt} - B_0^{Wbt}) + [(m_b^2 - m_t^2)^2 \right. \\
&\quad \left. + (m_b^2 + m_t^2)m_W^2] B_0^{Wbt} \right\}, \\
\delta m_Z^2 &= \frac{g^2 s_W^2}{18c_W^2 \pi^2} \left[\frac{m_b^2}{2} (3 - 2s_W^2) (B_0^{Zbb} + B_0^{0bb}) - m_t^2 (3 - 4s_W^2) (B_0^{Ztt} - B_0^{0tt}) \right] \\
&\quad + \frac{g^2}{32c_W^2 \pi^2} [m_b^2 (B_0^{Zbb} - 2B_0^{0bb}) - m_t^2 (B_0^{Ztt} + 2B_0^{0tt})], \\
\delta Z_{H^+} &= \frac{3}{16\pi^2} [2(h_t^2 \beta_{11}^2 + h_b^2 \beta_{21}^2) (B_1^{H^+bt} + m_b^2 B_0'^{H^+bt} + m_{H^+}^2 B_1'^{H^+bt}) \\
&\quad - 4h_b h_t m_b m_t \beta_{11} \beta_{21} B_0'^{H^+bt} + \sum_{i,j,i',j'} (\theta_{ii'}^b)^2 (\theta_{jj'}^t)^2 (h_b \Theta_{i'j'1}^5 + h_t \Theta_{i'j'1}^6)^2 B_0'^{H^+ \bar{b}_i \bar{t}_j}], \\
\delta m_{H_k}^2 &= \frac{3}{16\pi^2} \left\{ -2h_t^2 \alpha_{1k}^2 [m_t^2 (1 + B_0^{0tt} + 2B_0^{H_k tt}) + m_{H_k}^2 B_1^{H_k tt}] - 2h_b^2 \alpha_{2k}^2 [m_b^2 (1 \right. \\
&\quad \left. + B_0^{0bb} + 2B_0^{H_k bb}) + m_{H_k}^2 B_1^{H_k bb}] + \sum_{i,j,i',j'} [(h_t \theta_{ii'}^t \theta_{jj'}^t \Theta_{i'j'k}^1)^2 B_0^{H_k \bar{t}_i \bar{t}_j} \right. \\
&\quad \left. + (h_b \theta_{ii'}^b \theta_{jj'}^b \Theta_{i'j'k}^2)^2 B_0^{H_k \bar{b}_i \bar{b}_j}] + \sum_i h_b^2 m_{b_i}^2 \alpha_{2k}^2 (1 + B_0^{0\bar{b}_i \bar{b}_i}) \right. \\
&\quad \left. + \sum_i h_t^2 m_{t_i}^2 \alpha_{1k}^2 (1 + B_0^{0\bar{t}_i \bar{t}_i}) \right\}, \\
\delta Z_{H_k} &= \frac{3}{16\pi^2} \left\{ 2h_t^2 \alpha_{1k}^2 (B_1^{H_k tt} + 2m_t^2 B_0'^{H_k tt} + m_{H_k}^2 B_1'^{H_k tt}) + 2h_b^2 \alpha_{2k}^2 (B_1^{H_k bb} \right. \\
&\quad \left. + 2m_b^2 B_0'^{H_k bb} + m_{H_k}^2 B_1'^{H_k bb}) + \sum_{i,j,i',j'} [(h_t \theta_{ii'}^t \theta_{jj'}^t \Theta_{i'j'k}^1)^2 B_0'^{H_k \bar{t}_i \bar{t}_j} \right. \\
&\quad \left. + (h_b \theta_{ii'}^b \theta_{jj'}^b \Theta_{i'j'k}^2)^2 B_0'^{H_k \bar{b}_i \bar{b}_j}] \right\}, \\
\delta m_{A_k}^2 &= \frac{3}{16\pi^2} \left\{ 2h_t^2 \beta_{1k}^2 [m_t^2 (1 + B_0^{0tt}) + m_{A_k}^2 B_1^{A_k tt}] + 2h_b^2 \beta_{2k}^2 [m_b^2 (1 + B_0^{0bb}) \right. \\
&\quad \left. + m_{A_k}^2 B_1^{A_k bb}] - \sum_{i,j,i',j'} [(h_t \theta_{ii'}^t \theta_{jj'}^t \Theta_{i'j'k}^3)^2 B_0^{A_k \bar{t}_i \bar{t}_j} + (h_b \theta_{ii'}^b \theta_{jj'}^b \Theta_{i'j'k}^4)^2 B_0^{A_k \bar{b}_i \bar{b}_j}] \right. \\
&\quad \left. + \sum_i h_b^2 m_{b_i}^2 \beta_{2k}^2 (1 + B_0^{0\bar{b}_i \bar{b}_i}) + \sum_i h_t^2 m_{t_i}^2 \beta_{1k}^2 (1 + B_0^{0\bar{t}_i \bar{t}_i}) \right\}, \\
\delta Z_{A_k} &= \frac{3}{16\pi^2} \left\{ 2h_t^2 \beta_{1k}^2 (B_1^{A_k tt} + m_{A_k}^2 B_1'^{A_k tt}) + 2h_b^2 \beta_{2k}^2 (B_1^{A_k bb} + m_{A_k}^2 B_1'^{A_k bb}) \right. \\
&\quad \left. - \sum_{i,j,i',j'} [(h_t \theta_{ii'}^t \theta_{jj'}^t \Theta_{i'j'k}^3)^2 B_0'^{A_k \bar{t}_i \bar{t}_j} + (h_b \theta_{ii'}^b \theta_{jj'}^b \Theta_{i'j'k}^4)^2 B_0'^{A_k \bar{b}_i \bar{b}_j}] \right\}, \\
Z_{H^+W} &= \frac{-3g}{16\sqrt{2}\pi^2 m_{H^+}^2 m_W^2} [(h_t m_t \beta_{11} + h_b m_b \beta_{12}) ((m_b^2 - m_t^2) (B_0^{0bt} - B_0^{H^+bt}) - m_{H^+}^2 B_0^{H^+bt})]
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i,j,i',j'} \theta_{i1}^b \theta_{i'1}^b \theta_{j1}^t \theta_{j'1}^t (h_b \Theta_{i'j'1}^5 + h_t \Theta_{i'j'1}^6) (m_{\tilde{t}_j}^2 - m_{\tilde{b}_i}^2) (B_0^{0\tilde{b}_i \tilde{t}_j} - B_0^{H^+ \tilde{b}_i \tilde{t}_j}), \\
Z_{AZ} &= \frac{-i3gc_W}{16\sqrt{2}\pi^2 m_W^2} (h_t m_t \beta_{11} B_0^{Att} - h_b m_b \beta_{12} B_0^{Abb}) \\
& + \frac{igc_W}{32\pi^2 m_A^2 m_W^2} \sum_{i,j,i',j'} \{h_b \theta_{i'1}^b \theta_{j'1}^b \Theta_{i'j'1}^4 [(3 - 2s_W^2) \theta_{i1}^b \theta_{j1}^b - 2s_W^2 \theta_{i2}^b \theta_{j2}^b] (m_{\tilde{b}_i}^2 - m_{\tilde{b}_j}^2) (B_0^{0\tilde{b}_i \tilde{b}_j} \\
& - B_0^{A\tilde{b}_i \tilde{b}_j}) - h_t \theta_{i'1}^t \theta_{j'1}^t \Theta_{i'j'1}^3 [(3 - 4s_W^2) \theta_{i1}^t \theta_{j1}^t - 4s_W^2 \theta_{i2}^t \theta_{j2}^t] (m_{\tilde{t}_i}^2 - m_{\tilde{t}_j}^2) (B_0^{0\tilde{t}_i \tilde{t}_j} - B_0^{A\tilde{t}_i \tilde{t}_j})\}, \\
Z_{hH}^{1/2} &= \frac{3\alpha_{11}\alpha_{12}}{16\pi^2 (m_h^2 - m_H^2)} [2m_b^2 (1 + B_0^{0bb} + 2B_0^{Hbb}) - 2m_t^2 (1 + B_0^{0tt} + 2B_0^{Htt}) \\
& - m_H^2 (B_0^{Hbb} - B_0^{Htt})] \\
& + \frac{3}{16\pi^2 (m_h^2 - m_H^2)} \sum_{i,j,i',j'} [(h_b \theta_{i'1}^b \theta_{j'1}^b)^2 \Theta_{i'j'1}^2 \Theta_{i'j'2}^2 B_0^{H\tilde{b}_i \tilde{b}_j} + (h_t \theta_{i'1}^t \theta_{j'1}^t)^2 \Theta_{i'j'1}^1 \Theta_{i'j'2}^1 B_0^{H\tilde{t}_i \tilde{t}_j}] \\
& - \frac{3\alpha_{11}\alpha_{12}}{16\pi^2 (m_h^2 - m_H^2)} \sum_i [h_b^2 m_{\tilde{b}_i}^2 (1 + B_0^{0\tilde{b}_i \tilde{b}_i}) + h_t^2 m_{\tilde{t}_i}^2 (1 + B_0^{0\tilde{t}_i \tilde{t}_i})], \\
Z_{Hh}^{1/2} &= Z_{hH}^{1/2} |_{h \leftrightarrow H}, \tag{10}
\end{aligned}$$

with

$$B_n^{ijk} = (-1)^n \left\{ \frac{\Delta}{n+1} - \int_0^1 dy y^n \ln \left[\frac{m_i^2 y(y-1) + m_j^2 (1-y) + m_k^2 y}{\mu^2} \right] \right\}, \tag{11}$$

$$B_n'^{ijk} = (-1)^n \int_0^1 dy \frac{y^{n+1} (1-y)}{m_i^2 y(y-1) + m_j^2 (1-y) + m_k^2 y}. \tag{12}$$

The notations θ_{ij}^t and θ_{ij}^b used in above expressions are defined in Appendix A. A_i stands for A with $i = 1$ and G^0 with $i = 2$. H_i^+ stands for H^+ with $i = 1$ and G^+ with $i = 2$. $\frac{\delta m_b}{m_b}$, δZ_L^b , δZ_R^b can be obtained, respectively, from $\frac{\delta m_t}{m_t}$, δZ_L^t , δZ_R^t by the transformation:

$$h_b \leftrightarrow h_t, m_b \leftrightarrow m_t, m_{\tilde{b}_i} \leftrightarrow m_{\tilde{t}_i}, \alpha_{1i} \leftrightarrow \alpha_{2i}, \beta_{1i} \leftrightarrow \beta_{2i}, \theta_{ij}^b \leftrightarrow \theta_{ij}^t, N_{i4} \rightarrow N_{i3}, U_{i2} \rightarrow V_{i2}.$$

The corresponding amplitude squared is

$$\overline{\sum} |M_{ren}|^2 = \overline{\sum} |M_0^{(s)} + M_0^{(t)}|^2 + 2Re \overline{\sum} [(\sum \delta M)(M_0^{(s)} + M_0^{(t)})^\dagger]. \tag{13}$$

The cross section for the process $b\bar{b} \rightarrow W^\pm H^\mp$ is

$$\hat{\sigma} = \int_{\hat{t}_-}^{\hat{t}_+} \frac{1}{16\pi \hat{s}^2} \overline{\sum} |M_{ren}|^2 d\hat{t} \tag{14}$$

with

$$\hat{t}_{\pm} = \frac{m_W^2 + m_{H^-}^2 - \hat{s}}{2} \pm \frac{1}{2} \sqrt{(\hat{s} - (m_W + m_{H^-})^2)(\hat{s} - (m_W - m_{H^-})^2)}. \quad (15)$$

The total hadronic cross section for $pp \rightarrow b\bar{b} \rightarrow W^{\pm}H^{\mp}$ can be obtained by folding the subprocess cross section $\hat{\sigma}$ with the parton luminosity:

$$\sigma(s) = \int_{(m_W + m_{H^-})/\sqrt{s}}^1 dz \frac{dL}{dz} \hat{\sigma}(b\bar{b} \rightarrow W^{\pm}H^{\mp} \text{ at } \hat{s} = z^2 s). \quad (16)$$

Here \sqrt{s} and $\sqrt{\hat{s}}$ are the CM energies of the pp and $b\bar{b}$ states, respectively, and dL/dz is the parton luminosity, defined as

$$\frac{dL}{dz} = 2z \int_{z^2}^1 \frac{dx}{x} f_{b/P}(x, \mu) f_{\bar{b}/P}(z^2/x, \mu), \quad (17)$$

where $f_{b/P}(x, \mu)$ and $f_{\bar{b}/P}(z^2/x, \mu)$ are the bottom and anti-bottom quark parton distribution functions, respectively.

3. Numerical results and conclusion

We now present some numerical results for the SUSY EW corrections to $W^{\pm}H^{\mp}$ associated production at the LHC. The SM input parameters in our calculations were taken to be $\alpha_{ew}(m_Z) = 1/128.8$, $m_W = 80.375\text{GeV}$ and $m_Z = 91.1867\text{GeV}$ [17], and $m_t = 175.6\text{GeV}$ and $m_b = 4.7\text{GeV}$, which were taken according to Ref.[10] for comparison. We used the CTEQ5M parton distributions throughout the calculations[18]. The one-loop relations[19] between the Higgs boson masses $M_{h,H,A,H^{\mp}}$ and the parameters α and β in the MSSM were used, and m_{H^+} and β were chosen as the two independent input parameters. Other MSSM parameters were determined as follows:

(i) For the parameters M_1 , M_2 and μ in the chargino and neutralino matrix, we take M_2 and μ as the input parameters, and then used the relation $M_1 = (5/3)(g'^2/g^2)M_2 \simeq 0.5M_2$ [2] to determine M_1 .

(ii) For the parameters $m_{\tilde{Q},\tilde{U},\tilde{D}}^2$ and $A_{t,b}$ in squark mass matrices

$$M_{\tilde{q}}^2 = \begin{pmatrix} M_{LL}^2 & m_q M_{LR} \\ m_q M_{RL} & M_{RR}^2 \end{pmatrix} \quad (18)$$

with

$$\begin{aligned}
M_{LL}^2 &= m_{\tilde{Q}}^2 + m_q^2 + m_Z^2 \cos 2\beta (I_q^{3L} - e_q \sin^2 \theta_W), \\
M_{RR}^2 &= m_{\tilde{U}, \tilde{D}}^2 + m_q^2 + m_Z^2 \cos 2\beta e_q \sin^2 \theta_W, \\
M_{LR} &= M_{RL} = \begin{pmatrix} A_t - \mu \cot \beta & (\tilde{q} = \tilde{t}) \\ A_b - \mu \tan \beta & (\tilde{q} = \tilde{b}) \end{pmatrix},
\end{aligned} \tag{19}$$

to simplify the calculation we assumed $M_{\tilde{Q}} = M_{\tilde{U}} = M_{\tilde{D}}$ and $A_t = A_b$, and we used $M_{\tilde{Q}}$ and A_t as the input parameters except the numerical calculations as shown in Fig.6, where we took $m_{\tilde{t}_1}$, $m_{\tilde{b}_1}$ and $A_t = A_b$ as the input parameters.

Some typical numerical calculations of the Yukawa corrections and the genuine SUSY EW corrections are given in Fig.3-4 and Fig.5-9, respectively.

In Fig.3 we present the Yukawa corrections to the total cross sections relative to the tree-level values as a function of m_{H^+} for $\tan \beta = 1.5, 2, 6$ and 30 . For $\tan \beta = 1.5$ and 2 the corrections decrease the total cross sections significantly, which exceed -6% for $m_{H^+} < 500\text{GeV}$ and -12% for $m_{H^+} < 300\text{GeV}$. For $\tan \beta (= 6)$ these corrections also decrease the total cross sections, although relatively smaller, which exceed -2.5% for $m_{H^+} < 500\text{GeV}$ and exceed -5% for $m_{H^+} < 250\text{GeV}$. But for high $\tan \beta (= 30)$ these corrections become positive, which increase the total cross sections slightly. Note that there are the peaks at $m_{H^+} = 180.3\text{GeV}$, which arise from the singularity of the charged Higgs boson wavefunction renormalization constant at the threshold point $m_{H^+} = m_t + m_b$.

In Fig.4 we show the Yukawa corrections as a function of $\tan \beta$ for $m_{H^+} = 100, 150, 200$ and 300GeV . For $\tan \beta < 4$ the corrections reduce the total cross sections by more than 10% with $m_{H^+} = 100, 150$ and 200GeV . With $m_{H^+} = 300\text{GeV}$ the corrections are only significant for $1 < \tan \beta < 5$. For high $\tan \beta (> 10)$ the corrections become negligibly small for all above m_{H^+} values.

Fig.5 gives the genuine SUSY EW corrections as a function of m_{H^+} for $\tan \beta = 1.5, 2, 6$ and 30 , respectively, assuming $M_2 = 300\text{GeV}$, $\mu = -100\text{GeV}$, $A_t = A_b = 200\text{GeV}$, and $M_{\tilde{Q}} = M_{\tilde{U}} = M_{\tilde{D}} = 500\text{GeV}$. From this figure one sees that the corrections are very small and negligible, which is reasonable because the squark masses are now very large and also the couplings of the charged Higgs boson-squarks are small

for the values of $A_{t,b}$, $M_{\tilde{Q},\tilde{U},\tilde{D}}$ and μ used in those numerical calculations. In contrast, in Fig.6 when we take the lighter squarks masses: $m_{\tilde{t}_1} = 100\text{GeV}$ and $m_{\tilde{b}_1} = 150\text{GeV}$, and put $A_t = A_b = 1\text{TeV}$, which are relatively larger, assuming $M_2 = 200\text{GeV}$, $\mu = 100\text{GeV}$ and $M_{\tilde{Q}} = M_{\tilde{U}}$, the genuine SUSY EW corrections are enhanced significantly, especially for low $\tan\beta (= 1.5)$ and m_{H^+} below 250GeV , which can exceed -30% . But when $m_{H^+} > 250\text{GeV}$ the corrections increase the cross sections, which can exceed 10% . For $\tan\beta = 6$ and 30 the corrections are at most 10% and become small with an increase of m_{H^+} . The sharp dips at $m_{H^+} = 250\text{GeV}$ are again due to the singularity of the charged Higgs boson wavefunction renormalization constant at the threshold point $m_{H^+} = m_{\tilde{t}_1} + m_{\tilde{b}_1} = 250\text{GeV}$.

Fig.7, Fig.8 and Fig.9 give the genuine SUSY EW corrections versus $A_t = A_b$, $M_{\tilde{Q}} = M_{\tilde{U}} = M_{\tilde{D}}$ and μ , respectively, for $\tan\beta = 1.5$ and 30 . In each figure we fixed $m_{H^+} = 200\text{GeV}$ and $M_2 = 300\text{GeV}$.

Fig.7 shows that the corrections are negative for $\tan\beta = 1.5$ and positive for $\tan\beta = 30$, assuming $M_{\tilde{Q}} = M_{\tilde{U}} = M_{\tilde{D}} = 400\text{GeV}$ and $\mu = 100\text{GeV}$. For both $\tan\beta = 1.5$ and 30 the magnitude of the corrections increases with increasing $A_t = A_b$. When $A_t = A_b = 1\text{TeV}$ the corrections can reach -6% and 7.5% for $\tan\beta = 1.5$ and 30 , respectively. Otherwise, when $A_t = A_b$ decrease to 100GeV , the corrections become negligibly small. This result is due to the fact that large values of $A_t = A_b$ not only enhance the couplings, but also give a large splitting between the masses of $\tilde{t}_1(\tilde{b}_1)$ and $\tilde{t}_2(\tilde{b}_2)$, and in consequence lighter \tilde{t}_1 and \tilde{b}_1 .

Fig.8 also show that the corrections are negative for $\tan\beta = 1.5$ and positive for $\tan\beta = 30$, assuming $A_t = A_b = 500\text{GeV}$ and $\mu = 100\text{GeV}$. When $M_{\tilde{Q},\tilde{U},\tilde{D}} = 250\text{GeV}$ the corrections can reach -3.6% for $\tan\beta = 1.5$ and 7.3% for $\tan\beta = 30$. But the magnitude of the corrections drops below one percent when $M_{\tilde{Q},\tilde{U},\tilde{D}}$ increase to 750GeV . This is because for larger values of $M_{\tilde{Q},\tilde{U},\tilde{D}}$ the squarks have larger masses and their virtual effects decrease due to the decoupling effects.

In Fig.9 we present the genuine SUSY EW corrections as a function of μ , assuming $A_t = A_b = 500\text{GeV}$ and $M_{\tilde{Q}} = M_{\tilde{U}} = M_{\tilde{D}} = 400\text{GeV}$. For $\tan\beta = 30$ the magnitude

of the corrections increase with an increase of $|\mu|$, which varies from 0% to 5% when $|\mu|$ ranges between $0 \sim 500\text{GeV}$. For $\tan\beta = 1.5$ the corrections are relatively small and increase slowly from about 0% to 3.5% when μ ranges between $-500\text{GeV} \sim 500\text{GeV}$. This result indicates that large values of μ and $\tan\beta$ can enhance the corrections significantly since the couplings become stronger.

In conclusion, we have calculated the $O(\alpha_{ew}m_{t(b)}^2/m_W^2)$ and $O(\alpha_{ew}m_{t(b)}^4/m_W^4)$ SUSY EW corrections to the cross sections for $W^\pm H^\mp$ associated production at the LHC in the MSSM. The numerical results show that the Yukawa corrections arising from the Higgs sector can decrease the total cross sections significantly for low $\tan\beta$ ($= 1.5$ and 2) when $m_{H^+} (< 300)\text{GeV}$, which exceed -12% . For high $\tan\beta$ the Yukawa corrections become negligibly small. The genuine SUSY EW corrections can increase or decrease the total cross sections depending on the SUSY parameters, which can exceed -25% for the favorable SUSY parameter values. We also show that the genuine SUSY EW corrections depend strongly on the choice of $\tan\beta$, A_t , $M_{\tilde{Q}}$ and μ . For large values of A_t , or large values of μ and $\tan\beta$, one can get much larger corrections. The corrections can become very small, in contrast, for larger values of $M_{\tilde{Q}}$.

This work was supported in part by the National Natural Science Foundation of China, the Doctoral Program Foundation of Higher Education of China, the Post Doctoral Foundation of China and a grant from the State Commission of Science and Technology of China. S.H. Zhu also gratefully acknowledges the support of the K.C. Wong Education Foundation of Hong Kong.

Appendix A

We present some notations used in this paper here. We introduce an angle $\varphi = \beta - \alpha$, and for each angle $\alpha, \beta, \varphi, \theta^t$ or θ^b , we define

$$\alpha_{ij} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix}, \beta_{ij} = \begin{pmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{pmatrix}, \varphi_{ij} = \begin{pmatrix} \cos \varphi & \sin \varphi \\ -\sin \varphi & \cos \varphi \end{pmatrix},$$

$$\theta_{ij}^t = \begin{pmatrix} \cos \theta^t & \sin \theta^t \\ -\sin \theta^t & \cos \theta^t \end{pmatrix}, \theta_{ij}^b = \begin{pmatrix} \cos \theta^b & \sin \theta^b \\ -\sin \theta^b & \cos \theta^b \end{pmatrix}$$

We define six matrix $\Theta_{ij}^i, i = 1 - 6$ for the couplings between squarks and Higgses:

$$\Theta_{ij1}^1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 2m_t \cos \alpha & A_t \cos \alpha + \mu \sin \alpha \\ A_t \cos \alpha + \mu \sin \alpha & 2m_t \cos \alpha \end{pmatrix}$$

$$\Theta_{ij2}^1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 2m_t \sin \alpha & A_t \sin \alpha - \mu \cos \alpha \\ A_t \sin \alpha - \mu \cos \alpha & 2m_t \sin \alpha \end{pmatrix}$$

$$\Theta_{ij1}^2 = \frac{-1}{\sqrt{2}} \begin{pmatrix} 2m_b \sin \alpha & A_b \sin \alpha + \mu \cos \alpha \\ A_b \sin \alpha + \mu \cos \alpha & 2m_b \sin \alpha \end{pmatrix}$$

$$\Theta_{ij2}^2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 2m_b \cos \alpha & A_b \cos \alpha - \mu \sin \alpha \\ A_b \cos \alpha - \mu \sin \alpha & 2m_b \cos \alpha \end{pmatrix}$$

$$\Theta_{ij1}^3 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & A_t \cos \beta + \mu \sin \beta \\ -A_t \cos \beta - \mu \sin \beta & 0 \end{pmatrix}$$

$$\Theta_{ij2}^3 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & A_t \sin \beta - \mu \cos \beta \\ -A_t \sin \beta + \mu \cos \beta & 0 \end{pmatrix}$$

$$\Theta_{ij1}^4 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & A_b \sin \beta + \mu \cos \beta \\ -A_b \sin \beta - \mu \cos \beta & 0 \end{pmatrix}$$

$$\Theta_{ij2}^4 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -A_b \cos \beta + \mu \sin \beta \\ A_b \cos \beta - \mu \sin \beta & 0 \end{pmatrix}$$

$$\Theta_{ij1}^5 = \begin{pmatrix} m_b \sin \beta & 0 \\ A_b \sin \beta + \mu \cos \beta & m_t \sin \beta \end{pmatrix}$$

$$\Theta_{ij2}^5 = \begin{pmatrix} -m_b \cos \beta & 0 \\ -A_b \cos \beta + \mu \sin \beta & 0 \end{pmatrix}$$

$$\Theta_{ij1}^6 = \begin{pmatrix} m_t \cos \beta & A_t \cos \beta + \mu \sin \beta \\ 0 & m_b \cos \beta \end{pmatrix}$$

$$\Theta_{ij2}^6 = \begin{pmatrix} m_t \sin \beta & A_t \sin \beta - \mu \cos \beta \\ 0 & 0 \end{pmatrix}$$

Appendix B

The form factors defined in Eq.(9) are the following:

$$\begin{aligned}
f_1^{V_1(s)}(H) &= \sum_{i,j} \frac{gh_b^3 \alpha_{2i}^2 \alpha_{2j} \varphi_{j1}}{32\sqrt{2}\pi^2(\hat{s} - m_{H_j}^2)} \{B_0^{bbH_i} + [4m_b^2 C_0 \\
&\quad + (4m_b^2 + \hat{s})C_1](\hat{s}, m_b^2, m_b^2, m_b^2, m_b^2, m_{H_i}^2)\} \\
&\quad + \sum_{i,j} \frac{-gh_b^3 \beta_{2i}^2 \alpha_{2j} \varphi_{j1}}{32\sqrt{2}\pi^2(\hat{s} - m_{H_j}^2)} [B_0^{bbA_i} - (4m_b^2 - \hat{s})C_1(\hat{s}, m_b^2, m_b^2, m_b^2, m_b^2, m_{A_i}^2)] \\
&\quad + \sum_{i,j} \frac{gh_t \alpha_{1j} \varphi_{j1}}{16\sqrt{2}\pi^2(\hat{s} - m_{H_j}^2)} \{-h_b h_t \beta_{1i} \beta_{2i} B_0^{btH_i^+} + [(h_t^2 m_b m_t \beta_{1i}^2 \\
&\quad + 2h_b h_t m_t^2 \beta_{1i} \beta_{2i} + h_b^2 m_b m_t \beta_{2i}^2)C_0 + (2h_t^2 m_b m_t \beta_{1i}^2 + h_b h_t \hat{s} \beta_{1i} \beta_{2i} \\
&\quad + 2h_b^2 m_b m_t \beta_{2i}^2)C_1](\hat{s}, m_b^2, m_b^2, m_t^2, m_t^2, m_{H_i^+}^2)\} \\
&\quad + \sum_{i,j,k,l,i',j'} \frac{gh_t \varphi_{l1} \theta_{ii'}^t \theta_{jj'}^t \Theta_{j'i'l}^1}{16\pi^2(\hat{s} - m_{H_i}^2)} [h_b^2 m_b \theta_{i1}^t \theta_{j1}^t |U_{k2}|^2 (C_0 + C_1 + C_2) \\
&\quad + m_{\tilde{\chi}_k^+} h_b h_t \theta_{i2}^t \theta_{j1}^t U_{k2} V_{k2} C_0 - h_t^2 m_b \theta_{i2}^t \theta_{j2}^t |V_{j2}|^2 C_1](\hat{s}, m_b^2, m_b^2, m_{\tilde{t}_i}^2, m_{\tilde{t}_j}^2, m_{\tilde{\chi}_k^+}^2) \\
&\quad + \sum_{i,j,k,l,j',k'} \frac{gh_b^3 \varphi_{i1} \theta_{jj'}^b \theta_{kk'}^b N_{l3} \Theta_{j'k'i}^2}{16\pi^2(\hat{s} - m_{H_i}^2)} [m_b \theta_{j1}^b \theta_{k1}^b N_{l3}^* (C_0 + C_1 + C_2) \\
&\quad - m_b \theta_{j2}^b \theta_{k2}^b N_{l3}^* C_1 + m_{\tilde{\chi}_l^0} \theta_{j1}^b \theta_{k2}^b N_{l3} C_0](\hat{s}, m_b^2, m_b^2, m_{\tilde{b}_j}^2, m_{\tilde{b}_k}^2, m_{\tilde{\chi}_l^0}^2), \\
f_2^{V_1(s)}(H) &= f_1^{V_1(s)}(H) (h_b \theta_{n1}^t \leftrightarrow h_t \theta_{n2}^t, \theta_{n1}^b \leftrightarrow \theta_{n2}^b, U_{n2} \leftrightarrow V_{n2}^*, N_{n3} \leftrightarrow N_{n3}^*), \\
f_3^{V_1(s)}(H) &= f_1^{V_1(s)}(H), \\
f_4^{V_1(s)}(H) &= f_2^{V_1(s)}(H); \\
f_i^{V_1(s)}(A) &= f_i^{V_1(s)}(A)_a + f_i^{V_1(s)}(A)_b,
\end{aligned}$$

where

$$\begin{aligned}
f_1^{V_1(s)}(A)_a &= \sum_{i,j,k,i',j'} \frac{gh_t \theta_{ii'}^t \theta_{jj'}^t \Theta_{j'i'l}^3}{16\pi^2(\hat{s} - m_A^2)} [-h_b^2 m_b \theta_{i1}^t \theta_{j1}^t |U_{j2}|^2 (C_0 + C_1 + C_2) \\
&\quad - m_{\tilde{\chi}_k^+} h_b h_t \theta_{i2}^t \theta_{j1}^t U_{k2} V_{k2} C_0 + h_t^2 m_b \theta_{i1}^t \theta_{j2}^t |V_{j2}|^2 C_2](\hat{s}, m_b^2, m_b^2, m_{\tilde{t}_i}^2, m_{\tilde{t}_j}^2, m_{\tilde{\chi}_k^+}^2) \\
&\quad + \sum_{i,j,k,i',j'} \frac{gh_b^3 N_{k3} \theta_{ii'}^b \theta_{jj'}^b \Theta_{j'k'i}^4}{16\pi^2(\hat{s} - m_A^2)} [-m_b \theta_{i1}^b \theta_{j1}^b N_{k3}^* (C_1 + C_2 + C_3) \\
&\quad + m_b \theta_{i2}^b \theta_{j1}^b N_{k3}^* C_1 - m_{\tilde{\chi}_k^0} \theta_{j1}^b \theta_{i2}^b N_{k3} C_0](\hat{s}, m_b^2, m_b^2, m_{\tilde{b}_i}^2, m_{\tilde{b}_j}^2, m_{\tilde{\chi}_k^0}^2), \\
f_2^{V_1(s)}(A)_a &= f_1^{V_1(s)}(A)_a (h_b \theta_{n1}^t \leftrightarrow h_t \theta_{n2}^t, \theta_{n1}^b \leftrightarrow \theta_{n2}^b, U_{n2} \leftrightarrow V_{n2}^*, N_{n3} \leftrightarrow N_{n3}^*),
\end{aligned}$$

$$\begin{aligned}
f_3^{V_1(s)}(A)_a &= f_1^{V_1(s)}(A)_a, \\
f_4^{V_1(s)}(A)_a &= f_2^{V_1(s)}(A)_a, \\
f_1^{V_1(s)}(A)_b &= \sum_i \frac{gh_b^3 \alpha_{2i}^2 \beta_{21}}{32\sqrt{2}\pi^2(\hat{s} - m_A^2)} \{B_0^{bbH_i} - [4m_b^2 C_0 \\
&\quad + (4m_b^2 - \hat{s})C_1](\hat{s}, m_b^2, m_b^2, m_b^2, m_b^2, m_{H_i}^2)\} \\
&\quad + \sum_i \frac{-gh_b^3 \beta_{2i}' \beta_{21}}{32\sqrt{2}\pi^2(\hat{s} - m_A^2)} [B_0^{bbA_i} - (4m_b^2 - \hat{s})C_1(\hat{s}, m_b^2, m_b^2, m_b^2, m_b^2, m_{A_i}^2)] \\
&\quad + \sum_i \frac{gh_t \beta_{11}}{16\sqrt{2}\pi^2(\hat{s} - m_A^2)} \{h_b h_t \beta_{1i} \beta_{2i} B_0^{btH_i^+} \\
&\quad + [(h_t^2 m_b m_t \beta_{1i}^2 - 2h_b h_t m_b^2 \beta_{1i} \beta_{2i} + h_b^2 m_b m_t \beta_{2i}^2)C_0 \\
&\quad - h_b h_t (4m_b^2 - \hat{s})\beta_{1i} \beta_{2i} C_1](\hat{s}, m_b^2, m_b^2, m_t^2, m_t^2, m_{H_i^+}^2)\}, \\
f_2^{V_1(s)}(A)_b &= -f_3^{V_1(s)}(A)_b = f_4^{V_1(s)}(A)_b = -f_1^{V_1(s)}(A)_b; \\
f_1^{s(s)}(H) &= \sum_{i,j} \frac{-gh_b^3 \alpha_{2i}^2 \alpha_{2j} \varphi_{j1}}{8\sqrt{2}\pi^2(\hat{s} - m_{H_i}^2)(\hat{s} - m_{H_j}^2)} [m_b^2(1 + B_0^{0bb}) + (2m_b^2 B_0^{\hat{s}bb} + \hat{s}B_1^{\hat{s}bb})] \\
&\quad + \sum_{i,j} \frac{-gh_b h_t^2 \alpha_{1i} \alpha_{1j} \alpha_{2i} \varphi_{j1}}{8\sqrt{2}\pi^2(\hat{s} - m_{H_i}^2)(\hat{s} - m_{H_j}^2)} [m_t^2(1 + B_0^{0tt}) + (2m_t^2 B_0^{\hat{s}tt} + \hat{s}B_1^{\hat{s}tt})] \\
&\quad + \sum_{i,j,k,l,i',j'} \frac{gh_b^3 \alpha_{2k} \varphi_{l1} (\theta_{jj'}^b)^2 (\theta_{ii'}^b)^2 \Theta_{i'j'k}^2 \Theta_{j'l}^2}{16\sqrt{2}\pi^2(\hat{s} - m_{H_i}^2)(\hat{s} - m_{H_k}^2)} B_0^{\hat{s}\bar{b}_i \bar{b}_j} \\
&\quad + \sum_{i,j,k,l,i',j'} \frac{gh_b h_t^2 \alpha_{2l} \varphi_{k1} (\theta_{jj'}^t)^2 (\theta_{ii'}^t)^2 \Theta_{i'j'l}^1 \Theta_{j'k}^1}{16\sqrt{2}\pi^2(\hat{s} - m_{H_i}^2)(\hat{s} - m_{H_k}^2)} B_0^{\hat{s}\bar{t}_i \bar{t}_j} \\
&\quad + \sum_{i,j,k} \frac{3gh_b \alpha_{2i} \varphi_{j1}}{16\sqrt{2}\pi^2(\hat{s} - m_{H_i}^2)(\hat{s} - m_{H_j}^2)} [h_b^2 \alpha_{2i} \alpha_{2j} A_0(m_{b_k}^2) + h_t^2 \alpha_{1i} \alpha_{1j} A_0(m_{t_k}^2)], \\
f_2^{s(s)}(H) &= f_3^{s(s)}(H) = f_4^{s(s)}(H) = f_1^{s(s)}(H); \\
f_1^{s(s)}(A) &= \frac{gh_b^3 \beta_{21}^3}{8\sqrt{2}\pi^2(\hat{s} - m_A^2)^2} [m_b^2(1 + B_0^{0bb}) + \hat{s}B_1^{\hat{s}bb}] \\
&\quad + \frac{gh_b \beta_{11}^2 \beta_{21}}{8\sqrt{2}\pi^2(\hat{s} - m_A^2)^2} [m_t^2(1 + B_0^{0tt}) + \hat{s}B_1^{\hat{s}tt}] \\
&\quad - \sum_{i,j} \sum_{i',j'} \frac{gh_b^3 \beta_{21} (\theta_{jj'}^b)^2 (\theta_{ii'}^b)^2 (\Theta_{i'j'l}^4)^2}{16\sqrt{2}\pi^2(\hat{s} - m_A^2)^2} B_0^{\hat{s}\bar{b}_i \bar{b}_j} \\
&\quad - \sum_{i,j} \sum_{i',j'} \frac{-gh_b h_t^2 \beta_{21} (\theta_{jj'}^t)^2 (\theta_{ii'}^t)^2 (\Theta_{i'j'l}^3)^2}{16\sqrt{2}\pi^2(\hat{s} - m_A^2)^2} B_0^{\hat{s}\bar{t}_i \bar{t}_j} \\
&\quad - \sum_k \frac{3gh_b \beta_{21}}{16\sqrt{2}\pi^2(\hat{s} - m_A^2)^2} [h_b^2 \beta_{21}^2 A_0(m_{b_k}^2) + h_t^2 \beta_{11}^2 A_0(m_{t_k}^2)], \\
f_2^{s(s)}(A) &= -f_3^{s(s)}(A) = f_4^{s(s)}(A) = -f_1^{s(s)}(A);
\end{aligned}$$

$$\begin{aligned}
f_1^{V_2(s)}(H) = & \sum_i \frac{-gh_b^2 \alpha_{2i}^2}{16\sqrt{2}\pi^2(\hat{s} - m_{H_i}^2)} \left\{ \frac{3}{2} h_b \beta_{21} B_0^{\hat{s}bb} + [(h_t m_b m_t \beta_{11} \right. \\
& + h_b m_t^2 \beta_{21}) C_0 + h_b m_W^2 \beta_{21} C_1 - (h_b m_b^2 \beta_{21} + h_b m_t^2 \beta_{21} \\
& - 2h_t m_b m_t \beta_{11}) C_2 - 2h_b \beta_{21} C_{00} - h_b \beta_{21} (\hat{t} + \hat{u} - 2m_b^2) C_{12} \\
& \left. - 2h_b m_{H^+}^2 + \beta_{21} C_{22}] (\hat{s}, m_{H^+}^2, m_W^2, m_b^2, m_t^2, m_b^2) \right\} \\
& + \sum_i \frac{gh_b h_t \alpha_{1i} \alpha_{2i}}{16\sqrt{2}\pi^2(\hat{s} - m_{H_i}^2)} \left\{ \frac{3}{2} h_t \beta_{11} B_0^{\hat{s}tt} + [(h_t m_b^2 \beta_{11} \right. \\
& - h_b m_b m_t \beta_{21}) C_0 + h_t m_W^2 \beta_{11} C_1 - (h_t m_b^2 \beta_{11} + h_t m_t^2 \beta_{11} \\
& - 2h_b m_b m_t \beta_{21}) C_2 - 2h_t \beta_{11} C_{00} - h_t \beta_{11} (\hat{t} + \hat{u} - 2m_b^2) C_{12} \\
& \left. - 2h_t m_{H^+}^2 + \beta_{11} C_{22}] (\hat{s}, m_{H^+}^2, m_W^2, m_t^2, m_t^2, m_b^2) \right\} \\
& + \sum_{i,j,k,l,i',j',k'} \frac{gh_b^2}{16\pi^2(\hat{s} - m_{H_i}^2)} \alpha_{2l} (\theta_{ii'}^b)^2 \theta_{j1}^b \theta_{jj'}^b \theta_{k1}^b \theta_{kk'}^b \Theta_{i'j'l}^2 (h_b \Theta_{i'k'l}^5 \\
& + h_t \Theta_{i'k'l}^6) C_2 (\hat{s}, m_{H^+}^2, m_W^2, m_{b_j}^2, m_{b_i}^2, m_{t_k}^2) \\
& + \sum_{i,j,k,l,i',j',k'} \frac{gh_b h_t}{16\pi^2(\hat{s} - m_{H_i}^2)} \alpha_{2l} (\theta_{ii'}^t)^2 \theta_{j1}^t \theta_{jj'}^t \theta_{k1}^t \theta_{kk'}^t \Theta_{i'j'l}^1 (h_b \Theta_{i'k'l}^5 \\
& + h_t \Theta_{i'k'l}^6) C_2 (\hat{s}, m_{H^+}^2, m_W^2, m_{t_j}^2, m_{t_i}^2, m_{b_k}^2),
\end{aligned}$$

$$f_2^{V_2(s)}(H) = f_3^{V_2(s)}(H) = f_4^{V_2(s)}(H) = f_1^{V_2(s)}(H);$$

$$\begin{aligned}
f_1^{V_2(s)}(A) = & \frac{-gh_b^2 \beta_{21}^2}{16\sqrt{2}\pi^2(\hat{s} - m_A^2)} \left\{ \frac{3}{2} h_b \beta_{21} B_0^{\hat{s}bb} - [(h_t m_b m_t \beta_{11} - h_b m_t^2 \beta_{21}) C_0 \right. \\
& + h_b m_W^2 \beta_{21} C_1 + h_b \beta_{21} (m_b^2 - m_t^2) C_2 - 2h_b \beta_{21} C_{00} - h_b \beta_{21} (\hat{t} + \hat{u} \\
& - 2m_b^2) C_{12} - 2h_b m_{H^+}^2 + \beta_{21} C_{22}] (\hat{s}, m_{H^+}^2, m_W^2, m_b^2, m_b^2, m_t^2) \left. \right\} \\
& + \frac{-gh_b h_t \beta_{11} \beta_{21}}{16\sqrt{2}\pi^2(\hat{s} - m_A^2)} \left\{ \frac{3}{2} h_t \beta_{11} B_0^{\hat{s}tt} + [(h_t m_b^2 \beta_{11} - h_b m_b m_t \beta_{21}) C_0 \right. \\
& + h_t m_W^2 \beta_{11} C_1 - h_t \beta_{11} (m_b^2 - m_t^2) C_2 - 2h_t \beta_{11} C_{00} - h_t \beta_{11} (\hat{t} + \hat{u} \\
& - 2m_b^2) C_{12} - 2h_t m_{H^+}^2 + \beta_{11} C_{22}] (\hat{s}, m_{H^+}^2, m_W^2, m_t^2, m_t^2, m_b^2) \left. \right\} \\
& + \sum_{i,j,k,i',j',k'} \frac{-gh_b^2}{16\pi^2(\hat{s} - m_A^2)} \beta_{21} (\theta_{ii'}^b)^2 \theta_{j1}^b \theta_{jj'}^b \theta_{k1}^b \theta_{kk'}^b \Theta_{i'j'l}^4 (h_b \Theta_{i'k'l}^5 \\
& + h_t \Theta_{i'k'l}^6) C_2 (\hat{s}, m_{H^+}^2, m_W^2, m_{b_j}^2, m_{b_i}^2, m_{t_k}^2) \\
& + \sum_{i,j,k,i',j',k'} \frac{gh_b h_t}{16\pi^2(\hat{s} - m_A^2)} \beta_{21} (\theta_{ii'}^t)^2 \theta_{j1}^t \theta_{jj'}^t \theta_{k1}^t \theta_{kk'}^t \Theta_{i'j'l}^3 (h_b \Theta_{i'k'l}^5 \\
& + h_t \Theta_{i'k'l}^6) C_2 (\hat{s}, m_{H^+}^2, m_W^2, m_{t_j}^2, m_{t_i}^2, m_{b_k}^2),
\end{aligned}$$

$$f_2^{V_2(s)}(A) = -f_3^{V_2(s)}(A) = f_4^{V_2(s)}(A) = -f_1^{V_2(s)}(A);$$

$$\begin{aligned}
f_1^{V_1(t)} = & \sum_i \frac{-gh_b h_t^2 \alpha_{1i} \alpha_{2i} \beta_{11}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{-B_0^{Wbt} + [-m_{H_i}^2 C_0 + 2C_{00} + m_b^2 C_{11} \\
& + (m_b^2 + \hat{t})C_{12} + \hat{t}C_{22}](m_b^2, m_W^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2)\} \\
& + \sum_i \frac{-gh_b h_t^2 \beta_{1i} \beta_{2i} \beta_{11}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{-B_0^{Wbt} + [-m_{A_i}^2 C_0 + 2C_{00} \\
& + m_b^2 C_{11} + (m_b^2 + \hat{t})C_{12} + \hat{t}C_{22}](m_b^2, m_W^2, \hat{t}, m_{A_i}^2, m_b^2, m_t^2)\} \\
& + \sum_{i,j} \frac{-gh_t^2 \alpha_{1i} \beta_{11} \varphi_{ij}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} [(h_t m_b m_t \beta_{1j} - h_b m_t^2 \beta_{2j})C_0 + h_b \beta_{2j} (m_b^2 - m_t^2)C_1 \\
& + h_b \beta_{2j} (\hat{t} - m_t^2)C_2 + 2h_b \beta_{2j} C_{00} + (-h_t m_b m_t \beta_{1j} + h_b m_b^2 \beta_{2j} + h_b \hat{t} \beta_{2j})C_{12} \\
& + (h_b \hat{t} \beta_{2j} - h_t m_b m_t \beta_{1j})C_{22}](m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_j^+}^2, m_{H_i}^2) \\
& + \sum_i \frac{gh_t^2 \beta_{11} \beta_{1i}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} [h_t m_b m_t \beta_{1i} (C_0 + 2C_1 + 2C_2 + C_{11} + 2C_{12} + C_{22}) \\
& + 2h_b \beta_{2i} C_{00} + h_b m_t^2 \beta_{2i} (C_0 + C_1 + C_2) + h_b m_b^2 \beta_{2i} (C_1 + C_{11} + C_{12}) \\
& + h_b \beta_{2i} \hat{t} (C_2 + C_{12} + C_{22})](m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_i^+}^2, m_{A_i}^2) \\
& + \sum_{i,j} \frac{gh_b h_t \alpha_{2j} \beta_{11} \varphi_{ji}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} [h_t m_b^2 \beta_{1i} (C_0 + C_2 - C_{11} - C_{12}) \\
& - 2h_t \beta_{1i} C_{00} + h_b m_b m_t \beta_{2i} (-C_0 + C_{11} + 2C_{12} + C_{22}) - h_t \beta_{1i} \hat{t} (C_2 \\
& + C_{12} + C_{22})](m_b^2, m_W^2, \hat{t}, m_b^2, m_{H_j}^2, m_{H_i^+}^2) \\
& + \sum_{i,j,k} \frac{-gh_b h_t^2 \beta_{11} \theta_{j1}^b \theta_{i1}^t}{8\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{N_{k3} N_{k4}^* \theta_{j1}^b \theta_{i1}^t [m_b^2 (C_1 + C_{11} + C_{12}) + \hat{t} (C_2 + C_{12} \\
& + C_{22}) + 2C_{00}] - (N_{k3}^* N_{k4}^* \theta_{j2}^b \theta_{i1}^t m_b m_{\tilde{\chi}_k^0} + N_{k3} N_{k4} \theta_{j1}^b \theta_{i2}^t m_t m_{\tilde{\chi}_k^0}) (C_0 \\
& + C_1 + C_2) + N_{k3}^* N_{k4} m_b m_t \theta_{j1}^b \theta_{i2}^t (C_1 + C_2 + C_{11} + 2C_{12} \\
& + C_{22})\} (m_b^2, m_W^2, \hat{t}, m_{\tilde{\chi}_k^0}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2) \\
& + \sum_{i,j,k} \frac{gh_t^2 \beta_{11} N_{i4}^* \theta_{k1}^t}{8\pi^2(\hat{t} - m_t^2)} \{-h_b \theta_{k1}^t O_{ji}^{R*} U_{j2} B_0^{W\tilde{\chi}_j^+ \tilde{\chi}_i^0} + [h_b \theta_{k1}^t U_{j2} (m_{\tilde{\chi}_j^+} m_{\tilde{\chi}_i^0} O_{ji}^{L*} C_0 \\
& + O_{ji}^{R*} (2C_{00} + m_b^2 C_{11} + m_b^2 C_{12} + \hat{t} C_{12} + \hat{t} C_{22} - m_{\tilde{t}_k}^2 C_0)) + h_t m_b O_{ji}^{L*} V_{j2} \theta_{k2}^t (C_0 \\
& + C_1 + C_2)](m_b^2, m_W^2, \hat{t}, m_{\tilde{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2)\} \\
& + \sum_{i,j,k} \frac{gh_t^2 m_t \beta_{11} \theta_{k2}^t O_{ji}^{L*} N_{i4}}{8\pi^2(\hat{t} - m_t^2)} [h_b m_{\tilde{\chi}_j^+} \theta_{k1}^t (C_0 + C_1 + C_2) + h_t m_b \theta_{k2}^t V_{j2} (C_0 \\
& + 2C_1 + C_{11} + 2C_{12} + 2C_2 + C_{22})](m_b^2, m_W^2, \hat{t}, m_{\tilde{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2) \\
& + \sum_{i,j,k} \frac{gh_b h_t \beta_{11}}{8\pi^2(\hat{t} - m_t^2)} \{h_t (\theta_{k1}^b)^2 N_{j3} V_{i2} O_{ij}^{L*} B_0^{W\tilde{\chi}_i^+ \tilde{\chi}_j^0} - [h_t (\theta_{k1}^b)^2 N_{j3} V_{i2} (O_{ij}^{L*} (2C_{00}
\end{aligned}$$

$$\begin{aligned}
& +m_b^2(C_{11} + C_{12}) - m_{\tilde{b}_k}^2 C_0 + \hat{t}(C_{12} + C_{22})) + m_{\tilde{\chi}_i^+} m_{\tilde{\chi}_j^0} O_{ij}^{R*} C_0) \\
& + \theta_{k_1}^b \theta_{k_2}^b O_{ij}^{R*} (h_b m_t m_{\tilde{\chi}_j^0} N_{j_3} U_{i_2} + h_t m_b m_{\tilde{\chi}_i^+} N_{j_3}^* V_{i_2})(C_0 + C_1 + C_2) \\
& + h_b m_b m_t (\theta_{k_2}^b)^2 N_{j_3}^* U_{i_2} O_{ij}^{R*} (C_0 + 2C_1 + C_{11} + 2C_{12} \\
& + 2C_2 + C_{22}))(m_b^2, m_W^2, \hat{t}, m_{\tilde{b}_k}^2, m_{\tilde{\chi}_j^0}^2, m_{\tilde{\chi}_i^+}^2)\}, \\
f_2^{V_1(t)} = & \sum_i \frac{-gh_b^2 h_t m_b m_t \alpha_{1i} \alpha_{2i} \beta_{21}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} (4C_0 + 4C_1 + 4C_2 + C_{11} + 2C_{12} \\
& + C_{22})(m_b^2, m_W^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2) \\
& + \sum_i \frac{gh_b^2 h_t m_b m_t \beta_{1i} \beta_{2i} \beta_{21}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} (C_{11} + 2C_{12} + C_{22})(m_b^2, m_W^2, \hat{t}, m_{A_i}^2, m_b^2, m_t^2) \\
& + \sum_{i,j} \frac{-gh_b h_t \alpha_{1i} \beta_{21} \varphi_{ij}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} [(h_b m_b m_t \beta_{2j} - h_t m_t^2 \beta_{1j}) C_0 + h_t \beta_{1j} (m_b^2 - m_t^2) C_1 \\
& + h_t \beta_{1j} (\hat{t} - m_t^2) C_2 + 2h_t \beta_{1j} C_{00} + (-h_b m_b m_t \beta_{2j} + h_t m_b^2 \beta_{1j} + h_t \hat{t} \beta_{1j}) C_{12} \\
& + (h_t \hat{t} \beta_{1j} - h_b m_b m_t \beta_{2j}) C_{22}](m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_j^+}^2, m_{H_i}^2) \\
& + \sum_i \frac{-gh_b h_t \beta_{1i} \beta_{21}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} [h_b m_b m_t \beta_{2i} (C_0 + 2C_1 + 2C_2 + C_{11} + 2C_{12} + C_{22}) \\
& + 2h_t \beta_{1i} C_{00} + h_t m_t^2 \beta_{1i} (C_0 + C_1 + C_2) + h_t m_b^2 \beta_{1i} (C_1 + C_{11} + C_{12}) \\
& + h_t \beta_{1i} \hat{t} (C_2 + C_{12} + C_{22})](m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_i^+}^2, m_{A_i}^2) \\
& + \sum_{i,j} \frac{gh_b^2 \alpha_{2j} \beta_{21} \varphi_{ji}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} [h_b m_b^2 \beta_{2i} (C_0 + C_2 - C_{11} - C_{12}) \\
& - 2h_b \beta_{2i} C_{00} + h_t m_b m_t \beta_{1i} (-C_0 + C_{11} + 2C_{12} + C_{22}) - h_b \beta_{2i} \hat{t} (C_2 \\
& + C_{12} + C_{22})](m_b^2, m_W^2, \hat{t}, m_b^2, m_{H_j}^2, m_{H_i^+}^2) \\
& + \sum_{i,j,k} \frac{gh_b^2 h_t \beta_{21} \theta_{j_1}^b \theta_{i_1}^t}{8\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{N_{k_3} N_{k_4}^* \theta_{j_1}^b \theta_{i_1}^t m_b m_t (C_1 + C_2 + C_{11} + 2C_{12} + C_{22}) \\
& - (N_{k_3}^* N_{k_4}^* \theta_{j_2}^b \theta_{i_1}^t m_t m_{\tilde{\chi}_k^0} + N_{k_3} N_{k_4} \theta_{j_1}^b \theta_{i_2}^t m_b m_{\tilde{\chi}_k^0}) (C_0 + C_1 + C_2) \\
& + N_{k_3}^* N_{k_4} \theta_{j_2}^b \theta_{i_2}^t [m_b^2 (C_1 + C_{11} + C_{12}) + \hat{t} (C_2 + C_{12} + C_{22}) \\
& + 2C_{00}]\} (m_b^2, m_W^2, \hat{t}, m_{\tilde{\chi}_k^0}^2, m_{\tilde{b}_j}^2, m_{\tilde{t}_i}^2) \\
& + \sum_{i,j,k} \frac{gh_b h_t \beta_{12} \theta_{k_2}^t N_{i_4}}{8\pi^2(\hat{t} - m_t^2)} \{-h_t \theta_{k_2}^t O_{ji}^{L*} V_{j_2} B_0^{W\tilde{\chi}_j^+ \tilde{\chi}_i^0} + [h_b m_b m_{\tilde{\chi}_i^0} \theta_{k_1}^t O_{ji}^{R*} U_{j_2} (C_0 \\
& + C_1 + C_2) + h_t \theta_{k_2}^t O_{ji}^{L*} V_{j_2} (2C_{00} + m_b^2 C_{11} + m_b^2 C_{12} + \hat{t} C_{12} + \hat{t} C_{22} \\
& - m_{\tilde{t}_k}^2 C_0 + m_{\tilde{\chi}_i^0} m_{\tilde{\chi}_j^+} C_0)](m_b^2, m_W^2, \hat{t}, m_{\tilde{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2)\}
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i,j,k} \frac{gh_b h_t m_t \beta_{12} \theta_{k1}^t O_{ji}^{R*} N_{i4}^*}{8\pi^2(\hat{t} - m_t^2)} [h_b m_b \theta_{k1}^t U_{j2} (C_0 + 2C_1 + C_{11} + 2C_{12} + 2C_2 \\
& + C_{22}) + h_t m_{\tilde{\chi}_j^+} \theta_{k2}^t V_{j2} (C_0 + C_1 + C_2)] (m_b^2, m_W^2, \hat{t}, m_{\tilde{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2) \\
& - \sum_{i,j,k} \frac{gh_b^2 \beta_{21}}{8\pi^2(\hat{t} - m_t^2)} \{h_b (\theta_{k2}^b)^2 N_{j3}^* U_{i2} O_{ij}^{R*} B_0^{W\tilde{\chi}_i^+ \tilde{\chi}_j^0} - [h_b (\theta_{k2}^b)^2 N_{j3}^* U_{i2} (O_{ij}^{R*} (2C_{00} \\
& + m_b^2 (C_{11} + C_{12}) - m_{\tilde{b}_k}^2 C_0 + \hat{t} (C_{12} + C_{22})) + m_{\tilde{\chi}_i^+} m_{\tilde{\chi}_j^0} O_{ij}^{L*} C_0) \\
& + \theta_{k1}^b \theta_{k2}^b O_{ij}^{L*} (h_t m_t m_{\tilde{\chi}_j^0} N_{j3}^* V_{i2} + h_b m_b m_{\tilde{\chi}_i^+} N_{j3} U_{i2}) (C_0 + C_1 + C_2) \\
& + h_t m_b m_t (\theta_{k1}^b)^2 N_{j3} V_{i2} O_{ij}^{L*} (C_0 + 2C_1 + C_{11} + 2C_{12} \\
& + 2C_2 + C_{22})] (m_b^2, m_W^2, \hat{t}, m_{\tilde{t}_k}^2, m_{\tilde{\chi}_j^0}^2, m_{\tilde{\chi}_i^+}^2)\}, \\
f_5^{V_1(t)} & = \sum_i \frac{gh_b^2 h_t m_t \alpha_{1i} \alpha_{2i} \beta_{21}}{32\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{B_0^{Wbt} + [(4m_b^2 + m_{H_i}^2)C_0 + 4m_b^2 C_1 \\
& + 2(m_b^2 + \hat{t})C_2 - 2C_{00}] (m_b^2, m_W^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2)\} \\
& + \sum_i \frac{-gh_b^2 h_t m_t \beta_{1i} \beta_{2i} \beta_{21}}{32\sqrt{2}\pi^2(\hat{t} - m_t^2)} \Big|_0^{Wbt} + (m_{A_i}^2 C_0 - 2C_{00}) (m_b^2, m_W^2, \hat{t}, m_{A_i}^2, m_b^2, m_t^2) \\
& + \sum_{i,j} \frac{gh_b h_t \alpha_{1i} \beta_{21} \varphi_{ij}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} (h_t m_b \beta_{1j} + h_b m_t \beta_{2j}) C_{00} (m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_j^+}^2, m_{H_i}^2) \\
& + \sum_i \frac{gh_b h_t \beta_{1i} \beta_{21}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} (h_t m_b \beta_{1i} - h_b m_t \beta_{2i}) C_{00} (m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_i^+}^2, m_{A_i}^2) \\
& + \sum_{i,j} \frac{gh_b^2 \alpha_{2j} \beta_{21} \varphi_{ji}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} (h_t m_t \beta_{1i} + h_b m_b \beta_{2i}) C_{00} (m_b^2, m_W^2, \hat{t}, m_b^2, m_{H_j}^2, m_{H_i^+}^2) \\
& + \sum_{i,j,k} \frac{-gh_b^2 h_t \beta_{21} \theta_{j1}^b \theta_{i1}^t}{8\sqrt{2}\pi^2(\hat{t} - m_t^2)} (m_b N_{k3}^* N_{k4} \theta_{j2}^b \theta_{i2}^t - m_t N_{k3} N_{k4}^* \theta_{j1}^b \theta_{i1}^t) C_{00} \\
& (m_b^2, m_W^2, \hat{t}, m_{\tilde{\chi}_k^0}^2, m_{\tilde{t}_j}^2, m_{\tilde{t}_i}^2) \\
& + \sum_{i,j,k} \frac{gh_b h_t \beta_{12} \theta_{k2}^t N_{i4}^*}{16\pi^2(\hat{t} - m_t^2)} \{-h_t m_b \theta_{k2}^t O_{ji}^{L*} V_{j2} B_0^{W\tilde{\chi}_j^+ \tilde{\chi}_i^0} + [-h_b m_{\tilde{\chi}_j^+} \theta_{k1}^t O_{ji}^{L*} U_{j2} (m_b^2 C_1 \\
& + \hat{t} C_0 + \hat{t} C_2) + h_b m_{\tilde{\chi}_i^0} \theta_{k1}^t O_{ji}^{R*} U_{j2} (m_b^2 C_0 + m_b^2 C_1 + \hat{t} C_2) \\
& - h_t m_b \theta_{k2}^t O_{ji}^{L*} V_{j2} (-2C_{00} - m_b^2 C_1 + \hat{t} (C_0 + C_1 + 2C_2) + m_{\tilde{t}_k}^2 C_0 \\
& - m_{\tilde{\chi}_i^0} m_{\tilde{\chi}_j^+} C_0)] (m_b^2, m_W^2, \hat{t}, m_{\tilde{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2)\} \\
& + \sum_{i,j,k} \frac{gh_b h_t m_t \beta_{12} \theta_{k1}^t N_{i4}^*}{16\pi^2(\hat{t} - m_t^2)} \{h_b \theta_{k1}^t O_{ji}^{R*} U_{j2} B_0^{W\tilde{\chi}_j^+ \tilde{\chi}_i^0} + [-h_b m_{\tilde{\chi}_i^0} m_{\tilde{\chi}_j^+} \theta_{k1}^t O_{ji}^{L*} U_{j2} C_0 \\
& + h_b \theta_{k1}^t O_{ji}^{R*} U_{j2} (-2C_{00} + m_b^2 C_0 + 2m_b^2 C_1 + m_b^2 C_2 + \hat{t} C_2 + m_{\tilde{t}_k}^2 C_0) \\
& + h_t m_b \theta_{k2}^t V_{j2} (m_{\tilde{\chi}_j^+} O_{ji}^{R*} - m_{\tilde{\chi}_i^0} O_{ji}^{L*}) (C_0 + C_1
\end{aligned}$$

$$\begin{aligned}
& + C_2)](m_b^2, m_W^2, \hat{t}, m_{\hat{t}_k}^2, m_{\hat{\chi}_j^+}^2, m_{\hat{\chi}_i^0}^2) \} \\
& + \sum_{i,j,k} \frac{gh_b^2 \beta_{21}}{16\pi^2 (\hat{t} - m_{\hat{t}}^2)} \{ (-h_t (\theta_{k1}^b)^2 N_{j3} V_{i2} O_{ij}^{L*} + h_b (\theta_{k2}^b)^2 N_{j3}^* U_{i2} O_{ij}^{R*}) B_0^{W\hat{\chi}_i^+ \hat{\chi}_j^0} \\
& - [h_t m_t (\theta_{k1}^b)^2 N_{j3} V_{i2} (O_{ij}^{L*} (-2C_{00} + m_b^2 (C_0 + 2C_1 + C_2) + m_{b_k}^2 C_0 + \hat{t} C_2) \\
& - m_{\hat{\chi}_i^+} m_{\hat{\chi}_j^0} O_{ij}^{R*} C_0) + \theta_{k1}^b \theta_{k2}^b (h_b N_{j3} U_{i2} (m_{\hat{\chi}_i^+} O_{ij}^{L*} (m_b^2 C_0 + m_b^2 C_1 + \hat{t} C_2) \\
& - m_{\hat{\chi}_j^0} O_{ij}^{R*} (m_b^2 C_1 + \hat{t} C_0 + \hat{t} C_2)) + h_t m_b m_t N_{j3}^* V_{i2} (m_{\hat{\chi}_j^0} O_{ij}^{L*} (m_b m_t C_0 \\
& + m_b m_t C_1 + m_b m_t C_2) - m_{\hat{\chi}_i^+} O_{ij}^{R*} (C_0 + C_1 + C_2))] \\
& + h_b m_b (\theta_{k2}^b)^2 N_{j3}^* U_{i2} (m_{\hat{\chi}_i^+} m_{\hat{\chi}_j^0} O_{ij}^{L*} C_0 + O_{ij}^{R*} (2C_{00} - m_b^2 C_1 - m_{b_k}^2 C_0 \\
& - \hat{t} (C_0 + C_1 + C_2)))](m_b^2, m_W^2, \hat{t}, m_{b_k}^2, m_{\hat{\chi}_j^0}^2, m_{\hat{\chi}_i^+}^2) \}, \\
f_6^{V_1(t)} = & \sum_i \frac{gh_b h_{\hat{t}_i}^2 m_b \alpha_{1i} \alpha_{2i} \beta_{11}}{32\sqrt{2}\pi^2 (\hat{t} - m_{\hat{t}}^2)} \{ B_0^{Wbt} + [2(m_{\hat{t}}^2 + \hat{t} + m_{H_i}^2) C_0 \\
& + (2m_b^2 + m_{\hat{t}}^2 + \hat{t}) C_1 + (m_{\hat{t}}^2 + 3\hat{t}) C_2 - 2C_{00}] (m_b^2, m_W^2, \hat{t}, m_{H_i}^2, m_b^2, m_{\hat{t}}^2) \} \\
& + \sum_i \frac{-gh_b h_{\hat{t}_i}^2 m_b \beta_{1i} \beta_{2i} \beta_{11}}{32\sqrt{2}\pi^2 (\hat{t} - m_{\hat{t}}^2)} \{ B_0^{Wbt} + [m_{A_i}^2 C_0 - (m_{\hat{t}}^2 - \hat{t}) (C_1 + C_2) \\
& - 2C_{00}] (m_b^2, m_W^2, \hat{t}, m_{A_i}^2, m_b^2, m_{\hat{t}}^2) \} \\
& + \sum_{i,j} \frac{gh_{\hat{t}_i}^2 \alpha_{1i} \beta_{11} \varphi_{ij}}{16\sqrt{2}\pi^2 (\hat{t} - m_{\hat{t}}^2)} (h_t m_t \beta_{1j} + h_b m_b \beta_{2j}) C_{00} (m_b^2, m_W^2, \hat{t}, m_{\hat{t}_i}^2, m_{H_j^+}^2, m_{H_i}^2) \\
& + \sum_i \frac{-gh_{\hat{t}_i}^2 \beta_{1i} \beta_{11}}{16\sqrt{2}\pi^2 (\hat{t} - m_{\hat{t}}^2)} (h_b m_b \beta_{2i} - h_t m_t \beta_{1i}) C_{00} (m_b^2, m_W^2, \hat{t}, m_{\hat{t}_i}^2, m_{H_i^+}^2, m_{A_i}^2) \\
& + \sum_{i,j} \frac{gh_b h_{\hat{t}_i} \alpha_{2j} \beta_{11} \varphi_{ji}}{16\sqrt{2}\pi^2 (\hat{t} - m_{\hat{t}}^2)} (h_t m_b \beta_{1i} + h_b m_t \beta_{2i}) C_{00} (m_b^2, m_W^2, \hat{t}, m_b^2, m_{H_j}^2, m_{H_i^+}^2) \\
& + \sum_{i,j,k} \frac{-gh_b^2 h_{\hat{t}_i} \beta_{11} \theta_{j1}^b \theta_{i1}^t}{8\sqrt{2}\pi^2 (\hat{t} - m_{\hat{t}}^2)} (m_t N_{k3}^* N_{k4} \theta_{j2}^b \theta_{i2}^t - m_b N_{k3} N_{k4}^* \theta_{j1}^b \theta_{i1}^t) C_{00} \\
& (m_b^2, m_W^2, \hat{t}, m_{\hat{\chi}_k^0}^2, m_{b_j}^2, m_{\hat{t}_i}^2) \\
& + \sum_{i,j,k} \frac{gh_{\hat{t}_i}^2 \beta_{11} \theta_{k1}^t N_{i4}^*}{16\pi^2 (\hat{t} - m_{\hat{t}}^2)} \{ h_b m_b \theta_{k1}^t O_{ji}^{R*} U_{j2} B_0^{W\hat{\chi}_j^+ \hat{\chi}_i^0} + [-h_b m_b m_{\hat{\chi}_i^0} m_{\hat{\chi}_j^+} \theta_{k1}^t O_{ji}^{L*} U_{j2} C_0 \\
& + h_b m_b \theta_{k1}^t O_{ji}^{R*} U_{j2} (-2C_{00} + m_b^2 C_1 + \hat{t} C_0 + \hat{t} C_1 + 2\hat{t} C_2 + m_{\hat{t}_{k1}}^2 C_0) \\
& - h_t m_{\hat{\chi}_i^0} \theta_{k2}^t O_{ji}^{L*} V_{j2} (m_b^2 C_0 + m_b^2 C_1 + \hat{t} C_2) + h_t m_{\hat{\chi}_j^+} \theta_{k2}^t O_{ji}^{R*} V_{j2} (m_b^2 C_1 + \hat{t} C_0 \\
& + \hat{t} C_2)] (m_b^2, m_W^2, \hat{t}, m_{\hat{t}_k}^2, m_{\hat{\chi}_j^+}^2, m_{\hat{\chi}_i^0}^2) \} \\
& + \sum_{i,j,k} \frac{gh_{\hat{t}_i}^2 m_{\hat{t}_i} \beta_{11} \theta_{k2}^t N_{j4}}{16\pi^2 (\hat{t} - m_{\hat{t}}^2)} \{ -h_t \theta_{k2}^t O_{ji}^{L*} V_{j2} B_0^{W\hat{\chi}_j^+ \hat{\chi}_i^0} + [h_b m_b \theta_{k1}^t U_{j2} (m_{\hat{\chi}_i^0} O_{ji}^{R*} \\
& - m_{\hat{\chi}_j^+} O_{ji}^{L*}) (C_0 + C_1 + C_2) - h_t \theta_{k2}^t O_{ji}^{L*} V_{j2} (2C_{00} - m_b^2 C_0 - 2m_b^2 C_1 - m_b^2 C_2
\end{aligned}$$

$$\begin{aligned}
& -\hat{t}C_2 - m_{\hat{t}_k}^2 C_0 + m_{\tilde{\chi}_i^0} m_{\tilde{\chi}_j^+} C_0)(m_b^2, m_W^2, \hat{t}, m_{\hat{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2)\} \\
& - \sum_{i,j,k} \frac{gh_b h_t \beta_{11}}{16\pi^2(\hat{t} - m_{\hat{t}_k}^2)} \{[-h_b(\theta_{k2}^b)^2 N_{j3}^* U_{i2} O_{ij}^{R*} + h_t(\theta_{k1}^b)^2 N_{j3} V_{i2} O_{ij}^{L*}] B_0^{W\tilde{\chi}_i^+ \tilde{\chi}_j^0} \\
& - [h_b m_t (\theta_{k2}^b)^2 N_{j3}^* U_{i2} (O_{ij}^{R*} (-2C_{00} + m_b^2(C_0 + 2C_1 + C_2) + m_{b_k}^2 C_0 + \hat{t}C_2) \\
& - m_{\tilde{\chi}_i^+} m_{\tilde{\chi}_j^0} O_{ij}^{L*} C_0) + \theta_{k1}^b \theta_{k2}^b (h_t N_{j3}^* V_{i2} (m_{\tilde{\chi}_i^+} O_{ij}^{R*} (m_b^2 C_0 + m_b^2 C_1 + \hat{t}C_2) \\
& - m_{\tilde{\chi}_j^0} O_{ij}^{L*} (m_b^2 C_1 + \hat{t}C_0 + \hat{t}C_2)) + h_b m_b m_t N_{j3} U_{i2} (m_{\tilde{\chi}_j^0} O_{ij}^{R*} (m_b m_t C_0 \\
& + m_b m_t C_1 + m_b m_t C_2) - m_{\tilde{\chi}_i^+} O_{ij}^{R*} (C_0 + C_1 + C_2)))] \\
& + h_t m_b (\theta_{k1}^b)^2 N_{j3} V_{i2} (m_{\tilde{\chi}_i^+} m_{\tilde{\chi}_j^0} O_{ij}^{R*} C_0 + O_{ij}^{L*} (2C_{00} - m_b^2 C_1 - m_{\hat{t}_k}^2 C_0 \\
& - \hat{t}(C_0 + C_1 + C_2)))](m_b^2, m_W^2, \hat{t}, m_{\hat{t}_k}^2, m_{\tilde{\chi}_j^0}^2, m_{\tilde{\chi}_i^+}^2)\}, \\
f_7^{V_1(t)} &= \sum_i \frac{-gh_b^2 h_t m_t \alpha_{1i} \alpha_{2i} \beta_{21}}{32\sqrt{2}\pi^2(\hat{t} - m_{\hat{t}_k}^2)} (2C_2 + C_{12} + C_{22})(m_b^2, m_W^2, \hat{t}, m_{H_i}^2, m_b^2, m_{\hat{t}_k}^2) \\
& + \sum_i \frac{-gh_b^2 h_t m_t \beta_{1i} \beta_{2i} \beta_{21}}{16\sqrt{2}\pi^2(\hat{t} - m_{\hat{t}_k}^2)} (C_{12} + C_{22})(m_b^2, m_W^2, \hat{t}, m_{A_i}^2, m_b^2, m_{\hat{t}_k}^2) \\
& + \sum_{i,j} \frac{gh_b h_t \alpha_{1i} \beta_{21} \varphi_{ij}}{16\sqrt{2}\pi^2(\hat{t} - m_{\hat{t}_k}^2)} [h_t m_b \beta_{1j} (C_1 + C_{11} + C_{12}) + h_b m_t \beta_{2j} (C_0 + C_1 \\
& - C_{12} - C_{22})](m_b^2, m_W^2, \hat{t}, m_{H_i}^2, m_{H_j^+}^2, m_{H_i}^2) \\
& + \sum_i \frac{gh_b h_t \beta_{1i} \beta_{21}}{16\sqrt{2}\pi^2(\hat{t} - m_{\hat{t}_k}^2)} [h_t m_b \beta_{1i} (C_1 + C_{11} + C_{12}) + h_b m_t \beta_{2i} (C_0 + C_1 + 2C_2 \\
& + C_{12} + C_{22})](m_b^2, m_W^2, \hat{t}, m_{H_i}^2, m_{A_i}^2) \\
& + \sum_{i,j} \frac{-gh_b^2 \alpha_{2j} \beta_{21} \varphi_{ji}}{16\sqrt{2}\pi^2(\hat{t} - m_{\hat{t}_k}^2)} [h_t m_t \beta_{1i} (C_2 + C_{12} + C_{22}) + h_b m_b \beta_{2i} (C_0 + C_2 - C_{11} \\
& - C_{12})](m_b^2, m_W^2, \hat{t}, m_b^2, m_{H_j}^2, m_{H_i^+}^2) \\
& + \sum_{i,j,k} \frac{-gh_b^2 h_t \beta_{21} \theta_{j1}^b \theta_{i1}^t}{8\sqrt{2}\pi^2(\hat{t} - m_{\hat{t}_k}^2)} [m_b N_{k3}^* N_{k4} \theta_{j2}^b \theta_{i2}^t (C_2 + C_{11} + C_{12}) \\
& - N_{k3} N_{k4} \theta_{j1}^b \theta_{i2}^t m_{\tilde{\chi}_k^0} (C_0 + C_1 + C_2) + m_t N_{k3} N_{k4} \theta_{j1}^b \theta_{i1}^t (C_2 + C_{12} \\
& + C_{22})](m_b^2, m_W^2, \hat{t}, m_{\tilde{\chi}_k^0}^2, m_{b_j}^2, m_{\hat{t}_k}^2) \\
& - \sum_{i,j,k} \frac{gh_b h_t \beta_{12} \theta_{k2}^t N_{i4}}{8\pi^2(\hat{t} - m_{\hat{t}_k}^2)} [h_b m_{\tilde{\chi}_j^+} \theta_{k1}^t O_{ji}^{L*} U_{j2} C_1 + h_b m_{\tilde{\chi}_i^0} \theta_{k1}^t O_{ji}^{R*} U_{j2} C_2 \\
& + h_t m_b \theta_{k2}^t O_{ji}^{L*} V_{j2} (C_1 + C_{11} + C_{12})](m_b^2, m_W^2, \hat{t}, m_{\hat{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2) \\
& - \sum_{i,j,k} \frac{gh_b^2 h_t m_t \beta_{12} (\theta_{k1}^t)^2}{8\pi^2(\hat{t} - m_{\hat{t}_k}^2)} O_{ji}^{R*} U_{j2} N_{i4}^* (C_{12} + C_2 + C_{22})(m_b^2, m_W^2, \hat{t}, m_{\hat{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2)
\end{aligned}$$

$$\begin{aligned}
& - \sum_{i,j,k} \frac{gh_b^2 \beta_{21}}{8\pi^2(\hat{t} - m_t^2)} [h_t m_t (\theta_{k1}^b)^2 N_{j3} V_{i2} O_{ij}^{L*} (C_{12} + C_2 + C_{22}) \\
& + h_b \theta_{k1}^b \theta_{k2}^b N_{j3} U_{i2} (m_{\tilde{\chi}_i^+} O_{ij}^{L*} C_2 + m_{\tilde{\chi}_j^0} O_{ij}^{R*} C_1) \\
& + h_b m_b (\theta_{k2}^b)^2 N_{j3} U_{i2} O_{ij}^{R*} (C_{11} + C_1 + C_{12})] (m_b^2, m_W^2, \hat{t}, m_{\tilde{b}_k}^2, m_{\tilde{\chi}_j^0}^2, m_{\tilde{\chi}_i^+}^2), \\
f_8^{V_1(t)} = & \sum_i \frac{-gh_b h_t^2 m_b \alpha_{1i} \alpha_{2i} \beta_{11}}{32\sqrt{2}\pi^2(\hat{t} - m_t^2)} (2C_1 + C_{11} + C_{12}) (m_b^2, m_W^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2) \\
& + \sum_i \frac{gh_b h_t^2 m_b \beta_{1i} \beta_{2i} \beta_{11}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} (C_{11} + C_{12}) (m_b^2, m_W^2, \hat{t}, m_{A_i}^2, m_b^2, m_t^2) \\
& + \sum_{i,j} \frac{gh_t^2 \alpha_{1i} \beta_{11} \varphi_{ij}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} [h_b m_b \beta_{2j} (C_1 + C_{11} + C_{12}) + h_t m_t \beta_{1j} (C_0 + C_1 \\
& - C_{12} - C_{22})] (m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_j^+}^2, m_{H_i}^2) \\
& + \sum_i \frac{-gh_t^2 \beta_{1i} \beta_{11}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} [h_b m_b \beta_{2i} (C_1 + C_{11} + C_{12}) + h_t m_t \beta_{1i} (C_0 + C_1 + 2C_2 \\
& + C_{12} + C_{22})] (m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_i^+}^2, m_{A_i}^2) \\
& + \sum_{i,j} \frac{-gh_b h_t \alpha_{2j} \beta_{11} \varphi_{ji}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} [h_b m_t \beta_{2i} (C_2 + C_{11} + C_{22}) + h_t m_b \beta_{1i} (C_0 + C_2 - C_{11} \\
& - C_{12})] (m_b^2, m_W^2, \hat{t}, m_b^2, m_{H_j}^2, m_{H_i^+}^2) \\
& + \sum_{i,j,k} \frac{gh_b h_t^2 \beta_{11} \theta_{j1}^b \theta_{i1}^t}{8\sqrt{2}\pi^2(\hat{t} - m_t^2)} [m_t N_{k3}^* N_{k4} \theta_{j2}^b \theta_{i2}^t (C_2 + C_{12} + C_{22}) \\
& - N_{k3}^* N_{k4}^* \theta_{j2}^b \theta_{i1}^t m_{\tilde{\chi}_k^0} (C_0 + C_1 + C_2) + m_b N_{k3} N_{k4}^* \theta_{j1}^b \theta_{i1}^t (C_1 + C_{11} + C_{12})] \\
& (m_b^2, m_W^2, \hat{t}, m_{\tilde{\chi}_k^0}^2, m_b^2, m_{\tilde{t}_i}^2) \\
& - \sum_{i,j,k} \frac{gh_t^2 \beta_{11} \theta_{k1}^t N_{i4}^*}{8\pi^2(\hat{t} - m_t^2)} [h_t m_{\tilde{\chi}_j^+} \theta_{kw}^t O_{ji}^{R*} V_{j2} C_1 + h_t m_{\tilde{\chi}_i^0} \theta_{k2}^t O_{ji}^{L*} V_{j2} C_2 \\
& + h_b m_b \theta_{k1}^t O_{ji}^{R*} U_{j2} (C_1 + C_{11} + C_{12})] (m_b^2, m_W^2, \hat{t}, m_{\tilde{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2) \\
& - \sum_{i,j,k} \frac{gh_t^3 m_t \beta_{11} (\theta_{k2}^t)^2}{8\pi^2(\hat{t} - m_t^2)} O_{ji}^{L*} V_{j2} N_{i4}^* (C_2 + C_{12} + C_{22}) (m_b^2, m_W^2, \hat{t}, m_{\tilde{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2) \\
& + \sum_{i,j,k} \frac{gh_b h_t \beta_{11}}{8\pi^2(\hat{t} - m_t^2)} [h_b m_t (\theta_{k2}^b)^2 N_{j3}^* U_{i2} O_{ij}^{R*} (C_{12} + C_2 + C_{22}) \\
& + h_t \theta_{k1}^b \theta_{k2}^b N_{j3}^* V_{i2} (m_{\tilde{\chi}_i^+} O_{ij}^{R*} C_2 + m_{\tilde{\chi}_j^0} O_{ij}^{L*} C_1) \\
& + h_t m_b (\theta_{k1}^b)^2 N_{j3} V_{i2} O_{ij}^{L*} (C_{11} + C_1 + C_{12})] (m_b^2, m_W^2, \hat{t}, m_{\tilde{b}_k}^2, m_{\tilde{\chi}_j^0}^2, m_{\tilde{\chi}_i^+}^2), \\
f_{11}^{V_1(t)} = & \sum_i \frac{gh_b h_t^2 \alpha_{1i} \alpha_{2i} \beta_{11}}{32\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{B_0^{Wbt} + [m_{H_i}^2 C_0 + 2m_b^2 C_1 + (m_t^2 + \hat{t}) C_2 \\
& - 2C_{00}] (m_b^2, m_W^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2)\}
\end{aligned}$$

$$\begin{aligned}
& + \sum_i \frac{-gh_b h_t^2 m_b \beta_{1i} \beta_{2i} \beta_{11}}{32\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{B_0^{Wbt} + [m_{A_i}^2 C_0 - (m_t^2 - \hat{t})C_2 \\
& - 2C_{00}](m_b^2, m_W^2, \hat{t}, m_{A_i}^2, m_b^2, m_t^2)\} \\
& + \sum_{i,j} \frac{gh_b h_t^2 \alpha_{1i} \beta_{11} \beta_{2j} \varphi_{ij}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} C_{00}(m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_j^+}^2, m_{H_i}^2) \\
& + \sum_i \frac{-gh_b h_t^2 \beta_{1i} \beta_{2i} \beta_{11}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} C_{00}(m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_i^+}^2, m_{A_i}^2) \\
& + \sum_{i,j} \frac{gh_b h_t^2 \alpha_{2j} \beta_{11} \beta_{1i} \varphi_{ji}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} C_{00}(m_b^2, m_W^2, \hat{t}, m_b^2, m_{H_j}^2, m_{H_i^+}^2) \\
& + \sum_{i,j,k} \frac{gh_b h_t^2 \beta_{11} (\theta_{j1}^b)^2 (\theta_{i1}^t)^2}{8\sqrt{2}\pi^2(\hat{t} - m_t^2)} N_{k3} N_{k4}^* C_{00}(m_b^2, m_W^2, \hat{t}, m_{\tilde{\chi}_k^0}^2, m_{b_j}^2, m_{t_i}^2) \\
& + \sum_{i,j,k} \frac{gh_t^2 \beta_{11} \theta_{k1}^t N_{i4}^*}{16\pi^2(\hat{t} - m_t^2)} \{h_b \theta_{k1}^t O_{ji}^{R*} U_{j2} B_0^{Wbt} + [-h_b m_{\tilde{\chi}_i^0} m_{\tilde{\chi}_j^+} \theta_{k1}^t O_{ji}^{L*} U_{j2} C_0 \\
& + h_b \theta_{k1}^t O_{ji}^{R*} U_{j2} (-2C_{00} + m_b^2 C_1 + \hat{t} C_2 + m_{t_k}^2 C_0) + h_t m_b \theta_{k2}^t V_{j2} (m_{\tilde{\chi}_j^+} O_{ji}^{L*} \\
& - m_{\tilde{\chi}_i^0} O_{ji}^{R*}) C_1](m_b^2, m_W^2, \hat{t}, m_{t_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2)\} \\
& - \sum_{i,j,k} \frac{gh_t^2 \beta_{11} \theta_{k2}^t N_{i4}}{16\pi^2(\hat{t} - m_t^2)} [h_b m_{\tilde{\chi}_j^+} \theta_{k1}^t O_{ji}^{L*} U_{j2} (C_0 + C_2) - h_b m_{\tilde{\chi}_i^0} \theta_{k1}^t O_{ji}^{R*} U_{j2} C_2 \\
& + h_t m_b \theta_{k2}^t O_{ji}^{L*} V_{j2} (C_0 + C_1 + C_2)](m_b^2, m_W^2, \hat{t}, m_{t_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2) \\
& + \sum_{i,j,k} \frac{gh_b h_t \beta_{11}}{16\pi^2(\hat{t} - m_t^2)} \{-h_t (\theta_{k1}^b)^2 N_{j3} V_{i2} O_{ij}^{L*} B_0^{W\tilde{\chi}_i^+ \tilde{\chi}_j^0} + [-h_t (\theta_{k1}^b)^2 N_{j3} V_{i2} (m_{\tilde{\chi}_i^+} m_{\tilde{\chi}_j^0} O_{ij}^{R*} C_0 \\
& + O_{ij}^{L*} (-2C_{00} + m_b^2 C_1 + m_{b_k}^2 C_0 + \hat{t} C_2)) - \theta_{k1}^b \theta_{k2}^b (h_b m_t N_{j3} U_{i2} (m_{\tilde{\chi}_i^+} O_{ij}^{L*} C_2 \\
& - O_{ij}^{R*} (m_{\tilde{\chi}_j^0} C_0 + m_{\tilde{\chi}_i^+} C_2)) + h_t m_b N_{j3} V_{i2} (m_{\tilde{\chi}_j^0} O_{ij}^{L*} C_1 - O_{ij}^{R*} (m_{\tilde{\chi}_i^+} C_0 + m_{\tilde{\chi}_j^0} C_1)))] \\
& + h_b m_b m_t (\theta_{k2}^b)^2 N_{j3} U_{i2} O_{ij}^{R*} (C_0 + C_1 + C_2)](m_b^2, m_W^2, \hat{t}, m_{b_k}^2, m_{\tilde{\chi}_j^0}^2, m_{\tilde{\chi}_i^+}^2)\}, \\
f_{12}^{V_1(t)} & = \sum_i \frac{-gh_b^2 h_t \alpha_{1i} \alpha_{2i} \beta_{21}}{32\sqrt{2}\pi^2(\hat{t} - m_t^2)} m_b m_t (2C_0 + C_1 + C_2) (m_b^2, m_W^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2) \\
& + \sum_{i,j} \frac{gh_b h_t^2 \alpha_{1i} \beta_{21} \beta_{1j} \varphi_{ij}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} C_{00}(m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_j^+}^2, m_{H_i}^2) \\
& + \sum_i \frac{-gh_b h_t^2 \beta_{1i}^2 \beta_{21}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} C_{00}(m_b^2, m_W^2, \hat{t}, m_t^2, m_{H_i^+}^2, m_{A_i}^2) \\
& + \sum_{i,j} \frac{gh_b^3 \alpha_{2j} \beta_{21} \beta_{2i} \varphi_{ji}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} C_{00}(m_b^2, m_W^2, \hat{t}, m_b^2, m_{H_j}^2, m_{H_i^+}^2) \\
& + \sum_{i,j,k} \frac{-gh_b^2 h_t \beta_{21} \theta_{j1}^b \theta_{j2}^b \theta_{i1}^t \theta_{i2}^t}{8\sqrt{2}\pi^2(\hat{t} - m_t^2)} N_{k3}^* N_{k4} C_{00}(m_b^2, m_W^2, \hat{t}, m_{\tilde{\chi}_k^0}^2, m_{b_j}^2, m_{t_i}^2)
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i,j,k} \frac{gh_t^2 \beta_{11} \theta_{k1}^t N_{i4}^*}{16\pi^2 (\hat{t} - m_t^2)} \{ h_b \theta_{k1}^t O_{ji}^{R*} U_{j2} B_0^{Wbt} + [-h_b m_{\tilde{\chi}_i^0} m_{\tilde{\chi}_j^+} \theta_{k1}^t O_{ji}^{L*} U_{j2} C_0 \\
& + h_b \theta_{k1}^t O_{ji}^{R*} U_{j2} (-2C_{00} + m_b^2 C_1 + \hat{t} C_2 + m_{\tilde{t}_k}^2 C_0) + h_t m_b \theta_{k2}^t V_{j2} (m_{\tilde{\chi}_j^+} O_{ji}^{L*} \\
& - m_{\tilde{\chi}_i^0} O_{ji}^{R*}) C_1] (m_b^2, m_W^2, \hat{t}, m_{\tilde{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2) \} \\
& - \sum_{i,j,k} \frac{gh_b h_t \beta_{12} \theta_{k1}^t N_{i4}^*}{16\pi^2 (\hat{t} - m_t^2)} [h_t m_{\tilde{\chi}_j^+} \theta_{k2}^t O_{ji}^{R*} V_{j2} (C_0 + C_2) - h_t m_{\tilde{\chi}_i^0} \theta_{k2}^t O_{ji}^{L*} V_{j2} C_2 \\
& + h_b m_b \theta_{k1}^t O_{ji}^{R*} U_{j2} (C_0 + C_1 + C_2)] (m_b^2, m_W^2, \hat{t}, m_{\tilde{t}_k}^2, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2) \\
& - \sum_{i,j,k} \frac{gh_b^2 \beta_{21}}{16\pi^2 (\hat{t} - m_t^2)} \{ -h_b (\theta_{k2}^b)^2 N_{j3}^* U_{i2} O_{ij}^{R*} B_0^{W\tilde{\chi}_i^+ \tilde{\chi}_j^0} + [-h_b (\theta_{k2}^b)^2 N_{j3}^* U_{i2} (m_{\tilde{\chi}_i^+} m_{\tilde{\chi}_j^0} O_{ij}^{L*} C_0 \\
& + O_{ij}^{R*} (-2C_{00} + m_b^2 C_1 + m_{\tilde{b}_k}^2 C_0 + \hat{t} C_2)) - \theta_{k1}^b \theta_{k2}^b (h_t m_t N_{j3}^* V_{i2} (m_{\tilde{\chi}_i^+} O_{ij}^{R*} C_2 \\
& - O_{ij}^{L*} (m_{\tilde{\chi}_j^0} C_0 + m_{\tilde{\chi}_i^+} C_2)) + h_b m_b N_{j3}^* U_{i2} (m_{\tilde{\chi}_j^0} O_{ij}^{R*} C_1 - O_{ij}^{L*} (m_{\tilde{\chi}_i^+} C_0 + m_{\tilde{\chi}_j^0} C_1))] \\
& + h_t m_b m_t (\theta_{k1}^b)^2 N_{j3}^* V_{i2} O_{ij}^{L*} (C_0 + C_1 + C_2)] (m_b^2, m_W^2, \hat{t}, m_{\tilde{b}_k}^2, m_{\tilde{\chi}_j^0}^2, m_{\tilde{\chi}_i^+}^2) \}; \\
f_2^{s(t)} & = \sum_i \frac{-gh_b h_t^2 \alpha_{1i}^2 \beta_{21}}{16\sqrt{2}\pi^2 (\hat{t} - m_t^2)^2} [2m_t^2 B_0^{\hat{t}tH_i} - (m_t^2 + \hat{t}) B_1^{\hat{t}tH_i}] \\
& + \sum_i \frac{gh_b h_t^2 \beta_{1i}^2 \beta_{21}}{16\sqrt{2}\pi^2 (\hat{t} - m_t^2)^2} [2m_t^2 B_0^{\hat{t}tA_i} + (m_t^2 + \hat{t}) B_1^{\hat{t}tA_i}] \\
& + \sum_i \frac{gh_b \beta_{21}}{8\sqrt{2}\pi^2 (\hat{t} - m_t^2)^2} [2h_b h_t m_b m_t \beta_{1i} \beta_{2i} B_0^{\hat{t}bH_i^+} + (h_t^2 m_t^2 \beta_{1i} + h_b^2 \hat{t} \beta_{2i})^2 B_1^{\hat{t}bH_i^+}] \\
& + \sum_{i,j} \frac{-gh_b h_t^2 \beta_{21}}{8\sqrt{2}\pi^2 (\hat{t} - m_t^2)^2} \{ m_t \theta_{i1}^t \theta_{i2}^t (N_{j4}^2 + N_{j4}^{*2}) B_0^{\hat{t}\tilde{\chi}_j^0 \tilde{t}_i} - [m_t^2 (\theta_{i1}^t)^2 \\
& + \hat{t} (\theta_{i2}^t)^2] |N_{j4}|^2 B_1^{\hat{t}\tilde{\chi}_j^0 \tilde{t}_i} \} \\
& + \sum_{i,j} \frac{-gh_b \beta_{21}}{8\sqrt{2}\pi^2 (\hat{t} - m_t^2)^2} [-h_b^2 m_t^2 (\theta_{i2}^b)^2 |U_{j2}|^2 B_1^{\hat{t}\tilde{\chi}_j^+ \tilde{b}_i} + h_b h_t m_t \theta_{i1}^b \theta_{i2}^b (U_{j2} V_{j2} \\
& + U_{j2}^* V_{j2}^*) B_0^{\hat{t}\tilde{\chi}_j^+ \tilde{b}_i} - h_t^2 \hat{t} (\theta_{i1}^b)^2 |V_{j2}|^2 B_1^{\hat{t}\tilde{\chi}_j^+ \tilde{b}_i}], \\
f_5^{s(t)} & = -\frac{1}{2} m_b f_2^{s(t)}, \\
f_6^{s(t)} & = \sum_i \frac{-gh_t^3 m_t \alpha_{1i}^2 \beta_{11}}{32\sqrt{2}\pi^2 (\hat{t} - m_t^2)^2} [-(m_t^2 - \hat{t}) B_0^{\hat{t}tH_i} + 2\hat{t} B_1^{\hat{t}tH_i}] \\
& + \sum_i \frac{-gh_t^3 m_t \beta_{1i}^2 \beta_{11}}{32\sqrt{2}\pi^2 (\hat{t} - m_t^2)^2} [(m_t^2 + \hat{t}) B_0^{\hat{t}tA_i} + 2\hat{t} B_1^{\hat{t}tA_i}] \\
& + \sum_i \frac{-gh_t \beta_{11}}{16\sqrt{2}\pi^2 (\hat{t} - m_t^2)^2} [h_b h_t m_b \beta_{1i} \beta_{2i} (m_t^2 + \hat{t}) B_0^{\hat{t}bH_i^+} + (h_t^2 \beta_{1i}^2 + h_b^2 \beta_{2i}^2) m_t \hat{t} B_1^{\hat{t}bH_i^+}] \\
& + \sum_{i,j} \frac{-gh_t^3 \beta_{11}}{16\sqrt{2}\pi^2 (\hat{t} - m_t^2)^2} [-\theta_{i1}^t \theta_{i2}^t (m_t^2 N_{j4}^2 + \hat{t} N_{j4}^{*2}) B_0^{\hat{t}\tilde{\chi}_j^0 \tilde{t}_i} + m_t \hat{t} |N_{j4}|^2 B_1^{\hat{t}\tilde{\chi}_j^0 \tilde{t}_i}]
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i,j} \frac{-gh_t \beta_{11}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)^2} [h_b^2 m_t \hat{t} (\theta_{i2}^b)^2 |U_{j2}|^2 B_1^{\hat{t}\bar{\chi}_j^+ \bar{b}_i} - h_b h_t \theta_{i1}^b \theta_{i2}^b (m_t^2 U_{j2} V_{j2} \\
& + \hat{t} U_{j2}^* V_{j2}^*) B_0^{\hat{t}\bar{\chi}_j^+ \bar{b}_i} + h_t^2 m_t \hat{t} (\theta_{i1}^b)^2 |V_{j2}|^2 B_1^{\hat{t}\bar{\chi}_j^+ \bar{b}_i}], \\
f_{12}^{s(t)} &= -\frac{1}{2} f_2^{s(t)}; \\
f_2^{V_2(t)} &= \sum_i \frac{gh_b h_t \alpha_{1i} \alpha_{2i}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{h_t \beta_{11} B_0^{H+bt} + [h_t \beta_{11} (m_{H_i}^2 C_0 + 2m_b^2 C_1 + m_t^2 C_2 \\
& + \hat{t} C_2) - h_b m_b m_t \beta_{21} (4C_0 + 2C_1 + 2C_2)] (m_b^2, m_{H^+}^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2)\} \\
& + \sum_i \frac{gh_b h_t^2 \beta_{1i} \beta_{11} \beta_{2i}}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{B_0^{H+bt} + [m_{A_i}^2 C_0 \\
& + (\hat{t} - m_t^2) C_2] (m_b^2, m_{H^+}^2, \hat{t}, m_{A_i}^2, m_b^2, m_t^2)\} \\
& + \sum_{i,j,k} \sum_{i',j'} \frac{-gh_b h_t \theta_{jj'}^b \theta_{ii'}^t}{8\sqrt{2}\pi^2(\hat{t} - m_t^2)} (h_b \Theta_{j'i'1}^5 + h_t \Theta_{j'i'2}^6) (m_t N_{k3}^* N_{k4} \theta_{j1}^b \theta_{i1}^t C_2 \\
& - m_{\bar{\chi}_k^0} N_{k3}^* N_{k4}^* \theta_{j1}^b \theta_{j2}^t C_0 + m_b N_{k3} N_{k4}^* \theta_{j2}^b \theta_{i2}^t C_1) (m_b^2, m_{H^+}^2, \hat{t}, m_{\bar{\chi}_k^0}^2, m_{b_j}^2, m_{t_i}^2), \\
f_5^{V_2(t)} &= -\frac{m_b}{2} f_2^{V_2(t)}, \\
f_6^{V_2(t)} &= \sum_i \frac{-gh_b h_t \alpha_{1i} \alpha_{2i}}{32\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{h_b m_t \beta_{21} B_0^{H+bt} + [h_b m_t \beta_{21} (m_{H_i}^2 C_0 \\
& + 2\hat{t} C_2 + 2m_b^2 C_1) - h_t m_b \beta_{11} ((m_t^2 + \hat{t})(2C_0 + C_1) \\
& + 2\hat{t} C_2)] (m_b^2, m_{H^+}^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2)\} \\
& + \sum_i \frac{-gh_b h_t \beta_{1i} \beta_{2i}}{32\sqrt{2}\pi^2(\hat{t} - m_t^2)} \{h_b m_t \beta_{21} B_0^{H+bt} + [h_b m_t m_{A_i}^2 \beta_{21} C_0 \\
& + h_t m_b \beta_{11} (m_t^2 - \hat{t}) C_1] (m_b^2, m_{H^+}^2, \hat{t}, m_{A_i}^2, m_b^2, m_t^2)\} \\
& + \sum_{i,j,k} \sum_{i',j'} \frac{-gh_b h_t \theta_{jj'}^b \theta_{ii'}^t}{16\sqrt{2}\pi^2(\hat{t} - m_t^2)} (h_b \Theta_{j'i'1}^5 + h_t \Theta_{j'i'2}^6) (m_b m_t N_{k3}^* N_{k4} \theta_{j1}^b \theta_{i1}^t C_1 \\
& - m_t m_{\bar{\chi}_k^0} N_{k3} N_{k4} \theta_{j2}^b \theta_{j1}^t C_0 + \hat{t} N_{k3} N_{k4}^* \theta_{j2}^b \theta_{i2}^t C_2) (m_b^2, m_{H^+}^2, \hat{t}, m_{\bar{\chi}_k^0}^2, m_{b_j}^2, m_{t_i}^2), \\
f_{12}^{V_2(t)} &= -\frac{1}{2} f_2^{V_2(t)};
\end{aligned}$$

$$\begin{aligned}
f_1^{(b)} &= \sum_i \frac{gh_b^2 \alpha_{2i}^2}{16\sqrt{2}\pi^2} \{h_b \beta_{21} C_0 (m_{H^+}^2, m_W^2, \hat{s}, m_b^2, m_t^2, m_b^2) + [-h_t m_b m_t \beta_{11} (D_{13} \\
& + D_{23}) - h_b \beta_{21} 2D_{00} + h_b m_b^2 \beta_{21} (2D_3 - D_{11} - D_{12} + D_{13} + D_{23}) \\
& + h_b m_{H^+}^2 \beta_{21} (D_{13} + D_{23}) - h_b \hat{t} \beta_{21} (D_{12} + D_{13} + D_{22} + D_{23}) \\
& + h_b m_{H_i}^2 \beta_{21} D_0] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_t^2, m_b^2)\} \\
& + \sum_i \frac{gh_b^2 \beta_{2i}^2}{16\sqrt{2}\pi^2} \{-h_b \beta_{21} C_0 (m_{H^+}^2, m_W^2, \hat{s}, m_b^2, m_t^2, m_b^2) + [-h_t m_b m_t \beta_{11} (D_{13}
\end{aligned}$$

$$\begin{aligned}
& +D_{23}) + 2h_b\beta_{21}D_{00} + h_b m_b^2 \beta_{21}(D_{11} + D_{12} + D_{13} + D_{23}) \\
& - h_b m_{H^+}^2 \beta_{21}(D_{13} + D_{23}) + h_b \hat{t} \beta_{21}(D_{12} + D_{13} + D_{22} + D_{23}) \\
& - h_b m_{A_i}^2 \beta_{21} D_0 \{m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_t^2, m_b^2\} \\
& + \sum_i \frac{gh_b m_b \beta_{2i}}{8\sqrt{2}\pi^2} [h_t^2 m_b \beta_{11} \beta_{1i}(D_1 + D_{11} + D_{12} + D_{13}) - h_b h_t m_t \beta_{11} \beta_{2i}(D_1 \\
& + D_{12} + D_{13}) + h_b^2 m_b \beta_{21} \beta_{2i}(D_{12} + D_{13})] \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H^+}^2, m_t^2, m_b^2, m_t^2) \\
& + \sum_{i,j,k,l,i',k'} \frac{\sqrt{2}gh_b^2}{16\pi^2} N_{l3} \theta_{ii'}^b \theta_{j1}^b \theta_{k1}^t \theta_{kk'}^t (h_b \Theta_{i'k'1}^5 + h_t \Theta_{i'k'1}^6) [m_b \theta_{i1}^b \theta_{j1}^b N_{l3}^* (D_3 + D_{13} \\
& + D_{23}) - m_{\tilde{\chi}_l^0} N_{l3} \theta_{i2}^b \theta_{j1}^b (D_0 + D_1 + D_2) + m_b \theta_{i2}^b \theta_{j2}^b N_{l3}^* (D_1 + D_2 + D_{11} \\
& + 2D_{12} + D_{22})] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{\tilde{\chi}_l^0}^2, m_{\tilde{b}_i}^2, m_{\tilde{t}_k}^2, m_{\tilde{b}_j}^2) \\
& + \sum_{i,j,k,l,j',l'} \frac{g\theta_{l1}^b \theta_{l'l'}^b \theta_{i1}^t \theta_{jj'}^t}{8\sqrt{2}\pi^2} (h_b \Theta_{l'j'1}^5 + h_t \Theta_{l'j'1}^6) [h_b^2 m_b \theta_{i1}^t \theta_{j1}^t U_{k2}^2 (D_{12} + D_{23}) \\
& - h_b h_t m_{\tilde{\chi}_k^+} \theta_{i2}^t \theta_{j1}^t U_{k2} V_{k2} D_3 + h_t^2 m_b \theta_{i2}^t \theta_{j2}^t V_{k2}^2 (D_3 + D_{33})] \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{\tilde{\chi}_k^+}^2, m_{\tilde{t}_i}^2, m_{\tilde{b}_i}^2, m_{\tilde{t}_j}^2) \\
& + \sum_{i,j} \frac{gh_b \alpha_{2i} \varphi_{ij}}{16\sqrt{2}\pi^2} \{-h_b^2 \beta_{12} \beta_{2j} C_2(m_b^2, m_{H^+}^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2) \\
& + [h_t^2 m_b^2 \beta_{11} \beta_{1j}(D_{23} + 2D_3 + 2D_{33}) - h_b h_t m_b m_t \beta_{11} \beta_{2j}(D_{23} + 2D_3) \\
& + h_b h_t m_b m_t \beta_{12} \beta_{1j} D_{33} - h_b^2 \beta_{12} \beta_{2j}(m_b^2(D_{23} + D_{33}) + m_W^2 D_{13} + \hat{u} D_{23} \\
& + m_{H_j^+}^2 D_3)](m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H_j^+}^2, m_{H_i}^2, m_b^2, m_t^2)\} \\
& + \sum_i \frac{gh_b \beta_{2i}}{16\sqrt{2}\pi^2} \{h_b^2 \beta_{21} \beta_{2i}(C_0 + C_1 + C_2)(m_{H^+}^2, m_b^2, \hat{t}, m_t^2, m_b^2, m_{A_i}^2) \\
& + [h_b^2 \beta_{21} \beta_{2i}(m_b^2(D_{12} - D_{11}) + m_W^2 D_{13} - \hat{u} D_{12} - m_{H^+}^2 D_1) \\
& + h_b h_t m_b m_t (\beta_{21} \beta_{1i} D_{11} - \beta_{11} \beta_{2i} D_{12}) + h_t^2 m_b^2 \beta_{11} \beta_{1i} D_{12}] \\
& (m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H^+}^2, m_t^2, m_b^2, m_{A_i}^2)\} \\
& + \sum_{i,j,k,l,k',l'} \frac{gh_b \theta_{kk'}^b \theta_{ll'}^t}{8\pi^2} (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) [h_b \theta_{k1}^b \theta_{l1}^t U_{j2} (m_b N_{i3}^* O_{ij}^{R*} D_{23} \\
& + m_{\tilde{\chi}_j^+} N_{i3} O_{ij}^{L*} D_3) + h_t m_b \theta_{k2}^b \theta_{l2}^t N_{i3} V_{j2} O_{ij}^{L*} D_{33}] \\
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{\tilde{b}_k}^2, m_{\tilde{t}_i}^2), \\
& f_2^{(b)} = \sum_i \frac{gh_b^2 \alpha_{2i}^2}{16\sqrt{2}\pi^2} [-h_t m_b m_t \beta_{11}(2D_1 + 2D_2 + D_{11} + 2D_{12} + D_{22})
\end{aligned}$$

$$\begin{aligned}
& +h_b m_b^2 \beta_{21} (4D_0 + 6D_1 + 2D_{11} + 4D_{12} + D_{13} + 6D_2 + 2D_{22} \\
& + D_{23} + 2D_3)] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_t^2, m_b^2) \\
& + \sum_i \frac{-gh_b^2 \beta_{2i}^2}{16\sqrt{2}\pi^2} [h_t m_b m_t \beta_{11} (D_{11} + 2D_{12} + D_{22}) \\
& + h_b m_b^2 \beta_{21} (D_{13} + D_{23})] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_t^2, m_b^2) \\
& + \sum_i \frac{gh_b \beta_{2i}}{8\sqrt{2}\pi^2} \{h_t^2 \beta_{11} \beta_{1i} (C_0 + C_1 + C_2) (m_W^2, m_{H^+}^2, \hat{s}, m_t^2, m_b^2, m_t^2) \\
& + [-h_b h_t m_b m_t (\beta_{1i} \beta_{21} D_1 + \beta_{11} \beta_{2i} D_{11}) + h_b^2 m_b^2 \beta_{21} \beta_{2i}^2 (D_1 + D_{11}) \\
& + h_t^2 \beta_{11} \beta_{1i} (-m_b^2 D_{11} + m_W^2 D_{13} - \hat{u} D_{12} - \hat{u} D_{13} \\
& - m_{H_i^+}^2 D_1)] (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H_i^+}^2, m_t^2, m_b^2, m_t^2)\} \\
& + \sum_{i,j,k,l,i',k'} \frac{\sqrt{2}gh_b^2}{16\pi^2} N_{i3}^* \theta_{ii'}^b \theta_{j1}^b \theta_{k1}^t \theta_{kk'}^t (h_b \Theta_{i'k'1}^5 + h_t \Theta_{i'k'1}^6) [m_b \theta_{i1}^b \theta_{j1}^b N_{i3} (D_1 + D_2 \\
& + D_{11} + 2D_{12} + D_{22}) - m_{\tilde{\chi}_i^0} N_{i3}^* \theta_{i1}^b \theta_{j2}^b (D_0 + D_1 + D_2) + m_b \theta_{i2}^b \theta_{j2}^b N_{i3} (D_{13} \\
& + D_{23} + D_3)] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{\tilde{\chi}_i^0}^2, m_{b_i}^2, m_{t_k}^2, m_{b_j}^2) \\
& + \sum_{i,j,k,l,j',l'} \frac{g\theta_{i1}^b \theta_{l'j'}^b \theta_{i1}^t \theta_{jj'}^t}{8\sqrt{2}\pi^2} (h_b \Theta_{l'j'1}^5 + h_t \Theta_{l'j'1}^6) [h_b^2 m_b \theta_{i1}^t \theta_{j1}^t U_{k2}^2 D_{33} \\
& - h_b h_t m_{\tilde{\chi}_k^+} \theta_{i1}^t \theta_{j2}^t U_{k2} V_{k2} D_3 + h_t^2 m_b \theta_{i2}^t \theta_{j2}^t V_{k2}^2 (D_{13} + D_{23})] \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{\tilde{\chi}_k^+}^2, m_{t_i}^2, m_{b_l}^2, m_{t_j}^2) \\
& + \sum_{i,j} \frac{gh_b \alpha_{2i} \varphi_{ij}}{16\sqrt{2}\pi^2} \{h_t^2 \beta_{11} \beta_{1j} C_2 (m_b^2, m_{H^+}^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2) + [-h_b^2 m_b^2 \beta_{12} \beta_{2j} (D_{23} \\
& + 2D_3 + 2D_{33}) - h_b h_t m_b m_t \beta_{11} \beta_{2j} D_{33} + h_b h_t m_b m_t \beta_{12} \beta_{1j} (D_{23} + 2D_3) \\
& + h_t^2 \beta_{11} \beta_{1j} (m_b^2 (D_{23} + D_{33}) + m_W^2 D_{13} + \hat{u} D_{23} \\
& + m_{H_j^+}^2 D_3)] (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H_j^+}^2, m_{H_i}^2, m_b^2, m_t^2)\} \\
& - \sum_i \frac{gh_b \beta_{2i}}{16\sqrt{2}\pi^2} \{h_t^2 \beta_{11} \beta_{1i} (C_0 + C_1 + C_2) (m_{H^+}^2, m_b^2, \hat{t}, m_t^2, m_b^2, m_{A_i}^2) \\
& + [h_t^2 \beta_{11} \beta_{1i} (m_b^2 (D_{12} - D_{11}) + m_W^2 D_{13} - \hat{u} D_{12} - m_{H_i^+}^2 D_1) \\
& + h_b h_t m_b m_t (\beta_{11} \beta_{2i} D_{11} - \beta_{21} \beta_{1i} D_{12}) + h_b^2 \beta_{21} \beta_{2i} D_{12}] \\
& (m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_t^2, m_b^2, m_{A_i}^2)\} \\
& + \sum_{i,j,k,l,k',l'} \frac{gh_b \theta_{kk'}^b \theta_{ll'}^t}{8\pi^2} (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) [h_t \theta_{k2}^b \theta_{l2}^t V_{j2} (m_b N_{i3} O_{ij}^{L*} D_{23} \\
& + m_{\tilde{\chi}_j^+} N_{i3}^* O_{ij}^{R*} D_3) + h_b m_b \theta_{k1}^b \theta_{l1}^t N_{i3}^* U_{j2} O_{ij}^{R*} D_{33}]
\end{aligned}$$

$$\begin{aligned}
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{\tilde{t}_k}^2, m_{\tilde{t}_i}^2), \\
f_3^{(b)} = & \sum_i \frac{gh_b^2 \alpha_{2i}^2}{16\sqrt{2}\pi^2} \{ -h_b m_b^2 \beta_{21} C_2(m_{H^+}^2, m_W^2, \hat{s}, m_b^2, m_t^2, m_b^2) + [h_t m_b m_t \beta_{11} (2D_3 \\
& + D_{33}) - 2h_b m_b^2 \beta_{21} D_{33} + h_b m_W^2 \beta_{21} D_{13} - h_b \hat{t} \beta_{21} (D_{13} + D_{23}) \\
& - h_b m_{H_i}^2 \beta_{21} D_3] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_t^2, m_b^2) \} \\
& + \sum_i \frac{gh_b^2 \beta_{2i}^2}{16\sqrt{2}\pi^2} \{ h_b \beta_{21} C_2(m_{H^+}^2, m_W^2, \hat{s}, m_b^2, m_t^2, m_b^2) \\
& + [h_t m_b m_t \beta_{11} D_{33} - h_b m_W^2 \beta_{21} D_{13} + h_b \hat{t} \beta_{21} (D_{13} + D_{23}) \\
& + h_b m_{A_i}^2 \beta_{21} D_3] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_t^2, m_b^2) \} \\
& + \sum_i \frac{g\beta_{2i}}{8\sqrt{2}\pi^2} [h_t^3 m_b m_t \beta_{11} \beta_{1i} (D_0 + D_1 + D_2 + D_3) - h_b h_t^2 m_b^2 \beta_{11} \beta_{1i} (D_0 + D_1 \\
& + D_{12} + D_{13} + 2D_2 + D_{22} + 2D_{23} + 2D_3 + D_{33}) - h_b h_t^2 m_t^2 \beta_{11} \beta_{1i} (D_0 + D_2 \\
& + D_3) + h_b^2 h_t m_b m_t \beta_{1i} \beta_{21} (D_2 + D_3) + h_b^2 h_t m_b m_t \beta_{11} \beta_{2i} (D_1 + 2D_2 + D_{22} \\
& + 2D_{23} + 2D_3 + D_{33}) - h_b^3 m_b^2 \beta_{21} \beta_{2i} (D_2 + D_{22} + 2D_{23} + D_3 + D_{33})] \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H_i^+}^2, m_t^2, m_b^2, m_t^2) \\
& + \sum_{i,j,k,l,i',k'} \frac{-\sqrt{2}gh_b^2}{16\pi^2} N_{l3} \theta_{ii'}^b \theta_{j1}^b \theta_{k1}^t \theta_{kk'}^t (h_b \Theta_{i'k'1}^5 + h_t \Theta_{i'k'1}^6) [m_b \theta_{i1}^b \theta_{j1}^b N_{l3}^* D_{33} \\
& - m_{\tilde{\chi}_i^0} N_{l3} \theta_{i2}^b \theta_{j1}^b D_3 + m_b \theta_{i2}^b \theta_{j2}^b N_{l3}^* (D_{13} + D_{23})] \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{\tilde{\chi}_i^0}^2, m_{\tilde{t}_k}^2, m_{\tilde{t}_j}^2) \\
& + \sum_{i,j,k,l,j',l'} \frac{g\theta_{i1}^b \theta_{l'j'}^b \theta_{i1}^t \theta_{jj'}^t}{8\sqrt{2}\pi^2} (h_b \Theta_{l'j'1}^5 + h_t \Theta_{l'j'1}^6) [h_b h_t m_{\tilde{\chi}_k^+} \theta_{i2}^t \theta_{j1}^t U_{k2} V_{k2} (D_0 + D_1 + D_2) \\
& - h_t^2 m_b \theta_{i2}^t \theta_{j2}^t V_{k2}^2 (D_{13} + D_{33} + D_3) - h_b^2 m_b \theta_{i1}^t \theta_{j1}^t U_{k2}^2 (D_1 + D_{11} + 2D_{12} + D_2 \\
& + D_{22})] (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{\tilde{\chi}_k^+}^2, m_{\tilde{t}_i}^2, m_{\tilde{t}_l}^2, m_{\tilde{t}_j}^2) \\
& + \sum_{i,j} \frac{gh_b \alpha_{2i} \varphi_{ij}}{16\sqrt{2}\pi^2} \{ h_b^2 \beta_{12} \beta_{2j} C_1(m_b^2, m_{H^+}^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2) + [-h_t^2 m_b^2 \beta_{11} \beta_{1j} (2D_2 \\
& + D_{22} + 2D_{23}) + h_b h_t m_b m_t \beta_{11} \beta_{2j} (D_{22} + 2D_2) - h_b h_t m_b m_t \beta_{12} \beta_{1j} D_{23} \\
& + h_b^2 \beta_{12} \beta_{2j} (m_b^2 (D_{22} + D_{23}) + m_W^2 D_{12} + \hat{u} D_{22} + m_{H_j^+}^2 D_2)] \\
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H_j^+}^2, m_{H_i}^2, m_b^2, m_t^2) \} \\
& + \sum_i \frac{gh_b \beta_{2i}}{16\sqrt{2}\pi^2} \{ h_b^2 \beta_{21} \beta_{2i} C_1(m_{H^+}^2, m_b^2, \hat{t}, m_t^2, m_b^2, m_{A_i}^2) + [h_b^2 \beta_{21} \beta_{2i} (m_b^2 (D_{12} \\
& - D_{22}) + m_W^2 D_{23} + \hat{u} D_{22} + m_{H_i^+}^2 D_2) - h_b h_t m_b m_t (\beta_{11} \beta_{2i} D_{12} - \beta_{21} \beta_{1i} D_{22})
\end{aligned}$$

$$\begin{aligned}
& -h_t^2 m_b^2 \beta_{11} \beta_{1i} D_{22}) (m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_t^2, m_b^2, m_{A_i}^2) \} \\
& + \sum_{i,j,k,l} \sum_{k',l'} \frac{gh_b \theta_{kk'}^b \theta_{ll'}^t}{8\pi^2} (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) [h_b \theta_{k1}^b \theta_{l1}^t U_{j2} O_{ij}^{R*} (-m_b N_{i3}^* D_{22} \\
& + m_{\tilde{\chi}_i^0} N_{i3} D_2) - h_t m_b \theta_{k2}^b \theta_{l2}^t N_{i3} V_{j2} O_{ij}^{L*} D_{23}] \\
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{b_k}^2, m_{t_i}^2), \\
f_4^{(b)} = & \sum_i \frac{gh_b^2 \alpha_{2i}^2}{16\sqrt{2}\pi^2} [h_t m_b m_t \beta_{11} (D_{13} + D_{23}) - h_b m_b^2 \beta_{21} (2D_{13} + 2D_{23} \\
& + 2D_3 + D_{33})] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_t^2, m_b^2) \\
& + \sum_i \frac{gh_b^2 \beta_{2i}^2}{16\sqrt{2}\pi^2} [h_t m_b m_t \beta_{11} (D_{13} + D_{23}) + h_b m_b^2 \beta_{21} D_{33}] \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_t^2, m_b^2) \\
& + \sum_i \frac{gh_b \beta_{2i}}{8\sqrt{2}\pi^2} \{ -h_t^2 \beta_{11} \beta_{1i} C_0 (m_W^2, m_{H^+}^2, \hat{s}, m_t^2, m_b^2, m_t^2) + [h_b h_t m_b m_t \beta_{11} \beta_{2i} (D_1 \\
& + D_{12} + D_{13}) - h_b^2 m_b^2 \beta_{21} \beta_{2i} (D_{12} + D_{13}) - h_t^2 \beta_{11} \beta_{1i} (-2D_{00} + m_b^2 (D_1 - D_{23} \\
& - D_{33}) + m_{H^+}^2 (D_{12} + D_{13}) - \hat{u} (D_{12} + D_{13} + D_{22} + D_{23}) \\
& + m_{H_i^+}^2 D_0)] (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H_i^+}^2, m_t^2, m_b^2, m_t^2) \} \\
& + \sum_{i,j,k,l} \sum_{i',k'} \frac{-\sqrt{2} gh_b^2}{16\pi^2} N_{i3}^* \theta_{ii'}^b \theta_{j1}^b \theta_{k1}^t \theta_{kk'}^t (h_b \Theta_{i'k'1}^5 + h_t \Theta_{i'k'1}^6) [m_b \theta_{i2}^b \theta_{j2}^b N_{i3} D_{33} \\
& - m_{\tilde{\chi}_i^0} N_{i3}^* \theta_{i1}^b \theta_{j2}^b D_3 + m_b \theta_{i1}^b \theta_{j1}^b N_{i3} (D_{13} + D_{23})] \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{\tilde{\chi}_i^0}^2, m_{b_i}^2, m_{t_k}^2, m_{b_j}^2) \\
& + \sum_{i,j,k,l} \sum_{j',l'} \frac{g \theta_{l1}^b \theta_{ll'}^b \theta_{i1}^t \theta_{jj'}^t}{8\sqrt{2}\pi^2} (h_b \Theta_{l'j'1}^5 + h_t \Theta_{l'j'1}^6) [-h_b^2 m_b \theta_{i1}^t \theta_{j1}^t U_{k2}^2 (D_{13} + D_{23} + D_3) \\
& + h_b h_t m_{\tilde{\chi}_k^+} \theta_{i1}^t \theta_{j2}^t U_{k2} V_{k2} (D_0 + D_1 + D_2) - h_t^2 m_b \theta_{i2}^t \theta_{j2}^t V_{k2}^2 (D_1 + D_2 + D_{11} \\
& + 2D_{12} + D_{22})] (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{\tilde{\chi}_k^+}^2, m_{t_i}^2, m_{b_i}^2, m_{t_j}^2) \\
& + \sum_{i,j} \frac{gh_b \alpha_{2i} \varphi_{ij}}{16\sqrt{2}\pi^2} \{ -h_t^2 \beta_{11} \beta_{1j} C_1 (m_b^2, m_{H^+}^2, \hat{t}, m_{H_i}^2, m_b^2, m_t^2) + [h_b^2 m_b^2 \beta_{12} \beta_{2j} (2D_2 \\
& + D_{22} + 2D_{23}) + h_b h_t m_b m_t (\beta_{11} \beta_{2j} D_{23} + \beta_{12} \beta_{1j} (D_{22} + 2D_2)) \\
& - h_t^2 \beta_{11} \beta_{1j} (m_b^2 (D_{22} + D_{23}) + m_W^2 D_{12} + \hat{u} D_{22} + m_{H_j^+}^2 D_2)] \\
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H_j^+}^2, m_{H_i}^2, m_b^2, m_t^2) \} \\
& - \sum_i \frac{gh_b \beta_{2i}}{16\sqrt{2}\pi^2} \{ h_t^2 \beta_{11} \beta_{1i} C_1 (m_{H^+}^2, m_b^2, \hat{t}, m_t^2, m_b^2, m_{A_i}^2) + [h_t^2 \beta_{11} \beta_{1i} (m_b^2 (D_{12} \\
& - D_{22}) + m_W^2 D_{23} + \hat{u} D_{22} + m_{H_i^+}^2 D_2) - h_b h_t m_b m_t (\beta_{21} \beta_{1i} D_{12} - \beta_{11} \beta_{2i} D_{22}) \}
\end{aligned}$$

$$\begin{aligned}
& -h_b^2 m_b^2 \beta_{21} \beta_{2i} D_{22} \} (m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_t^2, m_b^2, m_{A_i}^2) \} \\
& + \sum_{i,j,k,l} \sum_{k',l'} \frac{gh_b \theta_{kk'}^b \theta_{ll'}^t}{8\pi^2} (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) [h_t \theta_{k2}^b \theta_{l2}^t V_{j2} O_{ij}^{L*} (-m_b N_{i3} D_{22} \\
& + m_{\tilde{\chi}_i^0} N_{i3}^* D_2) - h_b m_b \theta_{k1}^b \theta_{l1}^t N_{i3}^* U_{j2} O_{ij}^{R*} D_{23}] \\
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{b_k}^2, m_{t_l}^2), \\
f_5^{(b)} = & \sum_i \frac{gh_b^2 \alpha_{2i}^2}{32\sqrt{2}\pi^2} \{ [h_t m_t \beta_{11} C_0 - h_b m_b \beta_{21} (2C_0 + C_2)] (m_{H^+}^2, m_W^2, \hat{s}, m_b^2, m_t^2, m_b^2) \\
& + [2h_t m_b^2 m_t \beta_{11} (D_1 + D_2 + D_3) + 4h_b m_b \beta_{21} D_{00} - 2h_t m_t \beta_{11} D_{00} \\
& - h_b m_b^3 \beta_{21} (4D_0 + 6D_1 + D_{13} + 4D_2 + 4D_3 + D_{33}) + h_b m_b m_{H^+}^2 \beta_{21} (2D_3 \\
& + D_{33}) + h_b m_b m_W^2 \beta_{21} D_{13} - h_b m_b \hat{t} \beta_{21} (D_{13} + 2D_2 + 2D_{23} + 2D_3 + D_{33}) \\
& + h_t m_t m_{H_i}^2 \beta_{11} D_0 - h_b m_b m_{H_i}^2 \beta_{21} (2D_0 + D_3)] \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_t^2, m_b^2) \} \\
& + \sum_i \frac{gh_b^2 \beta_{2i}^2}{32\sqrt{2}\pi^2} \{ (h_t m_t \beta_{11} C_0 + h_b m_b \beta_{21} C_2) (m_{H^+}^2, m_W^2, \hat{s}, m_b^2, m_t^2, m_b^2) \\
& + [h_b m_b^3 \beta_{21} (D_{13} + D_{33}) - 2h_t m_t \beta_{11} D_{00} - h_b m_b m_{H^+}^2 \beta_{21} D_{33} \\
& - h_b m_b m_W^2 \beta_{21} D_{13} + h_b m_b \hat{t} \beta_{21} (D_{13} + 2D_{23} + D_{33}) + h_t m_t m_{A_i}^2 \beta_{11} D_0 \\
& + h_b m_b m_{A_i}^2 \beta_{21} D_3] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_t^2, m_b^2) \} \\
& + \sum_i \frac{g}{16\sqrt{2}\pi^2} \{ h_b \beta_{2i} [-h_b^2 m_b \beta_{21} \beta_{2i} C_0 + h_b h_t m_t \beta_{11} \beta_{2i} C_0 + h_t^2 m_b \beta_{11} \beta_{1i} (C_1 \\
& + C_2)] (m_W^2, m_{H^+}^2, \hat{s}, m_t^2, m_b^2, m_t^2) + [h_t^3 m_b^2 m_t \beta_{11} \beta_{1i}^2 (D_0 + D_1 + D_2 + D_3) \\
& + h_b^2 h_t m_b^2 m_t \beta_{1i} \beta_{21} \beta_{2i} (D_1 + D_2 + D_3) - h_b^3 m_b \beta_{21} \beta_{2i}^2 (-2D_{00} + m_b^2 (D_1 + D_2 \\
& + D_3) + m_{H_i^+}^2 D_0) + h_b^2 h_t m_t \beta_{11} \beta_{2i}^2 (-2D_{00} + m_b^2 (D_0 + D_2 + 2D_3) - m_{H^+}^2 D_1 \\
& + \hat{u} (D_1 + D_2) + m_{H_i^+}^2 D_0) + h_b h_t^2 m_b \beta_{11} \beta_{1i} \beta_{2i} (2D_{00} + m_{H^+}^2 (D_1 + D_{11}) \\
& - m_t^2 (D_0 + D_2 + D_3) + m_W^2 D_{13} - m_{H_i^+}^2 (D_0 + D_1) - m_b^2 (2D_1 + D_{11} \\
& + D_2 + 2D_3) - \hat{u} (D_1 + D_{11} + 2D_{12} + D_{13} + D_2)] \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H_i^+}^2, m_t^2, m_b^2, m_t^2) \} \\
& + \sum_{i,j,k} \sum_{i',k'} \frac{\sqrt{2}gh_b^2}{16\pi^2} |N_{l3}|^2 \theta_{ii'}^b \theta_{i1}^b (\theta_{j1}^b)^2 \theta_{k1}^t \theta_{kk'}^t (h_b \Theta_{i'k'1}^5 + h_t \Theta_{i'k'1}^6) D_{00} \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{\tilde{\chi}_l^0}^2, m_{b_i}^2, m_{t_k}^2, m_{b_j}^2)
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i,j,k,l} \sum_{j',l'} \frac{gh_b^2 \theta_{l1}^b \theta_{l'}^b (\theta_{i1}^t)^2 \theta_{j1}^t \theta_{jj'}^t U_{k2}^2}{8\sqrt{2}\pi^2} (h_b \Theta_{l'j'1}^5 + h_t \Theta_{l'j'1}^6) D_{00} \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{\tilde{\chi}_k^+}^2, m_{\tilde{t}_i}^2, m_{\tilde{b}_i}^2, m_{\tilde{t}_j}^2) \\
& + \sum_{i,j} \frac{gh_b \alpha_{2i} \varphi_{ij}}{16\sqrt{2}\pi^2} (h_t^2 m_b \beta_{11} \beta_{1j} - h_b h_t \beta_{11} \beta_{2j} - 2h_b^2 m_b \beta_{12} \beta_{2j}) D_{00} \\
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_{H_i}^2, m_b^2, m_t^2) \\
& + \sum_i \frac{gh_b h_t \beta_{11} \beta_{2i}}{16\sqrt{2}\pi^2} (h_t m_b \beta_{1i} - h_b m_t \beta_{2i}) D_{00} (m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_{\tilde{t}_i}^2, m_b^2, m_{A_i}^2) \\
& + \sum_{i,j,k,l} \sum_{k',l'} \frac{gh_b \theta_{kk'}^b \theta_{ll'}^t}{16\pi^2} (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) \{-h_b \theta_{k1}^b \theta_{l1}^t N_{i3}^* U_{j2} O_{ij}^{R*} \\
& C_0(m_b^2, m_{H^+}^2, \hat{t}, m_{\tilde{\chi}_i^0}^2, m_{\tilde{b}_k}^2, m_{\tilde{t}_i}^2) + [h_b m_{\tilde{\chi}_i^0} m_{\tilde{\chi}_j^+} \theta_{k1}^b \theta_{l1}^t N_{i3}^* U_{j2} O_{ij}^{L*} D_0 \\
& + h_b \theta_{k1}^b \theta_{l1}^t N_{i3}^* U_{j2} O_{ij}^{R*} (2D_{00} + m_b^2 D_2 - m_W^2 D_1 - \hat{u} D_2 - m_{\tilde{\chi}_j^+}^2 D_0) \\
& + h_b m_b \theta_{k2}^b \theta_{l1}^t N_{i3} U_{j2} (-m_{\tilde{\chi}_j^+} O_{ij}^{L*} + m_{\tilde{\chi}_i^0} O_{ij}^{R*}) D_2 + h_t m_b \theta_{k1}^b \theta_{l2}^t N_{i3}^* V_{j2} (m_{\tilde{\chi}_j^+} O_{ij}^{R*} \\
& - m_{\tilde{\chi}_i^0} O_{ij}^{L*}) D_3] (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{\tilde{b}_k}^2, m_{\tilde{t}_i}^2)\}, \\
f_6^{(b)} & = \sum_i \frac{gh_b^2 \alpha_{2i}^2}{32\sqrt{2}\pi^2} \{-h_b m_b \beta_{21} C_2(m_{H^+}^2, m_W^2, \hat{s}, m_b^2, m_{\tilde{t}_i}^2, m_b^2) + [-2h_t m_b^2 m_t \beta_{11} (2D_0 \\
& + D_1 + D_2 + D_3) + 4h_b m_b \beta_{21} D_{00} + h_b m_b^3 \beta_{21} (D_{11} + D_{12} + D_{13} + D_{23} + 2D_3) \\
& - h_b m_b m_{H^+}^2 \beta_{21} (D_{13} + D_{23}) - h_b m_b m_W^2 \beta_{21} (2D_1 + D_{11} + D_{12}) \\
& + h_b m_b \hat{t} \beta_{21} (2D_1 + D_{11} + 3D_{12} + D_{13} + 2D_2 + 2D_{22} + D_{33}) \\
& + h_b m_b m_{H_i}^2 \beta_{21} (D_0 + D_1 + D_2)] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_{\tilde{t}_i}^2, m_b^2)\} \\
& + \sum_i \frac{gh_b^3 m_b \beta_{21} \beta_{2i}^2}{32\sqrt{2}\pi^2} \{(2C_0 + C_2) (m_{H^+}^2, m_W^2, \hat{s}, m_b^2, m_{\tilde{t}_i}^2, m_b^2) + [-4D_{00} - m_b^2 (D_{11} \\
& + D_{12} + D_{13} + D_{23}) + m_{H^+}^2 (D_{13} + D_{23}) + m_W^2 (D_{11} + D_{12}) - \hat{t} (D_{11} + 3D_{12} \\
& + D_{13} + 2D_{22} + D_{23}) + m_{A_i}^2 (D_0 - D_1 - D_2)] \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_{\tilde{t}_i}^2, m_b^2)\} \\
& + \sum_i \frac{g}{16\sqrt{2}\pi^2} \{h_t^2 \beta_{11} \beta_{1i} [h_b m_b \beta_{2i} (C_1 + C_2) - h_t m_t \beta_{1i} C_0] \\
& (m_W^2, m_{H^+}^2, \hat{s}, m_{\tilde{t}_i}^2, m_b^2, m_{\tilde{t}_i}^2) + [h_b h_t^2 m_b m_{\tilde{t}_i}^2 \beta_{1i}^2 \beta_{21} D_0 - h_b^2 h_t m_b^2 m_t \beta_{1i} \beta_{21} \beta_{2i} (2D_0 \\
& + D_1 + D_2 + D_3) + h_b^3 m_b^3 \beta_{21} \beta_{2i}^2 (D_0 + D_1 + D_2 + D_3) - h_t^3 m_{\tilde{t}_i} \beta_{11} \beta_{1i}^2 (m_b^2 D_1 \\
& - m_W^2 D_3 + \hat{u} (D_2 + D_3) + m_{H_i^+}^2 D_0) + h_b h_t^2 m_b \beta_{11} \beta_{1i} \beta_{2i} (4D_{00} + m_b^2 (D_{12} \\
& + D_{13} + D_{23} + D_{33}) - m_{H^+}^2 D_{13} + m_{\tilde{t}_i}^2 D_1 - m_W^2 (D_{23} + D_3 - D_{33})
\end{aligned}$$

$$\begin{aligned}
& +\hat{u}(D_{12} + D_{13} + D_2 + 2D_{22} + 3D_{23} + D_3 + D_{33}) + m_{H_i^+}^2(D_2 + D_3)] \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H_i^+}^2, m_t^2, m_b^2, m_t^2) \} \\
& + \sum_{i,j,k,l,i',k'} \frac{\sqrt{2}gh_b^2}{16\pi^2} |N_{l3}|^2 \theta_{ii'}^b \theta_{i2}^b \theta_{j1}^b \theta_{j2}^b \theta_{k1}^t \theta_{kk'}^t (h_b \Theta_{i'k'1}^5 + h_t \Theta_{i'k'1}^6) D_{00} \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{\tilde{\chi}_i^0}^2, m_{\tilde{b}_i}^2, m_{\tilde{t}_k}^2, m_{\tilde{b}_j}^2) \\
& + \sum_{i,j,k,l,j',l'} \frac{gh_t^2 \theta_{l1}^b \theta_{l'l'}^b (\theta_{i2}^t)^2 \theta_{j1}^t \theta_{jj'}^t V_{k2}^2}{8\sqrt{2}\pi^2} (h_b \Theta_{l'j'1}^5 + h_t \Theta_{l'j'1}^6) D_{00} \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{\tilde{\chi}_k^+}^2, m_{\tilde{t}_i}^2, m_{\tilde{b}_l}^2, m_{\tilde{t}_j}^2) \\
& + \sum_{i,j} \frac{gh_b \alpha_{2i} \varphi_{ij}}{16\sqrt{2}\pi^2} (2h_t^2 m_b \beta_{11} \beta_{1j} + h_b h_t \beta_{12} \beta_{1j} - h_b^2 m_b \beta_{12} \beta_{2j}) D_{00} \\
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_{H_i}^2, m_b^2, m_t^2) \\
& - \sum_i \frac{gh_b h_t \beta_{21} \beta_{2i}}{16\sqrt{2}\pi^2} (h_b m_b \beta_{2i} - h_t m_t \beta_{1i}) D_{00} (m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_t^2, m_b^2, m_{A_i}^2) \\
& + \sum_{i,j,k,l,k',l'} \frac{gh_b \theta_{kk'}^b \theta_{ll'}^t}{16\pi^2} (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) \{ -h_t \theta_{k2}^b \theta_{l2}^t N_{i3} V_{j2} O_{ij}^{L*} \\
& C_0(m_b^2, m_{H^+}^2, \hat{t}, m_{\tilde{\chi}_i^0}^2, m_{\tilde{b}_k}^2, m_{\tilde{t}_i}^2) + [h_t m_{\tilde{\chi}_i^0} m_{\tilde{\chi}_j^+} \theta_{k2}^b \theta_{l2}^t N_{i3} V_{j2} O_{ij}^{R*} D_0 \\
& + h_t \theta_{k2}^b \theta_{l2}^t N_{i3} V_{j2} O_{ij}^{L*} (2D_{00} + m_b^2 D_2 - m_W^2 D_1 - \hat{u} D_2 - m_{\tilde{\chi}_j^+}^2 D_0) \\
& + h_t m_b \theta_{k1}^b \theta_{l2}^t N_{i3}^* V_{j2} (-m_{\tilde{\chi}_j^+} O_{ij}^{R*} + m_{\tilde{\chi}_i^0} O_{ij}^{L*}) D_2 + h_b m_b \theta_{k2}^b \theta_{l1}^t N_{i3} U_{j2} (m_{\tilde{\chi}_j^+} O_{ij}^{L*} \\
& - m_{\tilde{\chi}_i^0} O_{ij}^{R*}) D_3] (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{\tilde{b}_k}^2, m_{\tilde{t}_i}^2) \}, \\
f_7^{(b)} & = \sum_i \frac{gh_b^2 \alpha_{2i}^2}{16\sqrt{2}\pi^2} [h_t m_t \beta_{11} (D_{12} + D_{22}) - h_b m_b \beta_{21} (2D_{12} + D_{13} + 2D_2 + 2D_{22} \\
& + D_{23})] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_t^2, m_b^2) \\
& + \sum_i \frac{gh_b^2 \beta_{2i}^2}{16\sqrt{2}\pi^2} [h_t m_t \beta_{11} (D_{12} + D_{22}) + h_b m_b \beta_{21} (D_{13} + D_{23})] \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_t^2, m_b^2) \\
& + \sum_i \frac{gh_b \beta_{2i}^2}{8\sqrt{2}\pi^2} [h_t^2 m_b \beta_{11} \beta_{1i} (D_1 + D_{11} + D_{12}) - h_b h_t m_t \beta_{11} \beta_{2i} (D_1 + D_{12}) \\
& + h_b^2 m_b \beta_{21} \beta_{2i} D_{12}] (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H_i^+}^2, m_t^2, m_b^2, m_t^2) \\
& + \sum_{i,j,k,l,i',k'} \frac{\sqrt{2}gh_b^2}{16\pi^2} |N_{l3}|^2 \theta_{ii'}^b \theta_{i1}^b (\theta_{j1}^b)^2 \theta_{k1}^t \theta_{kk'}^t (h_b \Theta_{i'k'1}^5 + h_t \Theta_{i'k'1}^6) (D_{12} + D_2 + D_{22}) \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{\tilde{\chi}_i^0}^2, m_{\tilde{b}_i}^2, m_{\tilde{t}_k}^2, m_{\tilde{b}_j}^2)
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i,j,k,l} \sum_{j',l'} \frac{gh_b^2 \theta_{l1}^b \theta_{l'l'}^b (\theta_{i1}^t)^2 \theta_{j1}^t \theta_{jj'}^t U_{k2}^2}{8\sqrt{2}\pi^2} (h_b \Theta_{l'j'1}^5 + h_t \Theta_{l'j'1}^6) D_{23} \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{\chi_k^+}^2, m_{\tilde{t}_i}^2, m_{\tilde{b}_i}^2, m_{\tilde{t}_j}^2) \\
& + \sum_{i,j} \frac{gh_b \alpha_{2i} \varphi_{ij}}{16\sqrt{2}\pi^2} [h_t^2 m_b \beta_{11} \beta_{1j} (D_{13} + D_{23} + D_3 + D_{33}) - h_b h_t m_t \beta_{11} \beta_{2j} (D_{13} \\
& + D_{23} + D_3) - h_b^2 m_b \beta_{12} \beta_{2j} (2D_{13} + D_{23})] \\
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H_j^+}^2, m_{H_i}^2, m_b^2, m_t^2) \\
& + \sum_i \frac{gh_b \beta_{2i}}{16\sqrt{2}\pi^2} [h_b^2 m_b \beta_{21} \beta_{2i} D_{12} - h_b h_t m_t (\beta_{11} \beta_{2i} D_1 + \beta_{21} \beta_{1i} D_{12}) \\
& + h_t^2 m_b \beta_{11} \beta_{1i} (D_1 + D_{11} + D_{12} + D_{13})] (m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_{\tilde{t}_i}^2, m_b^2, m_{A_i}^2) \\
& + \sum_{i,j,k,l} \sum_{k',l'} \frac{gh_b^2}{8\pi^2} \theta_{k1}^b \theta_{kk'}^b \theta_{l1}^t \theta_{l'l'}^t (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) N_{i3}^* U_{j2} O_{ij}^{R*} (D_{13} \\
& + D_{23}) (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\chi_j^0}^2, m_{\chi_i^0}^2, m_{b_k}^2, m_{\tilde{t}_i}^2), \\
f_8^{(b)} & = \sum_i \frac{gh_b^3 m_b \alpha_{2i} \beta_{21}}{16\sqrt{2}\pi^2} (2D_1 + D_{11} + D_{12}) \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_{\tilde{t}_i}^2, m_b^2) \\
& + \sum_i \frac{-gh_b^3 m_b \beta_{2i}^2 \beta_{21}}{16\sqrt{2}\pi^2} (D_{11} + D_{12}) (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_{\tilde{t}_i}^2, m_b^2) \\
& + \sum_i \frac{-gh_b h_t^2 m_b}{8\sqrt{2}\pi^2} \beta_{11} \beta_{1i} \beta_{2i} D_{13} (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H_i^+}^2, m_{\tilde{t}_i}^2, m_b^2, m_t^2) \\
& + \sum_{i,j,k,l} \sum_{i',k'} \frac{\sqrt{2} gh_b^2}{16\pi^2} |N_{i3}|^2 \theta_{ii'}^b \theta_{i2}^b \theta_{j1}^b \theta_{j2}^b \theta_{k1}^t \theta_{kk'}^t (h_b \Theta_{i'k'1}^5 + h_t \Theta_{i'k'1}^6) (D_{12} + D_2 + D_{22}) \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{\chi_i^0}^2, m_{\tilde{b}_i}^2, m_{\tilde{t}_k}^2, m_{\tilde{b}_j}^2) \\
& + \sum_{i,j,k,l} \sum_{j',l'} \frac{gh_t^2 \theta_{l1}^b \theta_{l'l'}^b (\theta_{i2}^t)^2 \theta_{j1}^t \theta_{jj'}^t V_{k2}}{8\sqrt{2}\pi^2} (h_b \Theta_{l'j'1}^5 + h_t \Theta_{l'j'1}^6) D_{23} \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{\chi_k^+}^2, m_{\tilde{t}_i}^2, m_{\tilde{b}_i}^2, m_{\tilde{t}_j}^2) \\
& + \sum_{i,j} \frac{gh_b \alpha_{2i} \varphi_{ij}}{16\sqrt{2}\pi^2} [h_t^2 m_b \beta_{11} \beta_{1j} (2D_{13} + D_{23}) + h_b h_t m_t \beta_{11} \beta_{2j} (D_{13} + D_{23} \\
& + D_3) - h_b^2 m_b \beta_{12} \beta_{2j} (D_{13} + D_{23} + D_{33})] \\
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H_j^+}^2, m_{H_i}^2, m_b^2, m_t^2) \\
& - \sum_i \frac{gh_b \beta_{2i}}{16\sqrt{2}\pi^2} [h_t^2 m_b \beta_{11} \beta_{1i} D_{12} - h_b h_t m_t (\beta_{21} \beta_{1i} D_1 + \beta_{11} \beta_{2i} D_{12}) \\
& + h_b^2 m_b \beta_{21} \beta_{2i} (D_1 + D_{11} + D_{12} + D_{13})] (m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_{\tilde{t}_i}^2, m_b^2, m_{A_i}^2)
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i,j,k,l} \sum_{k',l'} \frac{gh_b h_t}{8\pi^2} \theta_{k2}^b \theta_{kk'}^b \theta_{l2}^t \theta_{l'l'}^t (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) N_{i3} V_{j2} O_{ij}^{L*} (D_{13} \\
& + D_{23}) (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{b_k}^2, m_{t_i}^2), \\
f_9^{(b)} & = \sum_i \frac{gh_b^2 \alpha_{2i}^2}{16\sqrt{2}\pi^2} [-h_t m_t \beta_{11} D_{23} + h_b m_b \beta_{21} (2D_{23} + 2D_3 + D_{33})] \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_t^2, m_b^2) \\
& + \sum_i \frac{-gh_b^2 \beta_{2i}^2}{16\sqrt{2}\pi^2} (h_t m_t \beta_{11} D_{23} + h_b m_b \beta_{21} D_{33}) \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_t^2, m_b^2) \\
& + \sum_i \frac{gh_b \beta_{2i}}{8\sqrt{2}\pi^2} [-h_t^2 m_b \beta_{11} \beta_{1i} (D_{12} + D_{13} + D_2 + D_{22} + D_{23}) \\
& + h_b h_t m_t \beta_{11} \beta_{2i} (D_2 + D_{22} + D_{23}) - h_b^2 m_b \beta_{11} \beta_{2i} (D_{22} \\
& + D_{23})] (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H_i^+}^2, m_t^2, m_b^2, m_t^2) \\
& + \sum_{i,j,k,l} \sum_{i',k'} \frac{\sqrt{2} gh_b^2}{16\pi^2} |N_{i3}|^2 \theta_{ii'}^b \theta_{i1}^b (\theta_{j1}^b)^2 \theta_{k1}^t \theta_{kk'}^t (h_b \Theta_{i'k'1}^5 + h_t \Theta_{i'k'1}^6) D_{23} \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{\tilde{\chi}_i^0}^2, m_{b_i}^2, m_{t_k}^2, m_{b_j}^2) \\
& - \sum_{i,j,k,l} \sum_{j',l'} \frac{gh_b^2}{8\sqrt{2}\pi^2} \theta_{l1}^b \theta_{l'l'}^b (\theta_{i1}^t)^2 \theta_{j1}^t \theta_{j'j'}^t U_{k2}^2 (h_b \Theta_{l'j'1}^5 + h_t \Theta_{l'j'1}^6) (D_2 + D_{12} + D_{22}) \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{\tilde{\chi}_k^+}^2, m_{t_i}^2, m_{b_l}^2, m_{t_j}^2) \\
& + \sum_{i,j} \frac{gh_b \alpha_{2i} \varphi_{ij}}{16\sqrt{2}\pi^2} [-h_t^2 m_b \beta_{11} \beta_{1j} (D_{12} + D_2 + D_{22} + D_{23}) \\
& + h_b h_t m_t \beta_{11} \beta_{2j} (D_{12} + D_2 + D_{22}) + h_b^2 m_b \beta_{12} \beta_{2j} (2D_{12} + D_{22})] \\
& (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H_j^+}^2, m_{H_i}^2, m_b^2, m_t^2) \\
& + \sum_i \frac{gh_b \beta_{2i}}{16\sqrt{2}\pi^2} [-h_b^2 m_b \beta_{21} \beta_{2i} D_{22} + h_b h_t m_t \beta_{11} \beta_{2i} (D_2 + D_{22} + D_{23}) \\
& - h_t^2 m_b \beta_{11} \beta_{1i} (D_{12} + D_2 + D_{22} + D_{23})] (m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_t^2, m_b^2, m_{A_i}^2) \\
& - \sum_{i,j,k,l} \sum_{k',l'} \frac{gh_b^2}{8\pi^2} \theta_{k1}^b \theta_{kk'}^b \theta_{l1}^t \theta_{l'l'}^t (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) N_{i3}^* U_{j2} O_{ij}^{R*} (D_{12} + D_2 \\
& + D_{22}) (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{b_k}^2, m_{t_i}^2), \\
f_{10}^{(b)} & = \sum_i \frac{-gh_b^3 m_b \alpha_{2i}^2 \beta_{21}}{16\sqrt{2}\pi^2} D_{13} (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_t^2, m_b^2) \\
& + \sum_i \frac{gh_b^3 m_b \beta_{2i}^2 \beta_{21}}{16\sqrt{2}\pi^2} D_{13} (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_t^2, m_b^2)
\end{aligned}$$

$$\begin{aligned}
& + \sum_i \frac{gh_t^2 \beta_{11} \beta_{1i}}{8\sqrt{2}\pi^2} [h_t m_t \beta_{1i} D_3 + h_b m_b \beta_{2i} (D_{23} + D_3 + D_{33})] \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H^+}^2, m_t^2, m_b^2, m_t^2) \\
& + \sum_{i,j,k,l,i',k'} \frac{\sqrt{2}gh_b^2}{16\pi^2} |N_{l3}|^2 \theta_{i'j'}^b \theta_{i2}^b \theta_{j1}^b \theta_{j2}^b \theta_{k1}^t \theta_{kk'}^t (h_b \Theta_{i'k'1}^5 + h_t \Theta_{i'k'1}^6) D_{23} \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{\tilde{\chi}_i^0}^2, m_{\tilde{b}_i}^2, m_{\tilde{t}_k}^2, m_{\tilde{b}_j}^2) \\
& - \sum_{i,j,k,l,j',l'} \frac{gh_t^2}{8\sqrt{2}\pi^2} \theta_{l1}^b \theta_{l'l'}^b (\theta_{i2}^t)^2 \theta_{j1}^t \theta_{j'j'}^t V_{k2}^2 (h_b \Theta_{l'j'1}^5 + h_t \Theta_{l'j'1}^6) (D_2 + D_{12} + D_{22}) \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{\tilde{\chi}_k^+}^2, m_{\tilde{t}_i}^2, m_{\tilde{b}_i}^2, m_{\tilde{t}_j}^2) \\
& + \sum_{i,j} \frac{gh_b \alpha_{2i} \varphi_{ij}}{16\sqrt{2}\pi^2} [-h_t^2 m_b \beta_{11} \beta_{1j} (D_{12} + D_{22}) - h_b h_t m_t \beta_{12} \beta_{1j} (D_{12} + D_2 + D_{22}) \\
& + h_b^2 m_b \beta_{12} \beta_{2j} (2D_{12} + D_{22})] (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H^+}^2, m_{H_i}^2, m_b^2, m_t^2) \\
& - \sum_i \frac{gh_b \beta_{2i}}{16\sqrt{2}\pi^2} [-h_t^2 m_b \beta_{11} \beta_{1i} D_{22} + h_b h_t m_t \beta_{21} \beta_{1i} (D_2 + D_{22} + D_{23}) \\
& - h_b^2 m_b \beta_{21} \beta_{2i} (D_{12} + D_2 + D_{22} + D_{23})] (m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H^+}^2, m_t^2, m_b^2, m_{A_i}^2) \\
& - \sum_{i,j,k,l,k',l'} \frac{gh_b h_t}{8\pi^2} \theta_{k2}^b \theta_{kk'}^b \theta_{l2}^t \theta_{l'l'}^t (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) N_{i3} V_{j2} O_{ij}^{L*} (D_{12} + D_2 \\
& + D_{22}) (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{\tilde{b}_k}^2, m_{\tilde{t}_i}^2), \\
f_{11}^{(b)} & = \sum_i \frac{gh_b^2 \alpha_{2i}^2}{32\sqrt{2}\pi^2} \{ -h_b \beta_{21} (C_0 - C_1) (m_{H^+}^2, m_W^2, \hat{s}, m_b^2, m_t^2, m_b^2) \\
& + [h_b \beta_{21} D_{00} - 2h_t m_b m_t \beta_{11} D_2 - h_b m_b^2 \beta_{21} (2D_1 - D_{12} - D_{23} + 2D_3) \\
& - h_b m_{H^+}^2 \beta_{21} D_{23} - h_b m_W^2 \beta_{21} D_{12} + h_b \hat{t} \beta_{21} (D_{12} + 2D_{22} + D_{23}) \\
& - h_b m_{H_i}^2 (D_0 - D_2)] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_t^2, m_b^2) \} \\
& + \sum_i \frac{gh_b^3 \beta_{21} \beta_{2i}^2}{32\sqrt{2}\pi^2} \{ (C_0 - C_1) (m_{H^+}^2, m_W^2, \hat{s}, m_b^2, m_t^2, m_b^2) + [-4D_{00} - m_b^2 (D_{12} \\
& + D_{23}) + m_{H^+}^2 D_{23} + m_W^2 D_{12} - \hat{t} (D_{12} + 2D_{22} + D_{23}) + m_{A_i}^2 (D_0 \\
& - D_2)] (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{A_i}^2, m_b^2, m_t^2, m_b^2) \} \\
& + \sum_i \frac{g}{16\sqrt{2}\pi^2} \{ h_t^3 m_b m_t \beta_{11} \beta_{1i}^2 (D_0 + D_1 + D_2) - h_b h_t^2 \beta_{11} \beta_{1i} \beta_{2i} [m_b^2 (D_0 + D_1 \\
& + D_2 + D_3) + m_t^2 (D_0 + D_2)] + h_b^2 h_t m_b m_t \beta_{2i} [\beta_{1i} \beta_{21} D_2 + \beta_{11} \beta_{2i} (D_0 + D_2 \\
& + D_3)] - h_b^3 m_b^2 \beta_{21} \beta_{2i}^2 D_2 \} (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H^+}^2, m_t^2, m_b^2, m_t^2) \\
& - \sum_{i,j} \frac{gh_b^3 \alpha_{2i} \beta_{12} \beta_{2j} \varphi_{ij}}{16\sqrt{2}\pi^2} D_{00} (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H^+}^2, m_{H_i}^2, m_b^2, m_t^2)
\end{aligned}$$

$$\begin{aligned}
& - \sum_i \frac{gh_b^3 \beta_{21} \beta_{2i}^2}{16\sqrt{2}\pi^2} D_{00}(m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_t^2, m_b^2, m_{A_i}^2) \\
& + \sum_{i,j,k,l} \sum_{k',l'} \frac{gh_b \theta_{kk'}^b \theta_{ll'}^t}{16\pi^2} (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) \{h_b m_b \theta_{k1}^b \theta_{l1}^t N_{i3}^* U_{j2} D_2 \\
& + h_b \theta_{k2}^b \theta_{l1}^t N_{i3} U_{j2} [-m_{\tilde{\chi}_j^+} O_{ij}^{L*}(D_0 + D_1 + D_2) + m_{\tilde{\chi}_i^0} O_{ij}^{R*}(D_1 + D_2)] \\
& + h_t m_b \theta_{k2}^b \theta_{l2}^t N_{i3} V_{j2} O_{ij}^{L*} D_3\} (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{b_k}^2, m_{t_i}^2), \\
f_{12}^{(b)} & = \sum_i \frac{gh_b^2 \alpha_{2i}^2}{16\sqrt{2}\pi^2} [h_t m_b m_t \beta_{11} D_2 - h_b m_b^2 \beta_{21} (2D_0 - D_1 - 2D_2 - D_3)] \\
& (m_b^2, m_{H^+}^2, m_W^2, m_b^2, \hat{t}, \hat{s}, m_{H_i}^2, m_b^2, m_t^2, m_b^2) \\
& + \sum_i \frac{g}{16\sqrt{2}\pi^2} \{-h_b h_t^2 \beta_{11} \beta_{1i} \beta_{2i} (C_0 - C_1) (m_W^2, m_{H^+}^2, \hat{s}, m_t^2, m_b^2, m_t^2) \\
& + [h_t^3 m_b m_t \beta_{11} \beta_{1i}^2 D_3 - h_b^2 h_t m_b m_t \beta_{2i} (\beta_{1i} \beta_{21} D_2 - \beta_{11} \beta_{2i} D_1) \\
& + h_b^3 m_b^2 \beta_{21} \beta_{2i}^2 D_2 - h_b h_t^2 \beta_{11} \beta_{1i} \beta_{2i} (-4D_{00} + m_b^2 (D_1 - D_{12} - D_{23} \\
& + D_3) + m_{H^+}^2 D_{12} + m_W^2 D_{23} - \hat{u} (D_{12} + D_{22} + D_{23}) + m_{H_i^+}^2 (D_0 - D_2))]\} \\
& (m_b^2, m_W^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{s}, m_{H_i^+}^2, m_t^2, m_b^2, m_t^2) \\
& + \sum_{i,j} \frac{gh_b^3 \alpha_{2i} \beta_{11} \beta_{1j} \varphi_{ij}}{16\sqrt{2}\pi^2} D_{00}(m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_{H_i}^2, m_b^2, m_t^2) \\
& + \sum_i \frac{gh_b h_t^2 \beta_{11} \beta_{1i} \beta_{2i}}{16\sqrt{2}\pi^2} D_{00}(m_b^2, m_{H^+}^2, m_b^2, m_W^2, \hat{u}, \hat{t}, m_{H_i^+}^2, m_t^2, m_b^2, m_{A_i}^2) \\
& + \sum_{i,j,k,l} \sum_{k',l'} \frac{gh_b \theta_{kk'}^b \theta_{ll'}^t}{16\pi^2} (h_b \Theta_{k'l'1}^5 + h_t \Theta_{k'l'1}^6) \{h_t m_b \theta_{k2}^b \theta_{l2}^t N_{i3} V_{j2} D_2 \\
& + h_t \theta_{k1}^b \theta_{l2}^t N_{i3}^* V_{j2} [-m_{\tilde{\chi}_j^+} O_{ij}^{R*}(D_0 + D_1 + D_2) + m_{\tilde{\chi}_i^0} O_{ij}^{L*}(D_1 + D_2)] \\
& + h_b m_b \theta_{k1}^b \theta_{l1}^t N_{i3}^* U_{j2} O_{ij}^{R*} D_3\} (m_W^2, m_b^2, m_{H^+}^2, m_b^2, \hat{u}, \hat{t}, m_{\tilde{\chi}_j^+}^2, m_{\tilde{\chi}_i^0}^2, m_{b_k}^2, m_{t_i}^2);
\end{aligned}$$

All other form factors f_i not listed above vanish.


Here A_0 , C_i , D_i and D_{ij} are the one-, three- and four-point Feynman integrals[20].

The definitions of U_{ij} , V_{ij} , N_{ij} , O_{ij}^L and O_{ij}^R can be found in Ref.[2].


References

- [1] For a review, see J.Gunion, H. Haber, G. Kane, and S.Dawson, *The Higgs Hunter's Guide*(Addison-Wesley, New York,1990).
- [2] H.E. Haber and G.L. Kane, *Phys. Rep.* **117**, 75(1985); J.F. Gunion and H.E. Haber, *Nucl. Phys.* **B272**, 1(1986).
- [3] CMS Technical Proposal. CERN/LHC94-43 LHCC/P1, December 1994.
- [4] CDF Collaboration, *Phys. Rev. Lett.* **79**, 35(1997); D0 Collaboration, *Phys. Rev.Lett.* **82**, 4975(1999).
- [5] Z.Kunszt and F. Zwirner, *Nucl. Phys.* **B385**, 3(1992), and references cited therein.
- [6] J.F. Gunion, H.E. Haber, F.E. Paige, W.-K. Tung, and S. Willenbrock, *Nucl. Phys.* **B294**,621(1987); R.M. Barnett, H.E. Haber, and D.E. Soper, *ibid.* **B306**, 697(1988); F.I. Olness and W.-K. Tung, *ibid.* **B308**, 813(1988).
- [7] V. Barger, R.J.N. Phillips, and D.P. Roy, *Phys. Lett.* **B324**, 236(1994).
- [8] C.S. Huang and S.H. Zhu, *Phys. Rev.* **D60**, 075012(1999). L.G. Jin, C.S.Li, R.J. Oakes, and S.H. Zhu, to appear in *Eur.Phys.J.C*. L.G. Jin, C.S.Li, R.J. Oakes, and S.H. Zhu, hep-ph/0003159.
- [9] D.A. Dicus, J.L.Hewett, C. Kao, and T.G. Rizzo, *Phys. Rev.* **D40**,789(1989).
- [10] A.A.Barrientos Bendezu and B.A. Kniehl, *Phys. Rev.* **D59**, 015009(1998).
- [11] S. Moretti and K. Odagiri, *Phys. Rev.* **D59**, 055008(1999).
- [12] K. Odagiri, hep-ph/9901432; *Phys. Lett.* **B452**, 327(1999).
- [13] D.P. Roy, *Phys. Lett.* **B459**, 607(1999).
- [14] S. Raychaudhuri and D.P.Roy, *Phys. Rev.* **D53**, 4902(1996).
- [15] S. Sirlin, *Phys. Rev.* **D22**, 971 (1980); W. J. Marciano and A. Sirlin,*ibid.* **22**, 2695(1980); **31**, 213(E) (1985); A. Sirlin and W.J. Marciano, *Nucl. Phys.* **B189**, 442(1981); K.I. Aoki et.al., *Prog. Theor. Phys. Suppl.* **73**, 1(1982).


- [16] A. Mendez and A. Pomarol, Phys.Lett.**B279**, 98(1992).
- [17] Particle Data Group, C.Caso *et al*, Eur.Phys.J.C 3, 1(1998).
- [18] H.L. Lai, et al.(CTEQ collaboration), hep-ph/9903282.
- [19] J.Gunion, A.Turski, Phys. Rev. **D39**, 2701(1989); **D40**, 2333(1990); J.R.Espinosa, M.Quiros, Phys. Lett. **B266**, 389(1991); M.Carena, M.Quiros, C.E.M.Wagner, Nucl. Phys. **B461**, 407(1996).
- [20] G.Passarino and M.Veltman, Nucl. Phys. **B160**, 151(1979); A.Axelrod, *ibid.* **B209**, 349 (1982); M.Clements *et al.*, Phys. Rev. **D27**, 570 (1983); A.Denner, Fortschr. Phys. **41**, 4 (1993); R. Mertig *et al.*, Comput. Phys. Commun. **64**, 345 (1991).




(a)




(b)




(c)




(d)




(e)




(f)




(g)




(h)




(i)




(j)




(k)




(l)



(m)



(n)



(o)





Figure 1: Feynman diagrams contributing to supersymmetric electroweak corrections to $b\bar{b} \rightarrow W^- H^+$: (a) and (b) are tree level diagrams; (c) – (x) are one-loop corrections. The dashed line 1 represents H, h, A ; the dashed line 2 represents H, h, A, G^0 ; the dashed line 3 represents H^+, G^+ . For diagram (r), the dashed line in the loop represents $H, h, A, G^0, H^+, G^+, \tilde{t}, \tilde{b}$.






Figure 2: Feynman diagrams contributing to renormalization constants: The dashed line represents $H, h, A, G^0, H^+, G^+, \tilde{t}, \tilde{b}$ for diagram (a), and H_i in diagrams (d) – (f) represents H, h, A .




Figure 3: The Yukawa corrections versus m_{H^+} for $\tan\beta = 1.5, 2, 6$ and 30 , respectively.




Figure 4: The Yukawa corrections versus $\tan\beta$ for $m_{H^+} = 100, 150, 200$ and 300 GeV, respectively.




Figure 5: The genuine SUSY EW corrections versus m_{H^+} for $\tan\beta = 1.5, 2, 6$ and 30 , respectively, assuming $M_2 = 300\text{GeV}$, $\mu = -100\text{GeV}$, $A_t = A_b = 200\text{GeV}$ and $M_{\tilde{Q}} = M_{\tilde{U}} = M_{\tilde{D}} = 500\text{GeV}$.




Figure 6: The genuine SUSY EW corrections versus m_{H^+} for $\tan\beta = 1.5, 6$ and 30 , respectively, assuming $M_2 = 200\text{GeV}$, $\mu = 100\text{GeV}$, $A_t = A_b = 1\text{TeV}$, $M_{\tilde{Q}} = M_{\tilde{U}}$, $m_{\tilde{t}_1} = 100\text{GeV}$ and $m_{\tilde{b}_1} = 150\text{GeV}$.




Figure 7: The genuine SUSY EW corrections versus $A_t = A_b$ for $\tan\beta = 1.5$ and 30, respectively, assuming $m_{H^+} = 200\text{GeV}$, $M_2 = 300\text{GeV}$, $\mu = 100\text{GeV}$, and $M_{\tilde{Q}} = M_{\tilde{U}} = M_{\tilde{D}} = 400\text{GeV}$.




Figure 8: The genuine SUSY EW corrections versus $M_{\tilde{Q}} = M_{\tilde{U}} = M_{\tilde{D}}$ for $\tan\beta = 1.5$ and 30, respectively, assuming $m_{H^+} = 200\text{GeV}$, $M_2 = 300\text{GeV}$, $\mu = 100\text{GeV}$, and $A_t = A_b = 500\text{GeV}$.




Figure 9: The genuine SUSY EW corrections versus μ for $\tan\beta = 1.5$ and 30, respectively, assuming $m_{H^+} = 200\text{GeV}$, $M_2 = 300\text{GeV}$, $A_t = A_b = 500\text{GeV}$ and $M_{\tilde{Q}} = M_{\tilde{U}} = M_{\tilde{D}} = 400\text{GeV}$.