

Optimal interior and boundary regularity for almost minimizers to elliptic variational integrals

FRANK DUZAAR *

KLAUS STEFFEN *

Abstract. We give a new proof of the small excess regularity theorems for integer multiplicity rectifiable currents of arbitrary dimension and codimension minimizing an elliptic parametric variational integral. This proof does not use indirect blow-up arguments, it covers interior and boundary regularity, it applies to almost minimizing currents, and it gives an explicit and often optimal modulus of continuity for the derivative, i.e. for the tangent plane field of the almost minimizing currents.

0 Introduction

The regularity theory for integer multiplicity rectifiable currents minimizing a parametric elliptic variational integral was initiated in the pioneering work of F.J. Almgren [Alm1], where the regularity is proved at interior support points with small cylindrical excess. Almgren's results are presented in F. Federer's monograph [Fe, Chap. 5]. E. Bombieri has given a somewhat simpler proof which also applies to approximately minimizing currents. This is important in order to be able to include problems with side conditions, such as volume constraints or obstacle conditions, in the regularity theory (see Section 1). In [Alm2] Almgren also treated the more general situation of almost minimizing currents and sets. A new and elegant proof of Almgren's original regularity theorem was presented by R. Schoen and L. Simon [SS]. This proof avoids the indirect blow-up arguments of the previous authors and it uses a direct P.D.E. argument instead. The interior regularity theory for (almost) minimizing currents has been thus quite well developed for some years.

With regard to boundary regularity the situation is different. One only has the work of R. Hardt [Ha1] which showed how to extend the reasoning from [Fe, Chap. 5.3] to the boundary situation. Apart from [Ha1], we are not aware of any published treatment of boundary regularity for (almost) minimizing currents and general elliptic integrands. (This problem is addressed, however, in the unpublished thesis [Li] of F.H. Lin.)

Our present work comes from an attempt to understand the question of boundary regularity better and to give a simpler proof of Hardt's boundary regularity result, and if possible to extend it to the case of almost minimizing currents. We first tried to carry over the simple method of [SS] to the boundary situation. This failed, however, for reasons explained below. Instead we have combined arguments from [Bo] with the idea of harmonic approximation as presented in [Si1, 21.1] and we have added some new estimates related to the integrands one obtains from a given smooth integrand by transforming it with diffeomorphisms that flatten the boundary. As a result, we have obtained small excess regularity theorems for almost minimizing currents which give an optimal modulus of continuity for the tangent plane field in many situations. It seems that even in the interior situation such

*We are grateful to the Volkswagen-Stiftung and to the Oberwolfach Mathematical Research Institute whose support through the RiP collaborative research program made the commencement of this work possible. We also acknowledge the support of the Sonderforschungsbereich 288 *Differential Geometry and Quantum Physics*, Berlin.



optimal regularity theorems are new and cannot be deduced from [Alm2], [Alm1], [Fe], [Bo], [SS] by simply giving quantitative versions of all the arguments involved.

We present our main result in Theorem 6.1 (the interior ϵ -regularity theorem), and Theorem 6.4 (the optimal boundary ϵ -regularity theorem). The precise technical assumptions are collected in Section 1 where we also explain the notion of almost minimizing currents. As an example for the boundary regularity statements one can deduce from Theorem 6.4 we state the following theorem:

0.1 Theorem (optimal boundary regularity). *Suppose F is a positive elliptic integrand of degree n and of class C^2 on the open subset D of \mathbb{R}^{n+k} and $0 < \beta < 1$. Then there exists $\vartheta > 0$ such that if T is an integer multiplicity rectifiable n -current which is almost minimizing for (F, ω) in D with modulus $\omega(r) = \text{const } r^{2\beta}$, if $a \in D \cap \text{spt}(\partial T)$ and $\text{spt}(\partial T)$ locally at a is an $(n-1)$ -dimensional submanifold of class $C^{1,\beta}$ in D , and if the lower density of $\|T\|$ at a satisfies*

$$\Theta_*^n(\|T\|, a) \leq \frac{1}{2} + \vartheta,$$

then $\text{spt } T$ is locally at a an n -dimensional submanifold of class $C^{1,\beta}$ up to the boundary.

It can be seen from the proof in Section 7 that the constant $\vartheta > 0$ depends only on the exponent β , the ellipticity constant of F , a bound for $F(x, \xi)$ and its derivatives up to second order with respect to the tangent n -plane variable ξ , and a modulus of continuity for the second derivative of F with respect to ξ . Moreover, a version of Theorem 0.1 holds for boundaries B satisfying a $C^{1,\kappa(\cdot)}$ Dini condition with modulus $\kappa(\cdot)$ such that $\mathbf{K}(r) = \int_0^1 \varrho^{-1} \kappa(\varrho) d\varrho < \infty$, and for moduli of almost minimality $\omega(\cdot)$ such that $\Omega(r) = \left[\int_0^1 \varrho^{-1} \sqrt{\omega(\varrho)} d\varrho \right]^2 < \infty$. Then $\text{spt } T$ is locally at a an n -dimensional submanifold of Dini class $C^{1,\gamma(\cdot)}$ up to the boundary with modulus $\gamma(r) = \text{const} \left[r^{2\beta} + \mathbf{K}(r)^2 + \Omega(r) \right]^{1/2}$ where $0 < \beta < 1$ can be prescribed. (The precise technical conditions on $\kappa(\cdot)$ and $\omega(\cdot)$ are described in Section 1.)

We should note that, in general, one cannot verify the density assumption for $\|T\|$ in Theorem 0.1 at least for some boundary points. In other words, it is not clear that the small excess boundary regularity theorem can be applied at all to minimizers of a given geometric variational problem. This unsatisfactory situation is in contrast to the interior theory where the ϵ -regularity theorem can always be applied at a dense set of points in the support and, hence, the regular set of an almost minimizing current is open and dense in its support.

Of course, if one makes additional assumptions concerning the prescribed boundary or the integrand, then one can say more. For example, in [Ha1, Sec. 4] it is proved that one has boundary regularity of codimension one currents near boundary points satisfying an appropriate barrier condition. For the area integrand, R. Hardt and L. Simon [HS] have even proved full boundary regularity for minimizing currents of codimension one. (This has been extended to minimizing currents with prescribed mean curvature in [DS2].) The interior singular set of area minimizing codimension one currents is known to be of codimension at least 7, by Federer's result [Fe1] (see [Ma], [MM], [DS1] for currents of prescribed mean curvature). With regard to the area integrand we also mention [HaL], Allard's work [All1], [All2] on regularity and boundary regularity of minimizers in the varifold setting, and Almgren's monumental theory [Alm3] which gives interior regularity up to codimension

two for area minimizing currents of arbitrary codimension. Finally, for codimension one minimizers of elliptic variational integrals we know that the singular set has codimension at least two, by the result of Schoen, Simon and Almgren [SSA]. In the present work, however, we are interested in general elliptic integrands and currents of arbitrary codimension, and we want to prove, starting from some weak initial assumptions (smallness of excess and a density condition) the best possible regularity for almost minimizing currents.

We now outline our proof of the regularity theorems. We begin in Section 2 with mass and height estimates for the rectifiable n -current T in \mathbb{R}^{n+k} which is almost minimizing with respect to an elliptic integrand F and has small excess over the n -plane $\mathbb{R}^n \times \{0\}^k$. These estimates are standard and extended to the boundary situation, where ∂T has support in $\mathbb{R}^{n-1} \times \{0\}^{k+1}$, without difficulties. The next step is to approximate T by the graph of a Lipschitz function g . The Lipschitz approximation is also well-known in the interior situation and easily established in the boundary case. We present in Section 3 a suitable version. Since T is almost minimizing for F it can be shown, along the lines of [Bo], that g is an approximate solution (in a certain sense) to the homogeneous system of P.D.E. associated with the constant coefficient elliptic quadratic form A that is obtained from F by freezing the coefficients at a point. We will call exact solutions of this system *A-harmonic* functions in what follows.

It is known from Almgren's work [Alm1] that the crucial step in the regularity proof is excess improvement, i.e. one must find a new n -plane \tilde{S} , obtained from $S = \mathbb{R}^n \times \{0\}^k$ by an isometry close to the identity, such that the excess of T over this *tilted* n -plane \tilde{S} is substantially smaller than the excess of T over S . In Section 5 we establish such an excess improvement (Lemma 5.2) and derive from it a growth condition for the cylindrical excess as a function of the radius of the cylinder (Lemma 5.4). From this excess growth condition the ϵ -regularity theorems of Section 6 then follow in a well-known manner, e.g. as in [Ha1], but we give full proofs in order to obtain an explicit modulus of continuity for the tangent plane field of T .

The essential point in which our proof differs from the previous ones, is the method of finding the tilted n -plane \tilde{S} with improved excess. In Almgren's and Bombieri's work [Alm1], [Alm2], [Bo], [Fe, 5.3] this is done using blow-up by a sequence of certain scaling transformations σ_ν for which the scaled Lipschitz approximations g_ν converge to an *A*-harmonic function u . The plane \tilde{S} is then obtained from the tangent plane to graph u (at the appropriate point) by the inverse scaling σ_ν^{-1} with $\nu \gg 1$. This proof of the excess improvement lemma is indirect and, hence, does not give an explicit constant in the estimate.

The idea of Schoen and Simon [SS] is to choose u as a solution of the Dirichlet problem associated with A on a ball, taking the values of the Lipschitz approximation g as boundary values. The plane \tilde{S} is then chosen as the tangent plane to graph u at the center of the ball. However, one needs a $C^{1,\delta}$ estimate for u up to the boundary of the ball in order to prove that the excess of T over \tilde{S} is substantially smaller, and therefore g has to be replaced by a smoothing g_δ of g with appropriately chosen smoothing radius (depending on the size of the excess over the original plane S). It is this point which creates problems when treating the boundary regularity problem. There one would have to consider half balls, and we could not find a way to smooth the half balls and the Lipschitz function g in such a way that the crucial estimates could be obtained along the lines of [SS]. Furthermore, as Schoen and Simon employ the Euler equation for the minimizing current T , it is not immediately clear

how to treat almost minimizing currents.

Our method here is to replace the Lipschitz function g , which is already known to be approximatively A -harmonic, by a nearest A -harmonic function, in a sense. This is made possible by a simple lemma (Lemma 3.3) analogous to [Si2, 21.1]. In order to show that the tangent plane \tilde{S} to the graph of this function u gives the desired excess improvement we need a reverse Poincaré inequality (Caccioppoli inequality). We establish this in Lemma 4.1 in a weak form, which can be proved in the boundary situation as in [Bo], and we use a covering argument from [Si2] to derive from this the required strong version (Lemma 4.2).

Of course, all the ingredients of our proof have already been used previously in regularity theory. In particular, the idea of approximating (almost) minimizing currents by the solution of a constant coefficient elliptic P.D.E. system is by now a standard technique, dating back to DeGiorgi [DeG]. We believe, however, that the way in which we have combined the ingredients in our proof is new and has definite conceptual advantages: It is the first unified treatment of interior and boundary regularity; it applies to almost minimizing currents; it does not use an indirect blow-up argument and gives constants which are (in principle) explicitly computable in the regularity estimates; it enables one to derive an explicit and frequently optimal modulus of continuity for the derivative (i.e. the tangent plane field); it uses only natural and quite weak assumptions on the elliptic integrand; it is very flexible and can be applied to regularity questions related to (almost) minimizers of variational integrals or to weak solutions of nonlinear elliptic P.D.E. systems in a variety of different situations. This last point is perhaps the most important one. In fact, variants of the method used in this paper play a role in [DGr], [DGa], and [DGG] to obtain new and sometimes optimal regularity results.

Finally, we want to point out that the fact that our proof only requires minimal assumptions on the integrand F is not merely an irrelevant technicality, but essential for optimal boundary regularity. Namely, if we study boundary regularity of an (almost) minimizing current T for a smooth elliptic integrand F , and we have small excess and a prescribed boundary of class $C^{1,\beta}$, say, then an almost unavoidable procedure is to flatten the boundary by transforming it into an affine subspace with the same dimension using suitable $C^{1,\beta}$ diffeomorphisms. This will give correspondingly transformed currents \tilde{T} and integrands \tilde{F} such that \tilde{T} is almost minimizing with respect to \tilde{F} and \tilde{T} has a flat boundary. However, $\tilde{F}(x, \xi)$ will only be $C^{1,\beta}$ with respect to the space variable x , and therefore we have to prove $C^{1,\beta}$ regularity for minimizers to integrands with this weak regularity in the case of a flat boundary in order to derive the expected $C^{1,\beta}$ regularity of T up to the boundary. In fact, with a straightforward adaptation of the interior regularity theory to the case of a flat boundary, with the given $C^{1,\beta}$ smoothness of \tilde{F} , we could only establish $C^{1,\beta/(2-\beta)}$ regularity of \tilde{T} losing something in the Hölder exponent. (We do not know whether this loss, which also occurs in a variety of other situations, is merely a technical problem or is indeed unavoidable.) In our estimates therefore we needed to keep distinct the continuity modulus of $x \mapsto \tilde{F}(x, \xi)$ from the continuity modulus of \tilde{F} of the same function in directions perpendicular to the flat boundary. For \tilde{F} as above, this second modulus is substantially smaller than the first one. Carefully tracking how the different moduli enter the various estimates (which is possible in our regularity proof), we were able to prove in the end $C^{1,\beta}$ regularity of T up to the boundary without any loss in the Hölder exponent. This way of proceeding is a new technique introduced in the present paper that may also be of interest with regard to other boundary regularity problems.

1 Rectifiable currents, parametric integrands, and almost minimizers

By an n -dimensional **integer multiplicity rectifiable current** T on \mathbb{R}^{n+k} we mean a generalized oriented surface which can be represented by integration as a linear functional on the space of smooth n -forms α with compact support in \mathbb{R}^{n+k} as follows

$$\langle T, \alpha \rangle = \int_{M_T} \langle \alpha, \vec{T} \rangle \vartheta_T d\mathcal{H}^n = \int_{\mathbb{R}^{n+k}} \langle \alpha, \vec{T} \rangle d\|T\|.$$

Here \mathcal{H}^n denotes the n -dimensional Hausdorff measure in \mathbb{R}^{n+k} , the **supporting set** $M_T \subset \mathbb{R}^{n+k}$ is countably (\mathcal{H}^n, n) rectifiable (i.e. it can be covered by countably many n -submanifolds of class C^1 in \mathbb{R}^{n+k} and a set of zero \mathcal{H}^n measure), ϑ_T is a positive integer valued \mathcal{H}^n summable **multiplicity function**, and \vec{T} is an n -measurable **orientation n -vectorfield** for M_T (i.e. $\vec{T}(x)$ equals, at \mathcal{H}^n almost every point $x \in M_T$, the exterior product of some orthonormal basis in the n -dimensional approximate tangent space $\text{Tan}^n(\mathcal{H}^n \llcorner M_T, x)$). As usual, we have written $\|T\|$ for the Radon measure $\mathcal{H}^n \llcorner \vartheta_T$ associated with T , understanding that ϑ_T is extended by the value zero on $\mathbb{R}^{n+k} \setminus M_T$, and we note that summability of ϑ_T is equivalent to finiteness of the **mass** (or **n -area**)

$$\mathbf{M}(T) = \|T\|(\mathbb{R}^{n+k}) = \int_{M_T} \vartheta_T d\mathcal{H}^n.$$

We refer to [Fe] and [Si1] for the basic notions and notations related to currents. Notations not explained in this paper are taken from [Fe]. Our usage is consistent with this reference, except that what we have called an integer multiplicity rectifiable current above would be a locally rectifiable current with finite mass in the language of [Fe]. The abelian group of these n -dimensional currents on \mathbb{R}^{n+k} is denoted by $\mathcal{R}_n(\mathbb{R}^{n+k})$, its elements will also be called rectifiable n -currents or simply n -currents. We will always assume dimension $n \geq 2$ and codimension $k \geq 1$.

A **parametric integrand** of degree n on an open set D in \mathbb{R}^{n+k} is a continuous map

$$F: D \times \bigwedge_n \mathbb{R}^{n+k} \rightarrow \mathbb{R}$$

which is positively homogeneous of degree one in the second variable, i.e. $F(x, r\xi) = rF(x, \xi)$ for $r > 0$. The **integral** of F over an n -current as above (with $\vartheta_T \equiv 0$ outside D) is then

$$\mathbf{F}(T) = \int_{M_T} F(x, \vec{T}(x)) \vartheta_T d\mathcal{H}^n x = \int_D F(x, \vec{T}(x)) d\|T\|_x,$$

provided the integral exists and is finite. This is the case, for example, if $|F|$ is bounded by the **area integrand**, i.e. $|F(x, \xi)| \leq \Lambda |\xi|$ holds for all (x, ξ) with a finite constant Λ . For the purpose of regularity theory it is no restriction to assume $D = \mathbb{R}^{n+k}$.

We now list the **assumptions on the integrand** which we shall use in our regularity theorems. The first set of assumptions requires bounds for $F(x, \xi)$ and its derivatives $D_{(2)}F$, $D_{(2)}^2 F$ with respect to the tangent plane variable $\xi \in \bigwedge_n \mathbb{R}^{n+k}$ as follows:

$$(1.1) \quad |\xi| \leq F(x, \xi) \leq \Lambda |\xi|,$$

$$(1.2) \quad \|D_{(2)}F(x, \xi)\| \leq \Lambda,$$

$$(1.3) \quad \|D_{(2)}^2F(x, \xi)\| \leq \Lambda|\xi|^{-1},$$

for all $x \in \mathbb{R}^{n+k}$ and $0 \neq \xi \in \Lambda_n \mathbb{R}^{n+k}$, where $\Lambda \geq 1$ is a finite constant. (1.1) means, in particular, that F is estimated from above and from below by the **area integrand** $M(x, \xi) = |\xi|$. (It is convenient to assume the constant 1 in the estimate from below; this is, of course, no essential restriction.) We also need a uniform modulus of continuity for $\xi \mapsto D_{(2)}^2F(x, \xi)$,

$$(1.4) \quad \left\| D_{(2)}^2F(x, \xi) - D_{(2)}^2F(x, \tilde{\xi}) \right\| \leq \nu(|\xi - \tilde{\xi}|)$$

for all $x \in \mathbb{R}^{n+k}$ and $\xi, \tilde{\xi} \in \Lambda_n \mathbb{R}^{n+k}$ with $|\xi| = 1 = |\tilde{\xi}|$, where $\nu(0+) = 0 = \nu(0)$. We assume continuity of F and $D_{(2)}F$ with respect to the space variable x as follows:

$$(1.5) \quad |F(x, \xi) - F(\tilde{x}, \xi)| \leq \kappa(|x - \tilde{x}|)|\xi|,$$

$$(1.6) \quad \left\| D_{(2)}F(x, \xi) - D_{(2)}F(\tilde{x}, \xi) \right\| \leq \kappa(|x - \tilde{x}|)$$

for all x, \tilde{x}, ξ , where $\kappa: [0, \infty[\rightarrow [0, \infty]$ is nondecreasing with $\kappa(0+) = 0 = \kappa(0)$ and $]0, \infty[\ni r \mapsto \frac{1}{r}\kappa(r)$ nonincreasing. We may also assume that $\kappa(1)$ is as small as we like; this can be achieved by scaling with homotheties. In the proof of the regularity theorem we need a restriction on the decay of $\kappa(r)$ as $r \downarrow 0$, cf. (1.15). This condition is satisfied when $\kappa(r) \leq r^\beta$ holds for some $\beta > 0$ and all sufficiently small $r > 0$. We also have to require that $r \mapsto r^{-\alpha}\kappa(r)$ is nonincreasing for some exponent $0 < \alpha < 1$.

In (1.1)–(1.6) we assume implicitly that all the appearing derivatives of F depend continuously on (x, ξ) , $\xi \neq 0$. With regard to the notation of norms we use $|\cdot|$ to denote natural Euclidean norms and $\|\cdot\|$ for operator type norms (largest Euclidean norm of values obtained from evaluating on unit vectors).

For the treatment of boundary regularity we also need the continuity modulus $\mu(\cdot)$ of $x \mapsto F(x, \xi)$ in directions perpendicular to the boundary tangent plane $\mathbb{R}^{n-1} \times \{0\}$, i.e.

$$(1.7) \quad |F(x, \xi) - F(\tilde{x}, \xi)| \leq \mu(|x - \tilde{x}|)|\xi|, \quad \text{if } x_i = \tilde{x}_i \text{ for } i = 1, \dots, n-1,$$

$$(1.8) \quad \left\| D_{(2)}F(x, \xi) - D_{(2)}F(\tilde{x}, \xi) \right\| \leq \mu(|x - \tilde{x}|), \quad \text{if } x_i = \tilde{x}_i \text{ for } i = 1 \dots, n-1,$$

where $0 \leq \mu(r) \leq \kappa(r)$ is a nondecreasing function of $r \geq 0$ with $]0, \infty[\ni r \mapsto \frac{1}{r}\mu(r)$ nondecreasing and a further condition stated in (1.15) below. The reason for introducing the modulus $\mu(\cdot)$ is the following: If F is obtained from a smooth integrand by transformation with a diffeomorphism Φ that maps a given C^1 submanifold, of \mathbb{R}^{n+k} to its tangent plane $\mathbb{R}^{n-1} \times \{0\} \subset \mathbb{R}^{n+k}$ at the origin and has the form $\Phi(x) = x + \Psi(x_1, \dots, x_{n-1})$, then the modulus $\kappa(\cdot)$ of F is essentially the continuity modulus of the tangent plane distribution of, while $\mu(r) \leq \text{const } r$ is a Lipschitz modul even when, is not of class $C^{1,1}$. We shall exploit this fact to obtain optimal moduli of continuity in the boundary regularity theorem.

With $\kappa(\cdot)$ and $\mu(\cdot)$ we associate the following functions $\widehat{\kappa}(\cdot)$ and $\widehat{\mu}(\cdot)$ which enter the regularity estimates:

$$(1.9) \quad \widehat{\kappa}(r) = \sup_{0 \leq 2s \leq 1} \left[\kappa(s) - \left(\frac{s}{r}\right)^2 \right] \quad \text{and} \quad \widehat{\mu}(r) = \sup_{0 \leq 2s \leq 1} \left[\mu(s) - \left(\frac{s}{r}\right)^2 \right] \quad \text{for } r > 0.$$

One readily verifies that $\widehat{\mu}(\cdot)$ is nondecreasing with $\widehat{\mu}(0+) = 0$ and $\widehat{\mu}(\frac{1}{2}) \leq \mu(\frac{1}{2}) \leq \frac{1}{4} + \widehat{\mu}(1)$. Furthermore, from the fact that $\mu(\vartheta^2 r) \geq \vartheta^2 \mu(r)$ holds for $0 < \vartheta \leq 1$ we deduce $\widehat{\mu}(\vartheta r) \geq \vartheta^2 \widehat{\mu}(r)$ and, hence, that $r \mapsto \frac{1}{r} \sqrt{\widehat{\mu}(r)}$ is nonincreasing. We also note that $\frac{1}{4} \mu(r)^2 \leq \widehat{\mu}(r) \leq \mu(r)$ holds for $0 < r \leq r_0$ whenever $0 < r_0 \leq \frac{1}{2}$ and $\mu(r_0) \leq 1$. The first inequality follows by taking $s = \frac{1}{\sqrt{2}} r \mu(\frac{1}{\sqrt{2}} r)$ in (1.9) and using $\mu(s) \geq \frac{1}{\sqrt{2}} \mu(\frac{1}{\sqrt{2}} r)^2 \geq \frac{1}{2} \mu(r)^2$, by the nonincreasing property of $r \mapsto \frac{1}{r} \mu(r)$. For the second inequality one observes that $\mu(s) - (\frac{s}{r})^2$ is nonpositive for $r \leq s \leq \frac{1}{2}$ and uses again (1.9) together with $\mu(s) \leq \frac{s}{r} \mu(r)$ for $0 < s \leq r$. As an example, we have $\widehat{\mu}(r) = \text{const } c^{\frac{2}{2-\beta}} r^{\frac{2\beta}{2-\beta}}$ for $0 < r \leq 1$, if $\mu(s) = cs^\beta$ with $0 < \beta \leq 1$ and $0 \leq c \leq 2^{\beta-2}$. The same statements hold for κ and $\widehat{\kappa}$, of course.

When dealing with the nonparametric situation where the n -currents are represented by graphs in $\mathbb{R}^n \times \mathbb{R}^k$ we will have to consider the associated **nonparametric integrand**

$$F^\S: \mathbb{R}^n \times \mathbb{R}^k \times \text{Hom}(\mathbb{R}^n, \mathbb{R}^k) \rightarrow \mathbb{R}$$

which is characterized by

$$\mathbf{F}(T_g) = \int_{\Omega} F^\S(y, g(y), Dg(y)) d\mathcal{L}^n y$$

whenever $g: \Omega \rightarrow \mathbb{R}^k$ is Lipschitz on a bounded domain Ω in \mathbb{R}^n and T_g is the n -current representing the graph of g with natural orientation and multiplicity one. We have

$$F^\S(y, z, p) = F\left((y, z), \bigwedge_n (\text{id}, p) e_1 \wedge \dots \wedge e_n\right),$$

where e_1, \dots, e_n is the canonical basis in \mathbb{R}^n and $(\text{id}, p)(y) = (y, py)$ for $y \in \mathbb{R}^n$, $p \in \text{Hom}(\mathbb{R}^n, \mathbb{R}^k)$. It is convenient to assume conditions analogous to (1.3), (1.4) also for the total second order derivative $D_{(3)}^2 F^\S$ of F^\S with respect to p , i.e.

$$(1.10) \quad \left\| D_{(3)}^2 F^\S(y, z, p) \right\| \leq \Lambda \quad \text{for all } y, z, p \text{ with } |p| \leq 1,$$

$$(1.11) \quad \left\| D_{(3)}^2 F^\S(y, z, p) - D_{(3)}^2 F^\S(y, z, \tilde{p}) \right\| \leq \nu(|p - \tilde{p}|)$$

for all y, z, p, \tilde{p} with $|p| \leq 2, |\tilde{p}| \leq 2$.

Actually these conditions follow from (1.1)–(1.4) with suitably modified constant Λ and modulus ν . For ν we will assume, as we may, that it is a nondecreasing function with $\nu(0+) = 0 = \nu(0)$.

We next come to the assumption of ellipticity for F . This is a condition on the constant coefficient integrands $F_a(x, \xi) = F(a, \xi)$ obtained from F by freezing the coefficients at $a \in \mathbb{R}^{n+k}$. Following [Fe, 5.1.2] we call F **elliptic** at a , if there exists a positive constant λ such that the inequality

$$(1.12) \quad \mathbf{F}_a(T) - \mathbf{F}_a(S) \geq \lambda [\mathbf{M}(T) - \mathbf{M}(S)]$$

holds whenever S, T are (compactly supported) rectifiable n -currents on \mathbb{R}^{n+k} with the same boundary $\partial S = \partial T$ and S is represented by an \mathcal{H}^n measurable subset of some n -dimensional affine subspace in \mathbb{R}^{n+k} with constant orientation n -vectorfield and \mathcal{H}^n summable positive integer valued multiplicity. (This is the ellipticity condition convenient for us. In Almgren's original definition [Alm1] only flat n -disks were considered. The ellipticity definition in [Bo] is an adaptation of this taking multiplicities into account; it assumes (1.12) only when S is an integer multiple of a flat n -disc with constant orientation. It can be shown that this is equivalent to the definition above.) We will suppose that F is **uniformly elliptic** in the sense that (1.12) holds with the same constant λ for all points $a \in \mathbb{R}^{n+k}$. This is no restriction as our considerations are of local nature. For the **ellipticity constant** we assume $0 < \lambda \leq 1$.

If F is elliptic at $a = (b, c) \in \mathbb{R}^n \times \mathbb{R}^k$, then the associated nonparametric integrand satisfies the **Legendre-Hadamard condition** in the sense that the Hessian of $F^\S(b, c, p)$ with respect to the variable $p = (p_\alpha^i) \in \text{Hom}(\mathbb{R}^n, \mathbb{R}^k)$ is rank one elliptic. In particular, if (1.12) holds, then with the same constant λ we have ([Fe, 5.1.10])

$$(1.13) \quad \sum_{\alpha, \beta=1}^n \sum_{i, j=1}^k \frac{\partial^2 F^\S}{\partial p_\alpha^i \partial p_\beta^j}(b, c, 0) \eta_\alpha \zeta^i \eta_\beta \zeta^j \geq \lambda |\eta|^2 |\zeta|^2$$

for all ζ in \mathbb{R}^k and η in (the dual space of) \mathbb{R}^n . The nonparametric Legendre-condition does not imply, however, ellipticity (1.12) if $n \geq 2, k \geq 2$ (see [Sv2]). On the other hand, the parametric Legendre condition [Fe, 5.1.3], which was adopted in [SS] as the definition of ellipticity, is presumably more restrictive than (1.12) if $n \geq 2, k \geq 2$. This is suggested by the analogous result for nonparametric variational integrals (e.g. [Sv1]).

Recalling [Fe, 5.1.1] for the transformation of integrands F by diffeomorphisms Φ and [Fe, 5.1.4] for the preservation of ellipticity under such transformations one readily verifies that the integrand \tilde{F} , obtained from the transformed integrand $\Phi^\# F$ by multiplying with a suitable positive factor and scaling with a suitable homothety of \mathbb{R}^{n+k} , satisfies the same set of conditions (1.1)–(1.13) with appropriate constants $\tilde{\Lambda}, \tilde{\lambda}$ and moduli $\tilde{\kappa}, \tilde{\nu}$, provided Φ is a biLipschitz diffeomorphism of \mathbb{R}^{n+k} with $D^2\Phi$ bounded or $D\Phi$ Lipschitz, at least. In this sense our assumptions on the integrand are invariant with respect to biLipschitz diffeomorphisms of class $C^{1,1}$.

A rectifiable n -current T is called F **minimizing** in an open subset D of \mathbb{R}^{n+k} if $\mathbf{F}(T) \leq \mathbf{F}(T + X)$ for all closed n -currents X with compact support in D . Following Almgren [Alm2] and Bombieri [Bo] we consider a relaxed minimizing condition which allows the existence of modifications $T + X$ of T with $\mathbf{F}(T + X)$ smaller than $\mathbf{F}(T)$, but only by a factor $1 + \omega(r)$ close to 1 if X has its support in a ball of small radius r . We adopt the definition from [Bo] which is as follows: For a given function $\omega:]0, \infty[\rightarrow [0, \infty]$ one says that T is **almost minimizing** for (F, ω) in D , or (F, ω) minimizing for short, if

$$(1.14) \quad \mathbf{F}(T) \leq \mathbf{F}(T + X) + \omega(r) \mathbf{M}(T \llcorner K + X)$$

holds for all closed n -currents X with support in a compact set $K \subset D$ which is contained in a ball of radius r in \mathbb{R}^{n+k} .

With regard to ω we will always assume that it is a nondecreasing function and finite for the values of r appearing in the following. For the regularity theory we need (1.14) only for small radii r , of course, i.e. we can admit $\omega(r) = \infty$ for $r \geq \delta$ and some $\delta > 0$.

Furthermore, it would be sufficient to know (1.14) for all X satisfying an additional mass bound $\mathbf{M}(X) \leq \gamma$ with some constant $0 < \gamma < \infty$. Additional assumptions on ω and on the moduli $\kappa, \widehat{\kappa}, \mu, \widehat{\mu}$ defined previously are needed for the iteration argument in the proof of the excess decay, Lemma 5.4. There we assume that $r^{-\alpha}\kappa(r), r^{-\alpha}\mu(r), r^{-2\alpha}\omega(r)$ are nonincreasing functions of $r > 0$ for some exponent $0 < \alpha < 1$ (this is not an essential restriction, but it excludes Lipschitz modules $\kappa(r) = \text{const}r$; as before, it follows that $r \mapsto r^{-2\alpha}\widehat{\kappa}(r)$ and $r \mapsto r^{-2\alpha}\widehat{\mu}(r)$ are also nonincreasing) and we require that

$$(1.15) \quad \begin{aligned} \mathbf{K}(r) &= \int_0^r \frac{1}{\varrho} \kappa(\varrho) d\varrho < \infty, & \widehat{\mathbf{K}}(r) &= \left[\int_0^r \frac{1}{\varrho} \sqrt{\widehat{\kappa}(\varrho)} d\varrho \right]^2 < \infty, \\ \widehat{\mathbf{M}}(r) &= \left[\int_0^r \frac{1}{\varrho} \sqrt{\widehat{\mu}(\varrho)} d\varrho \right]^2 < \infty, & \Omega(r) &= \left[\int_0^r \frac{1}{\varrho} \sqrt{\omega(\varrho)} d\varrho \right]^2 < \infty \end{aligned}$$

for small $r > 0$. This growth condition for $\omega(r)$ as $r \downarrow 0$ appears already in [Alm2], [Bo], [Ta1] and it is, in a sense, necessary for C^1 regularity of (F, ω) minimizers as shown by examples in [Ta1] (for the area integrand with codimension $k = 1$ and a somewhat different notion of almost minimality). Note that $\omega(r) = r^{2\beta}$ and $\kappa(r) = r^\beta$ with $0 < \beta < 1$ satisfies all the assumptions above with $\Omega(r) = \beta^{-2}r^{2\beta}$ and $\mathbf{K}(r) = \beta^{-1}r^\beta$. In view of $\widehat{\mu}(r) \leq \mu(r)$ for sufficiently small $r > 0$ the condition $\widehat{\mathbf{M}}(r) < \infty$ is satisfied for small $r > 0$ whenever $\int_0^r \frac{1}{\varrho} \sqrt{\mu(\varrho)} d\varrho < \infty$ holds, so that $\mu(r) = r^\beta$ with $0 < \beta < 1$ is again admissible here.

We now discuss some situations where (F, ω) minimizing currents arise. The first examples are minimizers for **functionals of mean curvature type**. Here we assume that the n -current T is minimizing for $\mathbf{F} + \mathbf{V}$ on \mathbb{R}^{n+k} where \mathbf{V} is some kind of additive volume functional. More precisely, suppose $|\mathbf{V}(X)| \leq \gamma\mathbf{M}(Q)$ whenever X is a closed rectifiable n -current and Q an $(n+1)$ -current of least mass with $\partial Q = X$. Then we have

$$(1.16) \quad \mathbf{F}(T) \leq \mathbf{F}(T + X) + \mathbf{V}(X) \leq \mathbf{F}(T + X) + \gamma\mathbf{M}(Q),$$

and if $\text{spt } X \subset K$, where K is contained in a ball of radius r , we can estimate $\mathbf{M}(Q)$ by the mass of the cone over X , to obtain

$$(1.17) \quad \mathbf{M}(Q) \leq \frac{r}{n+1}\mathbf{M}(X) \leq \frac{r}{n+1}[\mathbf{M}(T \llcorner K + X) + \mathbf{M}(T \llcorner K)].$$

With the help of (1.1) we readily deduce from (1.16), (1.17) that T is (F, ω) minimizing with a function $\omega(r) = O(r)$ as $r \downarrow 0$. Functionals of the type $\mathbf{M} + \mathbf{V}$ occur as energies in the variational approach to hypersurfaces with prescribed mean curvature or surfaces with prescribed mean curvature vector (e.g. [Du1], [DF1], [DF2], [DS3], [Ma]). In [DS1] currents with the almost minimizing property (1.16) for the area integrand $F = M$ were systematically studied.

A similar situation occurs when T minimizes \mathbf{F} subject to a **volume constraint** $\mathbf{V}(T) = \text{const}$. If the first variation of \mathbf{V} does not vanish and X, K, r are as above with r sufficiently small and $\mathbf{M}(X)$ bounded by a finite constant, then we can correct the volume of $T + X$ by a deformation h_t which is the identity on K to achieve $\mathbf{V}(h_{t\#}T + X) = \mathbf{V}(T)$ and $|\mathbf{F}(h_{t\#}T + X) - \mathbf{F}(T + X)| \leq \text{const}|\mathbf{V}(X)|$. Since $h_{t\#}T + X$ satisfies the volume constraint we have $\mathbf{F}(T) \leq \mathbf{F}(h_{t\#}T + X)$, and with $|\mathbf{V}(X)| \leq \gamma\mathbf{M}(Q)$ and (1.17) we see as before that T is (F, ω) minimizing (for the unconstrained problem) with $\omega(r) = O(r)$ as $r \downarrow 0$.

See [Du2], [GMT], [Ta3] for a more thorough discussion in the case $F = M$ and $k = 1$ and [Alm2] for general F and $k = 1$.

If T minimizes F subject to an **obstacle condition** $\text{spt} T \subset A$, where $A \subset \mathbb{R}^{n+k}$ is closed and $\mathbb{R}^{n+k} \setminus A$ is the ‘‘obstacle’’, then one can derive (F, ω) minimality of T (for the unconstrained problem), provided A is sufficiently regular at the boundary and the integrand F is compatible with the obstacle. By the latter condition we mean that for $0 < r$ sufficiently small and each point $a \in \partial A$ there is a Lipschitz retraction π_a of the ball $\mathbf{U}(a, r)$ in \mathbb{R}^{n+k} onto $A \cap \mathbf{U}(a, r)$ such that $\mathbf{F}_a(\pi_{a\#} S) \leq \mathbf{F}_a(S)$ holds for n -currents S with $\text{spt} S \subset \mathbf{U}(a, r)$ and $\text{spt} \partial S \subset A \cap \mathbf{U}(a, r)$. If X is closed with support in $\mathbf{U}(a, r)$, then we can use $\pi_{a\#}(T \llcorner \mathbf{U}(a, r) + X)$ as a comparison current for $T \llcorner \mathbf{U}(a, r)$, and with the help of (1.5) we deduce that T is (F, ω) minimizing with $\omega(r) = O(r)$ as $r \downarrow 0$ (assuming that the conditions on the obstacle hold uniformly for all $a \in \partial A$). More generally we assume that π_a retracts $\mathbf{U}(a, r)$ without decreasing \mathbf{F}_a onto a subset of $\mathbf{U}(a, r)$ which is diffeomorphic to $A \cap \mathbf{U}(a, r)$ by a C^1 diffeomorphism φ of $\mathbf{U}(a, r)$ that deviates in the C^1 norm from the identity by at most $\text{const } r^\beta$, $0 < \beta \leq 1$. Using then $(\varphi \circ \pi_a \circ \varphi^{-1})_{\#}(T + X) \llcorner \mathbf{U}(a, r)$ as comparison current for $T \llcorner \mathbf{U}(a, r)$ in the obstacle problem we obtain the (F, ω) -minimizing property for T with $\omega(r) = O(r^\beta)$ as $r \downarrow 0$. This can be applied, for example, if $F = M$ is the area integrand and A is the closure of a (uniform) $C^{1,\beta}$ domain with π_a the nearest point retraction onto the affine tangent half space to A at a . (Actually $\omega(r) = O(r^{2\beta})$ in this case by the argument below.) More generally we may take A locally uniformly $C^{1,\beta}$ equivalent, in a suitable sense, to a convex set. For general elliptic integrands F , codimension $k = 1$, and A the closure of a uniform $C^{1,\beta}$ domain we can again use the nearest point retraction π_a onto the tangent half space of A at a , as the \mathbf{F}_a nonincreasing property of $\pi_{a\#}$ is then a consequence of ellipticity. Thus, solutions of obstacle problems will be (F, ω) minimizing with $\omega(r) = O(r^\beta)$. We refer to [Alm2], [Li], and (for the area integrand with codimension 1 and multiplicity 1) to [Ta1], [Mi].

Finally, we want to indicate that every oriented uniform $C^{1,\beta}$ submanifold \mathcal{T} of dimension n in \mathbb{R}^{n+k} is (M, ω) minimizing with M the area integrand and $\omega(r) = O(r^{2\beta})$. More precisely, the n -current T defined by integration over \mathcal{T} (with multiplicity 1) has this property. Considering $a \in \mathcal{T}$ and a closed n -current X with support in a compact set $K \subset \mathbf{U}(a, r)$ and letting π_a be the orthonormal projection of \mathbb{R}^{n+k} onto $\text{Tan}_a \mathcal{T}$ we observe

$$\mathbf{M}(\pi_{a\#}(T \llcorner K)) = \mathbf{M}(\pi_{a\#}(T \llcorner K + X)) \leq \mathbf{M}(T \llcorner K + X),$$

and hence it suffices to prove

$$\mathbf{M}(T \llcorner K) \leq (1 + \omega(r))\mathbf{M}(\pi_{a\#}(T \llcorner K))$$

with $\omega(r)$ as asserted. This is easily done by writing $\mathbf{U}(a, r) \cap \mathcal{T} = \mathbf{U}(a, r) \cap (\text{graph } u)$ as a graph over the tangent plane $\text{Tan}_a \mathcal{T}$, if r is small enough, and then using the estimate $|Du(y)| \leq \text{const}|y|^\beta$ in the area integral of the corresponding function u . For further discussions of the (F, ω) minimizing property of solutions to parametric variational problems we refer to [Alm2] and (for the area integrand with codimension 1 and multiplicity one, and with a somewhat different definition of almost minimality) to [Ta2].

We will frequently use the following **scaling property**: If the rectifiable n -current T is (F, ω) minimizing in $D \subset \mathbb{R}^{n+k}$, $a \in \mathbb{R}^{n+k}$, $\vartheta > 0$, $\tau_a(x) = x + a$ denotes the translation in \mathbb{R}^{n+k} , $\mu_\vartheta(x) = \vartheta x$ denotes the homothety in \mathbb{R}^{n+k} , $\tilde{T} = \mu_{\vartheta\#}^{-1} \tau_{a\#}^{-1} T$, and $\tilde{F}(\tilde{x}, \xi) =$

$F(a + \vartheta \tilde{x}, \xi)$ for $\tilde{x} \in \tilde{D} = \boldsymbol{\mu}_\vartheta^{-1} \boldsymbol{\tau}_a^{-1} D$, then \tilde{T} is $(\tilde{F}, \tilde{\omega})$ minimizing in \tilde{D} with $\tilde{\omega}(\tilde{r}) = \omega(\vartheta \tilde{r})$. Here we have taken \tilde{F} instead of the transformed integrand $\tilde{F} = (\boldsymbol{\tau}_a \circ \boldsymbol{\mu}_\vartheta)^\# F = \vartheta^n \tilde{F}$, because then \tilde{F} has the same structure constants $\tilde{\Lambda} = \Lambda$, $\tilde{\lambda} = \lambda$ and modulus $\tilde{\nu} = \nu$ as F in (1.1)–(1.4). For the moduli appearing in (1.5)–(1.8) we then have $\tilde{\kappa}(r) = \kappa(\vartheta r)$, $\tilde{\mu}(r) = \mu(\vartheta r)$. Furthermore, $\hat{\tilde{\mu}}(r) \leq \hat{\mu}(\vartheta r)$ can be easily derived from the definition (1.9) if $0 < \vartheta \leq 1$. We also note, for biLipschitz maps G with biLipschitz constant L , that $G_\#^{-1} T$ is $(G^\# F, \tilde{\omega})$ minimizing with $\tilde{\omega}(\tilde{r}) = L^n \omega(L\tilde{r})$.

In the ϵ -regularity theorem (Theorem 6.1) one considers a situation where the (F, ω) minimizing n -current has special projection properties and small cylindrical excess with respect to an n -plane in \mathbb{R}^{n+k} . We fix this plane to be $\mathbb{R}^n \times \{0\}^k \subset \mathbb{R}^n \times \mathbb{R}^k$ and introduce the following **notations** to be used throughout this paper: $\mathbf{p}: \mathbb{R}^{n+k} \rightarrow \mathbb{R}^n$, $\mathbf{q}: \mathbb{R}^{n+k} \rightarrow \mathbb{R}^k$ are the projections, i.e. $\mathbf{p}(x) = y$, $\mathbf{q}(x) = z$ for $x = (y, z) \in \mathbb{R}^n \times \mathbb{R}^k$. Open balls in \mathbb{R}^n are denoted by $\mathbf{U}_\varrho(y)$, the center y is omitted if $y = 0$, the radius ϱ is omitted if $\varrho = 1$, so that \mathbf{U}_ϱ and \mathbf{U} are open balls in \mathbb{R}^n with radius ϱ and 1 respectively and centered at the origin. Closed balls are denoted similarly with \mathbf{U} replaced by \mathbf{B} . Cylinders in \mathbb{R}^{n+k} over open balls in \mathbb{R}^n will be written $\mathbf{C}_\varrho(y) = \mathbf{U}_\varrho(y) \times \mathbb{R}^k = \mathbf{p}^{-1}(\mathbf{U}_\varrho(y))$ with the same conventions for the suppressing of y or ϱ , in particular we will usually work in the unit cylinder $\mathbf{C} = \mathbf{U} \times \mathbb{R}^k$. Closed cylinders are denoted using $\overline{\mathbf{C}}$ instead of \mathbf{C} . The superscript or the subscript ”+” at open balls in \mathbb{R}^n indicates intersection with the half space $\mathbb{R}^{n-1} \times]0, \infty[$, so that e.g. \mathbf{U}_+ is the open unit half ball in \mathbb{R}^n . Similarly, $\mathbf{C}_\varrho^+ = \mathbf{C}_\varrho(y) \cap \mathbb{R}^{n-1} \times]0, \infty[\times \mathbb{R}^k$. The closure of $\mathbf{U}_\varrho^+(y)$ and $\mathbf{C}_\varrho^+(y)$ is written $\mathbf{B}_\varrho^+(y)$ and $\overline{\mathbf{C}}_\varrho^+(y)$ respectively, in particular \mathbf{B}_+ is the closed unit half ball in \mathbb{R}^n and $\overline{\mathbf{C}}_+ = \mathbf{B}_+ \times \mathbb{R}^k$. Superscripts or subscripts ”–” have analogous meaning with $] -\infty, 0[$ replacing $]0, \infty[$. Balls in a Euclidean space \mathbb{R}^l of arbitrary dimension we denote $\mathbf{U}_r^l(a)$, $\mathbf{B}_r^l(y)$ or $\mathbf{U}^l(a, r)$, $\mathbf{B}^l(a, r)$, indicating the dimension. However, in the case $l = n + k$ we omit the dimension writing $\mathbf{U}(x, r)$, $\mathbf{B}(x, r)$; no confusion should arise from this, as it can be seen from the context that $x \in \mathbb{R}^{n+k}$ in this case.

For the interior ϵ -regularity theorem we consider the class of **admissible currents in the interior situation**

$$\mathcal{T}_i(n, k, m)$$

which consists of the n -dimensional currents T on \mathbb{R}^{n+k} with the following properties:

$$(1.18) \quad T \in \mathcal{R}_n(\mathbb{R}^{n+k}), \quad T = T \llcorner \mathbf{C},$$

$$(1.19) \quad \partial T \llcorner \mathbf{C} = 0, \quad 0 \in \text{spt } T,$$

$$(1.20) \quad \mathbf{p}_\# T = m(\mathbf{E}^n \llcorner \mathbf{U}).$$

The first condition means that T is rectifiable and vanishes outside \mathbf{C} , by the second condition T has zero boundary in \mathbf{C} and does not vanish on a neighborhood of the origin, the third condition tells us that the projection of T onto \mathbb{R}^n is represented by the unit disc with standard orientation and multiplicity m . Here m is a given positive integer, and \mathbf{E}^n denotes the n -current on \mathbb{R}^n defined by integration of n -forms (with respect to the standard orientation of \mathbb{R}^n).

For the boundary regularity theorem we consider an analogous class of currents T with $\partial T \llcorner \mathbf{C}$ prescribed (and nonzero). Thinking of a boundary condition $\partial T = B$ where B

is represented by an oriented submanifold \mathcal{B} of class $C^{1,1}$ in \mathbb{R}^{n+k} it is no restriction, by the invariance of the structure conditions for the integrand F and the (F, ω) minimizing property of T with respect to transformations by biLipschitz diffeomorphisms of class $C^{1,1}$, to assume that \mathcal{B} is an $(n-1)$ -dimensional affine subspace which we take as $\mathbb{R}^{n-1} \times \{0\}^{k+1} \subset \mathbb{R}^{n+k}$ with the boundary orientation induced from the half space $\mathbb{R}^{n-1} \times]0, \infty[$ (this is $(-1)^n$ times the standard orientation of \mathbb{R}^{n-1}). We therefore replace (1.19) by

$$(1.21) \quad \partial T \llcorner \mathbf{C} = B \quad \text{with} \quad B = (-1)^{n-1} (\mathbf{E}^{n-1} \times \delta_0^{k+1}) \llcorner \mathbf{C}$$

and (1.20) by

$$(1.22) \quad \mathbf{p}_\# T = m (\mathbf{E}^n \llcorner \mathbf{U}_+) + (m-1) (\mathbf{E}^n \llcorner \mathbf{U}_-),$$

that is $\mathbf{p}_\# T$ is represented by the two half disks \mathbf{U}_+ , \mathbf{U}_- with standard orientation and multiplicities m , $m-1$ respectively, and

$$(1.23) \quad \mathbf{p}_\# B = \partial (\mathbf{E}^n \llcorner \mathbf{U}_+) \llcorner \mathbf{U}$$

is the oriented boundary of the half disc \mathbf{U}_+ in \mathbf{U} . The class of **admissible currents in the boundary situation**

$$\mathcal{T}_b(n, k, m)$$

is then defined as the set of all currents T satisfying (1.18), (1.21) and (1.22). We use the notation

$$\mathcal{T}_*(n, k, m) = \mathcal{T}_i(n, k, m) \cup \mathcal{T}_b(n, k, m)$$

so that statements made about $T \in \mathcal{T}_*(n, k, m)$ are valid in the interior situation and in the boundary situation as well. We notice that $(\mu_{\varrho\#}^{-1} \tau_{a\#}^{-1} T) \llcorner \mathbf{C} \in \mathcal{T}_i(n, k, m)$ if $T \in \mathcal{T}_i(n, k, m)$, $a \in \mathbf{C} \cap \text{spt } T$, and $0 < \varrho \leq 1 - |\mathbf{p}a|$. The analogous assertion holds for \mathcal{T}_b instead of \mathcal{T}_i , provided $a \in \mathbf{C} \cap \mathbb{R}^{n-1} \times \{0\}^{k+1}$.

A crucial role in the regularity theory for rectifiable n -currents T on \mathbb{R}^{n+k} is played by the **cylindrical excess** which is defined, for $r > 0$, by

$$\mathbf{E}(T, r) = r^{-n} [\mathbf{M}(T \llcorner \mathbf{C}_r) - \mathbf{M}(\mathbf{p}_\#(T \llcorner \mathbf{C}_r))] \geq 0,$$

and, for $y \in \mathbb{R}^n$,

$$\mathbf{E}(T, y, r) = \mathbf{E}(\tau_{(y,0)\#}^{-1} T, r).$$

If $\mathbf{p}_\#(T \llcorner \mathbf{C}_r) = \mathbf{E}^n \llcorner \vartheta$ holds with a nonnegative multiplicity function ϑ , then one can write

$$(1.24) \quad \mathbf{E}(T, r) = r^{-n} \int_{\mathbf{C}_r} (1 - \vec{T} \cdot \vec{\mathbf{e}}) d\|T\| = \frac{1}{2} r^{-n} \int_{\mathbf{C}_r} |\vec{T} - \vec{\mathbf{e}}|^2 d\|T\|,$$

where $\vec{\mathbf{e}}$ is the orientation vector of $\mathbf{E}^n \times \delta_0^k$ (i.e. the exterior product $\mathbf{e}_1 \wedge \dots \wedge \mathbf{e}_n$ of the first n canonical basis vectors of \mathbb{R}^{n+k}). In particular, we may use (1.24) for $T \in \mathcal{T}_*(n, k, m)$, and we have

$$\mathbf{E}(T, r) = r^{-n} \mathbf{M}(T \llcorner \mathbf{C}_r) - m\alpha(n) \quad \text{if } T \in \mathcal{T}_i(n, k, m),$$

$$\mathbf{E}(T, r) = r^{-n} \mathbf{M}(T \llcorner \mathbf{C}_r) - \frac{1}{2}(2m-1) \boldsymbol{\alpha}(n) \quad \text{if } T \in \mathcal{T}_b(n, k, m),$$

where $\boldsymbol{\alpha}(n) = \mathcal{L}^n(\mathbf{U})$ is the volume of the unit ball in \mathbb{R}^n . We note the mass bound $\mathbf{M}(T \llcorner \mathbf{C}_r) \leq 1 + m \boldsymbol{\alpha}(n) r^n$ as a consequence, if $\mathbf{E}(T, r) \leq 1$. We also note the monotonicity property $r^n \mathbf{E}(T, r) \leq s^n \mathbf{E}(T, s)$ for $0 < r \leq s$. (The main part in the regularity theory is to prove that also the scaled quantity $\mathbf{E}(T, r)$ is decreasing to zero with $r \downarrow 0$.)

Finally, with C_1, C_2, \dots we denote various constants which are "large", i.e. $C_i \in [1, \infty[$, while δ, ϵ, \dots are small positive constants with values in $]0, 1[$. We usually write $C_i = \text{const}(n, k, \Lambda, \nu, \dots)$ or $\delta = \delta(n, k, \Lambda, \nu, \dots)$ when the constants appear for the first time, in order to indicate their dependence on parameters or moduli $n, k, \Lambda, \nu, \dots$

2 Lower mass estimate and height bound

In this section we establish a lower bound for the mass in balls and an upper bound for the height of (F, ω) minimizing currents $T \in \mathcal{T}_*(n, k, m)$. The arguments are well known in the interior case (cf. [Fe, 5.1.6], [Fe, 5.3.4], [Bo, Sec.IV], [SS, Appendix]) and easily adapted to the boundary situation, we give proofs here only for the sake of completeness. For the mass estimate (Lemma 2.1) one only needs the structure condition (1.1) on F , i.e. $|\xi| \leq F(\cdot, \xi) \leq \Lambda |\xi|$, and the boundedness of ω . The height estimate (Lemma 2.2) then follows assuming smallness of the excess of T and, in the case of multiplicity $m > 1$, additional conditions on the density of $\|T\|$ (Remark 2.3).

2.1 Lemma (lower mass estimate). *Suppose $T \in \mathcal{T}_*(n, k, m)$ is (F, ω) minimizing in \mathbf{C} . Then*

$$(2.1) \quad C_1 \mathbf{M}(T \llcorner \mathbf{U}(x, r)) \geq [\Lambda + \omega(r)]^{1-n} r^n$$

holds for all $x \in \text{spt } T$ and $0 < r < 1 - |\mathbf{p}x|$, where $C_1 = \text{const}(n)$.

Proof. We use the Euclidean distance function d_x to x to obtain, for almost every r as above, a rectifiable slice $\langle T, d_x, r \rangle = \partial(T \llcorner \mathbf{U}(x, r)) - (\partial T) \llcorner \mathbf{U}(x, r)$ with $T \llcorner \mathbf{U}(x, r) = T \llcorner \mathbf{B}(x, r)$ (see [Fe, Sec. 4.3]). We then consider a rectifiable current S of least mass in \mathbb{R}^{n+k} with boundary $\partial S = \partial(T \llcorner \mathbf{U}(x, r))$. From the isoperimetric inequality [Fe, 4.4.2], [Alm4] we have $\mathbf{M}(S) \leq \gamma_n \mathbf{M}(\partial S)^{n/(n-1)}$, and the convexity of $\mathbf{U}(x, r)$ implies $\text{spt } S \subset \mathbf{B}(x, r)$. We may therefore use the (F, ω) minimality of T and the structure conditions on F to infer

$$\begin{aligned} \mathbf{M}(T \llcorner \mathbf{U}(x, r)) &\leq \mathbf{F}(T \llcorner \mathbf{U}(x, r)) \leq \mathbf{F}(S) + \omega(r) \mathbf{M}(S) \\ &\leq [\Lambda + \omega(r)] \mathbf{M}(S) \leq [\Lambda + \omega(r)] \gamma_n \mathbf{M}(\partial S)^{n/(n-1)} \\ &\leq [\Lambda + \omega(r)] \gamma_n [\mathbf{M}(\langle T, d_x, r \rangle) + \mathbf{M}((\partial T) \llcorner \mathbf{U}(x, r))]^{n/(n-1)}. \end{aligned}$$

In the case $(\partial T) \llcorner \mathbf{U}(x, r) \neq 0$ we use additionally the projection from \mathbb{R}^{n+k} onto the subspace $\mathbb{R}^{n-1} \times \{0\}^{k+1}$ to see $\mathbf{M}((\partial T) \llcorner \mathbf{U}(x, r)) \leq \mathbf{M}(\langle T, d_x, r \rangle)$, because this projection maps $\langle T, d_x, r \rangle$ onto $-(\partial T) \llcorner \mathbf{U}(x, r)$. In any case

$$(2.2) \quad \mathbf{M}(T \llcorner \mathbf{U}(x, r)) \leq [\Lambda + \omega(r)] \gamma_n 2^{n/(n-1)} \mathbf{M}(\langle T, d_x, r \rangle)^{n/(n-1)}$$

is valid.

Introducing $f(r) = \mathbf{M}(T \llcorner \mathbf{U}(x, r))$ and recalling $\mathbf{M}(\langle T, d_x, r \rangle) \leq f'(r)$ for almost all r ([Fe, 4.2.1]) we deduce from (2.2)

$$f(r) \leq [\Lambda + \omega(r)] \gamma_n 2^{n/(n-1)} f'(r)^{n/(n-1)}.$$

Noting $f(r) > 0$ for $r > 0$, since $x \in \text{spt } T$, we may rewrite this differential inequality as

$$(2.3) \quad \frac{d}{dr} [f(r)^{1/n}] \geq [\Lambda + \omega(r)]^{(1-n)/n} \gamma_n^{(1-n)/n} 2n^{-1},$$

valid for almost all r . Finally, we integrate (2.3) using the monotonicity of f and $f(0) = 0$ to obtain (2.1). \blacksquare

2.2 Lemma (height bound). *Suppose $T \in \mathcal{T}_*(n, k, 1)$ is (F, ω) minimizing in \mathbf{C} with a smallness condition for the excess*

$$(2.4) \quad C_2 \left[\Lambda + \omega\left(\frac{1}{2}\right) \right]^{n-1} \mathbf{E}(T, 1)^{1/2} \leq 1$$

where $C_2 = \text{const}(n)$. Then the estimate

$$(2.5) \quad \sup \left\{ |\mathbf{q}x|^n : x \in \mathbf{C}_{1/2} \cap \text{spt } T \right\} \leq C_3 \left[\Lambda + \omega\left(\frac{1}{2}\right) \right]^{n-1} \mathbf{E}(T, 1)^{1/2}$$

is valid with $C_3 = \text{const}(n)$.

Proof. We first treat the boundary situation $T \in \mathcal{T}_b(n, k, 1)$. Here we have $\mathbf{M}(T) = \mathbf{E}(T, 1) + \frac{1}{2}\alpha(n)$ and, consequently, $\mathbf{M}(T) \leq \frac{3}{4}\alpha(n)$ if C_2 in (2.4) is chosen to imply $\mathbf{E}(T, 1) \leq \frac{1}{4}\alpha(n)$. Fix a unit vector $u \in \{0\}^n \times \mathbb{R}^k \subset \mathbb{R}^{n+k}$ and define $H(r) = \{x \in \mathbf{C} : u \cdot x > r\}$, $T_r = T \llcorner H(r)$, and $H(r, s) = \{x \in \mathbf{C} : r < u \cdot x < s\}$ for $r < s$. Since $\mathbf{M}(\mathbf{p}_\# T_r) \leq \mathbf{M}(T)$ and thus

$$(2.6) \quad \mathbf{M}(\mathbf{p}_\# T_r) \leq \frac{3}{4}\alpha(n),$$

we may apply the relative isoperimetric inequality [Fe, 5.3.2] to obtain

$$(2.7) \quad \mathbf{M}(\mathbf{p}_\# T_r)^{1-1/n} \leq \text{const}(n) \mathbf{M}(\partial(\mathbf{p}_\# T_r) \llcorner \mathbf{U}) \leq \text{const}(n) \mathbf{M}((\partial T_r) \llcorner \mathbf{C}).$$

Suppose now $\mathbf{M}(T_s) \geq \sqrt{E}$ for some $s > 0$, where $E = \mathbf{E}(T, 1) \leq \frac{1}{4}$ and we may assume $E > 0$. Then, for $0 < r < s$ we get from (2.7)

$$(2.8) \quad \begin{aligned} \left(\sqrt{E} - E \right)^{1-1/n} &\leq [\mathbf{M}(T_r) - \mathbf{E}(T, 1)]^{1-1/n} \leq \mathbf{M}(\mathbf{p}_\# T_r)^{1-1/n} \\ &\leq \text{const}(n) \mathbf{M}((\partial T_r) \llcorner \mathbf{C}). \end{aligned}$$

Integrating this inequality for $r \in]0, s[$ and using the identity

$$\langle T, \ell_u, r \rangle \llcorner \mathbf{C} = \partial T \llcorner H(r) - (\partial T_r) \llcorner \mathbf{C} = -(\partial T_r) \llcorner \mathbf{C}$$

for the slices of T with respect to the linear function $\ell_u(x) = u \cdot x$ (note that $H(r) \cap \text{spt } \partial T = \emptyset$ for $r > 0$) we obtain from the standard slicing estimate [Fe, 4.3.2(2)] and $\mathbf{M}(T) \leq \text{const}(n)$

$$\begin{aligned}
\left(\frac{1}{2}\sqrt{E}\right)^{1-1/n} s &\leq (\sqrt{E} - E)^{1-1/n} s \leq \text{const}(n) \int_0^s \mathbf{M}(\langle T, \ell_{ur} \rangle) dr \\
&\leq \text{const}(n) \int_{H(0,s)} |\vec{T} \lrcorner d\ell_u| d\|T\| \\
&\leq \text{const}(n) \left[\int_{H(0,s)} |\vec{T} \lrcorner d\ell_u|^2 d\|T\| \right]^{1/2} \\
&\leq \text{const}(n) \left[\int_{H(0,s)} (1 - (\vec{T} \cdot \vec{e})^2) d\|T\| \right]^{1/2} \\
&\leq \text{const}(n) \left[2 \int_{H(0,s)} (1 - \vec{T} \cdot \vec{e}) d\|T\| \right]^{1/2} \\
&\leq \text{const}(n) \mathbf{E}(T \lrcorner H(0, s), 1)^{1/2} \leq \text{const}(n) \sqrt{E}.
\end{aligned}$$

This implies

$$(2.9) \quad s \leq \text{const}(n) E^{1/2n} \quad \text{when } s > 0 \text{ and } \mathbf{M}(T_s) \geq \sqrt{E}.$$

On the other hand, if $s > 0$ with $\mathbf{M}(T_s) \leq \sqrt{E}$ and $x \in \overline{\mathbf{C}}_{1/2} \cap \text{spt } T$ with $u \cdot x \geq s + \delta$ for some $0 < \delta < \frac{1}{2}$, then $\mathbf{U}(x, \delta) \subset H(s)$, and we see from Lemma 2.1

$$[\Lambda + \omega(\delta)]^{1-n} \delta^n \leq C_1 \mathbf{M}(T \lrcorner \mathbf{U}(x, \delta)) \leq C_1 \mathbf{M}(T_s) \leq C_1 \sqrt{E},$$

an impossibility if we choose $\delta^n = 2 \left[\Lambda + \omega(\frac{1}{2}) \right]^{n-1} C_1 \sqrt{E}$ and we require (2.4) with C_2 large enough to secure $\delta < \frac{1}{2}$. We deduce

$$(2.10) \quad \sup \left\{ u \cdot x : x \in \overline{\mathbf{C}}_{1/2} \cap \text{spt } T \right\} \leq s + \text{const}(n) \left[\Lambda + \omega(\frac{1}{2}) \right]^{1-1/n} E^{1/2n}$$

when $s > 0$ and $\mathbf{M}(T_s) \leq \sqrt{E}$.

Clearly, with (2.9) and (2.10) we have proved

$$(2.11) \quad \sup \left\{ u \cdot x : x \in \overline{\mathbf{C}}_{1/2} \cap \text{spt } T \right\} \leq \text{const}(n) \left[\Lambda + \omega(\frac{1}{2}) \right]^{1-1/n} E^{1/2n},$$

and (2.5) follows since $u \in \{0\}^n \times \mathbb{R}^k$ is arbitrary with $|u| = 1$.

In the interior situation $T \in \mathcal{T}_i(n, k, 1)$ one first chooses a median value r_0 such that both $\mathbf{M}(T \lrcorner H(r_0, \infty))$ and $\mathbf{M}(T \lrcorner H(-\infty, r_0))$ do not exceed $\frac{1}{2} \mathbf{M}(T)$. Then $\mathbf{M}(\mathbf{p}_{\#} T_{r_0}) \leq \frac{1}{2} \boldsymbol{\alpha}(n) + \frac{1}{2} E$ holds, and (2.6) is valid for $r > r_0$ assuming again $E \leq \frac{1}{4} \boldsymbol{\alpha}(n)$. Reasoning as in (2.9), (2.10) one obtains an estimate analogous to (2.11) for the oscillation of ℓ_u on $\overline{\mathbf{C}}_{1/2} \cap \text{spt } T$. The inequality (2.5) then follows from $0 \in \text{spt } T$. (Cf. the proof of Lemma 2 in [SS].) ■

2.3 Remark. In the case of multiplicity $m > 1$ one can prove a height estimate similar to (2.5), provided the additional assumption

$$(2.12) \quad \Theta^n(\|T\|, x) \geq m \quad \text{for } \|T\| \text{ almost all } x \in \mathbf{C}$$

is satisfied when $T \in \mathcal{T}_i(n, k, m)$, and

$$(2.13) \quad \begin{aligned} \Theta^n(\|T\|, x) &\geq m && \text{for } \|T\| \text{ almost all } x \in \mathbf{C}_+, \\ \Theta^n(\|T\|, x) &\geq m - 1 && \text{for } \|T\| \text{ almost all } x \in \mathbf{C}_- \end{aligned}$$

when $T \in \mathcal{T}_b(n, k, m)$.

For $T \in \mathcal{T}_i(n, k, m)$ and the median value r_0 from the proof above we deduce from (2.12) and $E \leq \frac{m}{4}\alpha(n)$ that $\mathcal{L}^n(P(r)) \leq \frac{3}{4}\alpha(n)$ for $r \geq r_0$ and the set $P(r) \subset \mathbf{U}$ of points y with $\Theta^n(\|\mathbf{p}_\#(T \llcorner H(r, \infty))\|, y) \geq 1$. Therefore, the relative isoperimetric inequality (2.7) is again valid, and we may argue as before to obtain (2.5) with $C_3 = m^{n/2}\text{const}(n)$.

For $T \in \mathcal{T}_b(n, k, m)$ we first choose median values r_0^+ , r_0^- for the half cylinders \mathbf{C}_+ , \mathbf{C}_- analogous to r_0 for \mathbf{C} . We then argue as in the proof of (2.9) to obtain, with $c = \text{const}(n)$ and $H_\pm(\dots) = \mathbf{C}_\pm \cap H(\dots)$,

$$(2.14) \quad \begin{aligned} \mathbf{M}(T \llcorner H_+(r, \infty)) &\leq \frac{1}{2}\sqrt{E} && \text{for } r \geq r_0^+ + cE^{1/2n}, \\ \mathbf{M}(T \llcorner H_+(-\infty, r)) &\leq \frac{1}{2}\sqrt{E} && \text{for } r \leq r_0^+ - cE^{1/2n}, \end{aligned}$$

and analogous statements for $T \llcorner H_-(\dots)$ and r_0^- .

We claim that $|r_0^+ - r_0^-| \leq 2cE^{1/2n} + \delta$ where $\delta = (2C_1)^{1/n} [\Lambda + \omega(\frac{1}{2})]^{1-1/n} E^{1/2n} < \frac{1}{2}$ is as in the proof of Lemma 2.2 and $E = \mathbf{E}(T, 1)$ is sufficiently small (and positive). With this estimate for $|r_0^+ - r_0^-|$ we can then derive the height estimate (2.5) with $C_3 = m^{n/2}\text{const}(n)$ exactly as in the proof of (2.10).

For $x \in \mathbf{C}_{1-\delta} \cap \text{spt } T$ we first note $\mathbf{M}(T \llcorner \mathbf{U}(x, \delta)) \geq 2\sqrt{E}$, by Lemma 2.1 and the definition of δ , and we infer

$$(2.15) \quad \mathbf{M}(T \llcorner (\mathbf{C}_+ \cap \mathbf{U}(x, \delta))) > \frac{1}{2}\sqrt{E} \quad \text{or} \quad \mathbf{M}(T \llcorner (\mathbf{C}_- \cap \mathbf{U}(x, \delta))) > \frac{1}{2}\sqrt{E},$$

since $\mathbf{M}(T \llcorner (\mathbf{C} \setminus (\mathbf{C}_+ \cup \mathbf{C}_-))) \leq E < \sqrt{E}$. Assuming $|r_0^+ - r_0^-| > 2cE^{1/2n} + 2\delta$ we deduce $u \cdot x \neq s$ for $s = \frac{1}{2}(r_0^+ + r_0^-)$, because otherwise $\mathbf{U}(x, \delta)$ would be disjoint from $H_+(r_0^+ - cE^{1/2n}, r_0^+ + cE^{1/2n})$ and from $H_-(r_0^- - cE^{1/2n}, r_0^- + cE^{1/2n})$ and then (2.15) would contradict (2.14). Since $0 \in \text{spt } \partial T$ and $u \cdot x = 0$ for $x \in \mathbf{C} \cap \text{spt } \partial T$ we conclude $s > 0$ and $\partial(T \llcorner H(s, \infty)) \llcorner \mathbf{C}_{1-\delta} = 0$, or $s < 0$ and $\partial(T \llcorner H(-\infty, s)) \llcorner \mathbf{C}_{1-\delta} = 0$. The second case being similar, we proceed with the first case where $[\partial(\mathbf{p}_\# T_s)] \llcorner \mathbf{U}_{1-\delta} = 0$ and $\mathbf{p}_\# T_s$ is constant on $\mathbf{U}_{1-\delta}$. But it follows from (2.14), $|r_0^+ - r_0^-| > 2cE^{1/2n}$, and the choice of s that

$$\mathbf{M}((\mathbf{p}_\# T_s) \llcorner \mathbf{U}_+) \leq \frac{1}{2}\sqrt{E} \quad \text{and} \quad \mathbf{M}(T \llcorner H_+(-\infty, s)) \leq \frac{1}{2}\sqrt{E},$$

or

$$\mathbf{M}((\mathbf{p}_\# T_s) \llcorner \mathbf{U}_-) \leq \frac{1}{2}\sqrt{E} \quad \text{and} \quad \mathbf{M}(T \llcorner H_+(-\infty, s)) \leq \frac{1}{2}\sqrt{E}.$$

For $\sqrt{E} < 2^{-n-1}\alpha(n)$ this contradicts the constancy of $\mathbf{p}_\#T_s$ on $\mathbf{U}_{1/2}$, because $\mathbf{p}_\#T$ has multiplicity $\geq m-1 > 0$ everywhere on \mathbf{U} and $\{x \in \mathbf{C}_{1/2} \cap \text{spt } T : u \cdot x = s\}$ is empty. The claim is proved.

We note that the *bad* hypotheses (2.12), (2.13) are needed only for the relative isoperimetric inequality (2.7) and somewhat weaker assumptions would suffice here. For example, it would be enough to know the inequalities (2.12), (2.13) on subcylinders of \mathbf{C} , \mathbf{C}_+ , \mathbf{C}_- with cross sections of positive \mathcal{L}^n measure (independent of T). However, we shall need scaled versions of the height estimate which we can prove, in the case $m \geq 2$, only with extra assumptions similar to (2.12), (2.13) (cf. [Fe, 5.3.17] and the discussion following Lemma 2 in [SS]).

3 Harmonic approximation

As a *general hypothesis* we assume in this section that $T \in \mathcal{T}_*(n, k, m)$ is (F, ω) minimizing in \mathbf{C} , where the integrand F satisfies $|\xi| \leq F(\cdot, \xi) \leq \Lambda|\xi|$ and $\omega:]0, 1[\rightarrow [0, \infty[$ is nondecreasing, and that the additional density conditions (2.12), (2.13) hold in the case of multiplicity $m \geq 2$. Using the height estimate Lemma 2.2 we first show that T can be approximated, with error estimates in terms of the excess $\mathbf{E}(T, 1)$, by the current T_g associated with the graph of a Lipschitz function $g: \mathbb{R}^n \rightarrow \mathbb{R}^k$ equipped with appropriately chosen multiplicities and orientation. In the interior case this was done in [SS, Lemma 3] modifying [All1, 8.12] (cf. [Si1, §20]), and it is not difficult to adapt the arguments to the boundary situation treated here. Since the estimates we need are somewhat different from those in [SS] we give complete proofs. We next show that the approximating function g is, again with an error dominated by the excess, an approximate solution to the homogeneous constant coefficient elliptic system determined by the nonparametric elliptic integrand associated with F frozen at the origin (i.e. with $\xi \mapsto F(0, \xi)$). This is similar to Lemma 12 and Lemma 13 in [Bo], in the boundary situation $T \in \mathcal{T}_b(n, k, m)$, however, we can prove such a statement only on the half ball \mathbf{U}_+ (and also on \mathbf{U}_- if $m \geq 2$). Finally, we state a simple Lemma following [Si1, §21], [Si2, §1.6] which asserts that the approximate solution g is indeed close, in the weak topology of the Sobolev space $W^{1,2}$, to an exact solution.

3.1 Lemma (Lipschitz approximation). *Given $0 < \varrho \leq \frac{1}{17}$ and $0 < \beta \leq 1$ there exists a Lipschitz map $g: \mathbf{U}_\varrho \rightarrow \mathbb{R}^k$ with $\text{Lip } g \leq \beta$ such that the following assertions are true:*

(i) *The n -current T_g represented by graph g , with orientation and multiplicities induced from $(\mathbf{p}_\#T) \llcorner \mathbf{U}_\varrho$ by the projection \mathbf{p} , satisfies*

$$(3.1) \quad \|T - T_g\|(\mathbf{C}_\varrho) \leq C_4 \beta^{-2n} \mathbf{E}(T, 1)$$

with a constant $C_4 = \text{const}(n, m) \left[\Lambda + \omega\left(\frac{1}{2}\right) \right]^{2n-2}$.

(ii) *Let A_g be the set of $y \in \mathbf{U}_\varrho$ satisfying $\{y\} \times \mathbb{R}^k \cap \text{spt } T \setminus \text{graph } g \neq \emptyset$ and in the case $T \in \mathcal{T}_b(n, k, m)$ additionally $y \in \mathbf{U}_+$; then*

$$(3.2) \quad \mathcal{L}^n(A_g) \leq C_5 \beta^{-2n} \mathbf{E}(T, 1)$$

holds with a constant $C_5 = \text{const}(n, m) \left[\Lambda + \omega\left(\frac{1}{2}\right) \right]^{2n-2}$.

(iii) In the case $T \notin \mathcal{T}_i(n, k, m)$

$$(3.3) \quad \int_{\mathbf{U}_\varrho \setminus A_g} |Dg|^2 d\mathcal{L}^n \leq 3\mathbf{E}(T, 1)$$

holds; in the case $T \in \mathcal{T}_b(n, k, m)$ this is valid with $\mathbf{U}_+ \cap \mathbf{U}_\varrho \setminus A_g$ as domain of integration.

(iv) For $\gamma > 0$ let $A_{g,\gamma} = A_g \cup \{y \in \mathbf{U}_\varrho : |Dg(y)| \geq \gamma\}$ in the case $T \in \mathcal{T}_i(n, k, m)$ and $A_{g,\gamma} = A_g \cup \{y \in \mathbf{U}_\varrho \cap \mathbf{U}_+ : |Dg(y)| \geq \gamma\}$ in the case $T \in \mathcal{T}_b(n, k, m)$; then

$$(3.4) \quad \mathcal{L}^n(A_{g,\gamma}) \leq (1 + C_5\beta^{-2n}) 3\gamma^{-2}\mathbf{E}(T, 1),$$

$$(3.5) \quad \|T\|(A_{g,\gamma} \times \mathbb{R}^k) \leq [1 + m(1 + C_5\beta^{-2n}) 3\gamma^{-2}] \mathbf{E}(T, 1).$$

(v) $g(0) = 0$ and in the boundary case $T \in \mathcal{T}_b(n, k, m)$ additionally $g(y) = 0$ for all $y \in \mathbf{U}_\varrho^{n-1} \times \{0\}$.

(vi) If $C_2 \left[\Lambda + \omega(\frac{1}{2})\right]^{n-1} \mathbf{E}(T, 1)^{1/2} \leq 1$ holds, then

$$(3.6) \quad \sup_{\mathbf{U}_\varrho} |g|^n \leq C_3 \left[\Lambda + \omega(\frac{1}{2})\right]^{n-1} \mathbf{E}(T, 1)^{1/2}$$

where C_2, C_3 are the constants from Lemma 2.2.

Proof. We only treat the boundary case $T \in \mathcal{T}_b(n, k, m)$ as the proof in the interior situation is similar, but simpler.

For $0 < \varrho \leq \frac{1}{4}$ and η to be fixed later define the *good set*

$$(3.7) \quad G = \{y \in \mathbf{U}_\varrho : y_n \neq 0 \text{ and } \mathbf{E}(T, y, \sigma) \leq \eta \text{ for } 0 < \sigma < 1 - \varrho\}.$$

Then for each $y \in \mathbf{U}_\varrho \setminus G$ there exists $\sigma(y) \in]0, 1 - \varrho[$ with

$$\frac{1}{2} \int_{\mathbf{C}_{\sigma(y)}(y)} |\vec{T} - \vec{e}|^2 d\|T\| = \sigma(y)^n \mathbf{E}(T, y, \sigma(y)) > \sigma(y)^n \eta.$$

By the Besicovitch covering theorem we can find $N = \text{const}(n)$ disjoint systems $\mathcal{B}_1, \dots, \mathcal{B}_N$ of balls $\mathbf{B}(y, \sigma(y))$, $y \in \mathbf{U}_\varrho \setminus G$, such that $\mathcal{B}_1 \cup \dots \cup \mathcal{B}_N$ covers $\mathbf{U}_\varrho \setminus G$. For $i \in \{1, \dots, N\}$ chosen suitably we infer the following estimate for the *bad set*:

$$(3.8) \quad \begin{aligned} \mathcal{L}^n(\mathbf{U}_\varrho \setminus G) &\leq N\mathcal{L}^n\left(\bigcup_{B \in \mathcal{B}_i} B\right) \leq N\alpha(n) \sum_{\mathbf{B}(y, \sigma(y)) \in \mathcal{B}_i} \sigma(y)^n \\ &\leq N\alpha(n) \frac{1}{2\eta} \int_{\mathbf{C}} |\vec{T} - \vec{e}|^2 d\|T\| = N\alpha(n) \frac{1}{\eta} \mathbf{E}(T, 1). \end{aligned}$$

We want to show that $\text{spt } T$ is a graph over G . For this we fix $y, \tilde{y} \in G$ and $x, \tilde{x} \in \text{spt } T$ with $\mathbf{p}x = y$, $\mathbf{p}\tilde{x} = \tilde{y}$, and we distinguish

Case 1: $\mathbf{B}(y, 2|y - \tilde{y}|) \subset \mathbf{U}_+$ (or $\subset \mathbf{U}_-$);

Case 2: $\mathbf{B}(y, 2|y - \tilde{y}|) \cap \mathbf{U}^{n-1} \times \{0\} \neq \emptyset$.

In the first case we consider the scaled current $\tilde{T} = \left((\boldsymbol{\mu}_{2|y-\tilde{y}|}^{-1} \circ \boldsymbol{\tau}_x^{-1}) \# T \right) \llcorner \mathbf{C} \in \mathcal{T}_i(n, k, m)$ which is $(\tilde{F}, \tilde{\omega})$ minimizing in \mathbf{C} , where \tilde{F} has the same structure constant Λ as F , and $\tilde{\omega}(r) = \omega(2|y-\tilde{y}|r) \leq \omega(r)$ (see Section 1; in the case $y = \tilde{y}$ we consider $\left((\boldsymbol{\mu}_t^{-1} \circ \boldsymbol{\tau}_x^{-1}) \# T \right) \llcorner \mathbf{C}$ with $t > 0$ arbitrary small.) We can therefore apply Lemma 2.2 (and Remark 2.3 in the case $m \geq 2$, as (2.13) for T implies (2.12) for \tilde{T}) to obtain

$$|\mathbf{q}x - \mathbf{q}\tilde{x}| \leq \tilde{C}_3 \mathbf{E}(\tilde{T}, 1)^{1/2n} 2|y - \tilde{y}|,$$

provided $\tilde{C}_2 \mathbf{E}(\tilde{T}, 1)^{1/2} \leq 1$. Here we have abbreviated $\tilde{C}_2 = C_2 \left[\Lambda + \omega(\frac{1}{2}) \right]^{n-1}$ and $\tilde{C}_3 = C_3^{1/n} \left[\Lambda + \omega(\frac{1}{2}) \right]^{1-1/n}$. From $\varrho \leq \frac{1}{5}$ we have $2|y - \tilde{y}| < 1 - \varrho$ and hence $\mathbf{E}(\tilde{T}, 1) = \mathbf{E}(T, y, 2|y - \tilde{y}|) \leq \eta$, by (3.7). Choosing η such that $\tilde{C}_2 \eta^{1/2} \leq 1$ the conclusion is therefore

$$(3.9) \quad |\mathbf{q}x - \mathbf{q}\tilde{x}| \leq \tilde{C}_3 \eta^{1/2n} 2|y - \tilde{y}| \quad \text{in case 1.}$$

If $\mathbf{B}(y, 2|y - \tilde{y}|) \subset \mathbf{U}_-$ and $m \geq 2$, then $\tilde{T} \in \mathcal{T}_i(n, k, m - 1)$, and the conclusion is the same.

In the second case we introduce $y' = (y_1, \dots, y_{n-1}, 0)$ in \mathbb{R}^n and observe $|y' - y| \leq 2|y - \tilde{y}|$, hence

$$\mathbf{B}(y, |y - \tilde{y}|) \subset \mathbf{B}(y', 3|y - \tilde{y}|) \subset \mathbf{B}(y', 6|y - \tilde{y}|) \subset \mathbf{B}(y, 8|y - \tilde{y}|) \subset \mathbf{U},$$

since $\varrho \leq \frac{1}{16}$. We now apply the height estimate from Section 2 to the scaled current $\tilde{T} = \left((\boldsymbol{\mu}_{6|y-\tilde{y}|}^{-1} \circ \boldsymbol{\tau}_{(y',0)}^{-1}) \# T \right) \llcorner \mathbf{C} \in \mathcal{T}_b(n, k, m)$ and obtain

$$\max \{ |\mathbf{q}x|, |\mathbf{q}\tilde{x}| \} \leq \tilde{C}_3 \mathbf{E}(T, y', 6|y - \tilde{y}|)^{1/2n} 6|y - \tilde{y}|,$$

provided $\tilde{C}_2 \mathbf{E}(T, y', 6|y - \tilde{y}|)^{1/2} \leq 1$. On account of $\varrho \leq \frac{1}{17}$ we have $8|y - \tilde{y}| < 1 - \varrho$ and hence, by (3.7), $\mathbf{E}(T, y', 6|y - \tilde{y}|) \leq \left(\frac{4}{3} \right)^n \eta$. Choosing $\eta > 0$ such that $\tilde{C}_2 \left(\frac{4}{3} \right)^{n/2} \eta^{1/2} \leq 1$ we therefore obtain

$$(3.10) \quad |\mathbf{q}x - \mathbf{q}\tilde{x}| \leq 2\tilde{C}_3 \sqrt{\frac{4}{3}} \eta^{1/2n} 6|y - \tilde{y}| \quad \text{in case 2.}$$

We also note

$$(3.11) \quad |\mathbf{q}x| \leq \tilde{C}_3 \sqrt{\frac{3}{2}} \eta^{1/2n} 2|y - y'|,$$

which follows from the same reasoning if we apply it with y' instead of \tilde{y} and use $\mathbf{B}(y, |y - y'|) \subset \mathbf{B}(y', 2|y - y'|) \subset \mathbf{B}(y, 3|y - y'|) \subset \mathbf{U}$.

Considering $y = \tilde{y}$ in (3.9), (3.10) we see that a function g is well defined on G in the case $m \geq 2$ and on $G \cap \mathbf{U}_+$ in the case $m = 1$ by letting $g(y) = \mathbf{q}x$ for $x \in \text{spt } T$ with $\mathbf{p}x = y$. From (3.9), (3.10) we also see $\text{Lip } g \leq \beta$ if we choose $\eta = C_5^{-1} \beta^{2n}$ with $C_5 = \text{const}(n, m) \left[\Lambda + \omega(\frac{1}{2}) \right]^{2n-2}$. From (3.11) we further recognize that this remains true if we extend g to $\mathbf{U}_\varrho^{n-1} \times \{0\}$ by zero. Finally, we extend g to all of \mathbf{U}_ϱ preserving the Lipschitz constant, then $\text{Lip } g \leq \beta \leq 1$ holds on \mathbf{U}_ϱ and (v) is valid.

To prove (i) we note that T_g is the n -current associated with graph g and with multiplicities m on $\mathbf{C}_+ \cap \text{graph } g$, $m-1$ on $\mathbf{C}_- \cap \text{graph } g$, oriented such that $(\partial T_g) \llcorner \mathbf{C}_\varrho = (\partial T) \llcorner \mathbf{C}_\varrho$. Writing $G_\pm = \mathbf{U}_\pm \cap G$ and $A = \mathbf{U}_\varrho \setminus (G_+ \cup G_-)$ we estimate, using also (2.13) if $m > 1$,

$$\begin{aligned}
\|T - T_g\|(G_+ \times \mathbb{R}^k) &= \int_{\text{graph } g|_{G_+}} [\Theta^n(\|T\|, \cdot) - m] d\mathcal{H}^n \\
(3.12) \qquad \qquad \qquad &\leq \|T\|(G_+ \times \mathbb{R}^k) - m\mathcal{L}^n(G_+) \\
&= \mathbf{E}(T \llcorner (G_+ \times \mathbb{R}^k), 1) ,
\end{aligned}$$

and similarly

$$(3.13) \quad \|T - T_g\|(G_- \times \mathbb{R}^k) \leq \mathbf{E}(T \llcorner (G_- \times \mathbb{R}^k), 1) \quad \text{in the case } m \geq 2,$$

while

$$(3.14) \quad \|T\|(\mathbf{C} \setminus \mathbf{C}_+) \leq \mathbf{E}(T, 1) \quad \text{in the case } m = 1,$$

because then $(\mathbf{p}_\# T) \llcorner (\mathbf{U} \setminus \mathbf{U}_+) = 0$. Furthermore,

$$\begin{aligned}
\|T - T_g\|(A \times \mathbb{R}^k) &\leq (\|T\| + \|T_g\|)(A \times \mathbb{R}^k) \\
(3.15) \qquad \qquad \qquad &\leq \mathbf{E}(T \llcorner (A \times \mathbb{R}^k), 1) + m \left[1 + (1 + (\text{Lip } g)^2)^{n/2} \right] \mathcal{L}^n(A) .
\end{aligned}$$

Adding (3.13), (3.14), (3.15), taking account of $\text{Lip } g \leq \beta \leq 1$, and using also (3.8) which, by the choice of η above, may be rewritten

$$(3.16) \quad \mathcal{L}^n(A) \leq C_5 N \alpha(n) \beta^{-2n} \mathbf{E}(T, 1) ,$$

we deduce (3.1) in the case $m \geq 2$ writing C_5 for $C_5 N \alpha(n)$. In the case $m = 1$ we obtain the same conclusion from (3.12), (3.14), (3.15), and (3.16), noting that $T_g \llcorner (\mathbf{C} \setminus \mathbf{C}_+) = 0$.

Assertion (ii) follows from (3.16) and (3.14), since $A_g \subset A \cup \mathbf{p}[\text{spt}(T \llcorner (\mathbf{C} \setminus \mathbf{C}_+))]$ if $m = 1$.

For (iii) we use $|Dg| \leq \text{Lip } g \leq 1$, the definition of A_g and (2.13) (if $m > 1$) to infer

$$\begin{aligned}
\int_{\mathbf{U}_+ \cap \mathbf{U}_\varrho \setminus A_g} |Dg|^2 d\mathcal{L}^n &\leq 3 \int_{\mathbf{U}_+ \cap \mathbf{U}_\varrho \setminus A_g} \left[\sqrt{1 + |Dg|^2} - 1 \right] d\mathcal{L}^n \\
&\leq 3 \int_{\mathbf{U}_+ \cap \mathbf{U}_\varrho \setminus A_g} [J_n(\text{id} \times g) - 1] d\mathcal{L}^n \\
&\leq \frac{3}{m} \|T\|(\mathbf{p}^{-1}(\mathbf{U}_+ \cap \mathbf{U}_\varrho \setminus A_g)) - 3\mathcal{L}^n(\mathbf{U}_+ \cap \mathbf{U}_\varrho \setminus A_g) \\
&\leq \frac{3}{m} \mathbf{E}(T \llcorner \mathbf{C}_+, 1) ,
\end{aligned}$$

and a similar inequality, with m replaced by $m-1$, holds for the integral on $\mathbf{U}_- \cap \mathbf{U}_\varrho \setminus A_g$ if $m \geq 2$.

With regard to (iv) we note that (iii) implies

$$\mathcal{L}^n(A_{g,\gamma}) \leq \left(1 + \frac{1}{\gamma^2} \right) \mathcal{L}^n(A_g) + \frac{3}{\gamma^2} \mathbf{E}(T, 1) ,$$

hence (3.4) follows from (3.2), and (3.5) from

$$\|T\|(\mathbf{p}^{-1}A_{g,\gamma}) \leq \mathbf{E}(T, 1) + m\mathcal{L}^n(A_{g,\gamma}).$$

Finally, (vi) follows from Lemma 2.2, because g can be chosen to satisfy (3.6) if we secure $\tilde{C}_2\mathbf{E}(T, 1)^{1/2} \leq 1$. \blacksquare

For the next lemma we introduce

$$A = (a_{ij}^{\alpha\beta}) = \left(\frac{\partial^2}{\partial p_\alpha^i \partial p_\beta^j} F_0^\S(0) \right)$$

for the Hessian matrix of the nonparametric integrand $F_0^\S(p)$ associated with the frozen integrand $F_0 = F(0, \cdot)$. We also write

$$A(\xi, \eta) = \sum_{\alpha, \beta=1}^n \sum_{i, j=1}^k a_{ij}^{\alpha\beta} \xi_\alpha^i \eta_\beta^j$$

for the quadratic form on $\text{Hom}(\mathbb{R}^n, \mathbb{R}^k)$ associated with A , i.e. $A = D^2F_0^\S(0)$. The constant Λ and the nondecreasing functions κ , μ , $\hat{\mu}$, and ν are from the structure conditions (1.1)-(1.9) on F .

3.2 Lemma (approximate A-harmonicity). *Suppose $\omega(\frac{1}{2}) + \mu(\frac{1}{2}) \leq 1$, $0 < \varrho \leq \frac{1}{17}$, $\mathbf{C}_\varrho \cap \text{spt } T \subset \mathbf{U}_\varrho^n \times \mathbf{B}_{1/3}^k$, and let g be the Lipschitz approximation associated with T , ϱ , and $\beta = 1$ by Lemma 3.1. Then, with $C_{10} = \text{const}(n, m, \Lambda)$, $E = \mathbf{E}(T, 1)$, and $\chi = \kappa(\frac{1}{2})^2 + \hat{\mu}(\frac{1}{2}) + \omega(\frac{1}{2})$, the inequality*

$$(3.17) \quad \left| \int_{\mathbf{U}_\varrho} A(Dg, D\zeta) d\mathcal{L}^n \right| \leq C_{10} \left[\gamma^{-2}E + \sqrt{\chi}(1 + E) + (\sqrt{E} + \sqrt{\chi}) \nu(\gamma + \sqrt{\chi}) \right] \sup |D\zeta|$$

holds for all parameters $0 < \gamma \leq 1$ and all functions $\zeta \in C^1(\mathbf{U}, \mathbb{R}^k)$ with compact support in \mathbf{U}_ϱ and satisfying additionally $\text{spt } \zeta \subset \mathbf{U}_+$ in the case $T \in \mathcal{T}_b(n, k, 1)$, $\text{spt } \zeta \subset \mathbf{U} \setminus \mathbf{U}^{n-1} \times \{0\}$ in the case $T \in \mathcal{T}_b(n, k, m)$, $m \geq 2$.

The essential point here is that χ/\sqrt{E} is small in the later application of this lemma so that, with the appropriate choice of γ , the constant on the right hand side of (3.17) is a small quantity times \sqrt{E} . Since $\hat{\mu}(\frac{1}{2}) \leq \mu(\frac{1}{2})$ we could replace $\hat{\mu}(\frac{1}{2})$ by $\mu(\frac{1}{2})$ in the hypothesis. However, replacing $\hat{\mu}(\frac{1}{2})$ by $\mu(\frac{1}{2})$ in the conclusion, i.e. in the definition of χ , would yield a weaker conclusion, in general. The hypothesis that T has height $\leq \frac{1}{3}$ in \mathbf{C}_ϱ is made only to secure that the (F, ω) minimizing property of T is applied in the proof with a modification of T inside a ball of radius $\frac{1}{2}$. An appropriate version of the lemma is true without this hypothesis.

Proof. We follow [Bo, Section IV] closely. We may assume $\sup |D\zeta| \leq 1$, $\sup |\zeta| \leq \varrho$, and we define the *vertical deformation*

$$h_t(x) = x + t(0, \zeta(\mathbf{p}x)) \quad \text{for } |t| \leq 1.$$

We then decompose

$$\mathbf{F}((h_t \# T) \llcorner \mathbf{C}_\varrho) = K(t) + L(t) + M(t)$$

where

$$\begin{aligned} K(t) &= \int_{A_\gamma \times \mathbb{R}^k} F\left(h_t(x), \bigwedge_n Dh_t(x) \vec{T}(x)\right) d\|T\|_x, \\ L(t) &= \int_{\mathbf{C}_\varrho \setminus A_\gamma \times \mathbb{R}^k} F\left(h_t(x), \bigwedge_n Dh_t(x) \vec{S}_g(x)\right) d\|S_g\|_x, \\ M(t) &= \int_{\mathbf{C}_\varrho \setminus A_\gamma \times \mathbb{R}^k} F\left(h_t(x), \bigwedge_n Dh_t(x) \vec{T}_g(x)\right) d\|T_g\|_x \end{aligned}$$

and we have written A_γ for the set $A_{g,\gamma} \subset \mathbf{U}_\varrho$ defined in Lemma 3.1, (iv) and S_g for $T - T_g$. (Note that $T - T_g = T \llcorner \vartheta$ in $\mathbf{C}_\varrho \setminus A_\gamma \times \mathbb{R}^k$ with a nonnegative function ϑ , cf. (3.13).)

We estimate $K(s) - K(0)$ using the continuity modulus $\mu(\cdot)$ for $x \mapsto F(x, \xi)$ from (1.7) and the bound Λ for the derivative $D_{(2)}F$ from (1.2). We have, for $|s| \leq 1$, and $\|T\|$ almost all $x \in \mathbf{C}_\varrho$,

$$\begin{aligned} & \left| F\left(h_s(x), \bigwedge_n Dh_s(x) \vec{T}(x)\right) - F(x, \vec{T}(x)) \right| \\ & \leq \left| F\left(h_s(x), \bigwedge_n Dh_s(x) \vec{T}(x)\right) - F\left(x, \bigwedge_n Dh_s(x) \vec{T}(x)\right) \right| \\ & \quad + \left| \int_0^s D_{(2)}F\left(x, \bigwedge_n Dh_t(x) \vec{T}(x)\right) \frac{d}{dt} \bigwedge_n Dh_t(x) \vec{T}(x) dt \right| \\ & \leq \mu(|s\zeta(\mathbf{p}x)|) \left| \bigwedge_n Dh_s(x) \vec{T}(x) \right| + \Lambda \left| \int_0^s \frac{d}{dt} \bigwedge_n Dh_t(x) \vec{T}(x) dt \right| \\ & \leq \mu(|s|\varrho)2^n + |s|\Lambda n2^{n-1}. \end{aligned}$$

Integrating with respect to $\|T\| \llcorner A_\gamma \times \mathbb{R}^k$ we obtain

$$(3.18) \quad |K(s) - K(0)| \leq \left(\mu(|s|\varrho)2^n + |s|\Lambda n2^{n-1} \right) \|T\|(A_\gamma \times \mathbb{R}^k) \leq C_6(\mu(|s|\varrho) + |s|)\gamma^{-2}E,$$

where $C_6 = \text{const}(n, m, \Lambda)$ and we have used (3.5).

In the same way, using (3.1) instead of (3.5), we derive

$$(3.19) \quad |L(s) - L(0)| \leq C_7(\mu(|s|\varrho) + |s|)E$$

with $C_7 = \text{const}(n, m, \Lambda)$.

Similarly as above we see, using also the continuity modulus $\kappa(\cdot)$ for $x \mapsto D_{(2)}F(x, \xi)$ from (1.6) and again assuming $|s| \leq 1$,

$$\begin{aligned} & \left| F\left(h_s(x), \bigwedge_n Dh_s(x) \vec{T}_g(x)\right) - F\left(x, \vec{T}_g(x)\right) - F\left(0, \bigwedge_n Dh_s(x) \vec{T}_g(x)\right) + F\left(0, \vec{T}_g(x)\right) \right| \\ & \leq \left| F\left(h_s(x), \bigwedge_n Dh_s(x) \vec{T}_g(x)\right) - F\left(x, \bigwedge_n Dh_s(x) \vec{T}_g(x)\right) \right| \end{aligned}$$

$$\begin{aligned}
& + \left| \int_0^s \left[D_{(2)}F \left(x, \bigwedge_n Dh_t(x) \vec{T}_g(x) \right) - D_{(2)}F \left(0, \bigwedge_n Dh_t(x) \vec{T}_g(x) \right) \right] \frac{d}{dt} \bigwedge_n Dh_t(x) \vec{T}_g(x) dt \right| \\
& \leq \mu(|s|\varrho)2^n + |s|\kappa(|x|)n2^{n-1}.
\end{aligned}$$

Integrating this with respect to $\|T_g\|_{\mathbf{L}(\mathbf{C}_\varrho \setminus A_\gamma \times \mathbb{R}^k)}$ we get

$$\begin{aligned}
(3.20) \quad & \left| \int_{\mathbf{C}_\varrho \setminus A_\gamma \times \mathbb{R}^k} \left[F \left(h_s(x), \bigwedge_n Dh_s(x) \vec{T}_g(x) \right) - F \left(x, \vec{T}_g(x) \right) \right] d\|T_g\|_x \right. \\
& \left. - \int_{\mathbf{C}_\varrho \setminus A_\gamma \times \mathbb{R}^k} \left[F \left(0, \bigwedge_n Dh_s(x) \vec{T}_g(x) \right) - F \left(0, \vec{T}_g(x) \right) \right] d\|T_g\|_x \right| \\
& \leq \text{const}(n, m) \left[\mu(|s|\varrho) + |s|\kappa(\sqrt{2}\varrho) \right],
\end{aligned}$$

since $\text{Lip } g \leq 1$ implies $\|T_g\|(\mathbf{C}_\varrho) \leq m \text{const}(n)$ and $|x| \leq \sqrt{2}\varrho$ for $x \in \mathbf{C}_\varrho \cap \text{spt } T_g$ (recall $g(0) = 0$).

Defining the multiplicity function $\boldsymbol{\nu}_m$ by $\boldsymbol{\nu}_m \equiv m$ on \mathbf{U} in the case $T \in \mathcal{T}_i(n, k, m)$ and $\boldsymbol{\nu}_m \equiv m$ on \mathbf{U}_+ , $\boldsymbol{\nu}_m \equiv m - 1$ on \mathbf{U}_- in the case $T \in \mathcal{T}_b(n, k, m)$ we can write

$$(3.21) \quad \int_{\mathbf{C}_\varrho \setminus A_\gamma \times \mathbb{R}^k} F \left(0, \bigwedge_n Dh_s(x) \vec{T}_g(x) \right) d\|T_g\|_x = \int_{\mathbf{U}_\varrho \setminus A_\gamma} F_0^\S(D(g + s\zeta)) \boldsymbol{\nu}_m d\mathcal{L}^n$$

and

$$(3.22) \quad \int_{\mathbf{C}_\varrho \setminus A_\gamma \times \mathbb{R}^k} F \left(0, \vec{T}_g(x) \right) d\|T_g\|_x = \int_{\mathbf{U}_\varrho \setminus A_\gamma} F_0^\S(Dg) \boldsymbol{\nu}_m d\mathcal{L}^n.$$

Using (1.11) in the Taylor expansion

$$\left\| DF_0^\S(p) - DF_0^\S(0) - DDF_0^\S(0)p \right\| \leq \nu(|p|)|p|$$

with $p = Dg(y) + tD\zeta(y)$, multiplying with $D\zeta(y)$, and integrating with respect to (t, y) we further obtain

$$\begin{aligned}
(3.23) \quad & \left| \int_{\mathbf{U}_\varrho \setminus A_\gamma} \left[F_0^\S(D(g + s\zeta)) - F_0^\S(Dg) \right] \boldsymbol{\nu}_m d\mathcal{L}^n - s \int_{\mathbf{U}_\varrho \setminus A_\gamma} DF_0^\S(0) D\zeta \boldsymbol{\nu}_m d\mathcal{L}^n \right. \\
& \left. - s \int_{\mathbf{U}_\varrho \setminus A_\gamma} D^2 F_0^\S(0)(Dg, D\zeta) \boldsymbol{\nu}_m d\mathcal{L}^n - \frac{1}{2}s^2 \int_{\mathbf{U}_\varrho \setminus A_\gamma} D^2 F_0^\S(0)(D\zeta, D\zeta) \boldsymbol{\nu}_m d\mathcal{L}^n \right| \\
& \leq \left| \int_0^s \int_{\mathbf{U}_\varrho \setminus A_\gamma} (\nu \circ |Dg + tD\zeta|) |Dg + tD\zeta| |D\zeta| \boldsymbol{\nu}_m d\mathcal{L}^n dt \right| \\
& \leq m \text{const}(n) |s| \nu(\gamma + |s|) \left(\sqrt{E} + |s| \right),
\end{aligned}$$

where we have used (3.3), $|D\zeta| \leq 1$ and $|Dg| \leq \gamma$ on $\mathbf{U}_\varrho \setminus A_\gamma$ (on $\mathbf{U}_+ \cap \mathbf{U}_\varrho \setminus A_\gamma$ in the case $T \in \mathcal{T}_b(n, k, 1)$) in the last step. The second integral appearing in (3.23) can be estimated

with (3.4), because the integral of $DF_0^\S(0)D\zeta\iota_m$ on \mathbf{U}_ϱ vanishes on account of the hypothesis on $\text{spt } \zeta$:

$$(3.24) \quad \left| \int_{\mathbf{U}_\varrho \setminus A_\gamma} DF_0^\S(0)D\zeta\iota_m d\mathcal{L}^n \right| = \left| \int_{A_\gamma} DF_0^\S(0)D\zeta\iota_m d\mathcal{L}^n \right| \leq \text{const}(n, m, \Lambda)\gamma^{-2}E.$$

Combining (3.20)–(3.24) and applying (3.4) once more we arrive finally at the following inequality, valid for $|s| \leq 1$:

$$(3.25) \quad \left| M(s) - M(0) - s \int_{\mathbf{U}_\varrho} D^2F_0^\S(0)(Dg, D\zeta)\iota_m d\mathcal{L}^n \right| \leq C_8\mu(|s|\varrho) + C_8|s| \left[\kappa(\sqrt{2}\varrho) + \gamma^{-2}E + |s| + (\sqrt{E} + |s|) \nu(\gamma + |s|) \right],$$

where $C_8 = \text{const}(n, m, \Lambda)$. With (3.18), (3.19), (3.25), and the inequality $\widehat{\mu}(\frac{1}{2}) \geq \mu(|s|\varrho) - 4s^2\varrho^2$ which follows from (1.9), we have proved

$$(3.26) \quad \left| \mathbf{F}((h_{s\#}T) \llcorner \mathbf{C}_\varrho) - \mathbf{F}(T \llcorner \mathbf{C}_\varrho) - s \int_{\mathbf{U}_\varrho} D^2F_0^\S(0)(Dg, D\zeta)\iota_M d\mathcal{L}^n \right| \leq C_9\mu(|s|\varrho) + C_9|s| \left[\kappa(\sqrt{2}\varrho) + \gamma^{-2}E + |s| + (\sqrt{E} + |s|) \nu(\gamma + |s|) \right] \leq C_9\widehat{\mu}(\frac{1}{2}) + C_9|s| \left[\kappa(\frac{1}{2}) + \gamma^{-2}E + 2|s| + (\sqrt{E} + |s|) \nu(\gamma + |s|) \right],$$

provided $|s| \leq 1$ and $\text{spt } \zeta$ is compact in \mathbf{U}_ϱ not meeting $\mathbf{U}^{n-1} \times \{0\}$ unless ι_m is constant.

Since, with this condition on $\text{spt } \zeta$, $h_{s\#}T$ is an admissible comparison current, we can employ the (F, ω) minimality of T :

$$(3.27) \quad \begin{aligned} \mathbf{F}(T \llcorner \mathbf{C}_\varrho) &\leq \mathbf{F}((h_{s\#}T) \llcorner \mathbf{C}_\varrho) + \omega(\frac{1}{2})\mathbf{M}(h_{s\#}T \llcorner \mathbf{C}_\varrho) \\ &\leq \mathbf{F}((h_{s\#}T) \llcorner \mathbf{C}_\varrho) + \omega(\frac{1}{2})\text{const}(n)(m + E), \end{aligned}$$

because $h_{s\#}T - T$ has support in $\mathbf{B}^{n+k}(0, \frac{1}{2})$, by our assumptions on $\mathbf{C}_\varrho \cap \text{spt } T$, ζ and ϱ . Choosing $|s| = \sqrt{\widehat{\mu}(\frac{1}{2}) + \omega(\frac{1}{2})}$ (we may assume that this is positive) and the sign of s carefully (as in [Bo], proof of Lemma 13) we deduce (3.17) from (3.26) and (3.27). The condition $|s| \leq 1$ is satisfied for this choice in view of the hypothesis $\mu(\frac{1}{2}) + \omega(\frac{1}{2}) \leq 1$ (and $\widehat{\mu}(\frac{1}{2}) \leq \mu(\frac{1}{2})$). \blacksquare

For $T \in \mathcal{T}_b(n, k, m)$ with multiplicity $m \geq 2$ it is reasonable to expect that the condition $\mathbf{U}^{n-1} \times \{0\} \cap \text{spt } \zeta = \emptyset$ in (3.17) is not necessary, i.e. g is approximately A -harmonic on \mathbf{U}_ϱ and not just on $\mathbf{U}_+ \cap \mathbf{U}_\varrho$ and $\mathbf{U}_- \cap \mathbf{U}_\varrho$ separately. However, we could not find a proof based on vertical deformations of the type used above, and this is one of the reasons why we can obtain a boundary regularity result only in the case $m = 1$. It seems that a different type of variations of T is needed to make progress here, variations which can separate “leaves” of T with zero boundary in \mathbf{C} from the “leaf” with boundary ∂T and multiplicity 1, speaking intuitively.

So far we have not used ellipticity of the integrand F . However, if $A = D^2F_0^\S(0)$ is elliptic in the sense of the Legendre-Hadamard condition (1.13), then we can find an

A-harmonic function, i.e. an exact solution to the homogeneous constant coefficient elliptic system associated with A , which is close in the weak topology of the Sobolev space $W^{1,2}$, to the approximatively A -harmonic function g exhibited in Lemma 3.2. This can be proved similarly as in [Si1, §21], [Si2, §1]. For the convenience of the reader we formulate in the next lemma the exact statement which will be used later, and we include the simple proof.

3.3 Lemma (A-harmonic approximation). *There exists a positive function $\delta(n, k, \lambda, \Lambda, \epsilon) \leq 1$ with the following property: Whenever $A \in \odot^2 \text{Hom}(\mathbb{R}^n, \mathbb{R}^k)$ is (rank one) elliptic with ellipticity constant $\lambda > 0$ and upper bound Λ , ϵ and ϱ are given positive numbers, and $f \in W^{1,2}(\mathbf{U}_\varrho, \mathbb{R}^k)$ with $\varrho^{2-n} \int_{\mathbf{U}_\varrho} |Df|^2 d\mathcal{L}^n \leq 1$ is approximatively A -harmonic in the sense that*

$$\left| \varrho^{2-n} \int_{\mathbf{U}_\varrho} A(Df, D\zeta) d\mathcal{L}^n \right| \leq \delta(n, k, \lambda, \Lambda, \epsilon) \varrho \sup |D\zeta|$$

holds for all $\zeta \in C^1(\mathbf{U}_\varrho, \mathbb{R}^k)$ with compact support in \mathbf{U}_ϱ , then there exists an A -harmonic function $u \in W^{1,2}(\mathbf{U}_\varrho, \mathbb{R}^k)$ such that

$$\varrho^{-n} \int_{\mathbf{U}_\varrho} |u - f|^2 d\mathcal{L}^n \leq \epsilon \quad \text{and} \quad \varrho^{2-n} \int_{\mathbf{U}_\varrho} |Du|^2 d\mathcal{L}^n \leq 1.$$

The same assertion is true if we replace, everywhere in the statement, the n -ball \mathbf{U}_ϱ by the half ball $\mathbf{U}_{\varrho,+} = \mathbf{U}_\varrho \cap \mathbb{R}^{n-1} \times]0, \infty[$ and the Sobolev space $W^{1,2}(\mathbf{U}_\varrho, \mathbb{R}^k)$ by $W_{(0)}^{1,2}(\mathbf{U}_{\varrho,+}, \mathbb{R}^k)$, the space of $W^{1,2}$ functions on $\mathbf{U}_{\varrho,+}$ with vanishing boundary trace on $\mathbf{U}_\varrho^{n-1} \times \{0\}$.

Proof. We may assume $\varrho = 1$, the general case follows by scaling $y \mapsto \varrho^{-1}f(\varrho y)$. Supposing the Lemma to be false we have existence of a sequence of quadratic forms $A_i \in \odot^2 \text{Hom}(\mathbb{R}^n, \mathbb{R}^k)$ with fixed ellipticity constant $\lambda > 0$ and bound $\Lambda < \infty$, and a sequence of $f_i \in W^{1,2}(\mathbf{U}, \mathbb{R}^k)$ with $\|Df_i\|_{L^2} \leq 1$ and

$$(3.28) \quad \left| \int_{\mathbf{U}} A_i(Df_i, D\zeta) d\mathcal{L}^n \right| \leq \frac{1}{i} \sup |D\zeta|$$

for all $\zeta \in C^1(\mathbf{U}, \mathbb{R}^k)$ with compact support in \mathbf{U} , such that for some $\epsilon > 0$ the inequality

$$(3.29) \quad \int_{\mathbf{U}} |f_i - u_i|^2 d\mathcal{L}^n > \epsilon$$

is valid for each i and each A_i -harmonic function $u_i \in W^{1,2}(\mathbf{U}, \mathbb{R}^k)$ with $\|Du_i\|_{L^2} \leq 1$.

Clearly, we may assume $\int_{\mathbf{U}} f_i d\mathcal{L}^n = 0$ and apply Poincaré's inequality and Rellich's theorem to obtain, after having passed to a subsequence, $f_i \rightarrow f$ strongly in $L^2(\mathbf{U}, \mathbb{R}^k)$ and weakly in $W^{1,2}(\mathbf{U}, \mathbb{R}^k)$ and $A_i \rightarrow A$ in $\odot^2 \text{Hom}(\mathbb{R}^n, \mathbb{R}^k)$. Then $\|Df\|_{L^2} \leq 1$, and from

$$\begin{aligned} \int_{\mathbf{U}} A(Df, D\zeta) d\mathcal{L}^n &= \int_{\mathbf{U}} A(Df - Df_i, D\zeta) d\mathcal{L}^n \\ &\quad + \int_{\mathbf{U}} (A - A_i)(Df_i, D\zeta) d\mathcal{L}^n + \int_{\mathbf{U}} A_i(Df_i, D\zeta) d\mathcal{L}^n \end{aligned}$$

and (3.28) we infer that f is A -harmonic.

Now, the ellipticity of A_i implies (by Fourier transform and Plancherel's theorem, i.e. Gårding's inequality in the case of constant coefficients) the coercivity condition

$$(3.30) \quad \int_{\mathbf{U}} A_i(D\eta, D\eta) d\mathcal{L}^n \geq \lambda \int_{\mathbf{U}} |D\eta|^2 d\mathcal{L}^n \quad \text{for } \eta \in W_0^{1,2}(\mathbf{U}, \mathbb{R}^k),$$

hence we can choose u_i as the solution to the Dirichlet problem corresponding to A_i and the boundary values of f , i.e.

$$(3.31) \quad u_i - f \in W_0^{1,2}(\mathbf{U}, \mathbb{R}^k), \quad \int_{\mathbf{U}} A_i(Du_i, D\zeta) d\mathcal{L}^n = 0 \quad \text{for } \zeta \in W_0^{1,2}(\mathbf{U}, \mathbb{R}^k).$$

Then

$$\begin{aligned} \lambda \int_{\mathbf{U}} |Du_i - Df|^2 d\mathcal{L}^n &\leq \int_{\mathbf{U}} A_i(Du_i - Df, Du_i - Df) d\mathcal{L}^n \quad \text{by (3.30),} \\ &= - \int_{\mathbf{U}} A_i(Df, Du_i - Df) d\mathcal{L}^n \quad \text{by (3.31),} \\ &= \int_{\mathbf{U}} (A - A_i)(Df, Du_i - Df) d\mathcal{L}^n \quad \text{by } A\text{-harmonicity of } f, \\ &\leq |A_i - A| \left[\int_{\mathbf{U}} |Du_i - Df|^2 d\mathcal{L}^n \right]^{1/2} \quad \text{on account of } \|Df\|_{L^2} \leq 1, \end{aligned}$$

and in view of $A_i \rightarrow A$ we have convergence $u_i \rightarrow f$ in $W^{1,2}$ norm. But this implies $\|u_i - f_i\|_{L^2} \leq \|u_i - f\|_{L^2} + \|f - f_i\|_{L^2} \rightarrow 0$ as $i \rightarrow \infty$ and the same assertion is true for $\tilde{u}_i = \min(1, \|Du_i\|_{L^2}^{-1})u_i$ contradicting (3.29).

The proof for $W_{(0)}^{1,2}(\mathbf{U}_+, \mathbb{R}^k)$ is the same. We need only to observe that Poincaré's inequality is valid in $W_{(0)}^{1,2}(\mathbf{U}_+, \mathbb{R}^k)$, Rellich's compactness theorem holds for the embedding $W_{(0)}^{1,2}(\mathbf{U}_+, \mathbb{R}^k) \rightarrow L^2(\mathbf{U}_+, \mathbb{R}^k)$, and $W_{(0)}^{1,2}(\mathbf{U}_+, \mathbb{R}^k)$ is a closed subspace of $W^{1,2}(\mathbf{U}_+, \mathbb{R}^k)$. ■

Although the proof of the previous lemma is by contradiction, this does not mean that our regularity proof is indirect. Our point of view is simply that the lemma guarantees the existence of a positive function $\delta(n, k, \lambda, \Lambda, \epsilon)$ which is, admittedly not in an explicit form, determined by a property of constant coefficient elliptic systems, and we will later use this function in the regularity proof.

4 A reverse Poincaré inequality

For solutions u to an elliptic equation by a reverse Poincaré inequality (also called Caccioppoli's inequality) one means an estimate of Dirichlet's integral of u (on a ball) in terms of the integral of $|u|^2$ (on a larger ball). The analogue for (F, ω) minimizing currents $T \in \mathcal{T}_*(n, k, m)$ would be an estimate of the excess $\mathbf{E}(T, \frac{1}{5})$ of T in $\mathbf{C}_{1/5}$, say, by the integral $\int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|$ of the squared height function. Up to some small extra term we prove here such an inequality. In Lemma 4.1 we first establish a *weak* version which has an additional term $\vartheta \mathbf{E}(T, 1)$ on the right-hand side, with an arbitrarily small parameter $\vartheta > 0$, however. The proof is a refinement of [Bo, Sec. 5] which requires some new estimates; this is crucial for the optimality of our regularity results. We then apply a covering argument which we have found in [Si2] to deduce the *strong* form of the reverse Poincaré inequality in Lemma 4.2. In the following we use the ellipticity constant $\lambda > 0$ of the integrand F from (1.12), the bound Λ from the structure conditions (1.1)–(1.3) for F , and the moduli $\kappa, \mu, \hat{\mu}$ for F from (1.5)–(1.9).

4.1 Lemma (weak reverse Poincaré inequality). *Suppose $T \in \mathcal{T}_i(n, k, m)$ with $m \geq 0$ or $T \in \mathcal{T}_b(n, k, m)$ with $m \geq 1$ is (F, ω) minimizing in \mathbf{C} and $\mathbf{C}_{7/8} \cap \text{spt } T$ is contained in the ball $\mathbf{B}^{n+k}(0, R)$. Then, with $\chi(R) = \kappa(R)^2 + \hat{\mu}(R) + \omega(R)$ the following inequality holds for all parameters $\vartheta > 0$,*

$$(4.1) \quad \mathbf{E}\left(T, \frac{1}{2}\right) \leq \vartheta \mathbf{E}(T, 1) + \frac{1}{\vartheta} C_{11} \left[\chi(R) \mathbf{M}(T) + (1 + \chi(R)) \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|x \right],$$

where $C_{11} = \text{const}(n, \lambda, \Lambda)$.

Here $\mathcal{T}_i(n, k, 0)$ is defined exactly as $\mathcal{T}_i(n, k, m)$ for $m \geq 1$ in Section 1, without the condition $0 \in \text{spt } T$, however. It will be clear from the proof that this latter condition is not used in the case $T \in \mathcal{T}_i(n, k, m)$ with $m \geq 1$.

Proof. For $0 < r \leq 1$ we write $T_r = T \llcorner \mathbf{C}_r$ and $\langle T, r \rangle = \partial T_r - (\partial T) \llcorner \mathbf{C}_r$ for the cylindrical slices of T . Recalling the multiplicity function $\iota_m \equiv m$ on \mathbf{U} in the case $T \in \mathcal{T}_i(n, k, m)$ and $\iota_m \equiv m$ on \mathbf{U}_+ , $\iota_m \equiv m - 1$ on \mathbf{U}_- in the case $T \in \mathcal{T}_b(n, k, m)$ we further denote by P_r the image of $(\mathbf{p}_\# T) \llcorner \mathbf{U}_r = (\mathbf{E}^n \llcorner \mathbf{U}_r) \llcorner \iota_m$ under the inclusion $\mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^k$. Using, for $0 < \vartheta \leq \frac{5}{8}$ and $r > \frac{1}{5}$, the homotopy

$$h^\vartheta(t, x) = x - t(\vartheta \mathbf{p}x, \mathbf{q}x) = (1 - t)x + t(1 - \vartheta)(\mathbf{p}x, 0)$$

we define, similarly as in [Bo, Section V],

$$\Sigma_{r, \vartheta} = P_{(1-\vartheta)r} - h^\vartheta_\#(\llbracket 0, 1 \rrbracket \times \langle T, r \rangle)$$

and verify

$$(4.2) \quad \begin{aligned} \Sigma_{r, \vartheta} &= \Sigma_{r, \vartheta} \llcorner \mathbf{C}_r, \quad \Sigma_{r, \vartheta} \llcorner \mathbf{C}_{1/2} = P_{1/2}, \\ \mathbf{p}_\# \Sigma_{r, \vartheta} &= \mathbf{p}_\# T_r = (\mathbf{E}^n \llcorner \mathbf{U}_r) \llcorner \iota_m, \quad \partial \Sigma_{r, \vartheta} = \partial T_r, \end{aligned}$$

so that $\Sigma_{r, \vartheta}$ may be used as a comparison current for T_r if $\langle T, r \rangle$ is rectifiable.

For such r we estimate the excess of $\Sigma_{r, \vartheta}$ with the standard estimate [Fe, 4.1.9] noting $\text{Lip } h^\vartheta(t, \cdot) \leq 1 - t\vartheta$:

$$(4.3) \quad \begin{aligned} \mathbf{E}(\Sigma_{r, \vartheta}, 1) &= \mathbf{M}(P_{(1-\vartheta)r}) + \mathbf{M}(h^\vartheta_\#(\llbracket 0, 1 \rrbracket \times \langle T, r \rangle)) - \mathbf{M}(P_r) \\ &\leq [(1 - \vartheta)^n - 1] \mathbf{M}(P_r) + \int_0^1 \int_{\partial \mathbf{C}_r} |(\vartheta \mathbf{p}x, \mathbf{q}x)| (1 - t\vartheta)^{n-1} d\|\langle T, r \rangle\|x dt \\ &\leq [1 - (1 - \vartheta)^n] \left[\frac{1}{n\vartheta} \int_{\partial \mathbf{C}_r} \left(\vartheta r + \frac{1}{2\vartheta r} |\mathbf{q}x|^2 \right) d\|\langle T, r \rangle\|x - \mathbf{M}(P_r) \right]. \end{aligned}$$

On account of (see [Fe, 4.3.2(2)])

$$\int_{5/8}^{7/8} \int_{\partial \mathbf{C}_r} |\mathbf{q}x|^2 d\|\langle T, r \rangle\|x dr \leq \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|x$$

and

$$\int_{5/8}^{7/8} \mathbf{M}(\langle T, r \rangle) dr \leq \mathbf{M}(T \llcorner (\mathbf{C}_{7/8} \setminus \mathbf{C}_{5/8})),$$

hence also (observe $\mathbf{M}(P_r) = r^n \mathbf{M}(P_1)$)

$$\begin{aligned} & \int_{5/8}^{7/8} \left[\mathbf{M}(\langle T, r \rangle) - \frac{n}{r} \mathbf{M}(P_r) \right] dr \\ & \leq \mathbf{M}(T \llcorner (\mathbf{C}_{7/8} \setminus \mathbf{C}_{5/8})) - \mathbf{M}(P_{7/8}) + \mathbf{M}(P_{5/8}) \leq \mathbf{E}(T, 1) , \end{aligned}$$

we may select $r \in]\frac{5}{8}, \frac{7}{8}[$ such that

$$\int_{\partial \mathbf{C}_r} |\mathbf{q}x|^2 d\|\langle T, r \rangle\|_x \leq 12 \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|_x ,$$

$$\mathbf{M}(\langle T, r \rangle) \leq 12 \mathbf{M}(T) ,$$

$$\mathbf{M}(\langle T, r \rangle) \leq \frac{n}{r} \mathbf{M}(P_r) + 12 \mathbf{E}(T, 1) ,$$

and $\langle T, r \rangle$ is rectifiable (cf. [Fe, 4.3.6]). With this choice of r , (4.3) and $[1 - (1 - \vartheta)^n] \leq n\vartheta$ give

$$(4.4) \quad \mathbf{E}(\Sigma_{r,\vartheta}, 1) \leq 12\vartheta r \mathbf{E}(T, 1) + \frac{6}{\vartheta r} \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|_x .$$

Similarly, denoting by \tilde{P}_ϑ the image of $(\mathbf{p}_\# T) \llcorner (\mathbf{U} \setminus \mathbf{U}_\vartheta) = (\mathbf{E}^n \llcorner (\mathbf{U} \setminus \mathbf{U}_\vartheta)) \llcorner \boldsymbol{\iota}_m$ under the inclusion $\mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^k$ and using the homotopy

$$\tilde{h}^\vartheta(t, x) = x + t(\vartheta \mathbf{p}x, -\mathbf{q}x) = (1-t)x + t(1+\vartheta)(\mathbf{p}x, 0)$$

we define, similarly as in [Bo, Sec. 5],

$$\tilde{\Sigma}_{r,\vartheta} = \tilde{P}_{(1+\vartheta)r} + \tilde{h}_\#^\vartheta([0, 1] \times \langle T, r \rangle) .$$

For r as above and $0 < \vartheta \leq \frac{1}{7}$, we then have

$$(4.5) \quad \begin{aligned} \tilde{\Sigma}_{r,\vartheta} &= \tilde{\Sigma}_{r,\vartheta} \llcorner (\mathbf{C} \setminus \mathbf{C}_r) , \quad \tilde{\Sigma}_{r,\vartheta} \llcorner (\mathbf{C} \setminus \mathbf{C}_{(1+\vartheta)r}) = P_1 \llcorner (\mathbf{C} \setminus \mathbf{C}_{(1+\vartheta)r}) , \\ \mathbf{p}_\# (T_r + \tilde{\Sigma}_{r,\vartheta}) &= \mathbf{p}_\# P_1 = (\mathbf{E}^n \llcorner \mathbf{U}) \llcorner \boldsymbol{\iota}_m , \quad \partial (T_r + \tilde{\Sigma}_{r,\vartheta}) = \partial P_1 , \end{aligned}$$

and obtain the excess estimate (noting $\text{Lip } \tilde{h}^\vartheta(t, \cdot) \leq 1 + t\vartheta$ and $[(1+\vartheta)^n - 1] \leq (1+\vartheta)^{n-1} n\vartheta$ this time)

$$(4.6) \quad \mathbf{E}(\tilde{\Sigma}_{r,\vartheta}, 1) \leq (1+\vartheta)^{n-1} \left[12\vartheta r \mathbf{E}(T, 1) + \frac{6}{\vartheta r} \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|_x \right] .$$

On account of (4.2) and the hypothesis $\text{spt } T_{7/8} \subset \mathbf{B}^{n+k}(0, R)$, which by the construction of $\Sigma_{r,\vartheta}$ implies $\text{spt}(\Sigma_{r,\vartheta} - T_r) \subset \mathbf{B}^{n+k}(0, R)$, we can apply the (F, ω) minimality of T to get

$$(4.7) \quad \mathbf{F}(T_r) \leq \mathbf{F}(\Sigma_{r,\vartheta}) + \omega(R) \mathbf{M}(\Sigma_{r,\vartheta}) .$$

On the other hand, in view of (4.5) we can make use of the ellipticity of $F_0(x, \xi) = F(0, \xi)$ (we also write $F_0(\xi)$ for $F(0, \xi)$ in the sequel) to obtain

$$\begin{aligned} \lambda 2^{-n} \mathbf{E} \left(T, \frac{1}{2} \right) &= \lambda \left[\mathbf{M} \left(T_{1/2} \right) - \mathbf{M} \left(P_{1/2} \right) \right] \\ &\leq \lambda \left[\mathbf{M} \left(T_r + \tilde{\Sigma}_{r, \vartheta} \right) - \mathbf{M} \left(P_1 \right) \right] \leq \mathbf{F}_0 \left(T_r + \tilde{\Sigma}_{r, \vartheta} \right) - \mathbf{F}_0 \left(P_1 \right) \\ &= \mathbf{F} \left(T_r + \tilde{\Sigma}_{r, \vartheta} \llcorner \mathbf{C}_{(1+\vartheta)r} \right) + (\mathbf{F}_0 - \mathbf{F}) \left(T_r + \tilde{\Sigma}_{r, \vartheta} \llcorner \mathbf{C}_{(1+\vartheta)r} \right) - \mathbf{F}_0 \left(P_{(1+\vartheta)r} \right). \end{aligned}$$

Introducing $S_{r, \vartheta} = \Sigma_{r, \vartheta} + \tilde{\Sigma}_{r, \vartheta} \llcorner \mathbf{C}_{(1+\vartheta)r}$ and recalling (4.2), (4.5) we deduce from (4.7) that the last expression does not exceed

$$\begin{aligned} &\mathbf{F} \left(S_{r, \vartheta} \right) + \omega(R) \mathbf{M} \left(\Sigma_{r, \vartheta} \right) + (\mathbf{F}_0 - \mathbf{F}) \left(T_r + \tilde{\Sigma}_{r, \vartheta} \llcorner \mathbf{C}_{(1+\vartheta)r} \right) - \mathbf{F}_0 \left(P_{(1+\vartheta)r} \right) \\ &= \mathbf{F}_0 \left(S_{r, \vartheta} \right) - \mathbf{F}_0 \left(P_{(1+\vartheta)r} \right) + (\mathbf{F}_0 - \mathbf{F}) \left(T_r \right) - (\mathbf{F}_0 - \mathbf{F}) \left(\Sigma_{r, \vartheta} \right) + \omega(R) \mathbf{M} \left(\Sigma_{r, \vartheta} \right), \end{aligned}$$

hence we arrive at

$$(4.8) \quad \begin{aligned} \lambda 2^{-n} \mathbf{E} \left(T, \frac{1}{2} \right) &\leq \mathbf{F}_0 \left(S_{r, \vartheta} \right) - \mathbf{F}_0 \left(P_{(1+\vartheta)r} \right) \\ &\quad + \omega(R) \mathbf{M} \left(\Sigma_{r, \vartheta} \right) + (\mathbf{F}_0 - \mathbf{F}) \left(T_r \right) - (\mathbf{F}_0 - \mathbf{F}) \left(\Sigma_{r, \vartheta} \right). \end{aligned}$$

Now using Taylor expansion of $F_0(\xi)$ at $\xi = \vec{\mathbf{e}}$ for $|\xi - \vec{\mathbf{e}}| < \sqrt{2}$ and recalling (1.1)–(1.3) we find

$$\begin{aligned} &\mathbf{F}_0 \left(S_{r, \vartheta} \right) - \mathbf{F}_0 \left(P_{(1+\vartheta)r} \right) \\ &= \int_{\mathbf{C}_{(1+\vartheta)r}} F_0 \left(\vec{S}_{r, \vartheta}(x) \right) d\|S_{r, \vartheta}\|x - \int_{\mathbf{C}_{(1+\vartheta)r}} F_0 \left(\vec{\mathbf{e}} \right) d\|P_{(1+\vartheta)r}\| \\ &\leq \int_{\mathbf{C}_{(1+\vartheta)r}} \left[F_0 \left(\vec{\mathbf{e}} \right) + DF_0 \left(\vec{\mathbf{e}} \right) \left(\vec{S}_{r, \vartheta} - \vec{\mathbf{e}} \right) \right] d\|S_{r, \vartheta}\|x \\ &\quad + \Lambda \int_{\mathbf{C}_{(1+\vartheta)r}} |\vec{S}_{r, \vartheta}(x) - \vec{\mathbf{e}}|^2 d\|S_{r, \vartheta}\|x - \int_{\mathbf{C}_{(1+\vartheta)r}} F_0 \left(\vec{\mathbf{e}} \right) d\|P_{(1+\vartheta)r}\|, \end{aligned}$$

where $DF_0(\vec{\mathbf{e}})\vec{\mathbf{e}} = F_0(\vec{\mathbf{e}})$, by homogeneity, and

$$\int_{\mathbf{C}_{(1+\vartheta)r}} DF_0(\vec{\mathbf{e}})\vec{S}_{r, \vartheta}(x) d\|S_{r, \vartheta}\|x = \int_{\mathbf{C}_{(1+\vartheta)r}} DF_0(\vec{\mathbf{e}})\vec{\mathbf{e}} d\|P_{(1+\vartheta)r}\|,$$

because $\partial S_{r, \vartheta} = \partial P_{(1+\vartheta)r}$, by (4.2) and (4.5), and the integrals are just the evaluations of $S_{r, \vartheta}$ and $P_{(1+\vartheta)r}$ on the constant (hence exact) differential n -form $DF_0(\vec{\mathbf{e}})$. Consequently, by (4.4), (4.6), and (1.24),

$$(4.9) \quad \begin{aligned} \mathbf{F}_0 \left(S_{r, \vartheta} \right) - \mathbf{F}_0 \left(P_{(1+\vartheta)r} \right) &\leq \Lambda \int_{\mathbf{C}_{(1+\vartheta)r}} |\vec{S}_{r, \vartheta} - \vec{\mathbf{e}}|^2 d\|S_{r, \vartheta}\| \\ &\leq \text{const}(n) \Lambda \left[\vartheta \mathbf{E} \left(T, 1 \right) + \frac{1}{\vartheta} \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|x \right]. \end{aligned}$$

To estimate the other terms appearing in (4.8) we introduce $f(x) = F(0, \vec{\mathbf{e}}) - F(x, \vec{\mathbf{e}})$ and observe

$$\begin{aligned}
& |(\mathbf{F}_0 - \mathbf{F})(T_r) - (\mathbf{F}_0 - \mathbf{F})(P_r)| \\
& \leq \int \int_0^1 \left\| D_{(2)}F(0, (1-t)\vec{\mathbf{e}} + t\vec{T}(x)) - D_{(2)}F(x, (1-t)\vec{\mathbf{e}} + t\vec{T}(x)) \right\| |\vec{T}(x) - \vec{\mathbf{e}}| dt d\|T_r\|x \\
& \quad + \left| \int f(x) d\|T_r\|x - \int f(x) d\|P_r\|x \right| \\
& \leq \int \kappa(|x|) |\vec{T}(x) - \vec{\mathbf{e}}| d\|T_r\|x + \int |f(x) - f(\mathbf{p}x, 0)| d\|T_r\|x \\
& \quad + \left| \int f(\mathbf{p}x, 0) d\|T_r\|x - \int f(\mathbf{p}x, 0) d\|P_r\|x \right| \\
& \leq \frac{1}{2}\vartheta \int |\vec{T}(x) - \vec{\mathbf{e}}|^2 d\|T_r\|x + \frac{1}{2\vartheta} \kappa(R)^2 \mathbf{M}(T_r) + \int \mu(|\mathbf{q}x|) d\|T_r\|x \\
& \quad + \kappa(R) [\mathbf{M}(T_r) - \mathbf{M}(P_r)] \\
& \leq (\vartheta + \kappa(R)) \mathbf{E}(T, 1) + \frac{1}{2\vartheta} \kappa(R)^2 \mathbf{M}(T) + \int \mu(|\mathbf{q}x|) d\|T_r\|.
\end{aligned}$$

Here we have used (1.6), (1.7), (1.5), i.e. $|f(x)| \leq \kappa(|x|) \leq \kappa(R)$, and the inequality

$$\left| \int \varphi(\mathbf{p}x) d\|T_r\|x - \int \varphi(\mathbf{p}x) d\|P_r\|x \right| \leq \sup |\varphi| \mathbf{E}(T, 1)$$

for continuous functions φ on \mathbb{R}^n , an inequality which one readily proves by approximating φ with finitely valued functions. Recalling the definition (1.9) of $\hat{\mu}$ we use the inequalities $\mu(s) - s^2 \leq \hat{\mu}(\frac{1}{2})$ for $0 \leq s \leq \frac{1}{2}$ and $\mu(s) \leq 2s\mu(\frac{1}{2}) \leq s^2 + \mu(\frac{1}{2})^2 \leq s^2 + \kappa(\frac{1}{2})^2$ for $s > \frac{1}{2}$ to estimate

$$\int \mu(|\mathbf{q}x|) d\|T_r\|x \leq (\kappa(R)^2 + \hat{\mu}(R)) \mathbf{M}(T) + \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|x.$$

Collecting the estimates we have proved

$$\begin{aligned}
(4.10) \quad & |(\mathbf{F}_0 - \mathbf{F})(T_r) - (\mathbf{F}_0 - \mathbf{F})(P_r)| \\
& \leq 2\vartheta \mathbf{E}(T, 1) + \left[\frac{2}{\vartheta} \kappa(R)^2 + \hat{\mu}(R) \right] \mathbf{M}(T) + \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|x.
\end{aligned}$$

Similarly as above we estimate

$$\begin{aligned}
& |(\mathbf{F}_0 - \mathbf{F})(\Sigma_{r,\vartheta}) - (\mathbf{F}_0 - \mathbf{F})(P_r)| \\
& \leq (1 + \kappa(R)) \mathbf{E}(\Sigma_{r,\vartheta}, 1) + \frac{1}{2} \kappa(R)^2 \mathbf{M}(\Sigma_{r,\vartheta}) + \int_{\mathbf{C}} \mu(|\mathbf{q}x|) d\|\Sigma_{r,\vartheta}\|x.
\end{aligned}$$

Here we apply (4.4) to estimate $\mathbf{E}(\Sigma_{r,\vartheta}, 1)$, and the inequality

$$\mathbf{M}(\Sigma_{r,\vartheta}) = \mathbf{E}(\Sigma_{r,\vartheta}, 1) + \mathbf{M}(P_r) \leq \mathbf{E}(\Sigma_{r,\vartheta}, 1) + \mathbf{M}(T)$$

to estimate $\mathbf{M}(\Sigma_{r,\vartheta})$. We use again $\mu(s) \leq \hat{\mu}(\frac{1}{2}) + s^2$ for $0 < s \leq \frac{1}{2}$ and $\mu(s) \leq 2s\mu(\frac{1}{2})$ for $s > \frac{1}{2}$ to get

$$\int_{\mathbf{C}} \mu(|\mathbf{q}x|) d\|\Sigma_{r,\vartheta}\|x \leq \hat{\mu}(\frac{1}{2}) \mathbf{M}(\Sigma_{r,\vartheta}) + \int_{|\mathbf{q}x| \leq \frac{1}{2}} |\mathbf{q}x|^2 d\|\Sigma_{r,\vartheta}\|x + 2\mu(\frac{1}{2}) \int_{|\mathbf{q}x| > \frac{1}{2}} |\mathbf{q}x| d\|\Sigma_{r,\vartheta}\|x.$$

Noting

$$|\mathbf{q}(h^\vartheta(t, x'))|^2 = (1-t)^2 |\mathbf{q}x'|^2 \leq \min(|\mathbf{q}x'|, |\mathbf{q}x'|^2)$$

if $|\mathbf{q}(h^\vartheta(t, x'))| \leq \frac{1}{2}$ and $x' \in \text{spt} \langle T, r \rangle$, we estimate

$$\begin{aligned} & \int_{|\mathbf{q}x| \leq \frac{1}{2}} |\mathbf{q}x|^2 d\|\Sigma_{r, \vartheta}\|_x \\ & \leq \int_0^1 \int_{\partial \mathbf{C}_r} \min(|\mathbf{q}x'|, |\mathbf{q}x'|^2) |(\vartheta \mathbf{p}x', \mathbf{q}x')| (1-\vartheta t)^{n-1} d\|\langle T, r \rangle\|_{x'} dt \\ & \leq (1+\vartheta r) \int_{\partial \mathbf{C}_r} |\mathbf{q}x'|^2 d\|\langle T, r \rangle\|_{x'} \leq 12(1+\vartheta r) \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|_x. \end{aligned}$$

Furthermore,

$$\begin{aligned} & 2\mu\left(\frac{1}{2}\right) \int_{|\mathbf{q}x| > \frac{1}{2}} |\mathbf{q}x| d\|\Sigma_{r, \vartheta}\|_x \\ & \leq 2\mu\left(\frac{1}{2}\right) \int_0^1 \int_{\partial \mathbf{C}_r} (1-t) |\mathbf{q}x'| |(\vartheta \mathbf{p}x', \mathbf{q}x')| (1-\vartheta t)^{n-1} d\|\langle T, r \rangle\|_{x'} dt \\ & \leq \mu\left(\frac{1}{2}\right)^2 \vartheta^2 r^2 \mathbf{M}(\langle T, r \rangle) + \left[1 + 2\mu\left(\frac{1}{2}\right)\right] \int_{\partial \mathbf{C}_r} |\mathbf{q}x'|^2 d\|\langle T, r \rangle\|_{x'} \\ & \leq \mu\left(\frac{1}{2}\right)^2 \mathbf{M}(T) + 24 \left[1 + \mu\left(\frac{1}{2}\right)\right] \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|_x. \end{aligned}$$

Collecting terms and using the monotonicity of κ , μ , $\hat{\mu}$ as well as $\mu \leq \kappa$ we arrive at

$$(4.11) \quad \begin{aligned} |(\mathbf{F}_0 - \mathbf{F})(\Sigma_{r, \vartheta}) - (\mathbf{F}_0 - \mathbf{F})(P_r)| & \leq 24\vartheta \mathbf{E}(T, 1) + 5 \left[\kappa(R)^2 + \hat{\mu}(R) \right] \mathbf{M}(T) \\ & + \frac{80}{\vartheta} \left(1 + \kappa(R)^2 + \hat{\mu}(R) \right) \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|_x. \end{aligned}$$

Combining (4.8)–(4.11) and estimating the term $\omega(R)\mathbf{M}(\Sigma_{r, \vartheta})$ similarly as above we finally conclude

$$\begin{aligned} \lambda 2^{-n} \mathbf{E}\left(T, \frac{1}{2}\right) & \leq \text{const}(n) \Lambda \vartheta \mathbf{E}(T, 1) + \frac{7}{\vartheta} \left(\kappa^2(R) + \hat{\mu}(R) + \omega(R) \right) \mathbf{M}(T) \\ & + \text{const}(n) \frac{\Lambda}{\vartheta} \left(1 + \kappa^2(R) + \hat{\mu}(R) + \omega(R) \right) \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|_x. \end{aligned}$$

This is true for $0 < \vartheta \leq \frac{1}{7}$, but increasing the constants if necessary we obtain the same inequality trivially for all $\vartheta > 0$, and (4.1) follows. \blacksquare

4.2 Lemma (reverse Poincaré inequality). *Suppose $T \in \mathcal{T}_*(n, k, m)$ is (F, ω) minimizing in \mathbf{C} with support in $\mathbf{B}^n \times \mathbf{B}^k(0, h)$, $h > 0$, and $r = \sqrt{1+h^2}$. Then the inequality*

$$(4.12) \quad \mathbf{E}\left(T, \frac{1}{5}\right) \leq C_{12} \left[\chi(r) \mathbf{M}(T) + (1 + \chi(r)) \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|_x \right]$$

holds with $C_{12} = \text{const}(n, \lambda, \Lambda)$ and $\chi(r) = \kappa(r)^2 + \hat{\mu}(r) + \omega(r)$.

The point here is that $\chi(r)$ is small in later applications so that (4.12) is indeed a reverse Poincaré inequality, up to a small extra term. Multiplicity $m = 0$ is admitted in Lemma 4.2 but irrelevant for later applications.

Again we point out that the condition $0 \in \text{spt } T$ from the definition of $\mathcal{T}_i(n, k, m)$ is not needed in the proof.

Proof. We consider the boundary situation $T \in \mathcal{T}_b(n, k, m)$ and n -balls $\mathbf{B}_\sigma(y) \subset \mathbf{U}$ such that $y \in \mathbf{U}^{n-1} \times \{0\}$ or $\mathbf{B}_\sigma(y) \cap \mathbf{U}^{n-1} \times \{0\} = \emptyset$. We want to apply Lemma 4.1 (with the comment following it) to the scaled currents $T_\sigma = \left(\mu_{\sigma\#}^{-1} \tau_{(y,0)\#}^{-1} T \right) \llcorner \mathbf{C}$. (Note that we may have $T_\sigma \in \mathcal{T}_i(n, k, 0)$ in the case $T \in \mathcal{T}_b(n, k, 1)$; this is the reason for admitting $m = 0$ in Lemma 4.1.) Clearly, T_σ satisfies the hypothesis of Lemma 4.1 with respect to the integrand $F_\sigma(x, \xi) = F((y, 0) + \sigma x, \xi)$ and the function $\omega_\sigma(\varrho) = \omega(\sigma\varrho)$. For the structure constants and moduli of F_σ we have $\lambda_\sigma = \lambda$, $\Lambda_\sigma = \Lambda$, $\kappa_\sigma(\varrho) = \kappa(\sigma\varrho)$, $\mu_\sigma(\varrho) = \mu(\sigma\varrho)$, $\widehat{\mu}_\sigma(\varrho) \leq \widehat{\mu}(\sigma\varrho)$, and T_σ has support in $\mathbf{B}^{n+k}(0, \sigma^{-1}r)$. Therefore, (4.1) implies

$$(4.13) \quad \mathbf{E} \left(T, y, \frac{1}{2}\sigma \right) \leq \vartheta \mathbf{E} (T, y, \sigma) + \frac{1}{\vartheta} \left[c\sigma^{-n-2} \int_{\mathbf{C}_\sigma(y)} |\mathbf{q}x|^2 d\|T\|x + c'\sigma^{-n} \mathbf{M} (T \llcorner \mathbf{C}_\sigma(y)) \right],$$

where we have abbreviated $c = C_{11}(1 + \chi(r))$ and $c' = C_{11}\chi(r)$. For balls $\mathbf{B}_\varrho(y) \subset \mathbf{U}$ which intersect $\mathbf{U}^{n-1} \times \{0\}$ we define $y' = (y_1, \dots, y_{n-1}, 0)$ and observe

$$\mathbf{B}_\varrho(y) \subset \mathbf{B}_{2\varrho}(y') \subset \mathbf{B}_{4\varrho}(y') \subset \mathbf{B}_{5\varrho}(y),$$

hence (4.13) applied to $\mathbf{B}_{2\varrho}(y')$ gives, provided $\mathbf{B}_{5\varrho}(y) \subset \mathbf{U}$,

$$\begin{aligned} 2^{-n} \mathbf{E} (T, y, \varrho) &\leq \mathbf{E} (T, y', 2\varrho) \\ &\leq \vartheta \left(\frac{4}{5} \right)^n \mathbf{E} (T, y, 5\varrho) + \frac{1}{\vartheta} \left[c(4\varrho)^{-n-2} \int_{\mathbf{C}_{5\varrho}(y)} |\mathbf{q}x|^2 d\|T\|x + c'(4\varrho)^{-n} \mathbf{M} (T \llcorner \mathbf{C}_{5\varrho}(y)) \right]. \end{aligned}$$

Increasing c and c' by a factor $\text{const}(n)$ we therefore have

$$(4.14) \quad \mathbf{E} \left(T, y, \frac{1}{5}\sigma \right) \leq \vartheta \mathbf{E} (T, y, \sigma) + \frac{1}{\vartheta} \left[c\sigma^{-n-2} \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|x + c'\sigma^{-n} \mathbf{M} (T) \right]$$

for all balls $\mathbf{B}_\sigma(y) \subset \mathbf{U}$ and parameters $\vartheta > 0$.

With (4.14) we are in a position to apply a covering argument as in [Si2, §2.8]. For this we set

$$\varphi(\mathbf{B}_\sigma(y)) = \sigma^n \mathbf{E} (T, y, \sigma) = \frac{1}{2} \int_{\mathbf{C}_\sigma(y)} |\vec{T}(x) - \vec{\mathbf{e}}|^2 d\|T\|x$$

and rewrite (4.14), using also $\sigma^2 < 1$,

$$(4.15) \quad \sigma^2 \varphi(\mathbf{B}_{\sigma/5}(y)) \leq \vartheta \sigma^2 \varphi(\mathbf{B}_\sigma(y)) + \gamma(\vartheta)$$

with

$$(4.16) \quad \gamma(\vartheta) = \frac{1}{\vartheta} \left[c5^{-n} \int_{\mathbf{C}} |\mathbf{q}x|^2 d\|T\|x + 5^{-n} c' \mathbf{M} (T) \right].$$

We define

$$(4.17) \quad Q = \sup \left\{ \varrho^2 \varphi(\mathbf{B}_\varrho(y)) : \mathbf{B}_{5\varrho}(y) \subset \mathbf{U} \right\}$$

and note $Q \leq 5^{-2} \mathbf{E}(T, 0, 1) < \infty$. Covering $\mathbf{B}_\varrho(y) \subset \mathbf{U}_{1-\varrho}$ by balls $\mathbf{B}_{\varrho/25}(y_i)$ with $y_i \in \mathbf{B}_\varrho(y)$, $i = 1, \dots, M$, and $M = \text{const}(n)$, we deduce from the subadditivity of the functional $\varphi(\cdot)$ on balls in \mathbf{U} and from (4.15)

$$(4.18) \quad \begin{aligned} \varrho^2 \varphi(\mathbf{B}_\varrho(y)) &\leq \varrho^2 \sum_{i=1}^M \varphi(\mathbf{B}_{\varrho/25}(y_i)) \\ &\leq 5^2 \sum_{i=1}^M \left[\vartheta \left(\frac{\varrho}{5} \right)^2 \varphi(\mathbf{B}_{\varrho/5}(y_i)) + \gamma(\vartheta) \right] \\ &\leq 5^2 \vartheta Q M + 5^2 \gamma(\vartheta) M. \end{aligned}$$

Taking the supremum over all balls $\mathbf{B}_\varrho(y)$ with $\mathbf{B}_{5\varrho}(y) \subset \mathbf{U}$ in this inequality we get

$$(4.19) \quad Q \leq 5^2 \vartheta Q M + 5^2 M \gamma(\vartheta).$$

Fixing now $\vartheta = (50M)^{-1}$ we see $Q \leq 50M\gamma(\vartheta)$ from (4.19), and from (4.17) we obtain

$$\varrho^2 \varphi(\mathbf{B}_\varrho(y)) \leq 50M\gamma(\vartheta),$$

i.e.

$$\mathbf{E}(T, y, \varrho) \leq \varrho^{-n-2} 50M\gamma(\vartheta).$$

Applying this with $y = 0$, $\varrho = \frac{1}{5}$ and recalling (4.16) we have proved (4.12).

The proof in the interior situation $T \in \mathcal{T}_i(n, k, m)$ is simpler, as one can use directly (4.13) for all balls $\mathbf{B}_\sigma(y) \subset \mathbf{U}$. ■

5 Excess decay

As is well known [Alm1], [Alm2], [Bo], [Fe, §5.3], [Ha1] the crucial step in the regularity proof is excess improvement, an idea introduced by Almgren. If T is (F, ω) minimizing with small excess $\mathbf{E}(T, 1)$ in the cylinder of radius 1 over the n -plane $\mathbb{R}^n \times \{0\}^k$ in \mathbb{R}^{n+k} , then one shows that the excess of T over some n -plane S close to $\mathbb{R}^n \times \{0\}^k$ and in a cylinder with smaller radius is substantially smaller than $\mathbf{E}(T, 1)$. By iteration of this argument one can prove that the excess of T in concentric cylinders of radius r over some n -plane decreases to zero with r , and from this excess decay one deduces finally the regularity of T in a small cylinder.

In this section we first prove (Lemma 5.1) that the square integral of the height of T over a *tilted* n -plane S , which is close to $\mathbb{R}^n \times \{0\}^k$, can be dominated by a small constant times $\mathbf{E}(T, 1) + \kappa(\frac{1}{2})^2 + \hat{\mu}(\frac{1}{2}) + \omega(\frac{1}{2})$, provided this quantity is small. This n -plane S is found as the graph of the differential at zero of the A -harmonic function u which is associated by Lemma 3.3 with the approximatively A -harmonic Lipschitz approximation g of T (Lemma 3.1 and Lemma 3.2). Then we use the reverse Poincaré inequality (Lemma 4.2) to replace the square integral of the height of T over the tilted plane S by the excess of T over this plane

in a cylinder of smaller radius $\vartheta \in]0, 1[$. The result is Lemma 5.2, an estimate of this latter excess by $\vartheta \mathbf{E}(T, 1)$ plus extra terms involving only the (small) quantity $\kappa(\frac{1}{2})^2 + \widehat{\mu}(\frac{1}{2}) + \omega(\frac{1}{2})$. The iteration of Lemma 5.2 then leads to the desired excess decay in Lemma 5.4.

We assume in this section that F is an elliptic integrand with ellipticity constant λ , bound Λ , and moduli $\kappa, \mu, \widehat{\mu}, \nu$ as in Section 1. In Lemma 5.4 we need the extra assumption (1.15) on $\kappa, \widehat{\mu}$ and the function ω which appears in the (F, ω) minimality condition for the rectifiable n -current T .

5.1 Lemma (improving the L^2 height estimate by tilting). *Suppose $T \in \mathcal{T}_b(n, k, 1)$ or $T \in \mathcal{T}_i(n, k, m)$ with (2.12), T is (F, ω) minimizing in \mathbf{C} , $\chi = \kappa(\frac{1}{2})^2 + \widehat{\mu}(\frac{1}{2}) + \omega(\frac{1}{2}) \leq 1$, $E = \mathbf{E}(T, 1)$, $0 < \epsilon \leq 1$, $0 < \sigma \leq 1$, $0 < \gamma \leq 1$ and a smallness condition*

$$(5.1) \quad C_{17} \left[\gamma^{-2} \sqrt{E} + \sqrt{\sigma} + \nu(\gamma + \sqrt{\chi}) \right] \leq \delta(n, k, \lambda, \Lambda, \epsilon)$$

is satisfied, where $\delta(n, k, \lambda, \Lambda, \epsilon) \in]0, 1[$ is from Lemma 3.3 and $C_{17} = \text{const}(n, m, \lambda, \Lambda)$. Then there exist $L \in \text{Hom}(\mathbb{R}^n, \mathbb{R}^k)$ and $v \in \mathbb{R}^{n+k}$ perpendicular to the n -plane $S = \text{graph}(L)$ such that for $0 < \eta \leq \frac{1}{17}$ the inequality

$$(5.2) \quad \int_{\mathbf{C}_{\eta/2}} \text{dist}(x - v, S)^2 d\|T\|x \leq C_{18} \left(\gamma^{-2} E^{1/n} + \eta^{n+4} + \epsilon \right) \left(E + \sigma^{-1} \chi \right)$$

is valid with a constant $C_{18} = \text{const}(n, m, \lambda, \Lambda)$. Moreover

$$(5.3) \quad \|L\| \leq C_{18} \sqrt{E + \sigma^{-1} \chi}$$

holds, and in the case $T \in \mathcal{T}_b(n, k, 1)$ we have $L = 0$ on $\mathbb{R}^{n-1} \times \{0\}$, $v = 0$, while in the case $T \in \mathcal{T}_i(n, k, m)$

$$(5.4) \quad |v| \leq C_{18} \left(\sqrt{E + \sigma^{-1} \chi} + E^{1/2n} \right).$$

Proof. We consider the boundary situation $T \in \mathcal{T}_b(n, k, 1)$ and indicate only some modifications of the proof for the case $T \in \mathcal{T}_i(n, k, m)$ satisfying (2.12). We fix $A = D^2 F_0^{\S}$, choose $\varrho = \frac{1}{17}$ in Lemma 3.2, and let g be the Lipschitz approximation for T corresponding to $\beta = 1$. Then we know from Lemma 3.1, (ii) and (iii)

$$\int_{\mathbf{U}_+ \cap \mathbf{U}_\varrho} |Dg|^2 d\mathcal{L}^n \leq C_{13} E,$$

where $C_{13} = \text{const}(n, m) \Lambda^{2n-2}$ in view of our assumption $\omega(\frac{1}{2}) \leq 1$. Scaling

$$f = \varrho^{n/2-1} C_{13}^{-1/2} \left(E + \sigma^{-1} \chi \right)^{-1/2} g$$

with a parameter $0 < \sigma \leq 1$, we have

$$\varrho^{2-n} \int_{\mathbf{U}_+ \cap \mathbf{U}_\varrho} |Df|^2 d\mathcal{L}^n \leq 1$$

and obtain from Lemma 3.2

$$\int_{\mathbf{U}_\varrho} A(Df, D\zeta) d\mathcal{L}^n \leq \varrho^{n/2-1} C_{14} \left[\gamma^{-2} \sqrt{E} + \sqrt{\sigma} + \nu (\gamma + \sqrt{\chi}) \right] \sup |D\zeta|$$

for all functions $\zeta \in \mathbb{C}^1(\mathbf{U}, \mathbb{R}^k)$ with compact support in $\mathbf{U}_+ \cap \mathbf{U}_\varrho$, where $0 < \gamma \leq 1$ can be chosen arbitrarily and $C_{14} = \max \{ 2C_{13}^{-1/2} C_{10}, 1 \}$. The hypothesis $\mathbf{C}_\varrho \cap \text{spt } T \subset \mathbf{U}_\varrho^n \times \mathbf{B}_{1/3}^k$ in Lemma 3.2 is satisfied, by Lemma 2.2, if we choose the constant in (5.1) large enough.

We now apply Lemma 3.3 to obtain, for given $\epsilon > 0$, an A -harmonic function $u \in W_{(0)}^{1,2}(\mathbf{U}_+ \cap \mathbf{U}_\varrho, \mathbb{R}^k)$ such that

$$(5.5) \quad \varrho^{-2} \int_{\mathbf{U}_+ \cap \mathbf{U}_\varrho} |u - g|^2 d\mathcal{L}^n \leq \epsilon C_{13} \left[E + \sigma^{-1} \chi \right],$$

and

$$(5.6) \quad \int_{\mathbf{U}_+ \cap \mathbf{U}_\varrho} |Du|^2 d\mathcal{L}^n \leq C_{13} \left[E + \sigma^{-1} \chi \right],$$

provided

$$(5.7) \quad \varrho^{-n/2} C_{14} \left[\gamma^{-2} \sqrt{E} + \sqrt{\sigma} + \nu (\gamma + \sqrt{\chi}) \right] \leq \delta(n, k, \lambda, \Lambda, \epsilon).$$

Noting that u is smooth on $\overline{\mathbf{U}}_+ \cap \mathbf{U}_\varrho$ with $u = 0$ on $\mathbf{U}_\varrho^{n-1} \times \{0\}$ we define $L = Du(0)$ and observe $Ly = y_n \ell$ for some $\ell \in \mathbb{R}^k$ and all $y \in \mathbb{R}^n$. From the a-priori-estimate for solutions to homogeneous constant coefficient elliptic systems

$$(5.8) \quad \varrho^2 \sup_{\mathbf{U}_+ \cap \overline{\mathbf{U}}_{\varrho/2}} \|Du\|^2 + \varrho^4 \sup_{\mathbf{U}_+ \cap \overline{\mathbf{U}}_{\varrho/2}} \|D^2u\|^2 \leq \text{const}(n, \lambda, \Lambda) \varrho^{2-n} \int_{\mathbf{U}_+ \cap \mathbf{U}_\varrho} |Du|^2 d\mathcal{L}^n$$

(which follows from Caccioppoli's inequality for u and its derivatives of any order, Sobolev's inequality, and Poincaré's inequality, see [Ca], [Gi], [Mo]) we deduce by Taylor expansion

$$(5.9) \quad \sup_{y \in \mathbf{U}_+ \cap \mathbf{U}_{\eta/2}} |u(y) - Ly|^2 \leq C_{15} \left(\frac{\eta}{\varrho} \right)^4 \varrho^{2-n} \int_{\mathbf{U}_+ \cap \mathbf{U}_\varrho} |Du|^2 d\mathcal{L}^n$$

for $0 < \eta \leq \varrho$, where $C_{15} = \text{const}(n, \lambda, \Lambda)$. (In the interior situation $T \in \mathcal{T}_i(n, k, m)$ we have the same estimate for $|u(y) - u(0) - Ly|^2$ on $\mathbf{U}_{\eta/2}$.) Combining (5.9) with (5.5) and (5.6) we get

$$(5.10) \quad \begin{aligned} & \int_{\mathbf{U}_+ \cap \mathbf{U}_{\eta/2}} |g(y) - Ly|^2 d\mathcal{L}^n y \\ & \leq 2 \int_{\mathbf{U}_+ \cap \mathbf{U}_{\eta/2}} |g(y) - u(y)|^2 d\mathcal{L}^n y + 2 \int_{\mathbf{U}_+ \cap \mathbf{U}_{\eta/2}} |u(y) - Ly|^2 d\mathcal{L}^n y \\ & \leq C_{16} (\epsilon + \eta^{n+4}) (E + \sigma^{-1} \chi) \end{aligned}$$

with $C_{16} = \text{const}(n, m, \lambda, \Lambda)$. (In the interior situation we get the same estimate for the integral of $|g(y) - u(0) - Ly|^2$ on $\mathbf{U}_{\eta/2}$.)

Next we estimate with (5.8) and $L = Du(0)$

$$\sup_{y \in \mathbf{U}_+ \cap \mathbf{U}_{\varrho/2}} \|Du(y) - L\|^2 \leq \text{const}(n, \lambda, \Lambda) \varrho^{-n} \int_{\mathbf{U}_+ \cap \mathbf{U}_{\varrho}} |Du|^2 d\mathcal{L}^n$$

and obtain, using (5.6) again,

$$(5.11) \quad \begin{aligned} \|L\| &\leq \int_{\mathbf{U}_+ \cap \mathbf{U}_{\varrho/2}} \|L - Du(y)\| d\mathcal{L}^n y + \int_{\mathbf{U}_+ \cap \mathbf{U}_{\varrho/2}} \|Du(y)\| d\mathcal{L}^n y \\ &\leq \text{const}(n, m, \lambda, \Lambda) \sqrt{E + \sigma^{-1}\chi}. \end{aligned}$$

In order to estimate the integral of $\text{dist}(x, S)^2$ for the n -plane $S = \text{graph}(L)$ we observe

$$\begin{aligned} &\int_{\mathbf{C}_{\eta/2}} \text{dist}(x, S)^2 d\|T\|x \\ &= \int_{\mathbf{C}_+ \cap \mathbf{C}_{\eta/2} \setminus (A_{g,\gamma} \times \mathbb{R}^k)} \dots d\|T_g\| + \int_{\mathbf{C}_{\eta/2}} \dots d\|T - T_g\| + \int_{\mathbf{C}_{\eta/2} \cap (A_{g,\gamma} \times \mathbb{R}^k)} \dots d\|T_g\| \\ &= \text{I} + \text{II} + \text{III}, \end{aligned}$$

where $A_{g,\gamma}$ is the *bad* set of g corresponding to the parameter γ as in Lemma 3.1, (iv). For $x \in \mathbf{C}_{\eta/2} \cap \text{spt } T$ we have

$$\begin{aligned} \text{dist}(x, S) &\leq |\mathbf{q}x - L\mathbf{p}x| \leq |\mathbf{q}x| + |L\mathbf{p}x| \\ &\leq \text{const}(n, m, \lambda, \Lambda) \left[E^{1/2n} + \eta \sqrt{E + \sigma^{-1}\chi} \right], \end{aligned}$$

by (5.11) and Lemma 2.2. (We have already noted that (5.7) implies the smallness condition (2.4) necessary to apply Lemma 2.2.) From Lemma 3.1, (i) and (iv) (with $\beta = 1$) we therefore obtain

$$|\text{II}| + |\text{III}| \leq \text{const}(n, m, \lambda, \Lambda) \left[E^{1/n} + \eta^2 (E + \sigma^{-1}\chi) \right] \gamma^{-2} E.$$

Letting $G(y) = (y, g(y))$ we also see from (5.10), taking $\text{dist}(G(y), S) \leq |g(y) - Ly|$ and $\text{Lip } g \leq \beta = 1$ into account,

$$\begin{aligned} |\text{I}| &\leq \int_{\mathbf{U}_+ \cap \mathbf{U}_{\eta/2}} |g(y) - Ly|^2 JG(y) d\mathcal{L}^n y \\ &\leq \text{const}(n, m, \lambda, \Lambda) (\epsilon + \eta^{n+4}) (E + \sigma^{-1}\chi). \end{aligned}$$

The result is

$$\int_{\mathbf{C}_{\eta/2}} \text{dist}(x, S)^2 d\|T\|x \leq \text{const}(n, m, \lambda, \Lambda) (\gamma^{-2} E^{1/n} + \eta^{n+4} + \epsilon) (E + \sigma^{-1}\chi).$$

With this inequality and (5.11) we have proved all the assertions of the Lemma in the boundary case under the smallness condition (5.7), i.e. (5.1). In the interior case we obtain the same result and note additionally

$$\begin{aligned} \frac{1}{3} \int_{\mathbf{U}_{\varrho/2}} |u(0) + Ly|^2 d\mathcal{L}^n y &\leq \int_{\mathbf{U}_{\varrho/2}} |u(0) + Ly - g(y)|^2 d\mathcal{L}^n y + \int_{\mathbf{U}_{\varrho/2}} |g|^2 d\mathcal{L}^n \\ &\leq \text{const}(n, m, \lambda, \Lambda) \left[(1 + \epsilon) (E + \sigma^{-1}\chi) + E^{1/n} \right], \end{aligned}$$

by the analogue of (5.10), by (5.5), and by Lemma 3.1, (vi). In view of

$$|u(0)| = \left| \int_{\mathbf{U}_{\varrho/2}} (u(0) + Ly) d\mathcal{L}^n \right|$$

this implies the estimate (5.4) for v which equals the orthogonal projection of $u(0)$ onto S^\perp . \blacksquare

We note that in the boundary case $T \in \mathcal{T}_b(n, k, m)$ with multiplicity $m \geq 2$ and with the additional density condition (2.13) the proof above does not work, because we could only prove in Lemma 3.2 that g is approximatively A -harmonic on $\mathbf{U}_+ \cap \mathbf{U}_\varrho$ and on $\mathbf{U}_- \cap \mathbf{U}_\varrho$ separately. Consequently we can find A -harmonic functions u_+ near g on $\mathbf{U}_+ \cap \mathbf{U}_\varrho$ and u_- on $\mathbf{U}_- \cap \mathbf{U}_\varrho$ but we do not know that the differentials $Du_+(0)$, $Du_-(0)$ coincide and, hence, cannot find, with the method above, a tilted n -plane S such that (5.2) holds.

In the following Lemma we use, in the interior situation, the modulus $\widehat{\kappa}$ associated with κ by (1.9), and we recall $\frac{1}{4}\kappa(\frac{1}{2})^2 \leq \widehat{\kappa}(\frac{1}{2})$ if $\kappa(\frac{1}{2}) \leq 1$.

5.2 Lemma (excess improvement by tilting). *For $0 < \vartheta \leq \frac{1}{340}$ there exists a positive constant $\delta_1(n, k, m, \lambda, \Lambda, \nu(\cdot), \vartheta) \leq 1$ such that the following is true: Whenever $T \in \mathcal{T}_b(n, k, 1)$ or $T \in \mathcal{T}_i(n, k, m)$ with (2.12) is (F, ω) minimizing in \mathbf{C} with $\text{spt } T \subset \mathbb{R}^n \times \mathbf{B}^k$ and the smallness conditions*

$$(5.12) \quad E = \mathbf{E}(T, 1) \leq \delta_1, \quad \chi \leq \delta_1,$$

are satisfied, where $\chi = \widehat{\kappa}(\frac{1}{2}) + \omega(\frac{1}{2})$ in the interior case $T \in \mathcal{T}_i(n, k, m)$ and $\chi = \kappa(\frac{1}{2})^2 + \widehat{\mu}(\frac{1}{2}) + \omega(\frac{1}{2})$ in the boundary case $T \in \mathcal{T}_b(n, k, m)$, then there exists a linear isometry \mathbb{I} of \mathbb{R}^{n+k} with

$$(5.13) \quad \|\mathbb{I} - \text{id}\| \leq C_{21} \sqrt{E + \chi},$$

and with $\mathbb{I} = \text{id}$ on $(\mathbb{R}^{n-1} \times \{0\}) \times \{0\}^k$ in the case $T \in \mathcal{T}_b(n, k, 1)$, such that

$$(5.14) \quad \mathbf{E}(\mathbb{I} \# T, \vartheta) \leq C_{22} \vartheta^2 \mathbf{E}(T, 1) + C_{21} \chi.$$

Here $C_{22} = \text{const}(n, m, \lambda, \Lambda)$, and $C_{21} = \text{const}(n, k, m, \lambda, \Lambda, \vartheta)$.

Proof. We assume that (5.1) is satisfied with parameters ϵ, γ to be specified later, $0 < \sigma \leq 1$, $\chi \leq 1$, and $E \leq 1$. Since (5.1) implies the smallness condition (2.1) in the hypotheses of Lemma 2.2 (by the choice of C_{17}), we can use the height estimate

$$(5.15) \quad \sup \{ |\mathbf{q}x| : x \in \mathbf{C}_{1/2} \cap \text{spt } T \} \leq \text{const}(n, \Lambda) E^{1/2n}.$$

Letting $S = \text{graph}(L)$ as in Lemma 5.1 we see from (5.3) that there is a linear isometry \mathbb{I} of \mathbb{R}^{n+k} satisfying $\mathbb{I} \# S = \mathbb{R}^n \times \{0\}^k$, $\mathbb{I} = \text{id}$ on $(\mathbb{R}^{n-1} \times \{0\}) \times \{0\}^k$ in the case $T \in \mathcal{T}_b(n, k, m)$, and

$$(5.16) \quad \|\mathbb{I} - \text{id}\| \leq \text{const}(n, m, \lambda, \Lambda) \sqrt{E + \sigma^{-1} \chi}.$$

Assuming $0 < \eta \leq \frac{1}{17}$ and a smallness condition

$$(5.17) \quad C_{19} \left(E^{1/2n} + \sqrt{E + \sigma^{-1}\chi} \right) \leq \eta$$

with a sufficiently large constant $C_{19} = \text{const}(n, m, \lambda, \Lambda)$, we verify by elementary geometric considerations, using the height bound (5.15) in $\mathbf{C}_{1/2}$, the height bound 1 in $\mathbf{C} \setminus \mathbf{C}_{1/2}$, (5.16), and (5.4), that

$$(5.18) \quad \left(\cdot, \tau^{-1} \mathbf{C}_{\eta/4} \right) \cap \text{spt } T \subset \mathbf{C}_{\eta/2}$$

and

$$(5.19) \quad \mathbf{C}_{\eta/4} \cap \tau_w^{-1} \left(\cdot, \text{spt } T \right) \subset \mathbf{U}_{\eta/4}^n \times \mathbf{B}^k(0, \eta),$$

where $w = \cdot, v$ and $v \in S^\perp$ is from Lemma 5.1. The conclusion (5.2) of this Lemma and (5.15) imply

$$(5.20) \quad \begin{aligned} \int_{\mathbf{C}_{\eta/4}} |\mathbf{q}x - w|^2 d\|\cdot, \#T\|x &= \int_{\Gamma^{-1}\mathbf{C}_{\eta/4}} \text{dist}(x - v, S)^2 d\|T\|x \\ &\leq \int_{\mathbf{C}_{\eta/2}} \text{dist}(x - v, S)^2 d\|T\|x \leq C_{18} \left(\gamma^{-2} E^{1/n} + \eta^{n+4} + \epsilon \right) \left(E + \sigma^{-1}\chi \right). \end{aligned}$$

We next apply the reverse Poincaré inequality Lemma 4.2 to the scaled current

$$\tilde{T} = \mu_{\eta/4\#}^{-1} \tau_w^{-1} \left(\cdot, \#T \llcorner \mathbf{C}_{\eta/4} \right).$$

Since $(\partial\tilde{T}) \llcorner \mathbf{C} = (\partial T) \llcorner \mathbf{C}$ it is easy to see, by a homotopy connecting \cdot to the identity, that \tilde{T} satisfies all the requirements from the definition of the class $\mathcal{T}_b(n, k, 1)$ or $\mathcal{T}_i(n, k, m)$ respectively (including (2.12)) with the exception $0 \in \text{spt } \tilde{T}$ in the latter case. (This condition, however, is not needed in Lemma 4.2 as we have pointed out there.) \tilde{T} is $(\tilde{F}, \tilde{\omega})$ minimizing in \mathbf{C} with $\tilde{\omega}(r) = \omega(\frac{\eta}{4}r)$, structure constants $\tilde{\lambda} = \lambda$, $\tilde{\Lambda} = \Lambda$, and moduli $\tilde{\kappa}(r) = \kappa(\frac{\eta}{4}r)$, $\tilde{\mu}(r) = \mu(\frac{\eta}{4}r)$, $\tilde{\hat{\mu}}(r) \leq \hat{\mu}(\frac{\eta}{4}r)$. The height h of \tilde{T} in \mathbf{C} does not exceed 4, by (5.19), hence $\tilde{\omega}(\sqrt{1+h^2}) \leq \omega(\frac{1}{2})$ and similarly for $\tilde{\kappa}$, $\tilde{\mu}$, $\tilde{\hat{\mu}}$. The conclusion of Lemma 4.2 therefore gives

$$\begin{aligned} \mathbf{E} \left(\cdot, \#T, \frac{1}{20}\eta \right) &\leq C_{12} \left[\chi \left(\frac{\eta}{4} \right)^{-n} \mathbf{M} \left(\cdot, \#T \llcorner \mathbf{C}_{\eta/4} \right) \right. \\ &\quad \left. + (1 + \chi) \left(\frac{\eta}{4} \right)^{-n-2} \int_{\mathbf{C}_{\eta/4}} |\mathbf{q}x - w|^2 d\|\cdot, \#T\|x \right], \end{aligned}$$

and combining this with (5.20) we get

$$(5.21) \quad \mathbf{E} \left(\cdot, \#T, \frac{1}{20}\eta \right) \leq C_{20} \left[\chi \eta^{-n} + \left(\eta^{-n-2} \gamma^{-2} E^{1/n} + \eta^2 + \eta^{-n-2} \epsilon \right) \left(E + \sigma^{-1}\chi \right) \right],$$

where $C_{20} = \text{const}(n, m, \lambda, \Lambda)$ and we have used

$$\mathbf{M} \left(\cdot, \#T \llcorner \mathbf{C}_{\eta/4} \right) \leq \mathbf{M} \left(T \llcorner \mathbf{C}_{\eta/2} \right) \leq E + m\alpha(n).$$

We now specify $\eta = 20\vartheta$ with $0 < \vartheta \leq \frac{1}{340}$, $\epsilon = \eta^{n+2}\vartheta^2$, $\sqrt{\sigma} = \frac{1}{2}C_{17}^{-1}\delta(n, k, \lambda, \Lambda, \epsilon)$, and $\gamma = E^{1/4n}$. If we require

$$(5.22) \quad E^{1/2n} \leq \vartheta^{n+4},$$

then (5.21) gives the desired conclusion (5.14). Moreover, with these specifications of ϵ and γ all the smallness assumptions (5.1), (5.17), and (5.22) are satisfied if we require (5.12) with a sufficiently small constant δ_1 . (5.13) finally follows from (5.3). \blacksquare

We note that in the boundary case $T \in \mathcal{T}_b(n, k, m)$ with (2.13) and multiplicity $m \geq 2$ the proof above does not work, because it is not clear that the tilted current \tilde{T} does again satisfy the density condition (2.13) which we needed for the height estimate. Thus, here is a second difficulty (besides that pointed out after Lemma 5.1) which prevents us from proving boundary regularity in the case of multiplicity $m \geq 2$. It seems that one would need an estimate of the height of \tilde{T} by the distance to $\text{spt } \partial T$ to overcome this difficulty.

We now iterate Lemma 5.2. We consider $0 < \vartheta \leq \frac{1}{340}$ such that $C_{22}\vartheta^2 < \frac{1}{2}\vartheta$. With δ_1 from (5.12) we choose positive numbers $E_* \leq \delta_1$ and $\chi_* \leq \delta_1$. Starting with $T_0 \in \mathcal{T}_b(n, k, 1)$ or $T_0 \in \mathcal{T}_i(n, k, m)$ satisfying (2.12) such that $\text{spt } T_0 \subset \mathbb{R}^n \times \mathbf{B}^k$ and T_0 is (F_0, ω_0) minimizing in \mathbf{C} for some integrand F_0 with structure constants resp. moduli $\lambda_0 = \lambda$, $\Lambda_0 = \Lambda$, $\nu_0 = \nu$, $\kappa_0 \leq \kappa$, $\mu_0 \leq \mu$ and for some nondecreasing $\omega_0 \leq \omega$, where $\kappa_0, \kappa, \mu_0, \mu, \omega_0, \omega$ satisfy the conditions of Section 1, we assume

$$(5.23) \quad \mathbf{E}(T_0, 1) \leq E_*, \quad \chi_0(1) \leq \chi_*.$$

Here we define

$$(5.24) \quad \chi(r) = \begin{cases} \widehat{\kappa}(r) + \omega(r) & \text{in the case } T \in \mathcal{T}_i(n, k, m), \\ \kappa(r)^2 + \widehat{\mu}(r) + \omega(r) & \text{in the case } T_0 \in \mathcal{T}_b(n, k, 1). \end{cases}$$

The function $\chi_0(r)$ is defined similarly using $\kappa_0, \widehat{\kappa}_0, \widehat{\mu}_0$, and ω_0 . We may then apply Lemma 5.2 to find a linear isometry ι_0 of \mathbb{R}^{n+k} , with $\iota_0 = \text{id}$ on $(\mathbb{R}^{n-1} \times \{0\}) \times \{0\}^k$ in the case $T_0 \in \mathcal{T}_b(n, k, 1)$, such that

$$(5.25) \quad \|\iota_0 - \text{id}\| \leq C_{21}\sqrt{\mathbf{E}(T_0, 1) + \chi_0(1)}$$

and

$$(5.26) \quad \mathbf{E}(T_1, 1) \leq \eta\mathbf{E}(T_0, 1) + C_{21}\chi_0(1)$$

hold for $T_1 = (\boldsymbol{\mu}_{\vartheta\#}^{-1}, {}_{0\#}T_0) \llcorner \mathbf{C}$ and $\eta = C_{22}\vartheta^2 < \frac{1}{2}\vartheta$. Moreover, T_1 is (F_1, ω_1) minimizing in \mathbf{C} for an integrand F_1 satisfying the same conditions as F_0 with constants and moduli $\lambda_1 = \lambda$, $\Lambda_1 = \Lambda$, $\nu_1 = \nu$, $\kappa_1(r) \leq \kappa(\vartheta r)$, $\widehat{\kappa}_1(r) \leq \widehat{\kappa}(\vartheta r)$, $\omega_1(r) \leq \omega(\vartheta r)$, and in the boundary case also $\mu_1(r) \leq \mu(\vartheta r)$, $\widehat{\mu}_1(r) \leq \widehat{\mu}(\vartheta r)$, i.e. $\chi_1(r) \leq \chi(\vartheta r)$. Hence, making use of (5.26) we see that (5.23) holds for $T_1, \chi_1(1)$ provided

$$(5.27) \quad C_{21}\chi_* \leq (1 - \eta)E_*.$$

If we impose an additional smallness condition

$$(5.28) \quad C_{23}E_* \leq \vartheta^{2n}$$

with sufficiently large constant $C_{23} = \text{const}(n, k, m, \lambda, \Lambda)$, then the height estimate Lemma 2.2 and (5.25) imply also the height condition $\text{spt } T_1 \subset \mathbb{R}^n \times \mathbf{B}^k$ for T_1 .

Thus, T_1 satisfies the same set of conditions as T_0 and we can proceed to obtain $T_i, \chi_i, \tilde{\nu}_i$ such that

$$(5.29) \quad \begin{aligned} T_{i+1} &= \left(\mu_{\vartheta\#}^{-1}, i\#T_i \right) \llcorner \mathbf{C}, \quad \chi_i(r) \leq \chi(\vartheta^i r), \\ \|\tilde{\nu}_i - \text{id}\| &\leq C_{21} \sqrt{\mathbf{E}(T_i, 1) + \chi_i(1)}, \end{aligned}$$

and

$$(5.30) \quad \mathbf{E}(T_{i+1}, 1) \leq \eta \mathbf{E}(T_i, 1) + C_{21} \chi_i(1).$$

From (5.30) and (5.29) we deduce

$$\mathbf{E}(T_i, 1) \leq \eta^i \mathbf{E}(T_0, 1) + C_{21} \sum_{j=1}^i \eta^{j-1} \chi(\vartheta^{i-j}).$$

To simplify this we use the assumption that $r^{-2\alpha} \kappa(r)^2, r^{-2\alpha} \widehat{\kappa}(r), r^{-2\alpha} \widehat{\mu}(r), r^{-2\alpha} \omega(r)$, hence also $r^{-2\alpha} \chi(r)$, are nonincreasing for some $0 < \alpha < 1$. Therefore $\chi(\vartheta^{i-j}) \leq \vartheta^{-2\alpha j} \chi(\vartheta^i)$ in the last sum and, consequently,

$$(5.31) \quad \mathbf{E}(T_i, 1) \leq \eta^i \mathbf{E}(T_0, 1) + 2C_{21} \vartheta^{-2\alpha} \chi(\vartheta^i),$$

provided we have chosen $\eta = C_{22} \vartheta^2 \leq \frac{1}{2} \vartheta^{2\alpha}$.

We next want to prove that $\tilde{\nu}_i = \tilde{\nu}_{i, i-1} \dots \tilde{\nu}_{1,0}$ converges to some limit isometry $\tilde{\nu}_\infty$. From (5.29) and (5.31) we have

$$\|\tilde{\nu}_i - \tilde{\nu}_{i-1}\| = \|\tilde{\nu}_i - \text{id}\| \leq C_{21} \left[\eta^i \mathbf{E}(T_0, 1) + (2C_{21} \vartheta^{-2\alpha} + 1) \chi(\vartheta^i) \right]^{1/2}.$$

It follows that $\sum_{i=1}^\infty \|\tilde{\nu}_i - \tilde{\nu}_{i-1}\|$ converges, provided $\sum_{i=1}^\infty \sqrt{\chi(\vartheta^i)} < \infty$. In view of

$$(5.32) \quad \frac{1}{\alpha} (1 - \vartheta^\alpha) \sqrt{\chi(\vartheta^i)} = \frac{\sqrt{\chi(\vartheta^i)}}{\vartheta^{\alpha i}} \int_{\vartheta^{i+1}}^{\vartheta^i} r^{\alpha-1} dr \leq \int_{\vartheta^{i+1}}^{\vartheta^i} \frac{1}{r} \sqrt{\chi(r)} dr$$

the convergence of $\sum_{i=1}^\infty \sqrt{\chi(\vartheta^i)}$ is implied by the condition (cf. (1.15))

$$(5.33) \quad \infty > \mathbf{X}(r) = \begin{cases} \widehat{\mathbf{K}}(r) + \Omega(r) & \text{in the case } T \in \mathcal{T}_i(n, k, m), \\ \mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r) & \text{in the case } T_0 \in \mathcal{T}_b(n, k, 1). \end{cases}$$

Assuming (5.33) we obtain a limit $\tilde{\nu}_\infty$ of the $\tilde{\nu}_i$ with the estimate (recalling $\eta \leq \frac{1}{2} \vartheta^{2\alpha} \ll 1$)

$$(5.34) \quad \|\tilde{\nu}_\infty - \tilde{\nu}_{i-1}\| \leq 2C_{21} \left[\sqrt{\eta^i \mathbf{E}(T_0, 1)} + \frac{\alpha}{1 - \vartheta^\alpha} \sqrt{2C_{21} \vartheta^{-2\alpha} + 1} \sqrt{\mathbf{X}(\vartheta^i)} \right],$$

where $i = 0$ gives an estimate for $\|, \infty - \text{id}\|$. Imposing a further smallness condition

$$(5.35) \quad C_{24} \left(\sqrt{E_*} + \sqrt{X_*} \right) \leq 1$$

with a sufficiently large constant $C_{24} = \text{const}(n, k, m, \lambda, \Lambda, \vartheta)$ and assuming $X(1) \leq X_*$, we see from (5.34) and $\text{spt } T_i \subset \mathbb{R}^n \times \mathbf{B}^k$ that

$$\left(, \infty_{\#}, \tilde{\cdot}_{i\#}^{-1} T_{i+1} \right) \llcorner \mathbf{C}_{\vartheta} = \left(\boldsymbol{\mu}_{\vartheta}^{-1}, \infty_{\#}, \tilde{\cdot}_{i-1\#}^{-1} T_i \right) \llcorner \mathbf{C}_{\vartheta}.$$

This implies, by induction,

$$(5.36) \quad \mathbf{E} \left(, \infty_{\#}, \tilde{\cdot}_{i-1\#}^{-1} T_i, \vartheta \right) = \mathbf{E} \left(, \infty_{\#} T_0, \vartheta^{i+1} \right).$$

We now invoke the following simple Lemma:

5.3 Lemma (excess comparison for tilting). *Suppose $T \in \mathcal{T}_*(n, k, m)$ with $\text{spt } T \subset \mathbb{R}^n \times \mathbf{B}^k$ and $,$ is a linear isometry of \mathbb{R}^{n+k} with $, = \text{id}$ on $(\mathbb{R}^n \times \{0\}) \times \{0\}^k$ in the case $T \in \mathcal{T}_b(n, k, m)$. If $\|, - \text{id}\| \leq \frac{1}{5}$, then*

$$\varrho^n \mathbf{E} \left(, \#T, \varrho \right) \leq 2\mathbf{E}(T, 1) + n^2 \|, - \text{id}\|^2 \mathbf{M}(T)$$

holds for $0 < \varrho \leq \frac{2}{3}$.

Proof. The hypotheses imply $(, \cdot^{-1} \mathbf{C}_{\varrho}) \cap \text{spt } T \cap \partial \mathbf{C} = \emptyset$, and with a homotopy argument one sees that $\mathbf{p}_{\#} \left(, \#T \llcorner \mathbf{C}_{\varrho} \right) = (\mathbf{E}^n \llcorner \mathbf{U}_{\varrho}) \llcorner \boldsymbol{\iota}_m$ with the same nonnegative multiplicity function $\boldsymbol{\iota}_m$ as in $\mathbf{p}_{\#} T = (\mathbf{E}^n \llcorner \mathbf{U}) \llcorner \boldsymbol{\iota}_m$. Therefore

$$\begin{aligned} \varrho^n \mathbf{E} \left(, \#T, \varrho \right) &= \frac{1}{2} \int_{\mathbf{C}_{\varrho}} \left| \overrightarrow{\cdot}_{\#T} - \bar{\mathbf{e}} \right|^2 d\|, \#T\| \leq \frac{1}{2} \int_{\mathbf{C}} \left| (\wedge_n, \cdot) \vec{T} - \bar{\mathbf{e}} \right|^2 d\|T\| \\ &\leq \int_{\mathbf{C}} \left| \vec{T} - \bar{\mathbf{e}} \right|^2 d\|T\| + \int_{\mathbf{C}} \left| (\wedge_n, \cdot) \vec{T} - \vec{T} \right|^2 d\|T\| \\ &\leq 2\mathbf{E}(T, 1) + n^2 \|, - \text{id}\|^2 \mathbf{M}(T). \end{aligned}$$

■

Applying this with $, = , \infty_{\cdot, i-1}^{-1}$ and $T = T_i$, and noting that (5.34), (5.35) imply $\|, - \text{id}\| \leq \frac{1}{5}$ if C_{24} is chosen large enough, we deduce

$$\mathbf{E} \left(, \infty_{\#} T_0, \vartheta^{i+1} \right) \leq 2\vartheta^{-n} \mathbf{E}(T_i, 1) + \text{const}(n, m) \left\| , \infty - \cdot_{i-1}^{-1} \right\|^2,$$

and with (5.31), (5.36), (5.34) we obtain, noting that $\chi(\vartheta^i) \leq \text{const}(\vartheta) X(\vartheta^i)$ by (5.32),

$$(5.37) \quad \mathbf{E} \left(, \infty_{\#} T_0, \vartheta^{i+1} \right) \leq C_{25} \left[\eta^i \mathbf{E}(T_0, 1) + X(\vartheta^i) \right].$$

This is the discrete version of the excess decay for T_0 over the tilted n -plane $, \cdot^{-1}(\mathbb{R}^n \times \{0\}^k)$. The following Lemma is an immediate consequence. We recall the definition of X in (5.33) and the assumed existence of $0 < \alpha < 1$ such that $r^{-\alpha} \kappa(r)$, $r^{-\alpha} \mu(r)$, $r^{-2\alpha} \omega(r)$, hence also $r^{-2\alpha} \widehat{\kappa}(r)$, $r^{-2\alpha} \widehat{\mu}(r)$ and $r^{-2\alpha} \chi(r)$, are nonincreasing.

5.4 Lemma (excess decay). For $\alpha \leq \beta < 1$ there exists a positive constant $\epsilon_0 = \epsilon_0(n, k, m, \lambda, \Lambda, \nu(\cdot), \beta)$ such that the following is true: Whenever $T \in \mathcal{T}_b(n, k, 1)$ or $T \in \mathcal{T}_i(n, k, m)$ with (2.12) is (F, ω) minimizing in \mathbf{C} with

$$(5.38) \quad \mathbf{E}(T, 1) + \mathbf{X}(1) \leq \epsilon_0,$$

then there exists a linear isometry ϑ of \mathbb{R}^{n+k} with $\vartheta|_{\mathbb{R}^{n-1} \times \{0\}} = \text{id}$ on $(\mathbb{R}^{n-1} \times \{0\}) \times \{0\}^k$ in the case $T \in \mathcal{T}_b(n, k, 1)$ such that

$$(5.39) \quad \|\vartheta - \text{id}\|^2 \leq C_{26} [\mathbf{E}(T, 1) + \mathbf{X}(1)]$$

and

$$(5.40) \quad \mathbf{E}\left(\vartheta\#(T \llcorner \mathbf{C}_{1/2}), r\right) \leq C_{26} \left[r^{2\beta} \mathbf{E}(T, 1) + \mathbf{X}(r)\right] \quad \text{for } 0 < r \leq \frac{1}{3},$$

$$(5.41) \quad \mathbf{E}(T, r) \leq C_{26} [\mathbf{E}(T, 1) + \mathbf{X}(1)] \quad \text{for } 0 < r \leq 1,$$

where $C_{26} = \text{const}(n, k, m, \lambda, \Lambda, \beta)$.

Proof. We define $T_0 = \mu_{2\#}(T \llcorner \mathbf{C}_{1/2})$ and observe that T_0 is (F_0, ω_0) minimizing with $\lambda_0 = \lambda$, $\Lambda_0 = \Lambda$, $\nu_0 = \nu$, $\chi_0 \leq \chi(\frac{1}{2}r)$, for χ defined in (5.24) and χ_0 analogously. Clearly, we may replace α by any larger exponent in $]0, 1[$ and hence assume $\alpha = \beta$. We define ϑ depending on n, m, λ, Λ , and α by $\vartheta = \min\left\{(2C_{22})^{-1/(2-2\alpha)}, \frac{1}{340}\right\}$ so that $C_{22} \leq \frac{1}{2}\vartheta^{2\alpha-2}$ and $\eta = C_{22}\vartheta^2 \leq \frac{1}{2}\vartheta^{2\alpha}$. Choosing ϵ_0 small enough we see that all our previous smallness conditions (5.23), (5.27), (5.28), and (5.35) are satisfied, and $\text{spt } T_0 \subset \mathbb{R}^n \times \mathbf{B}^k$ holds by Lemma 2.2. We let ϑ be the limit isometry ϑ_∞ produced above. Then (5.39) follows from (5.34) and $\mathbf{E}(T_0, 1) \leq 2^n \mathbf{E}(T, 1)$.

For $\vartheta^{i+2} < r \leq \vartheta^{i+1}$ with $i \geq 0$ we obtain from (5.37)

$$\begin{aligned} \mathbf{E}\left(\vartheta\#T_0, r\right) &\leq \vartheta^{-n} \mathbf{E}\left(\vartheta\#T_0, \vartheta^{i+1}\right) \leq \vartheta^{-n} C_{25} \left[\eta^i \mathbf{E}(T_0, 1) + \mathbf{X}(\vartheta^i)\right] \\ &\leq \vartheta^{-n} C_{25} \left[\vartheta^{-4\alpha} r^{2\alpha} \mathbf{E}(T_0, 1) + \mathbf{X}(\vartheta^{-2}r)\right]. \end{aligned}$$

This implies (5.40) for $0 < r \leq \frac{1}{2}\vartheta$, because we have $\mathbf{X}(t\rho) \leq t^{2\alpha}\mathbf{X}(\rho)$ for $t \geq 1$ and $\mathbf{E}\left(\vartheta\#(T \llcorner \mathbf{C}_{1/2}), \frac{1}{2}r\right) = \mathbf{E}\left(\vartheta\#T_0, r\right)$. For $\frac{1}{2}\vartheta < r \leq \frac{1}{3}$ we deduce (5.40) from (5.39) with Lemma 5.3, decreasing ϵ_0 if necessary.

With regard to (5.41) we note that by Lemma 2.2, (5.38), (5.39), and (5.40) we have the following, provided ϵ_0 is chosen sufficiently small:

$$\begin{aligned} \tilde{T}_r &= \left[\mu_{r\#}^{-1}, \#(T \llcorner \mathbf{C}_{1/2})\right] \llcorner \mathbf{C} \in \mathcal{T}_*(n, k, m), \\ \sup\left\{|\mathbf{q}x| : x \in \text{spt } \tilde{T}_r\right\} &\leq 1, \\ \|\tilde{T}_r - \text{id}\|^2 &\leq C_{26} [\mathbf{E}(T, 1) + \mathbf{X}(1)] \leq \frac{1}{25}, \\ \mathbf{E}(\tilde{T}_r, 1) &\leq C_{26} [\mathbf{E}(T, 1) + \mathbf{X}(1)], \\ (\vartheta\#^{-1}\tilde{T}_r) \llcorner \mathbf{C}_{1/2} &= (\mu_{r\#}^{-1}T) \llcorner \mathbf{C}_{1/2}, \end{aligned}$$

for $0 < r \leq \frac{1}{8}$. With Lemma 5.3 and $\mathbf{M}(\tilde{T}_r) \leq \mathbf{E}(\tilde{T}_r, 1) + m\alpha(n)$ we conclude

$$\begin{aligned} \mathbf{E}\left(T, \frac{r}{2}\right) &= \mathbf{E}\left(\mu_{r\#}^{-1}T, \frac{1}{2}\right) = \mathbf{E}\left(\cdot, \frac{1}{\#}\tilde{T}_r, \frac{1}{2}\right) \\ &\leq \text{const}(n, m) \left[\mathbf{E}(\tilde{T}_r, 1) + \|\cdot, \cdot^{-1} - \text{id}\|^2 \right] \\ &\leq 2\text{const}(n, m)C_{26} [\mathbf{E}(T, 1) + \mathbf{X}(1)] . \end{aligned}$$

Increasing C_{26} suitably we have therefore proved (5.41) for $0 < r \leq \frac{1}{16}$, and for $\frac{1}{16} \leq r \leq 1$ it holds trivially, by the scaling property of the excess, provided $C_{26} \geq 16^n$. \blacksquare

6 The ϵ -regularity theorem

In this section we use the excess decay (Lemma 5.4) to prove the ϵ -regularity theorem: An (F, ω) minimizing current T with sufficiently small excess in arbitrarily small balls centered at $a \in \text{spt } T$ is, locally at a , represented by a C^1 submanifold (with boundary $\text{spt } \partial T$ if ∂T is, locally at a , represented by a smooth submanifold). It is well known how to deduce this result from the excess decay (e.g. [All2], [Ha1]), but we give a complete proof, in order to control the dependence of constants and to derive an explicit modulus of continuity for the first order derivatives.

We assume here all the structure conditions from Section 1 on the integrand F and the function ω in the condition of (F, ω) minimality. In particular, $\lambda > 0$ is the ellipticity constant of F , Λ is a bound for $F(x, \xi)$ and its derivatives up to order 2 with respect to the tangent plane variable ξ , $\nu(\cdot)$ is a modulus of continuity for this second derivative, κ, μ are moduli for $F(x, \xi)$ and $D_{(2)}F(x, \xi)$ with respect to the space variable x as in (1.4)–(1.8), $\hat{\kappa}, \hat{\mu}$ are associated with κ, μ by (1.9), and $\mathbf{K}, \hat{\mathbf{K}}, \hat{\mathbf{M}}, \Omega$ are defined in (1.15). The exponent $0 < \alpha < 1$ is such that $r^{-\alpha}\kappa(r), r^{-\alpha}\mu(r), r^{-2\alpha}\omega(r)$ are nonincreasing functions of $r > 0$. Recall also the density condition (2.12), i.e. $\Theta^n(\|T\|, x) \geq m$ for $\|T\|$ almost all x , in the case $T \in \mathcal{T}_i(n, k, m)$ with $m \geq 2$. In the boundary situation $T \in \mathcal{T}_b(n, k, m)$ we must assume $m = 1$ for reasons explained in section 5. $\mathbf{E}(T, 1)$ is the cylindrical excess of T in the cylinder $\mathbf{C} = \mathbf{U}^n \times \mathbb{R}^k \subset \mathbb{R}^{n+k}$.

6.1 Theorem (interior and boundary ϵ -regularity theorem). *Corresponding to $\alpha \leq \beta < 1$ there exists a positive constant $\epsilon(n, k, m, \lambda, \Lambda, \nu(\cdot), \beta)$ such that the following is true: If $T \in \mathcal{T}_i(n, k, m)$ with (2.12) or $T \in \mathcal{T}_b(n, k, m)$ with $m = 1$ is (F, ω) minimizing in \mathbf{C} with*

$$(6.1) \quad \mathbf{E}(T, 1) + \hat{\mathbf{K}}(1) + \Omega(1) \leq \epsilon ,$$

then $T \llcorner \mathbf{C}_{1/34}$ is represented by the graph, taken with multiplicity m , of a continuously differentiable function g which is defined over the n -disk $\mathbf{U}_{1/34}$ in the case $T \in \mathcal{T}_i(n, k, m)$ and over the half n -disk $\mathbf{B}_+ \cap \mathbf{U}_{1/34}$ in the case $T \in \mathcal{T}_b(n, k, m)$, $m = 1$. Moreover, the derivative Dg has the following modulus of continuity:

$$(6.2) \quad |Dg(a) - Dg(b)| \leq C_{33} \left[|a - b|^{2\beta} \mathbf{E}(T, 1) + \hat{\mathbf{K}}(|a - b|) + \Omega(|a - b|) \right]^{1/2}$$

for a, b in the domain of g , where $C_{33} = \text{const}(n, k, m, \lambda, \Lambda, \beta)$.

Since one can always flatten the boundary by a diffeomorphism and achieve smallness of $\widehat{\mathbf{K}}(1)$ and $\Omega(1)$ by scaling, Theorem 6.1 can be applied to (F, ω) minimizing rectifiable n -currents T in \mathbb{R}^{n+k} , with F sufficiently regular, at all points $a \in \text{spt } T$ where $\mathbf{E}(T, a, \varrho_i) < \frac{1}{2}\epsilon$ for a sequence $\varrho_i \downarrow 0$ and where, in the case $a \in \text{spt } \partial T$, ∂T is represented locally by an oriented $C^{1,1}$ submanifold of dimension $n-1$ in \mathbb{R}^{n+k} an $\mathbf{p}_\# T$ has multiplicity 0 and 1 on the two sides of the projection of $\text{spt } \partial T$ near a . In this situation the theorem gives interior and boundary $C^{1,\beta}$ regularity of T locally at 0 for all $0 < \beta < 1$ when $\Omega \equiv 0$ (i.e. T is F minimizing), and regularity with continuity modulus $\text{const } \Omega$ for the derivative when $\Omega \not\equiv 0$ (which is optimal in the case $\omega(r) = \text{const } r^\beta$, cf. Section 1). The radius $\frac{1}{17}$ of the domain of g can, of course, be replaced by any radius $0 < r < 1$ (if one admits dependence of ϵ and C_{33} on r). This is easily seen from the scaling properties of the excess and the functions $\widehat{\mathbf{K}}(r)$, $\Omega(r)$.

Proof. We require $\epsilon < \epsilon_0$ with $\epsilon_0(n, k, m, \lambda, \Lambda, \nu(\cdot), \beta)$ from Lemma 5.4, and we will impose further smallness conditions on ϵ in the course of the proof.

Step 1. We prove that the excess of T is uniformly small in all cylinders $\mathbf{C}_r(y) \subset \mathbf{C}$ with $y \in \mathbf{B}_{1/2}$ and in the boundary case additionally $y \in \mathbf{B}_+$.

In the interior situation $T \in \mathcal{T}_i(n, k, m)$ we observe $\mathbf{E}(T, y, r_0) \leq 2^n \mathbf{E}(T, 0, 1) \leq 2^n \epsilon$ for $r_0 = 1 - |y|$, hence choosing $\epsilon \leq 2^{-n-1} \epsilon_0$ we see that the scaled current $(\mu_{r_0\#}^{-1} \tau_{x\#}^{-1} T) \llcorner \mathbf{C}$ with $x \in \mathbf{p}^{-1}\{y\} \cap \text{spt } T$ satisfies the hypotheses of Lemma 5.4. From (5.41) we therefore get a bound $\text{const} \cdot \epsilon$ on $\mathbf{E}(T, y, r)$ for all $0 < r \leq r_0$, and decreasing ϵ we achieve

$$(6.3) \quad \mathbf{E}(T, y, r) + \widehat{\mathbf{K}}(r) + \Omega(r) \leq \text{const}(n, k, m, \lambda, \Lambda, \beta) \epsilon \leq \epsilon_0$$

for all $y \in \mathbf{B}_{1/2}$ and $0 < r \leq 1 - |y|$.

In the boundary situation we apply Lemma 5.4 similarly, but estimate $\mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) \leq 5\widehat{\mathbf{K}}(r)$ for $0 < r \leq \frac{1}{2}$, which follows from $\frac{1}{4}\kappa(\varrho)^2 \leq \widehat{\kappa}(\varrho)$ and $\widehat{\mu}(\varrho) \leq \widehat{\kappa}(\varrho)$ for $0 < \varrho \leq \frac{1}{2}$. We then obtain (6.3) for $T \in \mathcal{T}_b(n, k, 1)$ and centers $y \in \mathbf{B}_{1/2}^{n-1} \times \{0\}$. If $y \in \mathbf{U}_+ \cap \mathbf{B}_{1/2}$ then we consider $y' = (y_1, \dots, y_{n-1}, 0) \in \mathbf{B}_{1/2}^{n-1} \times \{0\}$ and $r_0 = y_n \in]0, \frac{1}{2}]$. In the case $r_0 \geq \frac{1}{4}$ we deduce (6.3) for $0 < r \leq 1 - |y|$ as above. In the case $r_0 < \frac{1}{4}$ we have $\mathbf{U}_r(y) \subset \mathbf{U}_{2r}(y') \subset \mathbf{U}$ for $r_0 \leq r \leq \frac{1}{4}$, and using the smallness of $\mathbf{E}(T, y', 2r)$ we deduce again a corresponding estimate for $\mathbf{E}(T, y, r)$, $r_0 \leq r \leq \frac{1}{4}$. For $0 < r < r_0$ a similar estimate follows from the smallness of $\mathbf{E}(T, y, r_0)$ by the scaling argument above and Lemma 5.4.

Replacing T by $\mu_{2\#}(T \llcorner \mathbf{C}_{1/2})$ and choosing ϵ small enough we may therefore assume that (6.3) holds for all $y \in \mathbf{U}$ and $0 < r \leq 1 - |y|$, where $y_n \geq 0$ is required additionally in the boundary case. (In Theorem 6.1 we have stated the conclusion for the radius $\frac{1}{34}$ instead of $\varrho = \frac{1}{17}$, because we have expanded T by a factor 2.) This means that Lemma 5.4 can be applied to all the rescaled currents $(\mu_{r\#}^{-1} \tau_{x\#}^{-1} T) \llcorner \mathbf{C}$ with $x \in \mathbf{p}^{-1}\{y\} \cap \text{spt } T$, $0 < r \leq 1 - |y|$, and in the boundary case additionally $x = (y, 0)$ or $\mathbf{U}_r(y) \subset \mathbf{U}_+$. By Lemma 2.2 we may also assume that these currents have height ≤ 1 , i.e. support in $\mathbb{R}^n \times \mathbf{B}^k$.

Step 2. We prove that $\mathbf{C}_{1/17} \cap \text{spt } T$ is the graph of a Lipschitz function defined on $\mathbf{U}_{1/17}$ in the case $T \in \mathcal{T}_i(n, k, m)$ and on $\mathbf{B}_+ \cap \mathbf{U}_{1/17}$ in the case $T \in \mathcal{T}_b(n, k, 1)$. Moreover, $T \llcorner \mathbf{C}_{1/17}$ is represented by the graph of g with appropriate orientation and multiplicity.

For this we recall the definition of the *good points* y in the proof of the Lipschitz approximation Lemma 3.1 (see (3.7)): $y \in \mathbf{U}_\varrho$ (and $y_n \neq 0$ in the boundary case) with $\mathbf{E}(T, y, r) \leq \eta$ for $0 < r < 1 - \varrho$, where $\eta^{-1} = \text{const}(n, \Lambda)$ (since we may assume $\omega \leq 1$

here). Comparing with (6.3) we see that the bad set is empty, if ϵ is chosen sufficiently small. This means that $\mathbf{C}_\varrho \cap \text{spt } T$ in the interior case and $\mathbf{C}_+ \cap \mathbf{C}_\varrho \cap \text{spt } T$ in the boundary case equals graph g , where g denotes here restriction of the Lipschitz approximation to \mathbf{U}_ϱ or to $\mathbf{U}_+ \cap \mathbf{U}_\varrho$ respectively. According to Lemma 3.1 we have $\text{Lip } g \leq 1$, $g(0) = 0$ in the interior case, g has zero boundary values on $\mathbf{U}_\varrho^{n-1} \times \{0\}$ in the boundary case, and we may take $\varrho = \frac{1}{17}$. For the current T_g representing graph g with appropriate orientation and multiplicity we have $\mathbf{p}_\# T_g = (\mathbf{p}_\# T) \llcorner \mathbf{U}_\varrho$. Then $T \llcorner \mathbf{C}_\varrho = T_g$ or $T \llcorner (\mathbf{C}_+ \cap \mathbf{C}_\varrho) = T_g$ follows, because $y \mapsto (y, g(y))$ is biLipschitz.

In the boundary case $T \in \mathcal{T}_b(n, k, 1)$ we must also prove $(\mathbf{C}_\varrho \setminus \mathbf{C}_+) \cap \text{spt } T \subset (\mathbf{U}^{n-1} \times \{0\}) \times \{0\}^k$. Suppose $x = (y, z) \in \mathbf{C}_\varrho \cap \text{spt } T$ such that $y_n < 0$ or $y_n = 0$ and $z \neq 0$. Choose $r = |y_n|$ if $y_n < 0$ or $r = \frac{1}{\sqrt{2}}|z|$ else. On account of $\text{Lip } g \leq 1$ we then have $\mathbf{U}(x, r) \cap \text{graph } g = \emptyset$ and $\mathbf{E}(T, y', 2r) \geq 2^{-n} \mathbf{E}(T, y, r) \geq (2r)^{-n} \mathbf{M}(T \llcorner \mathbf{U}(x, r))$, because $\mathbf{p}_\# T_g = \mathbf{p}_\# T$ and hence $\mathbf{p}_\#(T \llcorner \mathbf{U}(x, r)) = 0$. But from Lemma 2.1 we have a lower bound $\tilde{C}_1 \mathbf{M}(T \llcorner \mathbf{U}(x, r)) \geq r^n$ with $\tilde{C}_1 = \text{const}(n, \Lambda)$, hence $\mathbf{E}(T, y', 2r) \geq 2^{-n} \tilde{C}_1^{-1}$ follows. On the other hand, by choosing ϵ small enough we can achieve $\mathbf{E}(T, y', 2r) < 2^{-n} \tilde{C}_1^{-1}$ in step 1, and with this contradiction step 2 is finished.

For the next step in the regularity proof we need

6.2 Lemma. *Suppose $g, h: \mathbb{R}^n \rightarrow \mathbb{R}^k$ are Lipschitz mappings and differentiable at $y \in \mathbb{R}^n$. Let $\vec{G}(y), \vec{H}(y)$ be the orientation vectors of graph g and graph h . Then*

$$(6.4) \quad |Dg(y) - Dh(y)|^2 \leq 4 \left[1 + (\text{Lip } g)^2\right]^n \left[1 + (\text{Lip } h)^2\right]^n \left|\vec{G}(y) - \vec{H}(y)\right|^2.$$

Proof. We have $\vec{G}(y) = \frac{\xi}{|\xi|}$ for $\xi = \vec{e} + \sum_{i=1}^n \sum_{j=1}^k \xi_i^j \vec{e}_j^i + \vec{r}$, where $\xi_i^j = \partial_i g^j(y)$, $\vec{e}_j^i = e_1 \wedge \dots \wedge e_{i-1} \wedge e_{n+j} \wedge e_{i+1} \wedge \dots \wedge e_n$ in terms of the canonical basis e_1, \dots, e_{n+k} of \mathbb{R}^{n+k} , and $\vec{r} \in \wedge_n \mathbb{R}^{n+k}$ is orthogonal to \vec{e} and all the \vec{e}_j^i . Similarly $\vec{H}(y) = \frac{\zeta}{|\zeta|}$ with $\zeta = \vec{e} + \sum_{i=1}^n \sum_{j=1}^k \zeta_i^j \vec{e}_j^i + \vec{s}$. The claim then follows from

$$|\xi - \zeta|^2 \geq |(\xi_j^i) - (\zeta_j^i)|^2 = |Dg(y) - Dh(y)|^2$$

and

$$\begin{aligned} |\xi - \zeta| &\leq |\xi| \left| \frac{\xi}{|\xi|} - \frac{\zeta}{|\zeta|} \right| + |\xi| |\zeta| \left| \frac{1}{|\xi|} - \frac{1}{|\zeta|} \right| \\ &\leq |\xi| (1 + |\zeta|) \left| \frac{\xi}{|\xi|} - \frac{\zeta}{|\zeta|} \right| \leq 2|\xi| |\zeta| \left| \frac{\xi}{|\xi|} - \frac{\zeta}{|\zeta|} \right|, \end{aligned}$$

where we have used $\xi \cdot \vec{e} = 1 = \zeta \cdot \vec{e}$, $|\zeta| \geq 1$, and $|\xi|^2 \leq [1 + (\text{Lip } g)^2]^n$ resp. $|\zeta|^2 \leq [1 + (\text{Lip } h)^2]^n$ in the last inequality. \blacksquare

We continue with the proof of the ϵ -regularity-theorem:

Step 3 (Interior regularity). Here we treat the case $T \in \mathcal{T}_i(n, k, m)$ and define $D^*g(y)$ as the Lebesgue value of Dg at $y \in \mathbf{U}_\varrho$ whenever it exists. We consider $a, b \in \mathbf{U}_\varrho$ (with $\varrho = \frac{1}{17}$) and let $T_a = \tau_{(a, g(a))\#}^{-1} T$. By (6.3) we can apply Lemma 5.4 to $\mu_{r_0\#} T_a \llcorner \mathbf{C}$, where $0 < r_0 \leq 1 - |a|$, and we obtain a linear isometry \cdot_a of \mathbb{R}^{n+k} such that, by (5.40),

$$(6.5) \quad \mathbf{E}(\cdot_a T_a, 0, r) \leq C_{26} \left[\left(\frac{r}{r_0}\right)^{2\beta} \mathbf{E}(T, a, r_0) + \hat{\mathbf{K}}(r) + \Omega(r) \right]$$

for $0 < r \leq \frac{1}{3}r_0$. (Note that we can write $,_{a\#}T_a$ instead of $,_{a\#}(T_a \llcorner \mathbf{C}_{r_0/2})$ in (6.5), because $\|,_{a} - \text{id}\|$ is small and T_a has height $\leq r_0$ in \mathbf{C}_{r_0} , cf. the end of step 1.) Representing $,_{a}^{-1}(\mathbb{R}^n \times \{0\}^k)$ as the graph of a linear function $L_a \in \text{Hom}(\mathbb{R}^n, \mathbb{R}^k)$, recalling $\text{Lip } g \leq 1$, and observing $\|L_a\| \leq 1$ on account of (5.39) and (6.1) (with ϵ small enough), we deduce from Lemma 6.2, for $0 < r \leq r_0$,

$$\begin{aligned}
& \int_{\mathbf{U}_{r/2(a)} \cap \mathbf{U}_\varrho} |Dg(y) - L_a|^2 d\mathcal{L}^n y \leq 2^{2n+2} \int_{\mathbf{U}_{r/2(a)} \cap \mathbf{U}_\varrho} \left| \vec{T}(y, g(y)) - (\wedge_n,_{a}^{-1})\vec{\epsilon} \right|^2 d\mathcal{L}^n y \\
(6.6) \quad & \leq 2^{2n+2} \int_{\mathbf{C}_{r/2(a)}} \left| \vec{T}(x) - (\wedge_n,_{a}^{-1})\vec{\epsilon} \right|^2 d\|T\|x \leq 2^{2n+2} \int_{\mathbf{C}_r} \left| \overrightarrow{,_{a\#}T_a} - \vec{\epsilon} \right|^2 d\|,_{a\#}T_a\|x \\
& = 2^{2n+3} r^n \mathbf{E}(,_{a\#}T_a, 0, r).
\end{aligned}$$

(We have used here $,_a(\mathbf{C}_{r/2} \cap \text{spt } T_a) \subset \mathbf{C}_r$ which is true, because T_a has height $\leq r$ in \mathbf{C}_r and $\|,_{a} - \text{id}\|$ is small.) Letting $r \downarrow 0$ we conclude with (6.5) that $D^*g(a) = L_a$ and

$$(6.7) \quad (\wedge_n,_{a}^{-1})\vec{\epsilon} = \vec{T}^*(a, g(a)),$$

the Lebesgue value of \vec{T} at $(a, g(a))$ with respect to the measure $\|T\| = m\mathcal{H}^n \llcorner \text{graph } g$.

From (6.5), (6.6) and the analogous inequality for $L_b = D^*g(b)$ with $r = 2|b - a| \leq \frac{4}{17}$ we also deduce

$$\begin{aligned}
& |D^*g(b) - D^*g(a)|^2 \leq \frac{2^{n+1}}{\alpha(n)|b - a|^n} \int_{\mathbf{U}_{r/2(a)} \cap \mathbf{U}_{r/2(b)} \cap \mathbf{U}_\varrho} |L_a - L_b|^2 d\mathcal{L}^n \\
(6.8) \quad & \leq \frac{2^{2n+2}}{\alpha(n)r^n} \left[\int_{\mathbf{U}_{r/2(a)} \cap \mathbf{U}_\varrho} |Dg(y) - L_a|^2 d\mathcal{L}^n y + \int_{\mathbf{U}_{r/2(b)} \cap \mathbf{U}_\varrho} |Dg(y) - L_b|^2 d\mathcal{L}^n y \right] \\
& \leq C_{27} \left[r^{2\beta} \mathbf{E}(T, 0, 1) + \widehat{\mathbf{K}}(r) + \Omega(r) \right]
\end{aligned}$$

with $C_{27} = \text{const}(n, k, m, \lambda, \Lambda, \nu(\cdot), \beta)$. Since $r = 2|b - a|$ and $\widehat{\mathbf{K}}(2r) + \Omega(2r) \leq 2^{2\alpha}(\widehat{\mathbf{K}}(r) + \Omega(r))$ the continuity modulus (6.2) for Dg is proved.

Step 4 (boundary regularity). Here we consider $T \in \mathcal{T}_b(n, k, 1)$ and first distinguish several special cases for $a, b \in \mathbf{B}_+ \cap \mathbf{U}_\varrho$ ($\varrho = \frac{1}{17}$).

The **special case 1** is $a, b \in \mathbf{U}_\varrho^{n-1} \times \{0\}$. In this case we can proceed exactly as in Step 3, using the boundary version of Lemma 5.4 and $\mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) \leq 5\widehat{\mathbf{K}}(r)$. The result is that the Lebesgue values of Dg at a, b exist (defined here with half balls $\mathbf{U}_+ \cap \mathbf{U}_r(a)$, of course, instead of balls $\mathbf{U}_r(a)$) and satisfy

$$(6.9) \quad |D^*g(b) - D^*g(a)|^2 \leq C_{27} \left[|b - a|^{2\beta} \mathbf{E}(T, 1) + \widehat{\mathbf{K}}(|b - a|) + \Omega(|b - a|) \right],$$

if we redefine C_{27} suitably.

The **special case 2** is $a \in \mathbf{U}_\varrho^{n-1} \times \{0\}$ and $b \in \mathbf{U}_\varrho$ with $b_n > 0$ and $|a - b| \leq 2b_n$. We define $T_a, ,_a$ as in step 3 and have (6.5) again for $r_0 \leq 1 - |a|$ and $0 < r < \frac{1}{3}r_0$. We next want to apply the interior regularity result from step 3 to $,_{a\#}T$ in $\mathbf{C}_{r_1}(\widehat{b})$ where

, $a(b, g(b)) = (\widehat{b}, \widehat{c})$ and $r_1 = \widehat{b}_n$. Recalling that $, a = \text{id}$ and $g = 0$ on $\mathbf{U}_\rho^{n-1} \times \{0\}$, and $\text{Lip } g \leq 1$, we see

$$|b_n - r_1| = |b - \widehat{b}| \leq |(b, g(b)) - , a(b, g(b))| \leq \sqrt{2} \| , a - \text{id} \| b_n.$$

Since $\| , a - \text{id} \|$ is small we conclude $\frac{4}{5}b_n \leq r_1 \leq \frac{6}{5}b_n$ and $\mathbf{U}_{r_1}(\widehat{b}) \subset \mathbf{U}_+ \cap \mathbf{U}_{4b_n}(a)$, hence

$$(6.10) \quad \begin{aligned} \mathbf{E} \left(, a_{\#}T, \widehat{b}, r_1 \right) &\leq 5^n \mathbf{E} \left(, a_{\#}T, a, 4b_n \right) \\ &\leq 5^n C_{26} \left[\left(\frac{17}{4}b_n \right)^{2\beta} \mathbf{E} \left(T, a, \frac{16}{17} \right) + \widehat{\mathbf{K}}(4b_n) + \Omega(4b_n) \right], \end{aligned}$$

by (6.5) applied with $r_0 = \frac{16}{17}$ and $r = 4b_n \leq \frac{4}{17}$. (From (6.3) we could not obtain a bound on $\mathbf{E} \left(, a_{\#}T, \widehat{b}, r_1 \right)$ which is small for small $b_n > 0$ as in (6.10). This is the reason why we must apply the interior excess growth lemma to $, a_{\#}T$ instead of T .)

Now, as in case 1 and step 3 the Lebesgue values $D^*g(a)$, $D^*g(b)$ exist with

$$(6.11) \quad \begin{aligned} r^{-n} \int_{\mathbf{U}_\rho^+ \cap \mathbf{U}_{r/2}(a)} |Dg(y) - D^*g(a)|^2 d\mathcal{L}^n y \\ \leq 2^{2n+3} C_{26} \left[\left(\frac{r}{r_0} \right)^{2\beta} \mathbf{E} \left(T, a, r_0 \right) + \widehat{\mathbf{K}}(r) + \Omega(r) \right] \end{aligned}$$

for $0 < r \leq \frac{1}{3}r_0 \leq \frac{1}{3}(1 - |a|)$, by the analogs of (6.5) and (6.6), and

$$(6.12) \quad \begin{aligned} \int_{\mathbf{U}_{r/3}(b) \cap \mathbf{U}_\rho} |Dg(y) - D^*g(b)|^2 d\mathcal{L}^n y \\ \leq 2^{2n+2} \int_{\mathbf{C}_{r/3}(b)} \left| \vec{T}(x) - \vec{T}^*(b, g(b)) \right|^2 d\|T\|x \\ \leq 2^{2n+2} \int_{\mathbf{C}_{r/2}(\widehat{b})} \left| \overrightarrow{ , a_{\#}T}(x) - \overrightarrow{ , a_{\#}T}(\widehat{b}, \widehat{c}) \right|^2 d\| , a_{\#}T\|x \end{aligned}$$

for $0 < r \leq b_n$, by the analogs of (6.6), (6.7) at b and the fact that $\| , a - \text{id} \|$ is small, $\text{spt } T$ is the graph a function vanishing on $\mathbf{U}_\rho^{n-1} \times \{0\}$ with Lipschitz constant ≤ 1 , and $, a(\mathbf{C}_{r/3}(b) \cap \text{spt } T) \subset \mathbf{C}_{r/2}(\widehat{b}) \cap , a \text{spt } T$.

Decreasing ϵ if necessary we see from (6.10) that $, a_{\#}T$, after translating $(\widehat{b}, \widehat{c})$ to the origin and applying the homothety $\boldsymbol{\mu}_{r_1}^{-1}$, satisfies the hypotheses of Lemma 5.4. The analogs of (6.6), (6.7), (6.5) for $, a_{\#}T$, \widehat{b} , $r \leq r_1$ instead of T , a , $r \leq r_0$ then give

$$(6.13) \quad \begin{aligned} 2^{2n+2} \int_{\mathbf{C}_{r/2}(\widehat{b})} \left| \overrightarrow{ , a_{\#}T}(x) - \overrightarrow{ , a_{\#}T}(\widehat{b}, \widehat{c}) \right|^2 d\| , a_{\#}T\|x \\ \leq C_{28} r^n \left[\left(\frac{r}{r_1} \right)^{2\beta} \mathbf{E} \left(, a_{\#}T, \widehat{b}, r_1 \right) + \widehat{\mathbf{K}}(r) + \Omega(r) \right], \end{aligned}$$

with $C_{28} = \text{const}(n, k, m, \lambda, \Lambda, \beta)$. Combinig this with (6.12) we have, for $0 < r \leq \frac{1}{3}r_1$,

$$(6.14) \quad \begin{aligned} r^{-n} \int_{\mathbf{U}_{r/3}(b) \cap \mathbf{U}_\rho} |Dg(y) - D^*g(b)|^2 d\mathcal{L}^n y \\ \leq C_{28} \left[\left(\frac{r}{r_1} \right)^{2\beta} \mathbf{E} \left(, a_{\#}T, \widehat{b}, r_1 \right) + \widehat{\mathbf{K}}(r) + \Omega(r) \right]. \end{aligned}$$

We now choose $r = \frac{1}{3}r_1$ in (6.14), $r = 2|b - a|$ and $r_0 = \frac{16}{17}$ in (6.11) and recall $\frac{2}{5}|b - a| \leq \frac{4}{5}b_n \leq r_1 = \widehat{b}_n \leq \frac{6}{5}b_n \leq \frac{6}{5}|b - a|$, $\Omega(t\varrho) \leq t^{2\alpha}\Omega(\varrho)$ and $\widehat{\mathbf{K}}(t\varrho) \leq t^{2\alpha}\widehat{\mathbf{K}}(\varrho)$ for $t \geq 1$, to deduce

$$\begin{aligned} & \int_{\mathbf{U}_{r_1/9}(b) \cap \mathbf{U}_\varrho} |Dg(y) - D^*g(b)|^2 d\mathcal{L}^n y + \int_{\mathbf{U}_\varrho^+ \cap \mathbf{U}_{|b-a|(a)}} |Dg(y) - D^*g(a)|^2 d\mathcal{L}^n y \\ & \leq |b - a|^n C_{29} \left[|b - a|^{2\beta} \mathbf{E}(T, 0, 1) + \widehat{\mathbf{K}}(|b - a|) + \Omega(|b - a|) \right], \end{aligned}$$

where $C_{29} = \text{const}(n, k, m, \lambda, \Lambda, \beta)$. Since $\mathbf{U}_{r_1/9}(b) \cap \mathbf{U}_\varrho^+ \cap \mathbf{U}_{|b-a|(a)}$ contains a half-ball of radius $\frac{1}{18}r_1 \geq \frac{1}{45}|b - a|$ we conclude as in (6.8)

$$(6.15) \quad |D^*g(b) - D^*g(a)|^2 \leq C_{30} \left[|b - a|^{2\beta} \mathbf{E}(T, 0, 1) + \widehat{\mathbf{K}}(|b - a|) + \Omega(|b - a|) \right]$$

with $C_{30} = \text{const}(n, k, m, \lambda, \Lambda, \beta)$.

The **special case 3** is $b, \tilde{b} \in \mathbf{U}_\varrho$ with $0 < b_n \leq \tilde{b}_n$ and $|b - \tilde{b}| \leq \frac{2}{15}b_n$. In this case we choose $a \in \mathbf{U}_\varrho^{n-1} \times \{0\}$ such that $|a - b| \leq 2b_n$ and $|a - \tilde{b}| \leq 2\tilde{b}_n$. From (6.14) with $r = 2|b - \tilde{b}|$ and from (6.10) we obtain, recalling $r_1 \geq \frac{4}{5}b_n$,

$$\begin{aligned} & |b - \tilde{b}|^{-n} \int_{\mathbf{U}_{2|b-\tilde{b}|/3}(b) \cap \mathbf{U}_\varrho} |Dg(y) - D^*g(b)|^2 d\mathcal{L}^n y \leq C_{31} \left[|b - \tilde{b}|^{2\beta} \mathbf{E}(T, 0, 1) \right. \\ & \quad \left. + \widehat{\mathbf{K}}(2|b - \tilde{b}|) + \Omega(2|b - \tilde{b}|) + \left| \frac{b - \tilde{b}}{b_n} \right|^{2\beta} \widehat{\mathbf{K}}(4b_n) + \left| \frac{b - \tilde{b}}{b_n} \right|^{2\beta} \Omega(4b_n) \right] \end{aligned}$$

with $C_{31} = \text{const}(n, k, m, \lambda, \Lambda, \beta)$, and the same estimate is valid with b, \tilde{b} interchanged. On account of $\beta \geq \alpha$ the exponent 2β can be replaced by 2α in the last two summands, and from $\widehat{\mathbf{K}}(t\varrho) + \Omega(t\varrho) \leq t^{2\alpha}(\widehat{\mathbf{K}}(\varrho) + \Omega(\varrho))$ for $t \geq 1$ and the fact that $\mathbf{U}_{2|b-\tilde{b}|/3}(b) \cap \mathbf{U}_{2|b-\tilde{b}|/3}(\tilde{b}) \cap \mathbf{U}_\varrho$ contains a half-ball of radius $\frac{1}{6}|b - \tilde{b}|$ we infer

$$(6.16) \quad |D^*g(b) - D^*g(\tilde{b})|^2 \leq C_{32} \left[|b - \tilde{b}|^{2\beta} \mathbf{E}(T, 0, 1) + \widehat{\mathbf{K}}(|b - \tilde{b}|) + \Omega(|b - \tilde{b}|) \right]$$

with $C_{32} = \text{const}(n, k, m, \lambda, \Lambda, \beta)$.

In the **general case** we consider $a, b \in \mathbf{B}_+ \cap \mathbf{U}_\varrho$ ($\varrho = \frac{1}{17}$) with the aim to prove the estimate (6.2). In the special cases this estimate follows from (6.9), (6.15), and (6.16). If $a \in \mathbf{U}_\varrho^{n-1} \times \{0\}$ and $b \in \mathbf{U}_+ \cap \mathbf{U}_\varrho$ with $|b - a| > 2b_n$, we apply the special case 1 to a and $b' = (b_1, \dots, b_{n-1}, 0)$ and the special case 2 to b' and b , and (6.2) follows because $|a - b'| \leq |a - b|$ and $|b - b'| \leq |a - b|$. The only remaining case is $a, b \in \mathbf{U}_\varrho$ with $0 < a_n \leq b_n$ and $|a - b| > \frac{2}{15}a_n$. In this case we apply the special case 1 to $a' = (a_1, \dots, a_{n-1}, 0)$ and $b' = (b_1, \dots, b_{n-1}, 0)$, and the special case 2 to a', a and to b', b . On account of $|a' - b'| \leq |a - b|$, $|a - a'| = a_n < \frac{15}{2}|a - b|$, and $|b - b'| \leq |b - a| + |a - a'| + |a' - b'| \leq \frac{19}{2}|a - b|$ the estimate (6.2) then follows again from (6.9) and (6.15), since $\widehat{\mathbf{K}}(t\varrho) + \Omega(t\varrho) \leq t^{2\alpha}(\widehat{\mathbf{K}}(\varrho) + \Omega(\varrho))$ for $t \geq 1$. \blacksquare

The preceding theorem is not completely satisfactory with regard to boundary regularity, for the following reason: if T is (F, ω) minimizing for a smooth integrand F and ∂T is

represented locally by an $(n-1)$ -dimensional oriented submanifold B of \mathbb{R}^{n+k} with multiplicity 1 and with continuity modulus κ for the tangent plane field of B , then we can flatten the boundary locally by transforming B into $(\mathbf{U}^{n-1} \times \{0\}) \times \{0\}^k$ with a diffeomorphism Φ^{-1} of class $C^{1,\kappa(\cdot)}$ and apply Theorem 6.1 to the transformed curret $\tilde{T} = \Phi_{\#}^{-1}T$ and to the transformed integrand $\tilde{F} = \Phi_{\#}F$ (provided the hypotheses of the theorem are satisfied by \tilde{T} and \tilde{F}). However, the continuity modulus of \tilde{F} and $D_{(2)}\tilde{F}$ with respect to the space variable will not be better than κ , in general, hence Theorem 6.1 will only give C^1 regularity for T with the modulus $(\hat{K} + \Omega)^{1/2}$. For example, if $\kappa(r) = \text{const } r^\beta$ with $0 < \beta < 1$ and $\omega(r) = \text{const } r^{2\beta}$ then we obtain only the modulus $\text{const } r^{\beta/(2-\beta)}$ for the tangent plane field of T , whereas the expected boundary regularity of T in this situation is $C^{1,\beta}$. Using the full strength of Lemma 5.4, however, we can prove the continuity modulus $(K^2 + \Omega)^{1/2}$ for the tangent plane field of T , i.e. the optimal $C^{1,\beta}$ boundary regularity in the example. For this we need the following variant of Lemma 5.3 for nonlinear mappings:

6.3 Lemma (excess comparison for diffeomorphisms). *Suppose Φ is a C^1 diffeomorphism between open sets of \mathbb{R}^{n+k} , $\sup \|D\Phi\| \leq L$ and $\sup \|D\Phi^{-1}\| \leq L$ with $1 \leq L < \infty$, and $T \in \mathcal{R}_n(\mathbb{R}^{n+k})$ such that $\mathbf{p}_{\#}(T \llcorner \mathbf{C}_r(y))$ and $\mathbf{p}_{\#}((\Phi_{\#}T) \llcorner \mathbf{C}_{\tilde{r}}(\tilde{y}))$ have nonnegative multiplicities and $\mathbf{C}_{\tilde{r}}(\tilde{y}) \cap \text{spt}(\Phi_{\#}T)$ is contained in $\Phi(\mathbf{C}_r(y) \cap \text{spt } T)$. Then we have*

$$\tilde{r}^n \mathbf{E}(\Phi_{\#}T, \tilde{y}, \tilde{r}) \leq 2L^n r^n \mathbf{E}(T, y, r) + n^2 L^{3n-2} \mathbf{M}(T \llcorner \mathbf{C}_r(y)) \sup_{\mathbf{C}_r(y) \cap \text{spt } T} \|D\Phi - \text{id}\|^2.$$

Proof. From the area formula [Fe, 3.2.20, 4.1.30] and (1.24) we obtain

$$\begin{aligned} 2\tilde{r}^n \mathbf{E}(\Phi_{\#}T, \tilde{y}, \tilde{r}) &= \int_{\mathbf{C}_{\tilde{r}}(\tilde{y})} |\overrightarrow{\Phi_{\#}T} - \tilde{\mathbf{e}}|^2 d\|\Phi_{\#}T\| \\ &\leq \int_{\mathbf{C}_r(y)} |\overrightarrow{\Phi_{\#}T}(\Phi(x)) - \tilde{\mathbf{e}}|^2 |(\wedge_n D\Phi(x))\vec{T}(x)| d\|T\|_x \\ &\leq 2L^n \int_{\mathbf{C}_r(y)} \left[|\vec{T} - \tilde{\mathbf{e}}|^2 + |(\wedge_n D\Phi)\vec{T} - |(\wedge_n D\Phi)\vec{T}|\vec{T}|^2 \right] d\|T\|, \end{aligned}$$

because $L^{-n} \leq |(\wedge_n D\Phi)\vec{T}| \leq L^n$ and $\overrightarrow{\Phi_{\#}T}(\Phi(x)) = |(\wedge_n D\Phi(x))\vec{T}(x)|^{-1} (\wedge_n D\Phi(x))\vec{T}(x)$. The claim then follows with

$$|(\wedge_n D\Phi(x))\vec{T}(x) - \vec{T}(x)| \leq \|D\Phi - \text{id}\| \sum_{j=0}^{n-1} \|D\Phi(x)\|^j.$$

■

It is clear that a variant of the Lemma holds for Φ only biLipschitz, where $D\Phi$ above has to be interpreted as the approximative differential with respect to the rectifiable set underlying T .

For the formulation of the optimal small excess boundary regularity theorem we now fix some **notations** and **assumptions**: We consider $T = T \llcorner \mathbf{C} \in \mathcal{R}_n(\mathbb{R}^{n+k})$ such that ∂T is represented in \mathbf{C} by the graph of a C^1 function $(\varphi_0, \dots, \varphi_k) = \varphi: \mathbf{U}^{n-1} \rightarrow \mathbb{R}^{1+k}$, taken with suitable orientation and multiplicity 1. We assume $\varphi(0) = 0$, $D\varphi(0) = 0$, and

$$\|D\varphi(y') - D\varphi(y'')\| \leq \kappa(|y' - y''|) \quad \text{for } y', y'' \in \mathbf{U}^{n-1},$$

where κ is a continuity modulus for the tangent plane field of the boundary $(\partial T)\llcorner \mathbf{C}$. We assume κ sufficiently small so that \mathbf{U} is divided into two components by graph $\varphi_0 = \mathbf{p}(\text{graph } \varphi)$, and for $\mathbf{p}\#T$ we assume that it has multiplicities 1 on $D_+ = \{(y', y_n) \in \mathbf{U} : y_n > \varphi_0(y')\}$ and multiplicity 0 on $D_- = \{(y', y_n) \in \mathbf{U} : y_n < \varphi_0(y')\}$, i.e. $\mathbf{p}\#T = \mathbf{E}^n \llcorner D_+$. For the integrand F we require all conditions from Section 1 in particular the bound Λ , the ellipticity constant λ , and the modulus ν for $\xi \mapsto D_{(2)}^2 F(x, \xi)$ from (1.4), but we denote the continuity modulus for $x \mapsto F(x, \xi)$ and $x \mapsto D_2 F(x, \xi)$ by μ this time, i.e. instead of (1.5), (1.6) we have

$$\begin{aligned} |F(x, \xi) - F(\tilde{x}, \xi)| &\leq \mu(|x - \tilde{x}|)|\xi|, \\ \|D_{(2)} F(x, \xi) - D_{(2)} F(\tilde{x}, \xi)\| &\leq \mu(|x - \tilde{x}|), \end{aligned}$$

and we do not make use of the special modulus introduced in (1.7), (1.8) (and denoted μ there). For the functions κ, μ, ω we assume all the conditions stated in Section 1. In particular there exists $0 < \alpha < 1$ such that $r^{-\alpha}\kappa(r), r^{-\alpha}\mu(r)$ and $r^{-\alpha}\sqrt{\omega(r)}$ are nonincreasing functions of $r > 0$. With μ we associate the function $\widehat{\mu}$ by (1.9) and with $\kappa, \widehat{\mu}, \omega$ the functions $\mathbf{K}, \widehat{\mathbf{M}}, \Omega$ by (1.15). We then have

6.4 Theorem (optimal boundary ϵ -regularity theorem). *For $\alpha \leq \beta < 1$ there exists a positive constant $\epsilon_1(n, k, \lambda, \Lambda, \nu(\cdot), \beta)$ such that if T, F satisfy the assumptions above, T is (F, ω) minimizing in \mathbf{C} and*

$$(6.17) \quad \mathbf{E}(T, 1) + \mathbf{K}(1)^2 + \widehat{\mathbf{M}}(1) + \Omega(1) \leq \epsilon_1,$$

then $T \llcorner \mathbf{C}_{1/50}$ is represented by the graph, taken with multiplicity 1, of a continuously differentiable function $g: \mathbf{U}_{1/50} \cap \mathbf{p}(\text{spt } T) \rightarrow \mathbb{R}^k$. Moreover, the derivative Dg has the following modulus of continuity:

$$(6.18) \quad |Dg(a) - Dg(b)| \leq C_{41} \left[r^{2\beta} \mathbf{E}(T, 1) + \mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r) \right]^{1/2}$$

for $a, b \in \mathbf{U}_{1/50} \cap \mathbf{p}(\text{spt } T)$ with $|a - b| \leq r$, where $C_{41} = \text{const}(n, k, \lambda, \Lambda, \beta)$.

For example, if the boundary is of class $C^{1,\beta}$, i.e. $\kappa(r) \leq \text{const } r^\beta$ for small $r > 0$, if the integrand is smooth ($\mu(r) \leq \text{const } r^{2\beta/(1+\beta)}$ is sufficient, as then $\widehat{\mathbf{M}}(r) \leq \text{const } r^{2\beta}$ follows), and if $\omega(r) \leq \text{const } r^{2\beta}$, then (6.18) gives Hölder continuity of Dg up to the boundary with exponent $\beta \in]0, 1[$. This is, of course, the best possible boundary regularity result, if the prescribed boundary is merely of class $C^{1,\beta}$. If the boundary is of Dini class $C^{1,\kappa(\cdot)}$, where $r \mapsto r^{-\beta}\kappa(r) > 0$ is decreasing for all $0 < \beta < 1$, then Theorem gives $C^{1,\mathbf{K}(\cdot)}$ Dini boundary regularity, provided $\mathbf{K}(1) = \int_0^1 \frac{1}{\varrho} \kappa(\varrho) d\varrho$ is finite and $\widehat{\mathbf{M}}(r), \Omega(r)$ are dominated by $\mathbf{K}(r)$ as $r \downarrow 0$ (e.g. $\mu(r) \leq \text{const } r^\beta, \omega(r) \leq \text{const } r^{2\beta}$ as above, for some $0 < \beta < 1$).

Proof. We define $\Phi(\tilde{x}) = \tilde{x} + (0, \varphi(\tilde{x}_1, \dots, \tilde{x}_{n-1})) \in \mathbb{R}^{n-1} \times \mathbb{R}^{1+k}$ for $\tilde{x} \in \mathbf{C}$ so that $\Phi^{-1}(x) = x - (0, \varphi(x_1, \dots, x_{n-1}))$. If ϵ_1 , and hence also $\kappa(1) \leq \mathbf{K}(1)$, is sufficiently small, we have $\text{Lip } \varphi \leq \kappa(1) \leq \frac{1}{7}$ and $\text{Lip } \Phi \leq \frac{8}{7}, \text{Lip } \Phi^{-1} \leq \frac{8}{7}$. Then $\Phi(\mathbf{C}_{7/8}) \subset \mathbf{C}$ follows because $\Phi(0) = 0$, and Φ acts on $\{y\} \times \mathbb{R}^n$ as a translation. Letting $\tilde{T} = (\mu_{7/8\#}^{-1} \Phi_{\#}^{-1} T) \llcorner \mathbf{C}$ and $\tilde{F}(\tilde{x}, \tilde{\xi}) = (\frac{8}{7})^n F(\Phi(\frac{7}{8}\tilde{x}), \wedge_n D\Phi(\frac{7}{8}\tilde{x})\tilde{\xi})$ we verify that $\tilde{T} \in \mathcal{T}_b(n, k, 1)$ is $(\tilde{F}, \tilde{\omega})$ minimizing in \mathbf{C} with $\tilde{\omega}(r) = \omega((\text{Lip } \Phi)^{\frac{7}{8}} r) \leq \omega(r)$. The transformed integrand \tilde{F} satisfies (1.1)–(1.4) and (1.12) with bound $\tilde{\Lambda} \leq C\Lambda$, modulus $\tilde{\nu}(r) \leq C[\nu(Cr) + \Lambda r]$, and ellipticity constant

$\tilde{\lambda} \geq C^{-1}\lambda$ (by [Fe, 5.1.4]), where $C = \text{const}(n, k)$. For the modulus $\tilde{\kappa}$ associated with \tilde{F} by (1.5), (1.6) we obtain $\tilde{\kappa}(r) \leq C[\mu((\text{Lip } \Phi)\frac{7}{8}r) + \Lambda\kappa(\frac{7}{8}r)] \leq C[\mu(r) + \Lambda\kappa(r)]$, and for the modulus $\tilde{\mu}$ associated with \tilde{F} by (1.7), (1.8) we find, using the special structure of Φ , i.e. the independence of $D\Phi(\tilde{x})$ from $\tilde{x}_n, \dots, \tilde{x}_{n+k}$, the estimate $\tilde{\mu}(r) \leq \mu((\text{Lip } \Phi)\frac{7}{8}r) \leq C\mu(r)$. It follows that $\tilde{X}(r)$, defined by (5.33) and (1.15), (1.9) in terms of $\tilde{\kappa}, \tilde{\mu}, \tilde{\omega}$, does not exceed $C[\mathbf{K}(r)^2 + \mathbf{M}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r)]$, where $\mathbf{K}, \mathbf{M}, \widehat{\mathbf{M}}, \Omega$ are defined in terms of κ, μ, ω as in (1.15), (1.9). If ϵ_1 , and hence also $\widehat{\mu}(1) \leq \widehat{\mathbf{M}}(1)$, is small enough, then we see $\mu(r)^2 \leq 16\widehat{\mu}(r)$ and $\mathbf{M}(r)^2 \leq 16\widehat{\mathbf{M}}(r)$ for $0 < r \leq 1$, recalling the discussion following (1.9). Consequently,

$$(6.19) \quad \tilde{X}(r) \leq \text{const}(n, k) \left[\mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r) \right] \quad \text{for } 0 < r \leq 1.$$

We also note that

$$(6.20) \quad \mathbf{E}(\tilde{T}, 1) \leq \text{const}(n) \left[\mathbf{E}(T, 1) + \kappa(\frac{7}{8})^2 \right],$$

by Lemma 6.3. It follows from (6.17), (6.19), (6.20) and $\kappa(1) \leq \mathbf{K}(1)$ that \tilde{T} satisfies the hypotheses of Lemma 5.4, provided ϵ_1 is chosen sufficiently small.

In the rest of the proof we will need additional smallness conditions for ϵ_1 in various arguments, and we will tacitly assume that ϵ_1 has been chosen to satisfy these conditions. By C_{34}, C_{35}, \dots we denote constants depending only on $n, k, \lambda, \Lambda, \beta$. As in Step 1 of the proof of Theorem 6.1 we next establish uniform smallness

$$(6.21) \quad \mathbf{E}(\tilde{T}, y, r) + \mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r) \leq C_{34}\epsilon_1,$$

for $y \in \mathbf{B}_{1/2}$ with $y_n \geq 0$ and for $r > 0$ with $r \leq 1 - |y|$. In the case $y_n = 0, \frac{1}{2} \leq r \leq 1 - |y|$ this follows from (6.17) and (6.20) by passing to the scaled current $(\mu_{r\#}^{-1}\tau_{(y,0)}^{-1}\tilde{T})\llcorner \mathbf{C}$, and from the excess bound (5.41) together with (6.19) we obtain (6.21) also when $y_n = 0, 0 < r < \frac{1}{2}$. In the case $y_n > 0$ we deduce (6.21) for $(\frac{7}{8})^2 y_n \leq r \leq 1 - |y|$ by applying (6.21) with center $y' = (y_1, \dots, y_{n-1}, 0)$ and radius $r' = 2y_n$ when $2y_n \leq 1 - |y|$ and with center 0 and radius 1 when $2y_n > 1 - |y|$ (which implies $r \geq (\frac{7}{8})^2 y_n > (\frac{7}{8})^2 \frac{1}{4}$). Using this with $r = y_n$ we see from Lemma 6.3 and

$$(6.22) \quad \mathbf{M}(\tilde{T}\llcorner \mathbf{C}_r(y)) \leq r^n \left[\alpha(n) + \mathbf{E}(\tilde{T}, y, r) \right]$$

that $((\mu_{49r/64}^{-1}\tau_x^{-1})\#T)\llcorner \mathbf{C}$ for $x \in \text{spt } T$ with $\mathbf{p}x = \mathbf{p}\Phi(\frac{7}{8}y, 0)$ satisfies the hypotheses of Lemma 5.4 in the interior situation (with respect to the corresponding transformation of the integrand F). From the conclusion (5.41) of this lemma we get (6.21) also for $0 < r < (\frac{7}{8})^2 y_n$ by using Lemma 6.3 again (and the analogue of (6.22)). (Note that we cannot apply the interior case of lemma 5.4 directly to \tilde{T} , because we did not assume that $\int_0^1 \frac{1}{\varrho} \sqrt{\tilde{\kappa}(\varrho)} d\varrho$ is finite.) Combining (6.21) with the height estimate of Lemma 2.2 we find that the height of $(\mu_{r\#}^{-1}\tau_{\tilde{x}\#}^{-1}\tilde{T})\llcorner \mathbf{C}$ does not exceed $\frac{1}{3}$, say, if $\tilde{x} \in \text{spt } \tilde{T}, \mathbf{p}\tilde{x} \in \mathbf{B}_{1/2} \cap \mathbf{B}_+$, and $0 < 2r < 1 - |\mathbf{p}\tilde{x}|$. Applying the diffeomorphism Φ we obtain corresponding height estimates for T , e.g. for $0 < r \leq \frac{7}{16}$ the height of $T\llcorner \mathbf{C}_r$ does not exceed $\frac{2}{3}(\frac{8}{7})^2 r < r$.

Exactly as in Step 2 of the proof for Theorem 6.1 we see from (6.21) that $\tilde{T}\llcorner \mathbf{C}_{1/34}$ is represented by the graph, taken with multiplicity 1, of a Lipschitz function $\tilde{g}: \mathbf{U}_{1/34} \cap \mathbf{B}_+ \rightarrow \mathbb{R}^k$ with $\text{Lip } \tilde{g} \leq 1$ and $\tilde{g} = 0$ on $\mathbf{U}_{1/34}^{n-1} \times \{0\}$. Consequently, T is represented in

$\mathbf{C}_{1/50}$ (since $(\frac{7}{8})^2 \frac{1}{34} < \frac{1}{50}$) by the graph of a function $g: \mathbf{U}_{1/50} \cap \mathbf{p}(\text{spt } T) \rightarrow \mathbb{R}^k$ such that $\text{graph } g = \mathbf{C}_{1/50} \cap \Phi \circ \boldsymbol{\mu}_{7/8}(\text{graph } \tilde{g})$, $g(y', \varphi_0(y')) = (\varphi_1(y'), \dots, \varphi_k(y'))$ for $(y', \varphi_0(y')) \in \mathbf{C}_{1/50} \cap \mathbf{p}(\text{spt } \partial T)$, and $\text{Lip } g \leq 2$.

From the excess decay Lemma 5.4 for \tilde{T} we obtain the existence of a linear isometry $,_0$ of \mathbb{R}^{n+k} with $,_0 = \text{id}$ on $(\mathbb{R}^{n-1} \times \{0\}) \times \{0\}^k$, $\|,_0 - \text{id}\| \leq C_{35}\epsilon_1$, and

$$(6.23) \quad \mathbf{E}\left(,_{0\#}(\tilde{T} \llcorner \mathbf{C}_{1/2}), r\right) \leq C_{35} \left[r^{2\beta} \mathbf{E}(\tilde{T}, 1) + \mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r) \right]$$

for $0 < r < \frac{1}{3}$. Letting $\Psi = ,_0 \circ \boldsymbol{\mu}_{8/7} \circ \Phi \circ \boldsymbol{\mu}_{7/8} \circ ,_0^{-1}$ we have $(\Psi_{\#},_{0\#}(\tilde{T} \llcorner \mathbf{C}_{1/2})) \llcorner \mathbf{C}_{1/3} = (,_{0\#} \boldsymbol{\mu}_{8/7\#}(T \llcorner \mathbf{C}_{7/16})) \llcorner \mathbf{C}_{1/3}$ and

$$\|D\Psi(\tilde{x}) - \text{id}\| = \left\| ,_0 D\Phi\left(\frac{7}{8}, ,_0^{-1}\tilde{x}\right), ,_0^{-1} - \text{id} \right\| = \left\| D\Phi\left(\frac{7}{8}, ,_0^{-1}\tilde{x}\right) - \text{id} \right\| \leq \kappa\left(\frac{7}{8}|\tilde{x}|\right).$$

On account of the smallness of $\|,_0 - \text{id}\|$ and the controlled height of $\tilde{T} \llcorner \mathbf{C}_{1/2}$ and $T \llcorner \mathbf{C}_{7/16}$ we can therefore apply Lemma 6.3 to deduce with (6.23), the analog of (6.22), and $\kappa(r) \leq \mathbf{K}(r)$, the estimate

$$(6.24) \quad \mathbf{E}\left(,_{0\#}(T \llcorner \mathbf{C}_{7/16}), r\right) \leq C_{36} \left[r^{2\beta} \mathbf{E}(T, 1) + \mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r) \right]$$

for $0 < r \leq \frac{1}{4}$.

Consider now $x \in \text{spt } T$ with $\mathbf{p}x = y = (0, \dots, 0, y_n)$ and $0 < y_n \leq \frac{1}{8}$. For $\hat{x} = ,_0 x$ and $\hat{y} = \mathbf{p}\hat{x}$ we then have $\hat{y} = (0, \dots, 0, \hat{y}_n)$ with $\frac{2}{3}y_n \leq \hat{y}_n \leq \frac{3}{2}y_n$, because $,_0 = \text{id}$ on $(\mathbb{R}^{n-1} \times \{0\}) \times \{0\}^k$, $\|,_0 - \text{id}\|$ is small, $T \llcorner \mathbf{C}_r$ has height at most r for $0 < r \leq \frac{7}{16}$. Therefore $\mathbf{C}_{y_n/2}(\hat{y})$ is contained in \mathbf{C}_{2y_n} , and from (6.24) we obtain

$$(6.25) \quad \begin{aligned} & \mathbf{E}\left(,_{0\#}(T \llcorner \mathbf{C}_{7/16}), \hat{y}, \frac{1}{2}y_n\right) \\ & \leq 4^n C_{36} \left[(2y_n)^{2\beta} \mathbf{E}(T, 1) + \mathbf{K}(2y_n)^2 + \widehat{\mathbf{M}}(2y_n) + \Omega(2y_n) \right]. \end{aligned}$$

Since $\mathbf{C}_{y_n/2}(\hat{y})$ does not intersect the support of $,_{0\#}(\partial T \llcorner \mathbf{C}_{7/16})$, the hypotheses of the interior excess decay estimate in Lemma 5.4 are satisfied for $(\boldsymbol{\mu}_{y_n/2}^{-1},_{0\#} \boldsymbol{\tau}_x^{-1})_{\#}(T \llcorner \mathbf{C}_{7/16}) \llcorner \mathbf{C}$ and for the corresponding transformation of the integrand F . Consequently, there exists a linear isometry $,_x$ of \mathbb{R}^{n+k} , close to the identity, such that for $T_x = \boldsymbol{\tau}_{x\#}^{-1}(T \llcorner \mathbf{C}_{7/16})$ we have

$$\mathbf{E}\left(,_{x\#}T_x, r\right) \leq C_{26} \left[\left(\frac{r}{y_n}\right)^{2\beta} \mathbf{E}\left(,_{0\#}(T \llcorner \mathbf{C}_{7/16}), r\right) + \widehat{\mathbf{M}}(r) + \Omega(r) \right]$$

for $0 < r \leq \frac{1}{6}y_n$. Combining this with (6.25), and using $(r/y_n)^{2\beta}\Omega(2y_n) \leq (r/y_n)^{2\alpha}\Omega(2y_n) \leq 2^{2\alpha}\Omega(r) \leq 2^{2\beta}\Omega(r)$ and the analogous inequality for $\widehat{\mathbf{M}}$ we infer

$$(6.26) \quad \mathbf{E}\left(,_{x\#}T_x, r\right) \leq C_{37} \left[r^{2\beta} \mathbf{E}(T, 1) + \mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r) \right]$$

for $x \in \text{spt } T$ with $\mathbf{p}x = (0, \dots, 0, y_n)$, $0 < y_n \leq \frac{1}{8}$, and $0 < r \leq \frac{1}{6}y_n$.

Now, choosing ϵ_1 sufficiently small we obtain linear isometries Δ_a corresponding to $a \in \mathbf{C}_{1/8} \cap \text{spt } \partial T$ such that $\|\Delta_a - \text{id}\| \leq \text{const}(n, k)\kappa(|a|)$ and the scaled currents $((\boldsymbol{\mu}_4 \Delta_a \boldsymbol{\tau}_a^{-1})_{\#})$

$(T \llcorner \mathbf{C}_{7/16})) \llcorner \mathbf{C}$ together with the correspondingly scaled version of F satisfy all the assumptions which we have used for T above to derive (6.24) and (6.26). Here, the smallness of the excess follows from (6.17) and Lemma 5.3. More precisely, with a mass estimate as in (6.22) we obtain the following estimate, valid for $a \in \mathbf{C}_{1/8} \cap \text{spt } \partial T$, $T_a = \tau_{a\#}^{-1}(T \llcorner \mathbf{C}_{7/16})$, some linear isometry τ_a of \mathbb{R}^{n+k} close to the identity, and $0 < r \leq \frac{1}{16}$:

$$(6.27) \quad \begin{aligned} \mathbf{E}(\tau_a, T_a, r) &\leq C_{38} \left[r^{2\beta} \left(\mathbf{E}(T, 1) + \kappa(|a|)^2 \right) + \mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r) \right] \\ &\leq C_{39} \left[r^{2\beta} \mathbf{E}(T, 1) + \mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r) \right]. \end{aligned}$$

Here we have used $r^\beta \kappa(|a|) \leq r^\alpha \kappa(1) \leq \kappa(r) \leq \mathbf{K}(r)$. Moreover, if $b \in \mathbf{C}_{1/32} \cap \text{spt } T \setminus \text{spt } \partial T$ and a is a nearest point to b in $\text{spt } \partial T$, then $|a| < \frac{1}{16}$ and $\mathbf{p}\mu_4 \Delta_a \tau_a^{-1}(b) = (0, \dots, 0, y_n)$ with $0 < y_n \leq 4|b - a| < \frac{1}{8}$. Hence (6.26) implies, by the same reasoning, the existence of a linear isometry τ_b of \mathbb{R}^{n+k} close to the identity such that $T_b = \tau_{b\#}^{-1}(T \llcorner \mathbf{C}_{7/16})$ satisfies

$$(6.28) \quad \mathbf{E}(\tau_b, T_b, r) \leq C_{40} \left[r^{2\beta} \mathbf{E}(T, 1) + \mathbf{K}(r)^2 + \widehat{\mathbf{M}}(r) + \Omega(r) \right]$$

for $0 < r \leq \frac{1}{24}y_n$. Since $4|b - a| \leq \sqrt{2}y_n$, because $\mu_{4\#} \Delta_{a\#} \tau_{a\#}^{-1}(T \llcorner \mathbf{C}_{7/16})$ has height in \mathbf{C}_r not exceeding r for $0 < r \leq \frac{7}{16}$, we see that (6.28) is valid in particular for $b \in \mathbf{C}_{1/32} \cap \text{spt } T \setminus \text{spt } (\partial T)$ and $0 < r \leq \frac{1}{9} \text{dist}(\mathbf{p}b, \mathbf{p}(\text{spt } (\partial T)))$.

With (6.27) and (6.28) we can now proceed exactly as in Step 4 of the proof of Theorem 6.1 in order to establish the asserted continuity modulus (6.18) for Dg . One uses the fact that $T \llcorner \mathbf{C}_{1/50}$ is represented by the graph of g with $\text{Lip } g \leq 2$ and estimates $|Dg(a) - Dg(b)|^2$ for $a, b \in \mathbf{C}_{1/50} \cap \mathbf{p}(\text{spt } T)$ distinguishing the three special cases $a, b \in \mathbf{p}(\text{spt } \partial T)$, resp. $a \in \mathbf{p}(\text{spt } \partial T)$ and $b \notin \mathbf{p}(\text{spt } \partial T)$, resp. $a, b \notin \mathbf{p}(\text{spt } \partial T)$ with $|a - b| \leq \frac{1}{9} \text{dist}(a, \mathbf{p}(\text{spt } \partial T)) \leq \frac{1}{9} \text{dist}(b, \mathbf{p}(\text{spt } \partial T))$. The general case is reduced to the special cases by the triangle inequality. As the reasoning is clear from the proof of Theorem 6.1 we can omit details. \blacksquare

7 Proof of the boundary regularity theorem

For completeness we indicate here how the boundary regularity theorem stated in the introduction follows from the ϵ -regularity theorem. The proof follows [Ha1, Section 3] closely, with simple modifications to treat almost minimizing currents. We first observe that the following compactness lemma for almost minimizing currents holds:

7.1 Lemma (compactness for sequences of almost minimizing currents). *Suppose $T_i \in \mathcal{R}_n(\mathbb{R}^{n+k})$ is (F_i, ω_i) minimizing in the open subset D of \mathbb{R}^{n+k} for $i = 1, 2, \dots$, where the parametric integrands F_i satisfy (1.1) uniformly and converge to F uniformly on compact subsets of $D \times \bigwedge_n \mathbb{R}^{n+k}$ as $i \rightarrow \infty$ and where $\limsup_{i \rightarrow \infty} \omega_i \leq \omega$ holds on $]0, \infty[$ with a nondecreasing function $\omega:]0, \infty[\rightarrow [0, \infty[$ which is continuous from the right with $\omega(0+) = 0$. If $T_i \rightarrow T_0$ holds locally on D with respect to the flat metric distance and $\sup_i \mathbf{M}(T_i) < \infty$ then $T_0 \in \mathcal{R}_n(\mathbb{R}^{n+k})$ is (F, ω) minimizing in D . Moreover, every subset A of D with positive distance to $(\text{spt } T_0) \cup \bigcup_{i=1}^{\infty} \text{spt } \partial T_i$ intersects only finitely many of the sets $\text{spt } T_i$.*

The proof is a straightforward adaptation of [Ha1, Lemma 3.5] which in turn is modelled after [Fe, 5.4.2]. It uses only the definition of almost minimality, standard properties of

convergence with respect to the flat distance [Fe, 4.1.24], and the lower mass bound from Lemma 2.1 which holds uniformly for the T_i with sufficiently large i by the hypotheses on the integrands F_i and the functions ω_i . (Note that $\omega_i(r) \leq 1 + \omega(r) < \infty$ for some $r > 0$ and all sufficiently large i .) We omit the details and proceed to the

Proof of the boundary regularity theorem. We argue by contradiction verifying also the assertions made in Section 0 about the dependence of the constant $\vartheta > 0$ on the data. If the theorem were false we would have, after applying linear isometries to transform the point a to the origin and the tangent space of the boundaries at a to $\mathbb{R}^{n-1} \times \{0\}^{k+1} \subset \mathbb{R}^{n+k}$ and after suitable scalings by homotheties of \mathbb{R}^{n+k} , sequences of currents $T_i \in \mathcal{R}_n(\mathbb{R}^{n+k})$, parametric integrands F_i , and $C^{1,\beta}$ functions $\varphi_i: \mathbf{U}^{n-1}(0, 1) \rightarrow \mathbb{R}^{1+k}$, such that no T_i satisfies the conclusion of Theorem 6.4 locally at 0 and the following assertions hold:

$$(7.1) \quad T_i \text{ is } (F_i, \omega_i) \text{ minimizing in } \mathbf{U}(0, 1) = \mathbf{U}^{n+k}(0, 1) \text{ with } \omega_i(r) = \sigma_i r^\beta, \\ \lim_{i \rightarrow \infty} \sigma_i = 0;$$

$(\partial T_i) \llcorner \mathbf{U}(0, 1)$ is represented by the graph of φ_i taken with suitable orientation and multiplicity 1; moreover $\varphi(0) = 0$, $D\varphi(0) = 0$, and

$$(7.2) \quad \|D\varphi(y') - D\varphi(y'')\| \leq \kappa_i |y' - y''|^\beta \quad \text{for } y', y'' \in \mathbf{U}^{n-1},$$

with $\lim_{i \rightarrow \infty} \kappa_i = 0$;

$$(7.3) \quad \mathbf{M}(T_i \llcorner \mathbf{U}(0, 1)) \leq \frac{1}{2} \boldsymbol{\alpha}(n) + \frac{1}{i};$$

$$(7.4) \quad F_i \text{ satisfies the structure conditions (1.1)–(1.12) with } \lim_{i \rightarrow \infty} \mu_i = 0 \\ \text{and with } \Lambda, \lambda, \nu(\cdot) \text{ independent of } i.$$

Moreover, as the F_i satisfy a uniform Lipschitz condition on each compact subset of $\mathbf{U}(0, 1) \times \bigwedge_n \mathbb{R}^{n+k}$ we may assume, after passing to a subsequence, that for some continuous integrand $F_0: \mathbf{U}(0, 1) \times \bigwedge_n \mathbb{R}^{n+k} \rightarrow [0, \infty[$ (which has constant coefficients, because $\mu_i \rightarrow 0$) we have

$$(7.5) \quad F_i \rightarrow F_0 \text{ uniformly on compact subsets of } \mathbf{U}(0, 1) \times \bigwedge_n \mathbb{R}^{n+k} \text{ as } i \rightarrow \infty.$$

Now, since $\mathbf{M}(T_i \llcorner \mathbf{U}(0, 1)) + \mathbf{M}((\partial T_i) \llcorner \mathbf{U}(0, 1))$ is bounded, by (7.3) and (7.2), we may apply the local version of the compactness theorem [Si1, 31.3] (cf. the reasoning in the proof of [Fe, 5.4.3]) to deduce, passing to a subsequence again, convergence $T_i \rightarrow T_0 \in \mathcal{R}_n(\mathbb{R}^{n+k})$ locally on $\mathbf{U}(0, 1)$ in the flat metric distance. Clearly,

$$(7.6) \quad \mathbf{M}(T_0) \leq \frac{1}{2} \boldsymbol{\alpha}(n)$$

holds, by (7.3), and

$$(7.7) \quad \mathbf{U}(0, 1) \cap \text{spt}(\partial T_0) = \mathbf{U}^{n-1}(0, 1) \times \{0\}^{1+k} \subset \mathbb{R}^{n+k}$$

by (7.2). Using the nonexpanding map $f(y, z) = (y_1, \dots, y_{n-1}, |(y_n, z)|)$ from $\mathbb{R}^n \times \mathbb{R}^k$ to $\mathbb{R}^{n-1} \times [0, \infty[$ one gets from (7.6)

$$(7.8) \quad \mathbf{M}(f_{\#} T_0) \leq \int_{\mathbf{U}(0,1)} \left| \left(\bigwedge_n Df \right) \vec{T}_0 \right| d\|T_0\| \leq \int_{\mathbf{U}(0,1)} d\|T_0\| \leq \frac{1}{2} \boldsymbol{\alpha}(n),$$

and since, for $\mathbf{U} = \mathbf{U}^n(0, 1)$,

$$\mathbf{U} \cap \text{spt}(f_{\#}T_0) \subset f(\mathbf{U}^{n-1} \times \{0\}^{1+k}) = \mathbf{U}^{n-1} \times \{0\} \subset \mathbf{U}$$

by (7.7) and the definition of f , one deduces with the constancy theorem [Fe, 4.1.7] that $f_{\#}T_0$ represents the half ball \mathbf{U}_+ with multiplicity $+1$ or -1 and that $|(\wedge_n Df)\vec{T}_0| = 1$ holds $\|T_0\|$ -almost everywhere. This means that $\vec{T}_0(y, z)$ is an orientation vector of the n -dimensional subspace $\mathbb{R}(y, z) + \mathbb{R}^{n-1} \times \{0\}$ of \mathbb{R}^{n+k} for $\|T_0\|$ -almost all $(y, z) \in \mathbb{R}^n \times (\mathbb{R}^k \setminus \{0\})$, and slicing with the map $(y, z) \mapsto |(y_n, z)|^{-1}(y_n, z) \in S^k$ one sees, using also (7.6), that T_0 represents an n -dimensional half ball in \mathbb{R}^{n+k} with equator $\mathbf{U}^{n-1} \times \{0\}$. (See [Br2, 4.5] where the argument of [Ha2, Corollary 1] is adapted to the present situation.) After a suitable orthogonal transformation of \mathbb{R}^{n+k} we may assume $T_0 = (\mathbf{E}^n \llcorner \mathbf{U}_+) \times \delta_0^k$.

By Lemma 7.1, T_0 is absolutely minimizing for F_0 , and for every $0 < \delta < 1$ the set $\{(y, z) \in \mathbf{B}(0, 1 - \delta) : |z| \geq \delta\}$ intersects only finitely many of the sets $\text{spt} T_i$. It is now clear from (7.1)–(7.4) and the constancy theorem [Fe, 4.1.7] that the currents

$$\tilde{T}_i = (\mu_{1-\delta}^{-1})_{\#}(T_i \llcorner \mathbf{U}^n(0, 1 - \delta) \times \mathbf{U}^k(0, \delta)) \in \mathcal{R}_n(\mathbf{C})$$

for $\delta > 0$ fixed sufficiently small and for i sufficiently large satisfy the hypotheses of the boundary case in Theorem 6.4 for the integrands \tilde{F}_i and functions $\tilde{\omega}_i$ obtained from F_i and ω_i by scaling with $\mu_{1-\delta}^{-1}$. Therefore, for i sufficiently large the currents \tilde{T}_i are regular at the origin as described in the conclusion of Theorem 6.4. This contradicts, however, our assumption on the sequence T_i . \blacksquare

In the interior situation $a \in \text{spt} T \setminus \text{spt}(\partial T)$ one can conclude with a similar reasoning that the (F, ω) minimizing current T is regular locally at a , provided the n -dimensional density of $\|T\|$ has a local minimum at a (on the complement of a set of $\|T\|$ measure zero) and $\text{Tan}(\text{spt} T, a)$ is an n -dimensional subspace of \mathbb{R}^{n+k} (cf. [Fe, 5.3.16] and [Bo, VIII]). The interior case of Theorem 6.4 then gives the same modulus of continuity for the field of the tangent n -planes to $\text{spt} T$ near a as in our boundary regularity theorem.

References

- [All1] *Allard, W.K.*: On the first variation of a varifold. *Annals of Math.* **95** (1972), 417–491.
- [All2] *Allard, W.K.*: On the first variation of a varifold: boundary behaviour. *Annals of Math.* **101** (1975), 418–446.
- [Alm1] *Almgren, F. J.*: Existence and regularity almost everywhere of solutions to elliptic variational problems among surfaces of varying topological type and singularity structure. *Annals of Math.* **87** (1968), 321–391.
- [Alm2] *Almgren, F. J.*: Existence and regularity almost everywhere of solutions to elliptic variational problems with constraints. *Mem. Am. Math. Soc.* **165** (1976).
- [Alm3] *Almgren, F. J.*: \mathbf{Q} -valued functions minimizing Dirichlet’s integral and the regularity of area minimizing rectifiable currents up to codimension two. Princeton Univ. 1984, 3745 g; see also *Bull. Amer. Math. Soc.* **8** (1983), 327–328.
- [Alm4] *Almgren, F. J.*: Optimal isoperimetric inequalities. *Indiana Univ. Math. J.* **35** (1986), 451–547.
- [Bo] *Bombieri, E.*: Regularity theory for almost minimal currents. *Arch. Rational Mech. Anal.* **7** (1982), 99–130.

- [Br1] *Brothers, J. E.*: The structure of solutions to Plateau's problem in the n -sphere. *J. Diff. Geometry* **11** (1976), 387–400.
- [Br2] *Brothers, J. E.*: Existence and structure of tangent cones at the boundary of an area minimizing integral current. *Indiana Univ. Math. J.* **26** (1977), 1027–1044.
- [Ca] *Campanato, S.*: Equazioni ellittiche del II^e ordine e spazi $\mathcal{L}^{2,\lambda}$. *Ann. Mat. Pura Appl.* **69** (1965), 321–380.
- [DeG] *DeGiorgi, E.*: Frontiere orientate di misura minima. *Sem. Mat. Norm. Sup. Pisa*, 1960–61.
- [Du1] *Duzaar, F.*: On the existence of surfaces with prescribed mean curvature and boundary in higher dimensions. *Analyse non linéaire* **10** (1993), 191–214.
- [Du2] *Duzaar, F.*: Boundary regularity for area minimizing currents with prescribed volume. *J. of Geom. Anal.* **7** (1997), 585–592.
- [DF1] *Duzaar, F., Fuchs, M.*: On the existence of integral currents with prescribed mean curvature vector. *Manuscripta math.* **67** (1990), 41–67.
- [DF2] *Duzaar, F., Fuchs, M.*: A general existence theorem for integral currents with prescribed mean curvature form. *Bolletino U.M.I. (7)* **6–B** (1992), 901–912.
- [DGa] *Duzaar, F., Gastel, A.*: Nonlinear elliptic systems with Dini continuous coefficients. Preprint SFB 288, Berlin.
- [DGr] *Duzaar, F., Grotowski, J.F.*: Partial regularity for nonlinear elliptic systems: The method of A -harmonic approximation. To appear in *Manuscripta. Math.*
- [DGG] *Duzaar, F., Gastel, A., Grotowski, J.F.*: Partial regularity for almost minimizing functions of quasiconvex integrands. Preprint SFB 288, Berlin.
- [DS1] *Duzaar, F., Steffen, K.*: λ minimizing currents. *Manuscripta math.* **80** (1993), 403–447.
- [DS2] *Duzaar, F., Steffen, K.*: Boundary regularity for minimizing currents with prescribed mean curvature. *Calc. Var.* **1** (1993), 355–406.
- [DS3] *Duzaar, F., Steffen, K.*: Existence of hypersurfaces with prescribed mean curvature in Riemannian manifolds. *Indiana Univ. Math. J.* **45** (1996), 1045–1093.
- [Fe] *Federer, H.*: Geometric measure theory. Springer–Verlag, Berlin Heidelberg New York, 1969.
- [Fe1] *Federer, H.*: The singular set of area minimizing rectifiable currents with codimension one and of area minimizing flat chains modulo two with arbitrary codimension. *Bull. Amer. Math. Soc.* **76** (1970), 767–771.
- [Gi] *Giaquinta, M.*: Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems. Princeton NJ, Princeton University Press, 1983.
- [GMT] *Gonzalez, E., Massari, U., Tamanini, I.*: On the regularity of boundaries of sets minimizing perimeter with a volume constraint. *Indiana Univ. Math. J.* **32** (1983), 25–37.
- [Ha1] *Hardt, R.*: On boundary regularity for integral currents or flat chains modulo two minimizing the integral of an elliptic integrand. *Comm. Partial Differ. Equations* **2** (1977), 1163–1232.
- [Ha2] *Hardt, R.*: Uniqueness of nonparametric area minimizing currents. *Indiana Univ. Math. J.* **26** (1977), 65–71.
- [HaL] *Hardt, R., Lin, F.-H.*: Tangential regularity near the C^1 boundary. *Proc. Symposia Pure Math.* **44** (1986), 245–253.
- [HS] *Hardt, R., Simon, L.*: Boundary regularity and embedded solutions for the oriented Plateau problem. *Annals of Math.* **110** (1979), 439–486.
- [Li] *Lin, F.-H.*: Regularity for a class of parametric obstacle problems. Thesis, Univ. of Minnesota, 1985.
- [Ma] *Massari, U.*: Esistenza e regolarità delle ipersuperfici di curvatura media assegnata in \mathbb{R}^n . *Arch. Rational Mech. Anal.* **55** (1974), 357–382.

- [MM] *Massari, U., Miranda, M.*: Minimal surfaces of codimension one. Mathematical studies, vol. 91, North-Holland, Amsterdam 1984.
- [Mi] *Miranda, M.*: Frontiere orientate con ostacoli. Ann. Univ. Ferrara **16** (1971), 29–37.
- [Mo] *Morrey, C.B.*: Multiple integrals in the calculus of variations. Springer–Verlag, Berlin Heidelberg New York, 1966.
- [SS] *Schoen, R., Simon, L.*: A new proof of the regularity for rectifiable currents which minimize parametric elliptic functionals. Indiana Univ. Math. J. **31** (1982), 413–434.
- [SSA] *Schoen, R., Simon, L., Almgren, F.J.*: Regularity and singularity estimates on hypersurfaces minimizing parametric elliptic variational integrals. Acta Math. **139** (1977), 217–265.
- [Si1] *Simon, L.*: Lectures on geometric measure theory. Proc. CMA, Vol. 3, ANU Canberra, 1983.
- [Si2] *Simon, L.*: Theorems on regularity and singularity of energy minimizing maps. Birkhäuser–Verlag, Basel Boston Berlin, 1996.
- [Sv1] *Šverák, V.*: Quasiconvex functions with subquadratic growth. Proc. Roy. Soc. London A **433** (1991), 223–225.
- [Sv2] *Šverák, V.*: Rank-one convexity does not imply quasiconvexity. Proc. Roy. Soc. Edinburgh **120** (1992), 185–189.
- [Ta1] *Tamanini, I.*: Regularity results for almost minimal hypersurfaces in \mathbb{R}^N . Quaderni Dip. Mat. Univ. Lecce, Q1 (1984).
- [Ta2] *Tamanini, I.*: Boundaries of Cacciopoli sets with Hölder-continuous normal vector. J. reine Angew. Math. **334** (1982), 27–39.
- [Ta3] *Tamanini, I.*: Variational problems of least area type with constraints. Ann. Univ. Ferrara **34** (1988), 183–217.

*Mathematisches Institut der Humboldt-Universität zu Berlin,
Unter den Linden 6, D-10099 Berlin, Germany*

and

*Mathematisches Institut der Heinrich–Heine–Universität Düsseldorf
Universitätstraße 1, D-40225 Düsseldorf, Germany.*