

United Nations Educational Scientific and Cultural Organization
and
International Atomic Energy Agency
THE ABDUS SALAM INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

**BOUNDED POINT EVALUATIONS FOR CYCLIC OPERATORS
AND LOCAL SPECTRA**

A. Bourhim¹

*Departement de Mathématiques - Informatique, Faculté des Sciences,
Université Mohammed V, B.P. 1014, Rabat, Morocco*

and

The Abdus Salam International Centre for Theoretical Physics, Trieste, Italy,

C.E. Chidume²

The Abdus Salam International Centre for Theoretical Physics, Trieste, Italy

and

E.H. Zerouali³

*Département de Mathématiques, Université Mohamed V, B.P. 1014, Rabat, Morocco
and*

The Abdus Salam International Centre for Theoretical Physics, Trieste, Italy.

MIRAMARE – TRIESTE

August 2000

¹E-mail: bourhim@ictp.trieste.it

²E-mail: chidume@ictp.trieste.it

³Regular Associate of the Abdus Salam ICTP. E-mail: zerouali@fsr.ac.ma

Abstract

We study in this paper the concept of bounded point evaluations for cyclic operators. We give a negative answer to a question of L.R. Williams *Dynamic Systems and Applications* 3(1994) 103-112. Furthermore, we generalize some results of Williams and give a simple proof of theorem 2.5 of [11] that nonnormal hyponormal weighted shifts have fat local spectra.

Throughout this paper, $L(H)$ will denote the algebra of all linear bounded operators on a Hilbert space H . For an operator $S \in L(H)$, let S^* denote its adjoint, $\sigma(S)$ its spectrum, $\sigma_p(S)$ its point spectrum, $\sigma_{ap}(S)$ its approximate spectrum, $\sigma_c(S)$ its compression spectrum, $r(S)$ its spectral radius, $m(S)$ its lower bound (i.e., $\inf \{\|Sx\| : \|x\| = 1\}$) and $r_1(S) = \sup_n m(S^n)^{\frac{1}{n}}$ which equals $\lim_n m(S^n)^{\frac{1}{n}}$. For $F \subset \mathbb{C}$, we denote by $\overline{F} = \{\bar{z} : z \in \mathbb{C}\}$ its conjugate set, $int(F)$ its interior and $cl(F)$ its closure.

Let $T \in L(H)$ be a cyclic operator on H with cyclic vector x that is the finite linear combinations of the vectors x, Tx, T^2x, \dots are dense. A complex number $\lambda \in \mathbb{C}$ is said to be a **bounded point evaluation** of T if there is a constant $M > 0$ such that

$$|p(\lambda)| \leq M \|p(T)x\|$$

for every complex polynomial p . The set of all bounded point evaluations of T will be denoted by $B(T)$. Note that it follows from the Riesz Representation Theorem that $\lambda \in B(T)$ if and only if there is a unique vector denoted $k(\lambda) \in H$ such that $p(\lambda) = \langle p(T)x, k(\lambda) \rangle$ for every complex polynomial p . By setting $q(z) = (z - \lambda)p(z)$, we obtain $\lambda \in \overline{\sigma_p(T^*)}$. Conversely if $k(\lambda)$ is an eigenvector of T^* associated to the eigenvalue $\bar{\lambda}$, we get $\langle p(T)x, k(\lambda) \rangle = p(\lambda) \langle x, k(\lambda) \rangle$. Hence $\langle x, k(\lambda) \rangle \neq 0$ and $\lambda \in \sigma_p(T)$. Which gives

Proposition 1.1. $B(T) = \sigma_p(T) = \overline{\sigma_p(T^*)}$.

An open subset O of \mathbb{C} is said to be an **analytic set** for T if it is contained in $B(T)$ and the vector valued function $k : \lambda \mapsto k(\lambda)$ is analytic on O . The largest analytic set for T will be denoted by $B_a(T)$ and every point of it will be called **analytic bounded point evaluation** for T . Remark that $B_a(T)$ coincides with conjugate of the usual spectral residuum S_{T^*} of T^* given by complex numbers λ such that there exist an open set $V \ni \lambda$ and $f : V \rightarrow H$ an analytic function satisfying $(T^* - z)f(z) = 0 \quad z \in V$ (see [12])

Tavan Trent proved in [9] that for cyclic subnormal operators, we have

$$\sigma_p(T) \setminus \sigma_{ap}(T) = B_a(T) \quad (*)$$

L.R. Williams proved in [10] proposition 1.3 that for every arbitrary cyclic operator $T \in L(H)$, $\sigma_p(T) \setminus \sigma_{ap}(T) \subset B_a(T)$ and asked if (*) remain valid for arbitrary cyclic operators ([10] Question A).

In this paper, section 2, we first give necessary and sufficient conditions for weighted shifts to satisfy (*) and we exhibit an operator which provides a negative answer to the question of Williams. Then, in section 3, we devote attention to operators with fat local spectra. Furthermore, we generalize theorem 2.1 of [10] to the case of operators satisfying Dunford Condition C and with no eigenvalues. Finally, we give a simple proof of theorem 2.5 of [11] using Zero-principle for analytic functions.

2. ANALYTIC BOUNDED POINT EVALUATIONS FOR UNILATERAL WEIGHTED SHIFT

In [7], A.L. Shields represented a weighted shift operator as ordinary shift operator (that is, as "multiplication by z ") on a Hilbert space of formal power series (in the unilateral case) or formal Laurent series (in the bilateral case). He defined the concept of bounded point evaluations of a weighted shift to examine which power series and Laurent series represent analytic functions. In fact, this concept of bounded point evaluations for injective unilateral weighted shift coincides with the one defined above by Williams (see [10]).

We now describe the set of bounded point evaluations and the set of analytic bounded point evaluations for an arbitrary injective unilateral weighted shift. Let S be a unilateral weighted shift on a Hilbert space H with a positive weight sequence $(\omega_n)_{n \geq 0}$, that is

$$S e_n = \omega_n e_{n+1}.$$

where $(e_n)_{n \geq 0}$ is an orthonormal basis of H . Let β be the following sequence given by:

$$\beta_n = \begin{cases} \omega_0 \dots \omega_{n-1} & \text{if } n > 0 \\ 1 & \text{if } n = 0 \end{cases}$$

The unilateral weighted shift S is cyclic with cyclic vector e_0 . It follows from Corollary 2 of [7] and Proposition 1.4 of [10] that the sets $B(S)$ and $B_a(S)$ have a circular symmetry about the origin. Also, it follows from Theorem 8 of [7] and Proposition 1.1 that $B(S) = \{0\}$ if $r_2(S) = 0$ otherwise

$$\{\lambda \in \mathbb{C} : |\lambda| < r_2(S)\} \subset B(S) \subset \{\lambda \in \mathbb{C} : |\lambda| \leq r_2(S)\},$$

where $r_2(S) = \liminf_{n \rightarrow +\infty} (\beta_n)^{\frac{1}{n}}$.

Theorem 2.1. *If $r_2(S) > 0$ then $B_a(S) = \{\lambda \in \mathbb{C} : |\lambda| < r_2(S)\}$.*

Proof. Let $\lambda \in B(S)$ then there is $k(\lambda) \in H$ such that $p(\lambda) = \langle p(S)e_0, k(\lambda) \rangle$ for every polynomial p . For every $n \geq 0$ we have

$$\begin{aligned} \langle e_n, k(\lambda) \rangle &= \left\langle \frac{1}{\beta_n} S^n e_0, k(\lambda) \right\rangle \\ &= \frac{\lambda^n}{\beta_n}. \end{aligned}$$

Hence, $k(\lambda) = \sum_{n \geq 0} \frac{\lambda^n}{\beta_n} e_n$ for every $\lambda \in B(S)$. And so, the desired result holds.

For the weighted shift S , the spectrum $\sigma(S)$ is known to be the disk $\{\lambda \in \mathbb{C} : |\lambda| \leq r(S)\}$, and the approximate point spectrum $\sigma_{ap}(S)$ is known to be the annulus $\{\lambda \in \mathbb{C} : r_1(S) \leq |\lambda| \leq r(S)\}$ where $r_1(S) = \lim_{n \rightarrow \infty} \left[\inf_{k \geq 0} \frac{\beta_{n+k}}{\beta_k} \right]^{\frac{1}{n}}$ (see [7]). Therefore,

$$, (S) \setminus \sigma_{ap}(S) = \sigma(S) \setminus \sigma_{ap}(S) = \{\lambda \in \mathbb{C} : |\lambda| < r_1(S)\}.$$

Hence we have the following theorem.

Theorem 2.2. *Let S be a weighted shift. Then, $, (S) \setminus \sigma_{ap}(S) = B_a(S)$ if and only if $r_1(S) = r_2(S)$.*

A negative answer to the question A (see [10]) can be given by a unilateral weighted shift S for which $r_1(S) < r_2(S)$.

Let us consider an example of a such weighted shift. For $s \in \mathbb{N}$ there are unique $n, k \in \mathbb{N}$ such that $s = n! + k$ with $0 \leq k \leq (n+1)! - n! - 1$. We set

$$\beta_s = \beta_{n!+k} = e^k.$$

And so, for every $k \in \mathbb{N}$, we have

$$\frac{\beta_{k+1}}{\beta_k} = \begin{cases} e & \text{if } n! \leq k < (n+1)! - 1 \\ \frac{1}{e^{(n+1)! - n! - 1}} & \text{if } k = (n+1)! - 1 \end{cases}$$

Hence, the unilateral weighted shift S corresponding to the weight $(\frac{\beta_{n+1}}{\beta_n})_{n \geq 0}$ is bounded. For every $n \geq 2$, set $k_n = n! - n$. Clearly, we have $(n-1)! \leq k_n < n!$ and $\beta_{k_n} = e^{(n! - (n-1)! - n)}$. And so,

$$\inf_{k \geq 0} \frac{\beta_{n+k}}{\beta_k} \leq \frac{\beta_{n+k_n}}{\beta_{k_n}} = \frac{1}{e^{(n! - (n-1)! - n)}}.$$

Hence,

$$r_1(S) = \lim_{n \rightarrow \infty} \left[\inf_{k \geq 0} \frac{\beta_{n+k}}{\beta_k} \right]^{\frac{1}{n}} = 0.$$

On the other hand, it is clear that

$$r_2(S) = \liminf_{n \rightarrow \infty} \left[\beta_n \right]^{\frac{1}{n}} = 1.$$

Therefore, $B_a(S) = \{\lambda \in \mathbb{C} : |\lambda| < 1\}$ and $(S) \setminus \sigma_{ap}(S) = \emptyset$.

For the unilateral weighted shift S the set of its analytic bounded point evaluations is exactly the interior of the set of its bounded point evaluations. This suggests the following question, which was posed by J.B. Conway in [2] page 65 for arbitrary cyclic subnormal operator.

Question: Does the interior of $B(T)$ always coincide with $B_a(T)$ for an arbitrary cyclic operator $T \in L(H)$?

3. LOCAL SPECTRA OF CYCLIC OPERATORS WITH DUNFORD'S CONDITION C

Let T be a bounded operator on a complex Hilbert space H . For an element $x \in H$, let $\sigma_T(x)$ be its local spectrum and $\rho_T(x)$ be its local resolvent (see [1]). For a closed subset F of the complex plane \mathbb{C} , let $H_T(F)$ be the set of elements $x \in H$ such that $\sigma_T(x) \subset F$; it is a linear subspace of H . The operator T is said to have the **single valued extension property (svep)** if zero is the unique vector $x \in H$ such that $\sigma_T(x) = \emptyset$, which is equivalent to that for every open set $U \subset \mathbb{C}$, the only analytic solution of the equation $(T - \lambda I)f(\lambda) = 0$ for $\lambda \in U$ is the zero function $f \equiv 0$. The operator T is said to satisfy **Dunford's Condition C (DCC)** if for every closed subset F of \mathbb{C} , the linear subspace $H_T(F)$ is closed. It is known that every operator which satisfies DCC has the single valued extension property (see [1]). The operator T is said to be **hyponormal** if $\|T^*x\| \leq \|Tx\|$ for every $x \in H$; J.G. Stampfli [8] and M. Radjabalipour [4] have shown that hyponormal operators satisfy DCC. Recall that T is said to be pure if the only

reducing subspace M of T such that $T|_M$ is normal is $M = \{0\}$. In [10] Theorem 2.1, it is shown that for a pure cyclic hyponormal operator $T \in L(H)$ with cyclic vector x , $\sigma_T(Sx) = \sigma(T)$ for every operator $S \in L(H)$ such that $ST = TS$ and $\ker(S^*)$ is finite dimensional. We generalize this result as follows.

Theorem 3.1. *Let $T \in L(H)$ be a cyclic operator on H with cyclic vector $x \in H$ and let $S \in L(H)$ be an operator on H which commutes with T such that $\ker(S^*)$ is finite dimensional. If T satisfies DCC and $\sigma_p(T) = \emptyset$ then $\sigma_T(Sx) = \sigma(T)$.*

We prove this theorem using several lemmas. The following two lemmas are given in [10].

Lemma 3.2. *Let $T \in L(H)$ be a cyclic operator on H with a cyclic vector $x \in H$. If T satisfies DCC then $\sigma_T(x) = \sigma(T)$.*

Lemma 3.3. *Suppose that $H = H_1 \oplus H_2$ where H_1 and H_2 are two Hilbert spaces such that H_2 is finite dimensional. If T has the single valued extension property and H_1 is an invariant subspace for T then $A = T|_{H_1}$ has the single valued extension property and $\sigma_A(x) = \sigma_T(x)$ for every $x \in H_1$.*

Lemma 3.4. *Suppose that $H = H_1 \oplus H_2$ where H_1 and H_2 are two Hilbert spaces such that H_2 is finite dimensional. If H_1 is an invariant subspace for an operator T which satisfies DCC then the operator $A = T|_{H_1}$ satisfies the DCC.*

Proof. It follows from Lemma 3.3 that for every closed subset F of \mathbb{C} , $H_{1_A}(F) = H_T(F) \cap H_1$. So, the desired result holds.

Lemma 3.5. *Suppose that $H = H_1 \oplus H_2$ where H_1 and H_2 are two Hilbert spaces such that H_2 is finite dimensional. If $T \in L(H)$ is an operator without point spectrum such that H_1 is an invariant subspace for T then $\sigma(T) = \sigma(A)$ where $A = T|_{H_1}$.*

Proof. We first write

$$T = \begin{bmatrix} A & B \\ 0 & C \end{bmatrix}$$

with respect to the decomposition $H = H_1 \oplus H_2$. Now, suppose that A is invertible in $L(H_1)$ then $TH_1 = H_1$. And so, $H_1 \cap TH_2 = \{0\}$ since T one to one. Let $x_2 \in H_2$ such that $Cx_2 = 0$ then $Tx_2 = Bx_2 \in H_1 \cap TH_2$, hence C is one to one and so, by the finite-dimensionality, C is invertible. Therefore the operator T is invertible with inverse given by:

$$\begin{bmatrix} A^{-1} & -A^{-1}BC^{-1} \\ 0 & C^{-1} \end{bmatrix}$$

Thus $\sigma(T) \subset \sigma(A)$. Conversely, It is known that $\sigma(A) = \bigcup_{x \in H_1} \sigma_A(x)$ (see [3] Theorem 1.9), it follows from Lemma 3.3 that $\sigma(A) = \bigcup_{x \in H_1} \sigma_T(x) \subset \sigma(T)$.

Proof of Theorem 3.1. Let H_1 be the closed linear subspace generated by $\{T^n Sx : n \geq 0\}$. Then it is clear that H_1 is an invariant subspace for T . Since $TS = ST$ and x is a cyclic vector for

T , then $H_1 = cl(Im(S))$. Therefore $H = H_1 \oplus H_2$ where $H_2 = \ker(S^*)$. It follows from Lemma 3.3 that $\sigma_T(Sx) = \sigma_A(Sx)$ where $A = T|_{H_1}$. Since A is a cyclic operator with cyclic vector Sx and satisfies DCC (Lemma 3.4) then it follows from Lemma 3.2 that $\sigma_A(Sx) = \sigma(A)$. Since $\sigma_p(T) = \emptyset$, it follows from Lemma 3.5 that $\sigma(T) = \sigma(A)$. And so, $\sigma(T) = \sigma(A) = \sigma_A(Sx) = \sigma_T(Sx)$. The proof is complete.

Remark 3.6. *Let $T \in L(H)$ be a cyclic operator with cyclic vector $x \in H$ and satisfies DCC such that $\sigma_p(T) = \emptyset$. For every non-zero polynomial p , $\sigma_T(p(T)x) = \sigma(T)$. Therefore, $\sigma_T(y) = \sigma(T)$ holds for all y in a dense subset of H . L.R. Williams proved in Theorem 2.5 of [11] that if T is a nonnormal hyponormal (unilateral or bilateral) weighted shift operator then $\sigma_T(x) = \sigma(T)$ for every non-zero element $x \in H$.*

We state and give a simple proof of Theorem 2.5 of [11] using the fact that a non-zero analytic function has isolated zeros.

Theorem 3.7. *Let T be a nonnormal hyponormal weighted shift on H . Then for every a non-zero element $x \in H$, $\sigma_T(x) = \sigma(T)$.*

Proof. First suppose that T is a nonnormal hyponormal unilateral weighted shift, then $r(T) = r_1(T) = r_2(T) = \|T\| > 0$ and by Theorem 2.1 we have $B_a(T) = int(\sigma(T))$. Now, let $x \in H$ such that there is $\lambda \in \sigma(T) \setminus \sigma_T(x)$. So, there is vector valued analytic function f on an open neighbourhood V of λ such that

$$(T - \mu I)f(\mu) = x \text{ for every } \mu \in V.$$

Since $\emptyset \neq V \cap int(\sigma(T)) \subset B_a(T)$, then for every $\mu \in V \cap int(\sigma(T))$ we have,

$$\begin{aligned} \hat{x}(\mu) &:= \langle x, k(\mu) \rangle \\ &= \langle (T - \mu I)f(\mu), k(\mu) \rangle \\ &= \langle f(\mu), (T - \mu I)^* k(\mu) \rangle \\ &= 0. \end{aligned}$$

Hence, $\hat{x} \equiv 0$. And so, $x = 0$.

The case of a nonnormal hyponormal bilateral weighted shift is similar using the bounded point evaluations in the sense of A.L. Shields [7].

Remark 3.8. *Using the same proof of this Theorem, one can see that for every injective unilateral weighted shift T , $cl(B_a(T)) \subset \sigma_T(x)$ for every non-zero element $x \in H$. The same holds for every bilateral weighted shift.*

Acknowledgments. This work has been partially supported by the Associate Scheme of the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy. The first and last authors are grateful to the mathematics section of the Abdus Salam ICTP.

REFERENCES

- [1] **I. Colojoara and C. Foias**, *Theory of Generalized Spectral Operators*, Gordon and Breach, New York 1968.
- [2] **J.B. Conway**, *The Theory of Subnormal Operators*, volume 36 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, R.I 1991.
- [3] **I. Erdelyi and R. Lange**, *Spectral Decomposition on Banach Spaces*, Lecture notes in Mathematics, Springer Verlag New York, V. 623 (1977).
- [4] **M. Radjabalipour**, *Ranges of Hyponormal Operators*, Illinois J. Math., 21(1977) 70-75.
- [5] **M. Raphael**, *Quasimilarity and Essential Spectra for Subnormal Operators*, Indiana Univ. Math. J., Vol 31. No.
- [6] **W.C. Ridge**, *Approximate Point Spectrum of a Weighted Shift*, Trans. Amer. Math. Soc. 147(1970) 349-356.
- [7] **A.L. Shields**, *Weighted Shift Operators and Analytic Function Theory*, in Topics in Operator Theory, Mathematical Surveys, N0 13 (ed. C. Pearcy), pp. 49-128. American Mathematical Society, Providence, Rhode Island 1974.
- [8] **J.G. Stampfli**, *A Local Spectral Theory for Operators. V: Spectral Subspaces for Hyponormal Operators*, Trans. Amer. Math. Soc., 21(1976) 285-296.
- [9] **T.T. Trent**, *$H^2(\mu)$ Spaces and Bounded Point Evaluations*, Pac. J. Math., 80(1979) 279-292.
- [10] **L.R. Williams**, *Bounded Point Evaluations and Spectra of Cyclic Hyponormal Operators*, Dynamic Systems and Applications 3(1994) 103-112.
- [11] **L.R. Williams**, *The Local Spectra of Pure Quasinormal Operators*, J. Math Anal. Appl. 187(1994) 842-850.
- [12] **F.H. Vasilescu**, *Analytic Functional Calculus and Spectral Decomposition*, Editura Academiei 1982.