

Electroweak fits at LEP

E. Lançon^{a*}

^aDAPNIA/SPP, CEA-Saclay,
91191 Gif-sur-Yvette Cedex, France

High precision electroweak measurements performed over ten years at LEP and SLC have allowed to constrain the Standard Model of electroweak interactions. The model have been used to predict the mass of the top quark and to set limits on the mass of the Higgs boson.

1. Why electroweak fits?

Electroweak fits have been extensively used at LEP and SLC for testing the Standard Model at the level of its quantum corrections, searching for deviations that may signal the presence of new physics. They have also successively predicted the top quark mass prior to its discovery and recently, allowed to set limit on the Higgs boson mass. Sensitivity to top quark and Higgs boson masses can be inferred from the following exemple relation :

$$m_W^2 \left(1 - \frac{m_W^2}{m_Z^2}\right) = \frac{\pi\alpha}{\sqrt{2}G_F} (1 + \Delta r)$$

where Δr contains several terms: $\Delta\alpha$ (due to light fermion masses), a negative term (several %) proportional to m_t^2 and a relatively small (below 1%) term proportional to $\ln(m_H)$. Therefore, the inferred constraints on m_H are much weaker than those on m_t .

Before LEP startup, the W and Z boson masses were measured to be 80.000 ± 0.360 GeV and 91.120 ± 0.160 GeV respectively. These measurements were equivalent to a precision on the weak mixing angle $\sin^2\theta_W$ of 3.7% worse than the direct measurement itself: 0.227 ± 0.006 corresponding to 2.6%. That is the reason why inputs to the Standard Model fits were α , G_F and $\sin^2\theta_W$. In 1995, at the end

*Presented at the Siena Conference, The Legacy of LEP and SLC, October 8-11 2001, on behalf of the LEP/SLD Collaborations and of the LEP Electroweak Working Group.

of LEP-I, the Z boson mass was known with an impressive precision of 2.210^{-5} hence replacing $\sin^2\theta_W$ as input parameter to electroweak fits. The Fermi constant G_F is determined from the μ lifetime, $G_F = 1.16637(1) \cdot 10^{-5} \text{GeV}^{-2}$ [1]. The relative error of G_F is comparable to that of m_Z ; both errors have negligible effects on the electroweak fit results.

More details can be found in the 2001 summary of the Electroweak Working Group [2].

1.1. The top quark mass

Many of the observables measured at LEP-I and SLC have sensitivity to the top quark mass (typically 20 GeV precision for the measurements of R_b , Γ_Z and $\sin^2\theta_W$), allowing for an impressive precision on the prediction of the top quark mass as the amount of data collected increased, as shown in Figure 1. In 2001, the top quark mass is measured to be 174.3 ± 5.1 GeV [3] and the electroweak fit yields 181_{-9}^{+11} GeV.

1.2. After m_t ?

Once the top quark mass is experimentally known, it is used to further constrain the electroweak fits allowing for predictions of m_W and of m_H . The first sensitivity curve to the Higgs boson mass (shown as a χ^2 curve on Figure 2) was published in the 1994 report of the Electroweak Working Group [4].




Figure 1. Time evolution of the top quark mass derived from electroweak fits, compared to direct searches results.

2. A word about a 3.3σ effect

At the time of this note and still for some time, it is difficult to present electroweak fit results without mentioning the so called “ 3.3σ ” effect.

The asymmetry measurements from LEP and SLD are combined into a single parameter, the effective electroweak mixing angle, $\sin^2 \theta_{\text{eff}}^{\text{lept}}$, defined as:

$$\sin^2 \theta_{\text{eff}}^{\text{lept}} \equiv \frac{1}{4} \left(1 - \frac{g_{V\ell}}{g_{A\ell}} \right),$$

without making strong model-specific assumptions. In Figure 3 are summarised the different measurements. As it can be seen, the combinations based on the leptonic results plus \mathcal{A}_ℓ (SLD) and on the hadronic forward-backward asymmetries differ by 3.3 standard deviations, mainly caused by the two most precise measurements of $\sin^2 \theta_{\text{eff}}^{\text{lept}}$, \mathcal{A}_ℓ (SLD) dominated by A_{LR} , and $A_{\text{FB}}^{0,b}$ (LEP). The averages of the two group of measurements are indicated in Table 1. The average of the six measurements leads to $\sin^2 \theta_{\text{eff}}^{\text{lept}} = 0.23152 \pm 0.00017$ with a χ^2 probability of 2.5%. The lepton based average prefers a low m_H value (below 100 GeV), while the hadronic based aver-




Figure 2. First $\Delta\chi^2 = \chi^2 - \chi_{\text{min}}^2$ vs m_H fit results plot including the CDF value of m_t .

age prefers a high m_H . Before any further considerations, one should keep in mind that 3σ effects exist in nature. Figure 4 shows the evolution of $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ determine at LEP from $A_{\text{FB}}^{0,b}$ (right) and SLD from \mathcal{A}_ℓ (left) as over seven years. As it can be seen, the significance of the discrepancy does not increase with time as the precision of the measurements does, suggesting a statistical fluctuation nature of the difference.

Table 1

Determinations of $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ from asymmetries.

	$\sin^2 \theta_{\text{eff}}^{\text{lept}}$
$A_{\text{FB}}^{0,\ell}, \mathcal{A}_\ell (\mathcal{P}_\tau), \mathcal{A}_\ell$ (SLD)	0.23113 ± 0.00021
$A_{\text{FB}}^{0,b}, A_{\text{FB}}^{0,c}, \langle Q_{\text{FB}} \rangle$	0.23230 ± 0.00029

3. Measurements

On figure 5, the various inputs to the electroweak fits are summarised, also shown the pulls (difference between measurement and fit in units of the total measurement error) of the various measurements. The pulls are derived from the Standard Model fit including all data with the Higgs mass treated as a free parameter. As it can be seen the largest pull (-2.9) is for $A_{\text{FB}}^{0,\ell}$ (reflecting the fact the it is the most precise measurement preferring a high m_H value). The pull distribution from all measurements follows a Gaussian law.




Figure 3. Comparison of several determinations of $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ from asymmetries. Also shown is the prediction of the Standard Model as a function of m_H . The width of the Standard Model band is due to the uncertainties in $\Delta\alpha_{\text{had}}^{(5)}$, m_Z and m_t . The total width of the band is the linear sum of these effects.

4. Theoretical and Parametric Uncertainties

Detailed studies of the theoretical uncertainties in the Standard Model predictions due to missing higher-order electroweak corrections and their interplay with QCD corrections can be found in [5].

The recently calculated complete fermionic two-loop corrections on m_W [6] are currently only used in the determination of the theoretical uncertainty. Their effect on m_W is small compared to the current experimental uncertainty on m_W , however, the naive propagation of this new m_W to $\sin^2 \theta_{\text{eff}}^{\text{lept}} = \kappa(1 - m_W^2/m_Z^2)$, keeping the electroweak form-factor κ unmodified, shows a more visible effect as $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ is measured very precisely. Thus the corresponding calculations for $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ (or κ) and for the partial Z widths are urgently needed; in particular since partial can-




Figure 4. Evolution of $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ determine at LEP from $A_{\text{FB}}^{0,b}$ (right) and SLD from \mathcal{A}_ℓ (left) as over seven years.

cellations of these new corrections in the product $\kappa(1 - m_W^2/m_Z^2) = \sin^2 \theta_{\text{eff}}^{\text{lept}}$ are expected [7].

The use of the new QCD corrections[8] increases the value of $\alpha_S(m_Z^2)$ by 0.001, as expected. The effects of missing higher-order QCD corrections on $\alpha_S(m_Z^2)$ covers missing higher-order electroweak corrections and uncertainties in the interplay of electroweak and QCD corrections and is estimated to be at least 0.002 [9]. The determination of the size of remaining theoretical uncertainties is under continued study.

At present the impact of theoretical uncertainties on the determination of Standard Model parameters from the precise electroweak measurements is small compared to the error due to the uncertainty in the value of $\alpha(m_Z^2)$, which is included in the results.

The uncertainty in $\alpha(m_Z^2)$ arises from the contribution of light quarks to the photon vacuum




Figure 5. Summary of measurements that enter the electroweak fits together with their pull.

polarisation ($\Delta\alpha_{\text{had}}^{(5)}(m_Z^2)$):

$$\alpha(m_Z^2) = \frac{\alpha(0)}{1 - \Delta\alpha_\ell(m_Z^2) - \Delta\alpha_{\text{had}}^{(5)}(m_Z^2) - \Delta\alpha_{\text{top}}(m_Z^2)},$$

where $\alpha(0) = 1/137.036$. The top contribution, $-0.00007(1)$, depends on the mass of the top quark, and is therefore determined inside the electroweak libraries. The leptonic contribution is calculated to third order[10] to be 0.03150, with negligible uncertainty.

The new evaluation of the hadronic contribution 0.02761 ± 0.00036 which takes into account the recently published results on electron-positron annihilations into hadrons at low centre-of-mass energies by the BES collaboration [11] is used. The uncertainty translates into an error of 0.00013 on the Standard Model prediction of $\sin^2\theta_{\text{eff}}^{\text{lept}}$, and errors of 0.2 GeV and 0.1 on the fitted values of m_t and $\log(m_H)$, all included in the results presented.

There are also several evaluations of $\Delta\alpha_{\text{had}}^{(5)}(m_Z^2)$ [12–19] which are more theory-driven. One of the most recent ones (Reference [18]) also includes the new results from BES, yielding 0.02738 ± 0.00020 . To show the effects of the uncertainty of $\alpha(m_Z^2)$, this evaluation of the hadronic vacuum polarisation is also used.

5. Fit results

Most observables have sensitivity to the Higgs boson mass as it can be seen in Figure 6. The most sensitive measurements are the asymmetries, *i.e.*, $\sin^2\theta_{\text{eff}}^{\text{lept}}$ and the W mass. A reduced uncertainty for the value of $\alpha(m_Z^2)$ would therefore result in an improved constraint on $\log m_H$ and thus m_H .




Figure 6. Constraints on $\log m_H$ from various observables.

Several different Standard Model fits to the data are discussed. The predictions are calculated with TOPAZ0 [20] and ZFITTER [21].

The data can first be used to determine the top quark and W masses indirectly, which can be compared to the direct measurements performed at the $p\bar{p}$ colliders and LEP-II, all the results ex-

cept the LEP-II and $p\bar{p}$ colliders m_W and m_t results are used. The indirect measurements of m_W and m_t are shown in Figure 7, compared with the direct measurements. Also shown are the Standard Model predictions for Higgs masses between 114 and 1000 GeV. As can be seen in the figure, the indirect and direct measurements of m_W and m_t are in good agreement, and both sets prefer a low value of the Higgs mass




Figure 7. The comparison of the indirect measurements of m_W and m_t (solid contour) and the direct measurements (dashed contour). In both cases the 68% CL contours are plotted. Also shown is the Standard Model relationship for the masses as a function of the Higgs mass.

From a fit to all data except m_W , the best indirect determination of m_W is obtained. The indirect determination of W boson mass 80.373 ± 0.023 GeV is in agreement with the combination of direct measurements from LEP-II and $p\bar{p}$ colliders [22] of $m_W = 80.451 \pm 0.033$ GeV.

Similarly, the indirect determination of the top quark mass : $m_t = 181_{-9}^{+11}$ GeV, is in very good agreement with the direct measurement of $m_t = 174.3 \pm 5.1$ GeV.

Finally, the best constraints on m_H are obtained when all data are used in the fit. In Figure 8 the observed value of $\Delta\chi^2 \equiv \chi^2 - \chi_{\min}^2$ as a function of m_H is plotted for the fit including all data. The χ^2 per degree of freedom of the fit is 23/15 corresponding to a probability of 8%. The solid curve is the result using ZFITTER. The result is $\log(m_H/\text{GeV}) = 1.94 \pm 0.21$, corresponding to $m_H = 88_{-35}^{+53}$ GeV. The shaded band represents the uncertainty due to uncalculated higher-order corrections, as estimated by ZFITTER and TOPAZ0. Compared to previous analyses, its width is enlarged towards lower Higgs-boson masses due to the effects of the complete fermionic two-loop calculation of m_W discussed above. The 95% confidence level upper




Figure 8. $\Delta\chi^2 = \chi^2 - \chi_{\min}^2$ vs. m_H curve. The line is the result of the fit using all data; the band represents an estimate of the theoretical error due to missing higher order corrections. The vertical band shows the 95% CL exclusion limit on m_H from the direct search.

limit on m_H (taking the band into account) is 196 GeV. The lower limit on m_H of approximately 114 GeV obtained from direct searches[23] is not used

in the determination of this limit. Also shown is the result (dashed curve) obtained when using $\Delta\alpha_{\text{had}}^{(5)}(m_Z^2)$ of Reference [18]. That fit results in $\log(m_H/\text{GeV}) = 2.03 \pm 0.19$ corresponding to $m_H = 106_{-38}^{+57}$ GeV and an upper limit on m_H of approximately 222 GeV at 95% confidence level.

The variation of the upper limit on m_H against experimental parametric errors is tested in Figure 9, where $\Delta\chi^2$ curves corresponding to one sigma variation of $\Delta\alpha_{\text{had}}^{(5)}(m_Z^2)$ and m_t are plotted. Although the position of the minimum changes because of the relative changes of the different measurement in the fit, the upper limit on m_H does not increase above 300 GeV.




Figure 9. $\Delta\chi^2$ for one sigma change of $\Delta\alpha_{\text{had}}^{(5)}(m_Z^2)$ or m_t and both.

6. Conclusions and Perspectives

The measurement of m_W at LEP-II is likely to reach a precision of ≈ 25 MeV, not too far from the uncertainty on the prediction obtained via the radiative corrections of the Z data, providing a further important test of the Standard Model. As the LEP energy final value is not yet known, the absolute m_W mass value from LEP might change. However to further, significantly, con-

strain m_H a more accurate measurement of m_t is mandatory. A precision of 2 GeV is likely to be obtained at FERMILAB run II. Figure 10 shows the $\Delta\chi^2$ curves that would be obtained with such precisions, without any change in central values of measurements. A precision of about 20 GeV on m_H is reachable.




Figure 10. $\Delta\chi^2$ curves for a precision of 25 MeV on m_W and 2 GeV on m_t . Central values of m_W and m_t are kept unchanged.

Figure 11 shows the evolution of both the lower limit on the Higgs boson mass from direct searches (hatched area) and the upper limit from electroweak fits (dashed area) as a function of time. The gap is not closed, a slightly longer run for LEP with a bit higher energy might have closed it.

Electroweak fits from LEP and SLC data have allowed to test the internal consistency of the Standard Model with great precision, three to five times better than anticipated. The top quark mass was predicted several years before it has been discovered. The measurements led to the prediction of a relatively light Higgs boson (around 100 GeV), with the same precision ($\approx 50\%$) as on the top quark mass before LEP and SLC startup.




Figure 11. Variation over seven years of the lower limit on the Higgs boson mass from direct searches (dashed area) compared to the upper limit derived from electroweak fits (shaded area).

REFERENCES

1. T. van Ritbergen and R.G. Stuart, Phys. Rev. Lett. **82** (1999) 488.
2. The LEP Collaborations ALEPH, DELPHI, L3 and OPAL, the LEP Electroweak Working Group, the SLD Heavy Flavour Group and Electroweak Working Group, *A Combination of Preliminary Electroweak Measurements and Constraints on the Standard Model*, hep-ex/0112021.
3. The Top Averaging Group, L. Demortier *et al.*, for the CDF and DØ Collaborations, FERMILAB-TM-2084 (1999).
4. The LEP Collaborations ALEPH, DELPHI, L3, OPAL and the LEP Electroweak Working Group, *Combined Preliminary Data on Z Parameters from the LEP Experiments and Constraints on the Standard Model*, CERN-PPE/94-187.
5. D. Bardin, M. Grünewald and G. Passarino, *Precision Calculation Project Report*, hep-ph/9902452.
6. A. Freitas, W. Hollik, W. Walter and G. Weiglein, Phys. Lett. **B495** (2000) 338.
7. D. Bardin, P. Gambino, G. Passarino, G. Weiglein, private communication, spring 2001.
8. A. Czarnecki and J. Kühn, Phys. Rev. Lett. **77** (1996) 3955;
- R. Harlander, T. Seidensticker and M. Steinhauser, Phys. Lett. **B426** (1998) 125.
9. T. Hebbeker, M. Martinez, G. Passarino and G. Quast, Phys. Lett. **B331** (1994) 165; P.A. Raczka and A. Szymacha, Phys. Rev. **D54** (1996) 3073; D.E. Soper and L.R. Surguladze, Phys. Rev. **D54** (1996) 4566.
10. M. Steinhauser, Phys. Lett. **B429** (1998) 158.
11. The BES Collaboration, J.Z. Bai *et al.*, *Measurements of the Cross Section for $e^+e^- \rightarrow$ hadrons at Center-of-Mass Energies from 2 to 5 GeV*, hep-ex/0102003.
12. M. L. Swartz, Phys. Rev. **D53** (1996) 5268.
13. A.D. Martin and D. Zeppenfeld, Phys. Lett. **B345** (1994) 558.
14. R. Alemany, *et al.*, Eur. Phys. J. **C2** (1998) 123.
15. M. Davier and A. Höcker, Phys. Lett. **B419** (1998) 419.
16. J.H. Kühn and M. Steinhauser, Phys. Lett. **B437** (1998) 425.
17. J. Erler, Phys. Rev. **D59**, (1999) 054008.
18. A. D. Martin, J. Outhwaite and M. G. Ryskin, Phys. Lett. **B492** (2000) 69.
19. F. Jegerlehner, *Hadronic Effects in (g-2) and $\alpha_{QED}(M_Z)$: Status and Perspectives*, Proc. of Int. Symp. on Radiative Corrections, Barcelona, Sept. 1998, page 75.
20. G. Montagna *et al.*, Comput. Phys. Commun. **117** (1999) 278; <http://www.to.infn.it/~giampier/topaz0.html> .
21. D. Bardin *et al.*, Z. Phys. **C44** (1989) 493; Comp. Phys. Comm. **59** (1990) 303; Nucl. Phys. **B351**(1991) 1; Phys. Lett. **B255** (1991) 290 and CERN-TH 6443/92 (May 1992); the most recent version of ZFIT-TER (6.21) is described in DESY 99-070, hep-ph/9908433 (Aug 1999) published in Comp. Phys. Comm. **133** (2001) 229.
22. http://www-cdf.fnal.gov/physics/ewk/wmass_new.htm.
23. P. Igo-Kemenes, these proceedings.