

Influence of dioxin intoxication on the human system and possibilities of limiting its negative effects on the environment and living organisms

Ireneusz Całkosiński¹, Joanna Rosińczuk-Tonderys¹, Justyna Bazan¹, Maciej Dobrzyński², Agnieszka Bronowicka-Szydełko³, Katarzyna Dzierżba³

¹ Department of Nervous System Diseases, Medical University, Wrocław, Poland

² Department of Conservative Dentistry and Pedodontics, Medical University, Wrocław, Poland

³ Department of Medical Biochemistry, Medical University in Wrocław, Poland

Całkosiński I, Rosińczuk-Tonderys J, Bazan J, Dobrzyński M, Bronowicka-Szydełko A, Dzierżba K. Influence of dioxin intoxication on the human system and possibilities of limiting its negative effects on the environment and living organisms. *Ann Agric Environ Med*. 2014; 21(3): 518–524. doi: 10.5604/12321966.1120594

Abstract

Introduction and objective. Despite the restrictive legal regulations related to the reduction of dioxins emission, their concentration in the environment is still too high. Mainly, this is related to the illegal utilisation of electronic equipment and combustion of wastes, and also to intensified activity and maintenance of ships, especially in developing countries. The most important remaining source in Europe is the metal industry. Studies on the mechanism of impact of dioxins are still being carried out. This review points at new possibilities for limiting the molecular mechanisms of dioxins activity, *inter alia*, through the application of high doses of tocopherol and acetylsalicylic acid while treating dioxins intoxication.

Brief description of the state of knowledge. Apart from the knowledge of dioxins affinity to the aryl hydrocarbon receptor (AhR), the multi-stage radical-form actions and the pro-inflammatory mechanism associated with cyclooxygenase-II enzyme (COX-2) are under intense investigation at the moment. Due to the high affinity of dioxins to animals adipose tissue and their ability to accumulate in it, they can enter the food chain. Furthermore, high dioxin doses can cause poisoning manifested as advanced clinical symptoms, whereas in smaller doses, when cumulated, can cause metabolic changes which are often difficult to associate with their presence. Recently, some serious food contaminations by dioxins have been demonstrated. Sea fish and products from contaminated aqueducts still constitute potential sources of dioxins pollution.

Conclusion. According to recent studies, dioxins are present in different concentrations in the environment and cause specific and long-time effects. These effects could be limited by the use of tocopherol and acetylsalicylic acid.

Key words

aryl hydrocarbon receptor (AhR), dioxin, inflammatory, persistent organic pollutants, 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (TCDD), tocopherol

INTRODUCTION

Dioxins organic pollutants belong to the group of persistent organic pollutants (POPs), the emission of which into the environment has been associated for many years with ecological and health problems. Although POPs are characterized by specific and known chemical and physical properties and are easily detected, their analysis (in both the environment and food) is still difficult and requires sophisticated equipment. Currently, their emission into the environment is limited by the provisions of The Stockholm Convention on Persistent Organic Pollutants of 22 May 2001, and the national legislation established by the regulatory authorities, agencies, or international organizations, such as the World Health Organization (WHO), United States Environmental Protection Agency (EPA), the European Food Safe Authority (EFSA) or the European Chemicals Agency (ECHA). Despite the rigorous limits in POPs emission established in legislation documents, some sources of POPs emission into the environment are still a huge problem, e.g. private waste combustion. Usually, they pollute soil and seas.

Although the impact of dioxins on the organism is inevitable, it can be limited. Unfortunately, people are not

aware that dioxins are produced while grilling, backyard barrel burning, or using some central heating systems, and how harmful these may be for the environment and every living organism. The economic and energy crisis also has a stimulating effect on this production.

Objective. This review focuses on determination of the influence of dioxins toxicity on the biochemical changes in the organism, and demonstrates the possibilities for reducing the negative effects of exposure to dioxin.

DESCRIPTION OF THE STATE OF KNOWLEDGE

The threat of dioxin poisoning. The organic pollutants group of POPs is divided into 12 groups of chemicals. One of them is polychlorinated dibenzo-*p*-dioxin (PCDDs), polychlorinated dibenzofurans (PCDFs) and polychlorinated biphenyls (PCBs) (Fig. 1). Polychlorinated dibenzo-*p*-dioxin (PCDDs), determined as dioxins, are a group of 75 compounds which contain benzene rings with varying numbers of chlorine atoms (Cl_x, Cl_y) in different positions of these rings. They can form congeners with varying numbers and localizations of chlorine atoms in the benzene frame. The 2, 3, 7, 8-tetrachlorodibenzo-*p*-dioxin (TCDD) is the standard compound of PCDDs.

According to epidemiological studies, dioxins are considered as cancerogenic factors, [2], defined by the

Address for correspondence: Justyna Bazan; Department of Nervous System Diseases, Medical University, Bartla 5, 51-618 Wrocław, Poland, tel. +48 71 784 18 27; e-mail: justyna.bazan@umed.wroc.pl;

Received: 04 January 2013; accepted: 06 May 2013






Figure 1. Structure of the individual groups POP representatives a) polychlorinated biphenyls (PCB), b) polychlorinated dibenzofurans (PCDF), c) polychlorinated dibenzodioxins (PCDD), d) 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD). On the basis of [82]

International Agency for Research on Cancer (IARC) as a group 1 carcinogenic compound of group 1, and their activity is still being investigated by experiments on animals [1]. Nevertheless, there are some reports that criticize the IARC classification of dioxins as proven human carcinogens [3].

The main dioxin and the best-examined compound is the TCDD which has a very high toxicity; it is therefore considered to be a reference for determining the toxicity of other toxic compounds of this group [4, 5]. The toxic equivalent factor determines TCDD toxicity and has a value of 1 [6]. The Total Toxic Equivalent (TEQ) presents an estimated value of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD)-like activity of the mixture by the sum of the individual PCDDs and PCDFs congeners in the sample, multiplied by the corresponding toxic equivalent factors (TEF). TEQ and TEF value indices were established and then systematically re-evaluated by the WHO to uniformity at the international level and provide recommendations for particular countries [7].

Indeed, the limitation of POPs emission that includes PCDD/PCDF (present in different proportions and having different toxicity) is regulated by the provisions of The Stockholm Convention on Persistent Organic Pollutants. According to this report, the main sources of POPs emissions are: open waste burning, including the burning of landfill sites and thermal processes in the metallurgical industry, specific chemical production processes releasing unintentionally formed persistent organic pollutants, and most of all, the production of chlorophenols and chloranil [8].

Dioxins are formed in every thermal process and are characterized by the presence of the organic matter, chlorine, high temperature (200–900 °C) and low concentration of oxygen. Dioxins are emitted not only during combustions and waste utilization, but also in various technological processes, such as textile and leather dyeing, oil refineries wastes, destruction of animal carcasses and products of the pesticide industry [8, 9]. It is noteworthy that dioxins may be converted from polychlorinated biphenyls (PCBs) or may be a result of sunshine interaction with plastic waste. Dioxins and PCBs are chemically similar and belong to the same group of POPs [10, 11]. Moreover, conflagrations are considered as one of the essential sources of PCBs and PCDD congeners emissions. The measurements in the conflagration region of World Trade Center (WTC) after the terrorist attack in 2001, showed a high concentration of polycyclic aromatic hydrocarbon, not only in the ground zero zone,

but also at a significant distance from it [12, 13, 14]. These observations have led to the conclusion that the risk of the tumour morbidity as a long-term effect of dioxin exposition in the population living near the WTC is high [14].

Interestingly, the research carried out on areas where, 30 years earlier, military operations using dioxins and herbicides were conducted, have shown a high elevation of PCDF concentration in the soil. These observations are important for indicating the correlation between food production in areas contaminated by polycyclic aromatic hydrocarbon and the capacity to deposition of these compounds in the food chain [15] (Fig. 2). It has been reported that dioxins are considered as xenobiotics with a hormone-like action. Their concentration, as a result of the utilization processes and individual heat production, is significant, especially in the urban environment [15, 16, 17].




Figure 2. Circulation of dioxins in food chain [83]

It is worth highlighting that prolonged, strong, biological actions of dioxins can be manifested in the second generation, even though only the first generation was exposed to this activity [18, 19].

Dioxin absorption. The important routes of dioxin entry into the organism include the digestion system, respiratory system and the skin. Dioxins have a high affinity to fats and can dissolve in them; thus, they can be delivered and stored in the organism in a simple way [5, 20].

Almost 90% of human exposure to dioxins is from food. These toxins appear in different food chains (Fig. 2) and are absorbed into the organism by the digestion system. For instance, dioxin accumulated in seals could be transferred to the Eskimos from Canada and Greenland, who eat these mammals. Moreover, studies of the metabolism of seals

living near the coast of Greenland have shown a correlation between the age of these mammals and dioxins concentration in their bodies [20]. What is more, a significantly higher concentration of PCDD/PCDF is present in the milk of Norwegian bears than in the milk of seals living in this area, suggesting that the amount of dioxins in the adipose tissue of the final consumers is directly proportional to the lifespan in the dioxins contaminated environment [21]. Additionally, consumption of high-fat foods containing dioxins results in dioxin accumulation in adipose tissue [1, 16, 20].

Although the air is contaminated by xenobiotic absorbing smokes and dusts, the absorption of dioxin by the organism through the respiratory system is insignificant (ca. 8%) and the absorption of dioxins by the skin lipids is even lower (ca. 2%). It should be noted that many different factors, for polluted air, such as soot are absorbed by the respiratory system and skin lipids, and they cause a significant rise of the dioxin level in an organism [17]. The knowledge of dioxin emission sources may help to distinguish occupational groups, such as chimney sweeps, firefighters, soldiers, members of Special Forces and anti-terrorist squads, for whom the risk of exposure to PCDD/PCDF is the greatest. It should be emphasized that the urea of chimney sweeps contains aromatic polychlorinated compounds, which are derivatives of POPs [22, 23].

Furthermore, employees of certain industries, as well as residents of industrialized zones, are also exposed to dioxins. The contamination of the population of Seveso in Italy is a prime example of this problem [24, 25, 26, 27].

Dioxin sensitivity. Due to their physical properties, i.e. no taste or smell and solubility in fats, dioxins may easily enter the digestion system. Because of these organoleptic features and the fact that the clinical symptoms of intoxication occur even after a long-time period, dioxins were considered as a potential chemical weapon. Studies conducted 20 and 30 years after the first exposure to these compounds have shown that their concentration in plasma lipids increases with age, since these compounds accumulate when the exposure is continuous. On the basis of these results, it was suggested that dioxin concentration is correlated with the body mass index (BMI). Due to the emerging dominance of catabolic processes in the body that appear with ageing or starvation, and because dioxin half-life is long, these compounds accumulated in the adipose tissue can be released into the plasma a long time after the first exposure [22, 24, 25].

So far, research on dioxins has been conducted only on a few animal models. The species most sensitive to TCDD activity, determined by the LD50 value, is the guinea pig (*Cavia porcellus*) [28]. Interestingly, rats and monkeys are 45 times more resistant to TCDD activity than the guinea pig. This species specificity is a result of changes in the regulation of gene expression, especially those responsible for the lipid metabolism [29].

The sensitivity to dioxins activity depends on gender, e.g. female rats are more sensitive to them than the male. The TCDD toxicity determined by LD50 for the female is between 22–50 µg/kg b.w., and for male rats it oscillates between 45–50 µg/kg b.w [30, 31]. Recent studies have shown that administration of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin at a dose of 25 µg/kg b.w in Buffalo strain rat females, causes their death on the third day after TCDD application, accompanied by liver damage and blood extravasation into the bowel.

Reducing the dose by half did not cause have these effects [32]. In connection with these studies, the need for the correction of toxicity studies in different animal species has been highlighted [31].

Reducing dioxin activity. Dioxins are transported from the external environment to the adipose tissue or liver through the plasma lipids, and TCDD binds to the aryl hydrocarbon receptor (AhR) in cytosol of hepatocytes [33–36].

The active dioxin-receptor complex is translocated to the cell nucleus and binds to the dioxin response element on DNA (dioxin responsive element, DRE). This complex is a signal for transcription of cytochromes P 450 and P 448 (aryl hydrocarbon hydroxylase – AHH) genes [37, 38]. The highest concentration of AHH, the enzyme responsible for dioxin conversions, occurs in endoplasmic reticulum of hepatocyte. In the initial phase, the hydrolytic degradation of dioxins by monooxygenase occurs and then the products of this reaction are conjugated with glucuronic acid or the glutathione molecule (GSH). After that, these complexes are released from organisms with urea or bile. Dioxins are metabolized more slowly than other high affinity AhR agonists, e.g. 6-formylindolo[3,2-b]carbazole (FICZ). For this reason, their action is much more prolonged in time and is not self-limited [39]. Nevertheless, it should be mentioned that AhR activation by FICZ (via IL-22) inhibited the experimental inflammation and colitis in the gastrointestinal tract of mice [40].

It was reported that certain cytokines play a significant role during dioxin elimination. Moreover, the AhR activation by IL-4 causes the expression of CYP1A1 gene, which is responsible for metabolism of xenobiotics such as TCDD [41, 42]. The AhR receptor inhibition by tocopherol as a possible therapeutic option has been recently described (Fig. 3). The inflammatory effects of dioxin can be also limited by reducing the level of TNF. Furthermore, the application of acetylsalicylic acid blocks cyclooxygenase-II enzyme (COX-2) and deactivates the AhR receptor [43].

The differences in dioxin intoxication effect depend on the type of tissue, subcutaneous adipose or liver tissue in which these compounds are accumulated. After binding to the AhR receptor, dioxins affect DNA stimulating mRNA cyclooxygenase synthesis (CYP 1 and CYP2) and influencing protein metabolism in hepatocytes and haematopoiesis. Accumulation of dioxins in the liver could modify the metabolism of cholesterol and hormones and is directly related to the synthesis of plasma proteins, thereby dioxins are able to change the peripheral blood picture and influence the biochemical response to an inflammatory reaction [44]. Studies have shown that changes in the smooth endoplasmic reticulum of hepatocyte occur between the 5th and 9th day after application of TCDD at a dose 5–25 µg/kg b.w. [45, 46]. The administered doses caused a decrease in bile secretion and an increase of the coproporphyrin concentration [47].

In the murine model, application of 100 µg/kg b.w. of TCDD caused an increase in liver mass during the first 72 hours, and a rise in alanine transaminase (ALT) activity in serum [48]. After this period, no histopathological changes in the liver were observed, whereas the macroscopic and histopathological changes in rat hepatocytes, such as hepatosteatosis and high cholesterol level in serum, were noticed after the 3rd week of TCDD application at a dose of 5 µg/kg b.w. [49]. This indicates that mice manifest a higher




Figure 3. Inflammatory mechanism of dioxin action by stimulation of COX-2 dual-receptor block by tocopherol and acetylsalicylic acid. Abbreviations: TCDD – 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin; AhR – Aryl hydrocarbon receptor; Ca²⁺– calcium ions; cPLA– cytosolic phospholipase A2; AA – arachidonic acid; COX-2 – Cyclooxygenase-II enzyme. On the basis of [82]

resistance to dioxin. Moreover, the studies of TCDD activity in monkeys and mice show the rise of lamine A concentration and intensive ALT activity in the liver [48, 50].

Pro-inflammatory character of dioxins. Currently, experiments designed to determine the effect of different doses of dioxins on the central nervous system (CNS) and cardiac muscle of animals are being conducted. These studies indicate the pro-inflammatory character of dioxins, which could be also correlated with oxidative stress and cardiotoxic interactions responsible for the proliferation of connective tissue compounds (Fig. 4) [51, 52].

The mechanisms of dioxin activity include processes related to the distribution of free radicals (hydroxylation, epoxidation), the appearance of free oxygen atoms that cause the formation of hypochlorites. A recently established mechanism is the stimulation of COX-2, which generates synthesis of prostaglandins (PGE 2) [53]. Furthermore, the effects of the AhR receptor contribute indirectly to the increased synthesis of monooxygenase (CYP1), which enhances the inflammatory processes. Recent studies have shown that inflammation generated in one of the organs – for example, the pleura – contributes, through increased pro-inflammatory interleukins and generation of free radicals, to the formation of destructive changes in the structures of the hippocampus. The observed changes are intensified in animals exposed to TCDD action. The mechanism of this process is related to the generation of free radicals by TCDD. These molecules enhance the existing inflammation by affecting AhR receptor, as well as contributing to the stimulation of COX-2. It has been reported recently that administration of high doses of tocopherol and acetylsalicylic acid, which could block the AhR receptor, contributes to the reduction of the changes in the hippocampus caused by administration of TCDD [54] (Fig. 5). One hypothesis states that this mechanism can have an influence on the uncontrollable emotional states




Figure 4. Mechanism of proinflammatory action of dioxins. Abbreviations: COX-2 – Cyclooxygenase-II enzyme; AhR – Aryl hydrocarbon receptor; CYP – Cytochrome P450.




Figure 5. Mechanism of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin, tocopherol and acetylsalicylic acid and their role in the development of pathological behavior. Abbreviations: TCDD – 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin; COX-2 – Cyclooxygenase-II enzyme; AhR – Aryl hydrocarbon receptor; ER – Estrogen receptor alpha; ROS – Reactive Oxygen Species; CYP – Cytochrome P450; CRH – Corticotiberin; AVP – Arginine vasopressin; IL-1 – Interleukin-1; IL-6 – Interleukin-6; TNF – Tumor Necrosis Factor alpha; INF – Interferon-gamma; 5-HT – Serotonin; NA – Noradrenaline; DA – Dopamine

observed in humans and animals, manifested as inexplicable aggression or depression [55].

The biochemical parameters and morphology of peripheral blood, as well as the character of inflammatory response in rats treated with dioxin, have also been analysed. Two routes of dioxin action for the change of the nature and dynamics of the inflammatory process have been shown. In the first route, the impact of dioxin on the synthesis of acute phase proteins in the liver and on the increases of catabolism, results in a change in biochemical parameters, whereas the second path of dioxin action in the inflammatory reaction is associated with change in the neutrophil cellular response of lymphocyte response. This observation has important diagnostic value due to indication of liver function in the regulation of inflammation [46, 56].

Dioxins can disturb the secretion of TSH and thyroid hormones (T3, T4) by increasing the metabolism of T4 by induction of glucuronidation, thus the concentration of TSH in blood increases and T4 decreases [57]. They also cause changes in the regularity of sex steroids secretion [57, 58, 59, 60], such as disorders in the menstrual cycle [61]. Moreover, it has been stated that dioxins are responsible for sequential loss of mass of genitalia and sex glands in polar bears (*Ursus maritimus*). Dioxins could cause disorders during spermatogenesis and oogenesis and could lead to intersexuality, miscarriages or perinatal mortality [19, 20]. Active dioxins may be released in milk during lactation [21, 27]. The offspring of bears fed on toxic milk have low survivability because their immunity is disrupted. These anomalies in the reproductive system and behavioural disorders in youthful individuals have also been observed in rats [61].

The suppressive action of dioxins on the estrogen gene transcription is the potential reason of changes in hormonal regulation [62]. Recent studies have shown that a single administration of TCDD contributes to a significant reduction of estrogen receptor (ER α) in the hippocampus, which is also related to the behaviour shown in the prolonged reaction time, as well as being the cause of apathy and depression in test animals [32].

Dioxins influence the immune system and are the main reason for chloracne disease [22, 25, 63]. It is noteworthy that POPs can affect the extracellular matrix (EM), which is a site of repair processes, such as connective tissue growth or destruction processes, and it is essential in the inflammatory reaction. Dioxins could cause oxidative stress, which is important during expression of liver gene responsible for collagen synthesis [64]. TCDD inhibits type 1 collagen synthesis, osteopontin and bones sialoproteins, and deactivates the alkaline phosphatase in osseous tissue culture [65]. Furthermore, POPs cause thymus involution [44], the rise of corticosteroids concentration and changes in the composition of plasma protein – the α - and β -globulins level increases and the immune response is weakened [66].

The interaction of dioxins with AhR contributes to the production of cytokines that influence blood cells development. TCDD inhibits mRNA IL-6 expression in the presence of lipopolysaccharides (LPS) and interacts with haematopoietic cells and lymphocytes B; it also decreases globulins synthesis [27, 44]. After administration of TCDD to lymphocytes B culture stimulated by LPS and IL-4, the secretion of immunoglobins, such as IgG1, IgE and IgM, was inhibited [67]. Other reports have shown that TCDD does not cause the increase of IgE synthesis by transformed lymphocytes B. Furthermore, POPs cause long-time immunosuppression of prolymphocytes B in the medulla, which is connected

with the stimulation of apoptosis [46]. The studies indicate that the AhR activation interacts with the haematopoiesis of immature lymphocytes as a result of low POPs doses action [68]. These immunosuppressive effects decrease the survivability of youthful individuals [69]. The experiments performed on monkeys (*Callithrix jacchus*) have shown the reduction of absolute and relative lymphocytes numbers with reference to general lymphocytes numbers three weeks after TCDD administration at a dose of 300 ng/kg b.w. Moreover, the reduction of 20% of lymphocytes CD4 was observed. These changes were the result of interaction of TCDD with the thymus [51]. Studies have shown that TCDD increase in the dose dependent manner the TGF- β and integrins concentration [51]. In the murine model immunization by SRBC and application of TCDD at dose of 5 μ g/kg b.w. has indicated the reduction of CD4 and CD8 lymphocytes in comparison to a control group in which these amount was increased [70]. The exposure to TCDD action causes changes in the innate and acquired immunity in the humoral and cell-mediated immune response. The behaviour of the immune system after dioxin application is a specific indicator of the adaptation of an organism to harmful conditions [69].

CONCLUSIONS

Dioxins are considered as pro-inflammatory environmental factors causing oxidative stress [71]. TCDD also inhibits the regeneration of destroyed fins in ray-finned fish *Archocentrus nigrofasciatus*, which contain connective tissue. These compounds interact with AhR and influence matrix creation [72]. TCDD administered hypodermically has shown cardiotoxic activity in monkeys [50, 52]. Moreover, the distinct character of the dioxins action in the heart has been shown, manifested by the growth in value of collagen, fibronectin and lamins. Dioxins influence prostaglandin and thromboxin synthesis, appearing in the first stage of the inflammatory reaction, and they have an influence on the second stage of the acute phase protein synthesis and may cause changes in the blood picture [73]. TCDD induces oxidative stress in which active forms of oxygen are released and the expression of antioxidant enzymes, e.g. superoxide dismutase, catalase, glutathione reductase and glutathione peroxidase, is decreased [74].

Symptoms of dioxins intoxication are divided, depending on their presence in early and late intoxication stages. The early symptoms occur few days after exposure to dioxin and include:

1. long-lasting Chloracne disease [25];
2. mental disorder (depression, apathy, long-time tiredness) [75];
3. emaciation, anorexia [76];
4. ascites [75];
5. often and long-lasting inflammatory reactions of airways [46];
6. atypical inflammatory reactions [77];
7. hypochromic anaemia – disorders in erythropoiesis [78];
8. susceptibility to infection – immunosuppression action (high mortality in young polar bears) [21];
9. long-lasting bleeding from the digestion system – disorders in thrombopoiesis and fibrinogen synthesis [56]
10. disorder in menstrual cycle, miscarriage, high mortality of infants, disorders in youthful individuals development [79]



Late symptoms appear many years after exposure to dioxin and may include:

1. embriotoxic actions (miscarriage) and teratogenic (cleft palate and backbone, acephalie) [80];
2. cancerogenic action – different cancers (especially in connective tissue) [81];
3. interaction on the hard tissue – disorders in glaze and teeth [7];

According to recent studies, dioxins are present in different concentrations in the environment and cause specific and long-time effects. They influence the main and multistage immune response, for instance, inflammatory reactions responsible for the oxidative stress.

REFERENCES

1. Viluksela M, Bager Y, Tuomisto JT, Scheu G, Unkila M, Pohjanvirta R, et al. Liver tumor-promoting activity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in TCDD-sensitive and TCDD-resistant rat strains. *Cancer Res.* 2000; 60: 6911–6920.
2. Kogevinas M. Epidemiological approaches in the investigation of environmental causes of cancer: the case of dioxins and water disinfection by-products. *Environ Health.* 2011; 10(1): 3.
3. Boffetta P, Mundt KA, Adami HO, Cole P, Mandel JS. TCDD and cancer: a critical review of epidemiologic studies. *Crit Rev Toxicol.* 2011; 41: 622–636.
4. Ahlborg UG, Brouwer A, Fingerhut MA, Jacobson JL, Jacobson SW, Kennedy SW, et al. Impact of polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls on human and environmental health, with special emphasis on application of the toxic equivalency factor concept. *Eur J Pharmacol.* 1992; 228: 179–199.
5. Krauthacker B, Reiner E, Lindstrom G, Rappe C. Residues of polychlorinated-dibenzodioxins, -dibenzofurans and -biphenyls in human milk samples collected in a continental town in Croatia, Yugoslavia. *Arh Hig Rada Toksikol.* 1989; 40: 9–14.
6. Harris M, Zacharewski T, Safe S. Comparative potencies of Aroclors 1232, 1242, 1248, 1254, and 1260 in male Wistar rats--assessment of the toxic equivalency factor (TEF) approach for polychlorinated biphenyls (PCBs). *Fundam Appl Toxicol.* 1993; 20: 456–463.
7. Dobrzyński M, Całkosiński I, Przywitowska I, Kobierska-Brzoza J, Czajczyńska-Waszkiwicz A, Sołtan E, et al. The effects of dioxins in environmental pollution on development of teeth disorders. *Polish J of Environ Stud.* 2009; 18: 319–323.
8. Fiedler H. National PCDD/PCDF release inventories under the Stockholm Convention on Persistent Organic Pollutants. *Chemosphere.* 2007; 67: 96–108.
9. Van den Berg M, Birnbaum LS, Denison M, De Vito M, Farland W, Feeley M, et al. The 2005 World Health Organization reevaluation of human and Mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol Sci.* 2006; 93: 223–241.
10. Liem AKD, Van Zorge JA. Dioxins and related compounds: status and regulatory aspects in selected countries. *Environ Sci Pollut Res.* 1995; 2: 46–56.
11. Liem AK, Furst P, Rappe C. Exposure of populations to dioxins and related compounds. *Food Addit Contam.* 2000; 17: 241–259.
12. Gaborek BJ, Mullikin JM, Pitrat AT, Cummings L, May LM. Pentagon surface wipe sampling health risk assessment. *Toxicol Ind Health.* 2001; 17: 254–261.
13. Rayne S. Using exterior building surface films to assess human exposure and health risks from PCDD/Fs in New York City, USA, after the World Trade Center attacks. *J Hazard Mater.* 2005; 127: 33–39.
14. Rayne S, Ikonomou MG, Butt CM, Diamond ML, Truong J. Polychlorinated dioxins and furans from the World Trade Center attacks in exterior window films from lower Manhattan in New York City. *Environ Sci Technol.* 2005; 39: 1995–2003.
15. Mai TA, Doan TV, Tarradellas J, de Alencastro LF, Grandjean D. Dioxin contamination in soils of Southern Vietnam. *Chemosphere.* 2007; 67: 1802–1807.
16. Grochowalski A, Chrzyszcz R, Piekło R, Gregoraszczyk EL. Estrogenic and antiestrogenic effect of in vitro treatment of follicular cells with 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Chemosphere.* 2001; 43: 823–827.
17. Grochowalski A, Chrzyszcz R. Determination of PCDFs/PCDDs in ambient air from Cracow city, Poland. *Organohal Comp.* 1995; 21: 321–326.
18. Całkosiński I, Stańda M, Borodulin-Nadzieja L, Wasilewska U, Majda J, Cegielski M, et al. The influence of 2,3,7,8-tetrachlorodibenzo-p-dioxin on changes of parenchymal organs structure and estradiol and cholesterol concentration in female rats. *Adv Clin Exp Med.* 2005; 14: 211–215.
19. Verreault J, Norstrom RJ, Ramsay MA, Mulvihill M, Letcher RJ. Composition of chlorinated hydrocarbon contaminants among major adipose tissue depots of polar bears (*Ursus maritimus*) from the Canadian high Arctic. *Sci Total Environ.* 2006; 370: 580–587.
20. Sonne C, Dietz R, Leifsson PS, Born EW, Kirkegaard M, Letcher RJ, et al. Are organohalogen contaminants a cofactor in the development of renal lesions in east Greenland polar bears (*Ursus maritimus*)? *Environ Toxicol Chem.* 2006; 25: 1551–1557.
21. Oehme M, Biseth A, Schlabach M, Wiig O. Concentrations of polychlorinated dibenzo-p-dioxins, dibenzofurans and non-ortho substituted biphenyls in polar bear milk from Svalbard (Norway). *Environ Pollut.* 1995; 90: 401–407.
22. Burton JE, Michalek JE, Rahe AJ. Serum dioxin, chloracne, and acne in veterans of Operation Ranch Hand. *Arch Environ Health.* 1998; 53: 199–204.
23. Wrbitzky R, Beyer B, Thoma H, Flatau B, Hennig M, Weber A, et al. Internal exposure to polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans (PCDDs/PCDFs) of Bavarian chimney sweeps. *Arch Environ Contam Toxicol.* 2001; 40: 136–140.
24. Baccarelli A, Pfeiffer R, Consonni D, Pesatori AC, Bonzini M, Patterson DG Jr, et al. Handling of dioxin measurement data in the presence of non-detectable values: overview of available methods and their application in the Seveso chloracne study. *Chemosphere.* 2005; 60: 898–906.
25. Neuberger M, Landvoigt W, Derntl F. Blood levels of 2,3,7,8-tetrachloro-dibenzo-p-dioxin in chemical workers after chloracne and in comparison groups. *Int Arch Occup Environ Health.* 1991; 63: 325–327.
26. Vezina CM, Walker NJ, Olson JR. Subchronic exposure to TCDD, PeCDF, PCB126, and PCB153: effect on hepatic gene expression. *Environ Health Perspect.* 2004; 112: 1636–1644.
27. Weiss J, Papke O, Bignert A, Jensen S, Greyerz E, Agostoni C, et al. Concentrations of dioxins and other organochlorines (PCBs, DDTs, HCHs) in human milk from Seveso, Milan and a Lombardian rural area in Italy: a study performed 25 years after the heavy dioxin exposure in Seveso. *Acta Paediatr.* 2003; 92: 467–472.
28. Schwetz BA, Norris JM, Sparschu GL, Rowe UK, Gehring PJ, Emerson JL, et al. Toxicology of chlorinated dibenzo-p-dioxins. *Environ Health Perspect.* 1973; 5: 87–99.
29. Boverhof DR, Burgoon LD, Tashiro C, Sharratt B, Chittim B, Harkema JR, et al. Comparative toxicogenomic analysis of the hepatotoxic effects of TCDD in Sprague Dawley rats and C57BL/6 mice. *Toxicol Sci.* 2006; 94: 398–416.
30. Kociba RJ, Schwetz BA. Toxicity of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (TCDD). *Drug Metab Rev.* 1982; 13: 387–406.
31. Całkosiński I, Dobrzyński M, Kobierska-Brzoza J, Majda J, Szymonowicz M, Całkosińska M, et al. The influence of strain, sex and age on selected biochemical parameters in blond serum of Buffalo and Wistar rats *Pol J Vet Sci.* 2010; 13: 293–299.
32. Rosińczuk-Tonderys J. Effect of dioxin on the structure of the central nervous system after tocopherol and acetylsalicylic acid administration in an experimental model [Habilitation Thesis]. Wrocław Medical University, Wrocław 2012.
33. Denison MS, Nagy SR. Activation of the aryl hydrocarbon receptor by structurally diverse exogenous and endogenous chemicals. *Annu Rev Pharmacol Toxicol.* 2003; 43: 309–334.
34. Denison MS, Pandini A, Nagy SR, Baldwin EP, Bonati L. Ligand binding and activation of the Ah receptor. *Chem Biol Interact.* 2002; 141: 3–24.
35. Franc MA, Pohjanvirta R, Tuomisto J, Okey AB. In vivo up-regulation of aryl hydrocarbon receptor expression by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in a dioxin-resistant rat model. *Biochem Pharmacol.* 2001; 62: 1565–1578.
36. Stevens EA, Mezrich JD, Bradfield CA. The aryl hydrocarbon receptor: a perspective on potential roles in the immune system. *Immunology.* 2009; 127: 299–311.
37. Goryo K, Suzuki A, Del Carpio CA, Siizaki K, Kuriyama E, Mikami Y, et al. Identification of amino acid residues in the Ah receptor involved in ligand binding. *Biochem Biophys Res Commun.* 2007; 354: 396–402.
38. Tijet N, Boutros PC, Moffat ID, Okey AB, Tuomisto J, Pohjanvirta R. Aryl hydrocarbon receptor regulates distinct dioxin-dependent and dioxin-independent gene batteries. *Mol Pharmacol.* 2006; 69: 140–153.



39. Bergander L, Wincent E, Rannug A, Foroozesh M, Alworth W, Rannug U. Metabolic fate of the Ah receptor ligand 6-formylindolo[3,2-b]carbazole. *Chem Biol Interact.* 2004; 149: 151–164.
40. Monteleone I, Rizzo A, Sarra M, Sica G, Sileri P, Biancone L, et al. Aryl hydrocarbon receptor-induced signals up-regulate IL-22 production and inhibit inflammation in the gastrointestinal tract. *Gastroenterology.* 2011; 141: 237–248, 248–231.
41. Beischlag TV, Luis Morales J, Hollingshead BD, Perdew GH. The aryl hydrocarbon receptor complex and the control of gene expression. *Crit Rev Eukaryot Gene Expr.* 2008; 18: 207–250.
42. Tanaka G, Kanaji S, Hirano A, Arima K, Shinagawa A, Goda C, et al. Induction and activation of the aryl hydrocarbon receptor by IL-4 in B cells. *Int Immunol.* 2005; 17: 797–805.
43. Kloser E, Böhmendorfer S, Brecker L, Kählig H, Netscher T, Mereiter K, et al. Synthesis of 5-(Fluorophenyl)tocopherols as Novel Dioxin Receptor Antagonists. *European Journal of Organic Chemistry.* 2011; 13: 2450–2457.
44. Ivens IA, Loser E, Rinke M, Schmidt U, Neupert M. Toxicity of 2,3,7,8-tetrabromodibenzo-p-dioxin in rats after single oral administration. *Toxicology.* 1992; 73: 53–69.
45. Fowler BA, Lucier GW, Brown HW, McDaniel OS. Ultrastructural changes in rat liver cells following a single oral dose of TCDD. *Environ Health Perspect.* 1973; 5: 141–148.
46. Calkosinski I, Rosinczuk-Tonderys J, Bazan J, Dzierzba K, Calkosinska M, Majda J, et al. The influence of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on hematological parameters during experimentally induced pleuritis in rats. *Inflammation.* 2013; 36: 387–404.
47. Hwang SW. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the biliary excretion of indocyanine green in rat. *Environ Health Perspect.* 1973; 5: 227–231.
48. Yoon CY, Park M, Kim BH, Park JY, Park MS, Jeong YK, et al. Gene expression profile by 2,3,7,8-tetrachlorodibenzo-p-dioxin in the liver of wild-type (AhR+/+) and aryl hydrocarbon receptor-deficient (AhR-/-) mice. *J Vet Med Sci.* 2006; 68: 663–668.
49. Kociba RJ, Keyes DG, Beyer JE, Carreon RM, Wade CE, Dittenber DA, et al. Results of a two-year chronic toxicity and oncogenicity study of 2,3,7,8-tetrachlorodibenzo-p-dioxin in rats. *Toxicol Appl Pharmacol.* 1978; 46: 279–303.
50. Oberemm A, Meckert C, Brandenburger L, Herzig A, Lindner Y, Kalenberg K, et al. Differential signatures of protein expression in marmoset liver and thymus induced by single-dose TCDD treatment. *Toxicology* 2005; 206: 33–48.
51. Nottebrock C, Riecke K, Kruse M, Shakibaei M, Stahlmann R. Effects of 2,3,7,8-tetrachloro-dibenzo-p-dioxin on the extracellular matrix of the thymus in juvenile marmosets (*Callithrix jacchus*). *Toxicology* 2006; 226: 197–207.
52. Riecke K, Grimm D, Shakibaei M, Kossmehl P, Schulze-Tanzil G, Paul M, et al. Low doses of 2,3,7,8-tetrachlorodibenzo-p-dioxin increase transforming growth factor beta and cause myocardial fibrosis in marmosets (*Callithrix jacchus*). *Arch Toxicol.* 2002; 76: 360–366.
53. Gilroy DW, Tomlinson A, Willoughby DA. Differential effects of inhibitors of cyclooxygenase (cyclooxygenase 1 and cyclooxygenase 2) in acute inflammation. *Eur J Pharmacol.* 1998; 355: 211–217.
54. MacDonald CJ, Ciolino HP, Yeh GC. The drug salicylamide is an antagonist of the aryl hydrocarbon receptor that inhibits signal transduction induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Cancer Res.* 2004; 64: 429–434.
55. Geusau A, Abraham K, Geissler K, Sator MO, Stingl G, Tschachler E. Severe 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) intoxication: clinical and laboratory effects. *Environ Health Perspect.* 2001; 109: 865–869.
56. Calkosinski I, Rosinczuk-Tonderys J, Bronowicka-Szydelko A, Dzierzba K, Bazan J, Dobrzyński M, et al. Effect of tocopherol on biochemical blood parameters in pleuritis-induced rats treated with 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicol Ind Health.* 2013.
57. Gregoraszczyk EL. Dioxin exposure and porcine reproductive hormonal activity. *Cad Saude Publica.* 2002; 18: 453–462.
58. Chaffin CL, Trewin AL, Hutz RJ. Estrous cycle-dependent changes in the expression of aromatic hydrocarbon receptor (AHR) and AHR-nuclear translocator (ARNT) mRNAs in the rat ovary and liver. *Chem Biol Interact.* 2000; 124: 205–216.
59. Chaffin CL, Heimler I, Rawlins RG, Wimpee BA, Sommer C, Hutz RJ. Estrogen receptor and aromatic hydrocarbon receptor in the primate ovary. *Endocrine.* 1996; 5: 315–321.
60. Gregoraszczyk EL, Zabiłny E, Ochwat D. Aryl hydrocarbon receptor (AhR)-linked inhibition of luteal cell progesterone secretion in 2,3,7,8-tetrachlorodibenzo-p-dioxin treated cells. *J Physiol Pharmacol.* 2001; 52: 303–311.
61. Całkosiński I, Borodulin-Nadzieja L, Stańda M, Wasilewska U, Cegielski M. Influence of a single dose of TCDD on estrogen levels and reproduction in female rats. *Med Wet.* 2003; 59.
62. Tian Y, Ke S, Thomas T, Meeker RJ, Gallo MA. Transcriptional suppression of estrogen receptor gene expression by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *J Steroid Biochem Mol Biol.* 1998; 67: 17–24.
63. Crow KD. Chloracne. *Br J Dermatol.* 1970; 83: 599–600.
64. Chojkier M, Houghlum K, Lee KS, Buck M. Long- and short-term D-alpha-tocopherol supplementation inhibits liver collagen alpha(I) gene expression. *Am J Physiol.* 1998; 275: G1480–1485.
65. Singh SU, Casper RF, Fritz PC, Sukhu B, Ganss B, Girard B, Jr., et al. Inhibition of dioxin effects on bone formation in vitro by a newly described aryl hydrocarbon receptor antagonist, resveratrol. *J Endocrinol.* 2000; 167: 183–195.
66. Vos JG, Moore JA, Zinkl JG. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the immune system of laboratory animals. *Environ Health Perspect.* 1973; 5: 149–162.
67. Rodriguez-Sosa M, Elizondo G, Lopez-Duran RM, Rivera I, Gonzalez FJ, Vega L. Over-production of IFN-gamma and IL-12 in AhR-null mice. *FEBS Lett.* 2005; 579: 6403–6410.
68. Near RI, Matulka RA, Mann KK, Gogate SU, Trombino AF, Sherr DH. Regulation of preB cell apoptosis by aryl hydrocarbon receptor/transcription factor-expressing stromal/adherent cells. *Proc Soc Exp Biol Med.* 1999; 221: 242–252.
69. Karras JG, Conrad DH, Holsapple MP. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on interleukin-4-mediated mechanisms of immunity. *Toxicol Lett.* 1995; 75: 225–233.
70. Kerkvliet NI, Brauner JA. Flow cytometric analysis of lymphocyte subpopulations in the spleen and thymus of mice exposed to an acute immunosuppressive dose of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Environ Res.* 1990; 52: 146–154.
71. Aly HA, Domenech O. Cytotoxicity and mitochondrial dysfunction of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in isolated rat hepatocytes. *Toxicol Lett.* 2009; 191: 79–87.
72. Andreasen CH, Stender-Petersen KL, Mogensen MS, Torekov SS, Wegner L, Andersen G, et al. Low physical activity accentuates the effect of the FTO rs9939609 polymorphism on body fat accumulation. *Diabetes* 2008; 57: 95–101.
73. Pohjanvirta R, Korkalainen M, McGuire J, Simanainen U, Juvonen R, Tuomisto JT, et al. Comparison of acute toxicities of indolo [3,2-b]carbazole (ICZ) and 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in TCDD-sensitive rats. *Food Chem Toxicol.* 2002; 40: 1023–1032.
74. Kovacic P, Somanathan R. Integrated approach to immunotoxicity: electron transfer, reactive oxygen species, antioxidants, cell signaling, and receptors. *J Recept Signal Transduct Res.* 2008; 28: 323–346.
75. Hochstein MS, Jr., Render JA, Bursian SJ, Aulerich RJ. Chronic toxicity of dietary 2,3,7,8-tetrachlorodibenzo-p-dioxin to mink. *Vet Hum Toxicol.* 2001; 43: 134–139.
76. Lensu S, Miettinen R, Pohjanvirta R, Linden J, Tuomisto J. Assessment by c-Fos immunostaining of changes in brain neural activity induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and leptin in rats. *Basic Clin Pharmacol Toxicol.* 2006; 98: 363–371.
77. Xu G, Li Y, Yoshimoto K, Chen G, Wan C, Iwata T, et al. 2,3,7,8-Tetrachlorodibenzo-p-dioxin-induced inflammatory activation is mediated by intracellular free calcium in microglial cells. *Toxicology* 2013.
78. Całkosiński I, Rosińczuk-Tonderys J, Bazan J, Dzierzba K, Całkosińska M, Majda J, et al. The Influence of 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) on Hematological Parameters During Experimentally Induced Pleuritis in Rats. *Inflammation* 2012.
79. Yang CY, Huang TS, Lin KC, Kuo P, Tsai PC, Guo YL. Menstrual effects among women exposed to polychlorinated biphenyls and dibenzofurans. *Environ Res.* 2011; 111: 288–294.
80. Weber H, Harris MW, Haseman JK, Birnbaum LS. Teratogenic potency of TCDD, TCDF and TCDD-TCDF combinations in C57BL/6N mice. *Toxicol Lett.* 1985; 26: 159–167.
81. La Merrill M, Harper R, Birnbaum LS, Cardiff RD, Threadgill DW. Maternal dioxin exposure combined with a diet high in fat increases mammary cancer incidence in mice. *Environ Health Perspect.* 2010; 118: 596–601.
82. Całkosiński I, Rosińczuk-Tonderys J, Szopa M, Dobrzyński M, Gamian A. High doses of tocopherol in the prevention and potentiation of dioxin in experimental inflammation-potential application. *Postepy Hig Med Dosw.* 2011; 65: 143–157.
83. Całkosiński I. The course of experimentally induced acute pleuritis with use of nitrogranulogen (NTG) and 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) [Habilitation Thesis]. Wrocław Medical University, Wrocław 2005.