

UDC 575: 632.4

DOI:10.2298/GENSR0802095B

Original scientific paper

TESTING OF WHEAT TO DURABLE RESISTANCE AND SURVEY APPROACH FOR *Puccinia recondita tritici*

Jelena BOŠKOVIĆ¹, Momčilo BOŠKOVIĆ² and Željana PRIJIĆ¹

¹Faculty of biofarming, Sombor

²Faculty of Agriculture, Novi Sad

Bošković J., M. Bošković and Ž. Prijić (2008): *Testing of wheat to durable resistance and survey approach for Puccinia recondita tritici.* – Genetika, Vol. 40, No. 2, 95 -107.

The main objective within new approach in international pathogenicity survey of *Puccinia recondita tritici* was to provide genetically diverse sources of resistance (wheat lines with pyramiding resistant genes) to be used in a survey of wheat leaf rust pathogen in European-Mediterranean regions and to search for and document pathogenicity of *P. recondita tritici* cultures useful in differentiating sources of resistance. Emphasis is placed on sources of resistance and their usefulness rather than on description of fungus populations. In this international survey new methods have been applied containing Central Field Nursery, Central Seedling Tests, Cooperative Seedling Tests and Regional Field Nurseries (ELRWN–European Leaf Rust of Wheat Nursery). The results have been reported from one year of investigations. ELRWN contained 20 winter wheat hybrid lines with pyramiding resistant genes including strong ones

Lr9, Lr19 and Lr24. In addition, 16 spring wheat lines were included, as control lines were Lr9, Lr18, Lr19, Lr24 and Lr14. In that year ELRWN have been realized in 13 countries and cooperative seedling test in 8 countries using 22 pathotypes of *P. recondita tritici*. The best results obtained by the winter wheat lines NS-66/5×Lr24, NS-77/2×Lr19, NS-37/2×Lr19 and spring wheat lines 647-CMA-14793 and 26TH-ESWYT-10. The results have shown losing almost complete resistance of Lr9 and Lr24, but much less Lr19.

Key words: International survey, hybrid lines, leaf rust, resistant hybrid wheat lines.

INTRODUCTION

Leaf rust caused by *Puccinia recondita* Rob. ex Desm. f.sp. *tritici* Eriks. is the most widespread and regularly occurring of the three rusts found on wheat. It is found nearly wherever wheat is grown. Leaf rust fungus is adapted to a range of different climates, and the disease can be found in diverse wheat growing areas throughout the world.

Genetic resistance is the most economical and preferable method of reducing yields losses due to leaf rust. Wheat cultivars that are susceptible regularly suffer yield reductions of 5–15% or greater, depending on the stage of crop development when the initial rust infections occur. To date, 51 leaf rust resistance genes have been designated and mapped in wheat. Resistance gene expression is dependent on the genetics of host-parasite interaction, temperature conditions, plant developmental stage, and interaction between resistance genes with suppressors or other resistance genes in the wheat genomes. The essential orientation for the international studies of the rust pathogens where their long distance dissemination as well established phenomenon. Wind is a great uncontrolled carrier of inoculum and urediospores of rust fungi are recognized as international travelers (ROELFS, 1985). This was the main reason why the best method of rust pathogen control was a network of international cooperative studies which would cover large epidemiological areas (BOŠKOVIĆ and BOŠKOVIĆ 1988, 2000, 2001; STUBBS *et al.* 1974).

Various wheat breeding programs throughout the world have had mixed results in producing cultivars with long-lasting, effective resistance to leaf rust. Spring and winter wheat breeding programs in the world have generally been very successful in producing cultivars that have had high levels of durable and effective resistance.

The importance and necessity of cooperative international investigations of the wheat rusts was especially emphasized by the European and Mediterranean Cereal Rusts Foundation. That was included first time in resolutions of Cereal Rust Conferences in Cambridge, 1964 and later on the others. Cooperative research of yellow rust of wheat for Europe had been organized in Netherlands, for stem rust in

Portugal and Italy, and for leaf rust in Yugoslavia. The European Project of Wheat Leaf Rust Research had been started in Novi Sad dealing primarily with pathogenicity surveys of *Puccinia recondita tritici* in European-Mediterranean regions and breeding for resistance (BOŠKOVIĆ, 1966). From that time in International surveys for European-Mediterranean regions different sets of Lr lines have been used (BOŠKOVIĆ, 1980; BOŠKOVIĆ and BOŠKOVIĆ, 1988; BOŠKOVIĆ *et al.* 2000). The most of these lines had not shown satisfactory efficiency for the surveys. A comparison of pathogenicity of *Puccinia recondita* in Europe, The United States and Canada has shown big differences (BOŠKOVIĆ and BROWDER, 1976). The same Lr lines used hadn't any value for European-Mediterranean regions. It was clear, even years ago, that these regions needed new more efficient resistance genes and large testing and crossing program started in that time. At the beginning 18 donors of resistance had been selected after extensive screening tests of several International rusts nurseries, for crossing with varieties Princ and Starke (BOŠKOVIĆ and BOŠKOVIĆ, 1993). Later on, eight of these hybrid lines with the most interesting donor, 66, 77, 26, 32, 46, 94 and 146, have been crossed with only effective genes Lr9, Lr19 and Lr24 (BOŠKOVIĆ and BOŠKOVIĆ, 1997, 2001; BOŠKOVIĆ *et al.*, 2006).

The main objective within new approach in international pathogenicity survey of *Puccinia recondita tritici* was to provide genetically diverse sources of resistance (wheat lines with pyramiding resistant genes) to be used in a survey of wheat leaf rust pathogen in European-Mediterranean regions and to search for and document pathogenicity of *P. recondita tritici* cultures useful in differentiating sources of resistance. Emphasis is placed on sources of resistance and their usefulness rather than on description of fungus populations.

MATERIALS AND METHODS

The methods are applied according to the following approaches and procedure:

Central Field Nursery Each year in this field nursery numerous field materials from International rust nurseries as well as numerous breeding wheat lines from our program have been tested in the condition of artificial inoculations.

Central Seeding Test *P. recondita tritici* collections from regional nurseries (ELRWN) have been sent to Novi Sad where has been cultured and there virulence to the source lines confirmed. When virulence to a given line is found and confirmed by greenhouse tests, that line should be removed from the field nursery and replaced by another line with potential value. This procedure is based on the concept of maximizing the number of sources of resistance to be studied. It is assumed that once virulent cultures are available, these cultures can be used to separate that line from other sources of resistance. Analysis of infection-type data has been done to distinguish between sources of resistance and to evaluate the usefulness of different sources of resistance in various places of the European-Mediterranean regions.

Cooperative Seeding Tests Uniform sets in European Leaf Rust Wheat Nursery (ELRWN) and possibly some other potentially useful sources of resistance,

should be inoculated with several prevalent cultures by 6-8 cooperators in several countries well-disposed on European-Mediterranean territory.

Regional Field Nurseries (ELRWN) This approach should involve testing of a uniform set of wheat lines to naturally occurring *P. recondita tritici* populations at 20-30 sites in Europe and Mediterranean regions. The materials included should emphasize only wheat lines previously tested and shown to be highly resistant, and for which there is indication of diverse resistance genotype. Observations of leaf rust severity should be made by cooperators and sent to Novi Sad for assembling and summarization. The materials in these nurseries will also provide a basis for collecting uredial cultures which are virulent to some or all of the wheat lines. These cultures are used in further greenhouse and laboratory studies for differentiation sources of resistance. The seedlings in the greenhouse were scored for infection type according to a scale 0-9 and variations were classified for easier computerization (BROWDER and YOUNG, 1975). Reaction classes (R, I and S) comprised the following variation of infections types »R« - 1, 2, 3, 4, (0, 0; 1, 2) »I« - 5, 6, (X⁻, X⁺) and »S« - 7, 8, and 9 (3⁻, 3⁺, 4). Since the segregation was very frequent in the seedlings and in the field, that was designated by »«,« For leaf rust and other rusts the reactions are recorded by severity (0-99) and response (VR-S). In the field are recorded disease severity, the parentage of the surface of the plant tillers and leaves affected, using the modified Cobb scale (PETERSON et al. 1984). Host response, the type of infections observed (R - resistant, I - all intermediate types and S - susceptible).

Severity is reduced to a single digit as follows: 0=0; 10=1; 11-25=2; 26-35=3; 36-45=4; 46-55=5; 56-65=6; 66-75=7; 76-85=8; 86-100=9. Host response is changed from R, I and S to 0-9 scale to computerization and deriving coefficient of infection. R= 0-3 or 2; I=4-6 or 5; S=7-9 or 8.

As a material have been used our hybrid lines with pyramiding resistant genes and other highly resistant wheat genotypes in ELRWN selected according to above explained procedures. In Central Field Nursery are included complete International Rust Nurseries and numerous of our breeding lines.

RESULTS

In Central Field Nursery have been tested in the field eight International Rust Nurseries with total of 410 entries and seven spring wheat - CIMMYT Nurseries with 708 entries. In addition to Central Nursery have been tested hybrid progenies from the breeding program of accumulation, or pyramiding resistant genes. In breeding material were included 834 hybrid lines. Some selected of all these material have been tested in the greenhouse (seedling stage) to twenty-two international cultures of *P. recondita tritici* from Regional Field Nurseries (ELRWN). Genetic differentiation in resistance within these selected wheat hybrids or cultivars have been realized applying computerized program with a Boolean algebraic approach to modeling the gene-for-gene relationship in different environments (BROWDER, 1985).

Cooperative Seedling Tests in the second year included selected 36 winter and spring wheat entries in ELRWN. Seedling tests to particular pathotypes of *P. recondita tritici* have been realized in the following countries: Germany (one pathotype), Czechoslovakia (two pathotypes), Sweden (one path.), China (three path.), France (four path.), Italy (two path.), Bulgaria (four path.) and Israel (five path.) – in total 22 pathotypes. Average reactions to these pathotypes are summarized in Table 1.

A Regional Field Nursery (ELRWN) comprised in second year twenty of winter wheat hybrid lines with pyramiding resistant genes from our breeding program and sixteen highly resistant spring wheat lines, again selected from tested and analyzed International Wheat Rust Nurseries. Field ELRWN nurseries with 36 entries have been realized in 13 countries and evaluated to *P. recondita tritici* and some other wheat pathogens: Germany (3 sites), Austria, Holland, Bulgaria, Israel, Sweden, Switzerland, Italy, Poland, Czechoslovakia, Spain, France and Chile. Summarized data of this ELRWN are presented in Table 1.

All winter wheat hybrid lines with accumulation of resistant genes containing strong resistant genes Lr9, Lr 19 and Lr24 have shown very good results. But, there is a very slight difference between them in degree of resistance. The best were the lines NS-66/5×Lr24, NS-77/2×Lr19, NS-37/2×Lr19, then NS-66/2×Lr19, NS-77/3×Lr24, NS-66/4×Lr19, NS-26/2×Lr19, and NS-26/1×Lr9, NS-32/2×Lr19, NS-94/4×Lr19. These hybrid lines have had a little better combining ability from the genes of the donors and strong resistant genes Lr9, Lr19 and Lr24, which resulted, with some higher degree of resistance. Within spring wheat lines in ELRWN, the best results obtained were the lines 647-CMA-14793 and 26TH-ESWYT-10. Less resistance have had 26TH-ESWYT-36, 11TH-ESWYT-20 and 26TH-ESWYT-3. For these spring lines it can be supposed that they contain several resistant genes. Other spring lines have had insufficient resistance or quite susceptible reactions. The most typical were the lines Lr9, Lr19 and Lr24 which had been used in our breeding program for accumulation of resistant genes. It is clear that these lines loosed almost complete resistance as Lr9 and Lr24, but much less Lr19.

It is important to compare these results of twenty wheat lines containing accumulated resistant genes with the same lines where have been reported the segregation ratios of F₂ generations, that is presented in Table 2 (BOŠKOVIĆ *et al.* 1999, 2000, 2006). The number of resistant genes of these twenty lines in the table is very good correlated with results obtained in the seedlings and adult plants in ELRWN nurseries in Table 1. That means, correlation of degree of resistance of cooperative seedling tests to particularly pathotypes of *P. recondita tritici*, as well as to degree of resistance in the field of the ELRWN in corresponding countries to the number of resistant genes in F₂ generations (Table 2) of each breeding combination.

Table 1. Seedling and field response in the second year ELRWN to *P. recondita tritici*

Winter wheat lines		Cooperative seedling tests			Field response	
		Reaction to 22 pathotypes of <i>P. recondita tritici</i>			Reactions in 13 ELRWN	
		R	Seg.	S	R	S
1	NS-66/5×Lr24	22	-	-	13	-
2	NS-66/2×Lr9	21	1	-	13	-
3	NS77/2×Lr19	22	-	-	13	-
4	NS-77/3×Lr24	21	1	-	13	-
5	NS-26/1×Lr9	20	2	-	12	1
6	NS-32/2×Lr19	20	2	-	13	-
7	NS-37/2×Lr9	22	-	-	13	-
8	NS-66/4×Lr19	21	1	-	13	-
9	NS-26/2×Lr19	21	1	-	13	-
10	NS-26/2×Lr24	19	3	-	11	2
11	NS-32/1×Lr9	18	2	2	12	1
12	NS-32/3×Lr24	19	3	-	12	1
13	NS-46/2×Lr9	16	4	2	11	2
14	NS-46/3×Lr19	17	3	2	10	3
15	NS-46/3×Lr24	18	2	2	11	2
16	NS-94/2×Lr9	18	4	-	12	1
17	NS-94/4×Lr19	20	2	-	12	1
18	NS-94/5×Lr24	19	2	1	13	-
19	NS-146/1×Lr9	16	4	2	11	2
20	NS-146/3×Lr19	18	4	-	12	1
Spring wheat lines						
1	81-ND-582	14	1	7	10	3
2	417-ND-660	9	4	9	8	5
3	647-CMA-14793	22	-	-	12	1
4	11TH-ESWYT-20	18	4	-	8	5
5	11TH-ESWYT-25	15	5	2	10	3
6	11TH-ESWYT-30	6	7	9	7	6
7	26TH-ESWYT-3	18	2	2	10	3
8	26TH-ESWYT-10	22	-	-	12	1
9	26TH-ESWYT-36	21	1	-	10	3
10	26TH-ESWYT-49	12	6	4	8	5
11	26TH-ESWYT-50	12	3	7	9	4
12	Lr9	4	9	9	6	7
13	Lr18	2	5	15	5	8
14	Lr19	12	6	4	11	2
15	Lr24	6	2	14	8	5
16	Lr14	2	8	12	3	10

Legend: R – resistant; Seg. – segregations; S – susceptible.

Table.2 – The segregation ratios in the F_2 generation of crosses between eight sources of resistance and *Lr* lines *Lr9*, *Lr19* and *Lr24* using three pathotypes of *Puccinia recondita tritici*.

Pathotype	B.g.s.1289			Is.w.889			Chl.w.1489		
	Exp. Ratio	χ^2	P	Exp. Ratio	χ^2	P	Exp. Ratio	χ^2	P
NS-66/5× <i>Lr24</i>	57:7	0.35	0.55	57:7	0.53	0.48	54:7	0.16	0.82
NS-66/2× <i>Lr9</i>	57:7	0.24	0.62	9:7	0.15	0.70	45:19	0.77	0.37
NS77/2× <i>Lr19</i>	57:7	0.67	0.40	13:3	0.01	0.92	54:10	0.01	0.92
NS-77/3× <i>Lr24</i>	15:1	0.36	0.53	45:19	2.65	0.12	57:7	0.35	0.55
NS-26/1× <i>Lr9</i>	9:7	3.20	0.08	54:10	0.01	0.99	57:7	0.01	0.92
NS-32/2× <i>Lr19</i>	45:19	2.65	0.12	15:1	0.59	0.44	54:10	0.17	0.65
NS-37/2× <i>Lr9</i>	54:10	0.17	0.65	3:1	0.08	0.75	54:10	0.17	0.65
NS-66/4× <i>Lr19</i>	13:3	0.35	0.55	15:1	0.60	0.53	54:10	0.16	0.70
NS-26/2× <i>Lr19</i>	9:7	0.02	0.90	57:7	0.04	0.82	15:1	1.44	0.24
NS-26/2× <i>Lr24</i>	13:3	0.14	0.70	3:1	1.16	0.32	45:19	0.33	0.58
NS-32/1× <i>Lr9</i>	3:1	0.95	0.75	57:7	0.05	0.82	9:7	0.02	0.90
NS-32/3× <i>Lr24</i>	15:1	1.60	0.24	15:1	1.07	0.32	57:7	0.01	0.92
NS-46/2× <i>Lr9</i>	9:7	0.02	0.90	57:7	0.27	0.58	15:1	1.44	0.24
NS-46/3× <i>Lr19</i>	15:1	0.09	0.75	51:13	0.01	0.92	13:3	0.01	0.92
NS-46/3× <i>Lr24</i>	15:1	0.18	0.65	57:7	0.27	0.58	9:7	0.02	0.02
NS-94/2× <i>Lr9</i>	9:7	0.07	0.80	15:1	1.12	0.32	54:10	1.87	0.16
NS-94/4× <i>Lr19</i>	9:7	1.47	0.30	15:1	0.32	0.58	45:19	2.65	0.12
NS-94/5× <i>Lr24</i>	15:1	0.65	0.42	9:7	1.02	0.32	54:10	0.15	0.70
NS-146/1× <i>Lr9</i>	15:1	3.15	0.08	54:10	0.19	0.65	15:1	0.82	0.27
NS-146/3× <i>Lr19</i>	15:1	0.18	0.94	45:19	0.01	0.92	15:1	1.60	0.24

DISCUSSION

It is well known in the last time that combining or pyramiding of resistance genes into individual cultivars has had considerable success in reducing the rate of evolution of pathogens particularly in the situations where the pathogen does not reproduce sexually, as in the case of *P. recondita tritici*. Considerable arguments for durability of cultivars with pyramided race-specific resistance genes have been already reported (KOLMER *et al.* 1991; MUNDT, 1991).

Same differences in F_2 generation (Table2) concerning the number of resistance genes related to particular pathotypes of *P. recondita tritici* was already reported by other authors stating that differences can depend from different donors and pathotypes used (SAMBORSKI and DYCK, 1982; SINGH and MCINTOSH, 1984).

In the time when we used the lines with strong genes Lr9, Lr19 and Lr24 in our breeding program that lines have had very high resistance on the large epidemiological territory (BOŠKOVIĆ and BOŠKOVIĆ, 1988, 1993, 1995; BOŠKOVIĆ *et al.* 2000; BOŠKOVIĆ *et al.* 2006; PARADIES, 1980; STATLER *et al.* 1985). Meanwhile, these lines loosed almost complete resistance, as Lr9 and Lr24 and much less Lr19 (Table 1). Recently has been reported that pathogenicity studies of European populations of *Puccinia recondita tritici* using pathogenicity and molecular markers resulted in 35 pathotypes identified from 68 isolates examined, all of which were avirulent for the genes Lr9, Lr19 and Lr24, as well as to the other Lr's - Lr21, Lr25 and Lr29 (PARK *et al.* 1996; HYSING, 2007).

That demonstrated the complexity of genetic parasite: host interactions of these hybrids, which differ from the other hybrid combinations.

As we mentioned, further pyramiding resistant genes was realized by selection of the eight best hybrids with same resistant donors and crossed with only effective isogenic lines containing the strong genes Lr9, Lr19 and Lr24. Twenty progenies of F₂ generation with different crossing combinations have been tested with three different international pathotypes of *P. recondita tritici* (Bg.s.12/89, Is.w.8/89 and Chl.w.14/89). Differentiation of the pathotypes that used was from International survey of *P. recondita tritici* on differential Lr single gene lines with different virulence/avirulence formulae from distinct regions. The first culture was from Bulgaria, second from Israel and third was from Chile. Genetic analyses of twenty crossing combinations tested in the seedling stage with the first culture are presented in Table 2.

Different segregation ratios of resistant and susceptible plants in all crossing combinations proved that no one of the hybrids used did not possess strong genes Lr9, Lr19 and Lr24. Trigenic control of resistance was present in the hybrids: NS-66/2 × Lr9, NS-66/5 × Lr24 and NS-77/2 × Lr19. In crossing combinations NS-66/4 × Lr19 and NS-26/2 × Lr24 segregation's ratio 13R : 3S showed epistatic dominant gene on one locus and epistatic recessive gene on the other. In F₁b₁ (BOŠKOVIĆ and BOŠKOVIĆ J. 2007; BOŠKOVIĆ *et al.* 2008) of the same donors have had two or three resistance genes depending of the pathotype applied. The mentioned hybrid with strong genes proved the accumulation of these genes by progenies segregation to resistance susceptible plants. Incorporations of the strong Lr genes with segregation ratio 9R : 7S and two dominant complementary genes have shown the hybrids: NS-77/1 × Lr9, NS-26/1 × Lr9, NS-26/2 × Lr19, NS-46/2 × Lr9, NS-94/2 × Lr9 and NS-94/4 × Lr19, since the majority of these hybrids in F₁b₁ have had only one resistant gene. Two isoeipistatic resistant genes according to ratios 15R : 1S were present in the hybrids: NS-77/3 × Lr24, NS-32/3 × Lr24, NS-37/3 × Lr24, NS-46/3 × Lr19, NS-46/3 × Lr24, NS-94/5 × Lr24, NS-146/1 × Lr9 and NS-146/3 × Lr19. It is typical that these ratios were mostly with Lr24 and again the majority in F₁b₁ has had only one resistant gene, which confirms accumulation of the strong resistant gene.

The same hybrid combinations tested with second culture of *P. recondita tritici* in Table 2. According to segregation ratios have proved successful incorporation of Lr9, Lr19 and Lr24. Comparing segregation ratios with the first

pathotype, it is expressed considerable influence of the culture used, which is related to the segregation ratio and number of resistant genes. Thus, the segregation ratios and number of the genes in all crossing combinations were different, except in the hybrids NS-94/2 × Lr9, NS-94/4 × Lr19, NS-94/5 × Lr24, NS-66/4 × Lr19, NS-77/1 × Lr9, and NS-32/3 × Lr24 where digenic resistant genes were found with both pathotypes and tregenic combination only in NS-66/5 × Lr24.

Segregation ratios with the third culture (Chl.w.14/89) of *P. recondita tritici* in Table 2. have confirmed transferring of the strong resistance genes, with corresponding differences related to the pathotype applies, as with the previous two cultures.

As it is presented (BOŠKOVIĆ and BOŠKOVIĆ J., 2007; BOŠKOVIĆ *et al.* 2008) segregation ratios in F₁b₁ of eight resistant donors in the same recurrent parents Princ and Starke tested with three International cultures of *P. recondita tritici* have shown presence of one, two and three resistance genes. Inheritance of resistance was mostly dominant, as it was reported by many authors (ROELFS *et al.* 1992; JERKOVIĆ, 1992). Yet, the first investigations found that inheritance of resistance to *P. recondita tritici* could be dominant, recessive and intermediate, depending of parents combinations (JERKOVIC, 1992; SCHACHERMAYR *et al.* 1996).

Complementary acting resistance genes have been found in the hybrids NS-66/4 × Starke² and NS-32/2 × Starke² containing three resistance genes, similar with investigations of (SINGH and HUERTA-ESPINO, 2003). They reported in some crosses three resistance genes and two of them complementary acting. In identification of resistance genes Lr33 and Lr34 (DYCK, 1987; DYCK *et al.* 1987; SINGH and HUERTA-ESPINO, 2003) it was found complementary effect of Lr34 with Lr33 or LrT3. It was expressed the influence of recurrent parent to the number resistance genes. Resistant donors 26 and 94 with Princ have had one resistant gene, but with Starke two genes. Similar influence of different recurrent parents was already reported (BOSKOVIC, *et al.* 2006).

Analyses of F₂ progenies in “pyramiding genes” and transferring of the strong Lr genes (Lr9, Lr19 and Lr24) tested with three different cultures of *P. recondita tritici*, it was proved successful incorporation of these genes (Table 2). That was on the base of essential genetic rule that segregation ratios of all crossing combinations were to the resistant and susceptible plants in the progenies. If all progenies of the corresponding crossing combinations would expressed highly resistant reactions, as in the lines Lr9, Lr19 and Lr24, that would means the identity of resistant genes of the both parents used (BARTOŠ *et al.* 1969). Inheritance of resistance in F₂ progenies was similar as in F₁b₁. The dominance of resistance prevailed, as in the papers (BOSKOVIC, *et al.* 1999) and complementary effects (JERKOVIĆ, 1992; BOŠKOVIĆ *et al.* 2001a, 2001b, 2006). Considerable influence of the pathotype used to different segregation ratios and number of resistant genes in F₂ (Table 2), was reported by some other authors (GUPTA *et al.* 1995).

Our further investigations in Serbia need to use reliable MAS which may help in resistance breeding of wheat and resistance gene pyramiding, providing durable resistance. However, the resistance gene that is actually present in the

material may be inactive or inefficient. Available markers should be verified with a possibly wide selection of wheat lines and, if possible, combined with studies of field resistance to rust fungi. The use of DNA markers helps to identify desirable genes more precisely and will facilitate transfer of R genes into wheat and even their pyramiding to derive cultivars with durable resistance to diseases. However, DNA gene marker reliability should be verified in diverse genetic backgrounds. It is particularly important in wheat accessions, considering the complex structure of their hexaploid genome. Gene bank accessions and cultivars of wheat often have additional numerous translocations from wild species. The development of new gene transfer techniques in the last decade allows direct biolistic or Agrobacterium-mediated wheat transformation. New biotechnological methods complement the conventional wheat breeding have opened new ways of transferring resistance genes into wheat cultivars and these new methods enrich conventional breeding methods, but cannot replace them (LUKASZ *et al.* 2003; BOŠKOVIĆ and BOSKOVIĆ J., 2007; BOŠKOVIĆ J. *et al.* 2008).

ACKNOWLEDGEMENTS

The authors are grateful for financial support by international and national projects and cooperators in projects.

Received May 17th, 2008

Accepted July 10th, 2008

REFERENCES

- BARTOŠ, P., P. L. DYCK, and D. J. SAMBORSKI (1969): Adult-plant leaf rust resistance in Thatcher and Marquis wheat: a genetic analysis of the host-parasite interaction. *Canadian Journal of Botany* 47, 267-269.
- BOŠKOVIĆ, J. and M. BOŠKOVIĆ (1993): Expansion of gene-for-gene models in wheat hybrids resistant to *Puccinia recondita tritici*. VIII International Wheat Genetics Symposium, Beijing, p. 101.
- BOŠKOVIĆ, J. and M. BOŠKOVIĆ (1997): New approach for resistance of wheat to leaf rust pathogen in European-Mediterranean regions. 10th Congress of the Mediterranean Phytopathological Union, Montpellier, Le Corum), June 1-5, 1997, p. 553-557.
- BOŠKOVIĆ, J., M. BOŠKOVIĆ and G. PRODANOVIĆ (1995): Enhancement of resistance to *Puccinia recondita tritici* of wheat by incorporation the strong genes in sources of resistance. *European Journal of Plant Pathology XIII International Plant Protection Congress, The Hague, The Netherlands*, 2-7 July 1995, Paper No. 1002.

- BOŠKOVIĆ, J., M. BOŠKOVIĆ, Z. JERKOVIĆ, G. PRODANOVIĆ and V. PEŠIĆ (1999): Durable resistance to wheat leaf rust pathogen with combinations of resistance genes, XIVth International Plant Protection Congress, Tel Aviv, Israel, Abstract, p. 76.
- BOŠKOVIĆ, J. and M. BOŠKOVIĆ (2001a): Primena sistema gen za gen u analizi populacije patogena i selekciji izvora otpornosti. Monografija. Izdavač: Poljoprivredni fakultet, Novi Sad, str. 260.
- BOŠKOVIĆ, J., M. BOŠKOVIĆ, M. BABOVIĆ, Z. JERKOVIĆ, and V. PEŠIĆ (2001b): Pyramiding strategy for durable resistance to wheat leaf rust pathogen. Wheat in a Global Environment. Developments in Plant Breeding, Volume 9, Eds. Z. Bedo and L. Lang. Kluwer Academic Publishers, p.804. Hardcover, May 31, 2001, 1 edition ISBN: 0792367227; p. 337-343.
- BOŠKOVIĆ, J., M. BOŠKOVIĆ, and Ž. PRIJIĆ (2006): Strategija akumulacije gena za dugotrajnu otpornost prema patogenu lisne rđe pšenice. Treći simpozijum sekcije za oplemenjivanje organizama. Zlatibor, 16-20 maja 2006. Zbornik apstarkta, str.15, I-Usm-10.
- BOŠKOVIĆ, J., M. BOŠKOVIĆ and Ž. PRIJIĆ (2008): Pyramiding major genes for resistance to leaf rust pathogen of wheat. Proc. Of the International Scientific Conference on Multifunctional agriculture. University of Szeged, Faculty of agriculture, Hungary. p. 24.
- BOŠKOVIĆ, M. M. (1966): The European Project of Wheat Leaf Rust Research. Contemporary Agriculture, Vol. 14, No. 11-12, p. 607-611.
- BOŠKOVIĆ, M. M. (1980): The use of some new differentials in international pathogenicity surveys of *Puccinia recondita* f. sp. *tritici*. Proc. of the fifth Europ. and Mediter. Cereal Rusts Conf., Bary, Italy, p. 185-190.
- BOŠKOVIĆ, M. and L. E. BROWDER (1976): A comparison of pathogenicity of *Puccinia recondita* f. sp. *tritici* in Europe, the United States and Canada, Plant Dis. Report 60: 278-280.
- BOŠKOVIĆ, M. M. and J. BOŠKOVIĆ (1988): International Pathogenicity Survey of Wheat Leaf Rust Pathogen and Sources of Resistance. Cereal Rust and Powdery Mildews Bulletin Vol. 16, Part 1, 33-47.
- BOŠKOVIĆ, M., L. E. BROWDER, J. BOŠKOVIĆ and Z. JERKOVIĆ, (2000): International survey approach for *Puccinia recondita* *tritici* of wheat using multilocation testing. Acta Phytopathologica et Entomologica Hungarica 35 (1-4), p. 351-358.
- BOŠKOVIĆ, M. and J. BOŠKOVIĆ (2007): Sistem nomenklature u analizi populacije *Puccinia recondita* *tritici*. Str. 287-321. Poglavlje. Naučna publikacija. NAUKA OSNOVA ODRŽIVOG RAZVOJA. Izdavač Društvo genetičara Srbije. ISBN 978-86-87109-00-1. Str. 339.
- BROWDER, L. E. (1985): Parasite: host: environment specificity in the cereal rusts. Ann. Rev. Phytopathology. 23: 201-222.
- DYCK, P. L., E. R. KERBER and O. M. LUKOW (1987): Chromosome location and linkage of a new gene (Lr33) for reaction to *Puccinia recondita* f. sp. *tritici*, Genome 29: 463-466.
- GUPTA, S, R. G. SAINI and A. K. GUPTA (1995): Genetical analysis of resistance of leaf-rust (*Puccinia recondita* f. sp. *tritici*) pathotypes in the durum wheat 'PBW 34' and 'DWL 5023'. Plant Breeding 114 (2) 176-178.
- HYSING, S.C. (2007): Genetic resources for disease resistance breeding in wheat: characterization and utilization. Ph.D. ISSN. 1652-6880; ISBN 978-91-576-7308-4. p. 55.
- JERKOVIĆ, Z. (1992): Identifying genes for *Puccinia recondita* f. sp. *Tritici* resistance in wheat lines. J. Genet. Breed. 46: 163-166.

- KOLMER, J.A, P.L DYCK and A.P. ROELFS (1991): An Appraisal of stem and leaf rust resistance in North American hard red spring wheats and the probability of multiple mutations to virulence in populations of cereal rust fungi. *Phytopathology* 81: 237-239.
- LUKASZ, S., L.GOLKA and J.CHEŁKOWSKI (2003): Leaf rust resistance genes of wheat: identification in cultivars and resistance sources. *J. Appl Genet.* 44 (2):139-49
- MUNDT, C. C. (1991): Probability of mutation to multiple virulence and durability of resistance gene pyramids: further comments. *Phytopathology* 81: 240-242.
- PARADIES, M. (1980): Standard races and variants (biotypes) of *Puccinia recondita* sp. *tritici* identified in Italy in 1977: effectiveness of some resistance genes and greenhouse tests of varieties of durum wheat (*Triticum durum* Des.) and bread wheat (*Triticum aestivum* L.). *Cereal Rusts Bull.* 8 (2): 3-9.
- PARK, R., F., AJAHOOR, and F. G FELSENSTEIN (1996): Genetic variation in European populations of *Puccinia recondita* f. sp. *tritici* using pathogenicity and molecular markers. Proc. of the 9th European and Mediterranean Cereal Rusts&Powdery Mildews Conference. Lunteren, The Netherlands. 92-94.
- PETERSON, R. F, A. B. CAMPBELL, and A. E HANNAH (1984): A diagrammatic scale for estimating rust intensity of leaves and stem of cereals. *Can. J. Res. Sect.* 26: 496-500.
- ROELFS, A. P. (1985): Epidemiology in North America. p. 403-434 in A. P. Roelfs and W. R. Bushnell. eds. *The Cereal Rusts Vol. II; Diseases, Distribution, Epidemiology and Control.* Academic Press, Orlando.
- ROELFS, A. P, R. P. SINGH and E. E. SAARI (1992): *Rust Diseases of Wheat: Concepts and methods of disease management.* Mexico, D. F. CIMMYT. 81 pages.
- SAMBORSKI, D. J. and P. L. DYCK (1982): Enhancement of resistance to *Puccinia recondita tritici* by interaction of resistance genes in wheat. *Canadian journal of Plant Pathology* 4, 152-156.
- SCHACHERMAYR, G., C.FEUILLET, M.MESSMER, M. WINZELER, and B.KELLER (1996): Development of molecular markers linked to the Lr1, Lr9 and Lr24 leaf rust resistance genes in wheat. Proc of the 9th European and Mediterranean Cereal Rusts&Powdery Mildews Conference, Lunteren, The Netherlands: 188-191.
- SINGH, R. P. and R. A. MCINTOSH (1984): Complementary genes for resistance to *Puccinia recondita tritici* in *Triticum aestivum* II. Cytogenic studies. *Can. J. Genet. Cytol.* 26:736-742.
- STATLER, G. D, J. D. MILLER and D. C. HIRSCH (1985): Wheat leaf rust in North Dakota during 1982-1984. *Plant dis.* 69:720-721.
- STUBBS, R. W, E. FUCHS, H. VECHT and E. J. W. BASSEST (1974): The international survey of factors of virulence of *Puccinia striiformis* Westend. in 1969, 1970, 1971. *Ned. Graan Centrum Tech. Ber.* 21; 1-88.

**TESTIRANJE PŠENICE NA DUGOTRAJNU OTPORNOST I ANALIZA
POPULACIJE *Puccinia recondita tritici***Jelena BOŠKOVIĆ¹, M. BOŠKOVIĆ² i Željana PRIJIĆ¹¹Fakultet za biofarming, Sombor²Poljoprivredni fakultet, Novi Sad

I z v o d

Glavni cilj u okviru novog pristupa međunarodne analize populacije patogeniteta *Puccinia recondita tritici* satojao se u uključenju genetički različiti izvora otpornosti (linije pšenice sa akumulacijom gena otpornosti) koji su upotrebljeni u analizi populacije patogena lisne rđe pšenice Evropsko-Mediterranskog područja kao i analiza dobijenih podataka patogeniteta kultura *P. recondita tritici* korisnih u diferencijaciji izvora otpornosti. Istaknut je značaj izvora otpornosti i njihova korisna primena pre nego opisivanje populacije. U ovim međunarodnim analizama primenjene su nove metode u koje se uključuju testiranje centralnog poljskog rasadnika pšenice, centralno testiranje sejanaca, kooperativna testiranja sejanaca i regionalni poljski rasadnici (ELRWN - Evropski Rasadnik za Patogena Lisne Rđe Pšenice). Rezultai su saopšteni za jednogodišnja istraživanja. ELRWN je sadržavao 20 ozimih hibrida linija pšenice sa akumulacijom gena otpornosti uključujući jake gene Lr9, Lr19 i Lr24. Dodatno je uključeno 16 jarin linija pšenice kao i kontrolne linije Lr9, Lr18, Lr19, Lr24 i Lr14. U toj godini rasadnik ELRWN je bio realizovan u 13 zemalja, a kooperativno testiranje sejanaca u 8 zemalja upotrebom 22 patotipa *P. recondita tritici*. Dobijeni su najbolji rezultati sa ozimim linijama pšenice NS-66/5×Lr24, NS-77/2×Lr19, NS-37/2×Lr19 i jarim linijama pšenice 647-CMA-14793 i 26TH-ESWYT-10. Ovi rezultati su pokazali gubitak skore potpune otpornosti linija Lr9 i Lr24, ali mnogo manje linije Lr19.

Primljeno 17. V. 2008.

Odobreno 10. VII. 2008.