

VERKALETS N.B., ZAGORODNYUK A.V.

ON GEOMETRIC EXTENSION OF POLYNOMIALS ON BANACH SPACES

We consider some questions related to Aron-Berner extensions of polynomials on infinitely dimensional complex Banach spaces, using natural extensions of their zeros.

Key words and phrases: homogeneous polynomials on Banach spaces, zeros of polynomials, Aron-Berner extension.

Vasyl Stefanyk Precarpathian National University, 57 Shevchenka str., 76018, Ivano-Frankivsk, Ukraine
E-mail: nadyaverkalets@ukr.net (Verkalets N.B.), andriyzag@yahoo.com (Zagorodnyuk A.V.)

INTRODUCTION AND PRELIMINARIES

Let X be a complex Banach space, $\mathcal{P}(X)$ be the algebra of all continuous complex valued polynomials on X and $\mathcal{P}^n(X)$ be a subspace of $\mathcal{P}(X)$ of n -homogeneous polynomials. Since every polynomial $P \in \mathcal{P}(X)$ admits a unique (up to a multiplicative constant) factorization $P = P_1^{m_1} \cdots P_k^{m_k}$ in $\mathcal{P}(X)$ into irreducible polynomials, where $m_1, \dots, m_k \in \mathbb{N}$, $\deg P_j > 0$, $j = 1, \dots, k$, we can denote the *radical* of P as $\text{Rad}(P) = P_1 \cdots P_k$.

In the general case, let $J = (P_1, \dots, P_n)$ be the ideal generated by $P_1, \dots, P_n \in \mathcal{P}(X)$, that is

$$J = \left\{ \sum_{i=1}^n P_i Q_i : Q_i \in \mathcal{P}(X) \right\}, \quad (1)$$

then the set $\text{Rad } J$ is called the *radical* of J , if $P^k \in J$ for some positive integer k implies $P \in \text{Rad } J$. For a given ideal $J \subset \mathcal{P}(X)$, $V(J)$ denotes the *zero* of J , that is, the common set of zeros of all polynomials in J .

Let G be a subset of X . Then $I(G)$ denotes the hull of G , that is the set of all polynomials in $\mathcal{P}(X)$ which vanish on G .

Ideals of the form (1) is called *finitely generated* in $\mathcal{P}(X)$. The following theorem is an analogue of the well known Hilbert Nulstellensatz for the infinite-dimensional case (see [4, 5]).

Theorem 1. *Let J be a finitely generated in $\mathcal{P}(X)$. Then $I[V(J)] = \text{Rad } J$.*

This theorem implies, in particular, that every irreducible polynomial in $\mathcal{P}(X)$ is defined (up to a multiplicative constant) by its zeros.

In this paper we consider questions related to extensions of polynomials, using their zeros. Here we will use the fact in [3] that zero set of every homogeneous polynomial on a complex infinite-dimensional Banach space X contains an infinite-dimensional linear subspace. Also, we will use the well known Aron-Berner extension of polynomials in $\mathcal{P}(X)$ to the second dual X'' and the fact that the extension operator $P \rightsquigarrow \tilde{P}$ is a topological homomorphism of algebras $\mathcal{P}(X)$ and $\mathcal{P}(X'')$ (see [1, 2]).

УДК 517.98

2010 *Mathematics Subject Classification:* 46A20, 46A22.

1 GEOMETRIC EXTENSION OF POLYNOMIALS

Let $P \in \mathcal{P}(X)$, $\dim X = \infty$. Since $\text{Ker } P$ consists of infinite-dimensional closed linear subspaces ([3]), we can present $\text{Ker } P = \cup_{\alpha} V_{\alpha}$, where V_{α} are *maximal* closed linear subspaces in $\text{Ker } P$ and α goes over a set of indexes. As usually,

$$V_{\alpha}^{\perp} = \{f \in X' : f(x) = 0, \forall x \in V_{\alpha}\}.$$

So, $V_{\alpha}^{\perp\perp}$ naturally contains V_{α} and

$$V_{\alpha}^{\perp\perp} = \{\varphi \in X'' : \varphi(f) = 0, \forall f \in V_{\alpha}^{\perp}\}.$$

Proposition 1. $V_{\alpha}^{\perp\perp} = V_{\alpha}''$.

Proof. $V_{\alpha}^{\perp\perp} \supset V_{\alpha}''$. By the Goldstain's theorem for every $\varphi \in V_{\alpha}''$ there is a net $(x_{\beta}) \in V_{\alpha}$ such that $x_{\beta} \xrightarrow{*weak} \varphi$ that is $f(x_{\beta}) \rightarrow \varphi(f)$ for all $f \in X'$. Hence if $f \in V_{\alpha}^{\perp}$, then $f(x_{\beta}) = 0$ for all β . So, we have $\varphi(f) = 0$ and $V_{\alpha}^{\perp\perp} \supset V_{\alpha}''$.

On the other hand, if we have $\varphi \in V_{\alpha}^{\perp\perp}$, then $\varphi \in X''$. For any $g \in V_{\alpha}'$ by the Hahn-Banach theorem there exists an extension $\tilde{g} \in X'$. Let \tilde{g}_1, \tilde{g}_2 be extensions of g to some elements in X' , then $\tilde{g}_1 - \tilde{g}_2 = 0$ on V_{α} , i.e. $\tilde{g}_1 - \tilde{g}_2 \in V_{\alpha}^{\perp}$. Since $\varphi \in V_{\alpha}^{\perp\perp}$, then $\varphi(\tilde{g}_1 - \tilde{g}_2) = 0$ and so, $\varphi(\tilde{g}_1) = \varphi(\tilde{g}_2)$.

Therefore we can define

$$\varphi(g) := \varphi(\tilde{g}),$$

where \tilde{g} is an arbitrary extension.

We have proved that $\varphi \in V_{\alpha}''$, which means that $V_{\alpha}^{\perp\perp} \subset V_{\alpha}''$. It follows that $V_{\alpha}^{\perp\perp} = V_{\alpha}''$. \square

The main question of this paper is: *does $\text{Ker } \tilde{P} = \cup_{\alpha} V_{\alpha}^{\perp\perp}$, where \tilde{P} is the Aron-Berner extension of P ?*

We have an affirmative answer for some partial cases.

Proposition 2. $\text{Ker } \tilde{P} \supset \cup_{\alpha} V_{\alpha}^{\perp\perp}$.

Proof. Since each $V_{\alpha}^{\perp\perp}$ is naturally identified with V_{α}'' , then the restriction of \tilde{P} on $V_{\alpha}^{\perp\perp}$ coincide with the Aron-Berner extension of restriction of P onto V_{α} . But P vanishes on V_{α} and so, $V_{\alpha}^{\perp\perp} \subset \text{Ker } \tilde{P}$. \square

Corollary 1. *If $\cup_{\alpha} V_{\alpha}^{\perp\perp}$ is a zero-set of a polynomial R on X'' , then $\text{Ker } \tilde{P} = \cup_{\alpha} V_{\alpha}^{\perp\perp}$.*

Proof. Without lost of the generality, we can assume that R is radical. By the Proposition 2, $\text{Ker } \tilde{P} \supset \text{Ker } R$, that is $\tilde{P} \in I[V(R)]$. By Theorem 1, $\tilde{P} = RQ$ for some $Q \in \mathcal{P}(X'')$. Note that $\text{Ker } P = \text{Ker } R|_X$ and $\text{Ker } P = \text{Ker } R|_X \cup \text{Ker } Q|_X$. So, $\text{Ker } Q|_X \subset \text{Ker } P|_X$ and we have that $\text{Ker } Q \subset \text{Ker } R$. Hence $\text{Ker } \tilde{P} = \text{Ker } RQ = \text{Ker } R = \cup_{\alpha} V_{\alpha}^{\perp\perp}$. \square

Corollary 2. *If X is a complemented subspace in X'' , then $\text{Ker } \tilde{P} = \cup_{\alpha} V_{\alpha}^{\perp\perp}$.*

Proof. Let $T : X'' \rightarrow X$ be a projection. Then T maps $V_{\alpha}^{\perp\perp}$ onto V_{α} . Let $R = P \circ T$. So $\cup_{\alpha} V_{\alpha}^{\perp\perp} \subset \text{Ker } R$.

On the other hand, if $z \notin \cup_{\alpha} V_{\alpha}^{\perp\perp}$, then $T(z) \notin \cup_{\alpha} V_{\alpha}$ and so $R(z) \neq 0$, that is $\cup_{\alpha} V_{\alpha}^{\perp\perp} = \text{Ker } R$. By the Corollary 1, $\text{Ker } \tilde{P} = \cup_{\alpha} V_{\alpha}^{\perp\perp}$. \square

Since every dual Banach space is complemented in its second dual we have the following corollary.

Corollary 3. *If X is a dual space to a Banach space, then $\text{Ker } P = \cup_{\alpha} V_{\alpha}^{\perp\perp}$.*

REFERENCES

- [1] Aron R.M., Berner P.D. *A Hahn-Banach extension theorem for analytic mappings*. Bull. Soc. Math. France 1978, **106**, 3–24.
- [2] Carne T.K., Cole B., Gamelin T.W. *A uniform algebra of analytic functions on a Banach spaces*. Trans. Amer. Math. Soc. 1989, **314** (2), 639–659. doi:10.1090/S0002-9947-1989-0986022-0
- [3] Plichko A., Zagorodnyuk A. *On automatic continuity and three problems of “The Scottish Book” concerning the boundedness of polynomials functionals*. J. Math. Anal. Appl. 1998, **220** (2), 477–494. doi:10.1006/jmaa.1997.5826
- [4] Zagorodnyuk A.V. *Spectra of algebras of analytic functions and polynomials on Banach spaces*. Contemp. Math. 2007, **435**, 381–394.
- [5] Zagorodnyuk A.V. *The Nulstellensatz on infinite-dimensional complex spaces*. J. Math. Sci. 1999, **96** (2), 2951–2956. doi:10.1007/BF02169686

Received 22.08.2013

Веркалець Н.Б., Загороднюк А.В. *Геометричне продовження поліномів на банахових просторах* // Карпатські математичні публікації. — 2013. — Т.5, №2. — С. 196–198.

У роботі розглянуті деякі питання, пов'язані з продовженням Арона-Бернера поліномів на нескінченно вимірних комплексних банахових просторах, використовуючи природне продовження їхніх нулів.

Ключові слова і фрази: однорідні поліноми на банахових просторах, нулі поліномів, продовження Арона-Бернера.

Веркалец Н.Б., Загороднюк А.В. *Геометрическое продолжение полиномов на банаховых пространствах* // Карпатские математические публикации. — 2013. — Т.5, №2. — С. 196–198.

У работе рассмотрены некоторые вопросы, связанные с продолжением Арона-Бернера полиномов на бесконечно измеримых комплексных банаховых пространствах, используя естественное продолжение их нулей.

Ключевые слова и фразы: однородные полиномы на банаховых пространствах, нули полиномов, продолжение Арона-Бернера.