

Differential Spatiotemporal Patterns of Galectin Expression are a Hallmark of Endotheliochorial Placentation

Melanie L. Conrad^{1,a}, Nancy Freitag^{1,a}, Mónica E. Diessler², Rocío Hernandez², Gabriela Barrientos^{1,3}, Matthias Rose¹, Luciano A. Casas⁴, Claudio G. Barbeito^{2,3}, Sandra M. Blois¹

¹Division of General Internal and Psychosomatic Medicine, Reproductive Medicine Research Group, Charité Center for Internal Medicine and Dermatology, Charité-Universitätsmedizin Berlin, Berlin, Germany;

²Cátedra de Histología y Embriología, Facultad de Ciencias Veterinarias, Universidad Nacional de La Plata, La Plata, Buenos Aires, Argentina;

³CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas), Buenos Aires, Argentina;

⁴Clínica Veterinaria Berizoo, Berisso, Argentina

Keywords

Canine, endotheliochorial, feline, galectins, placenta, pregnancy

Correspondence

Sandra M. Blois, Charité Universitätsmedizin Berlin, CC12 für Innere Medizin und Dermatologie, Medizinische Klinik mit Schwerpunkt Psychosomatik, AG Reproduktionsmedizin, Haus 037, Forum 4, Raum 2.0549, Campus Virchow Klinikum (CVK), Augustenburger Platz 1, 13353 Berlin, Germany.
E-mail: sandra.blois@charite.de

^aM.L.C. and N.F. contributed equally to this work.

Submission August 20, 2015;
accepted October 20, 2015.

Citation

Conrad ML, Freitag N, Diessler ME, Hernandez R, Barrientos G, Rose M, Casas LA, Barbeito CG, Blois SM. Differential spatiotemporal patterns of galectin expression are a hallmark of endotheliochorial placentation. *Am J Reprod Immunol* 2016; 75: 317–325

doi:10.1111/aji.12452

Introduction

In eutherian mammals, the placenta is a transitory organ that invades the maternal uterus to provide

Problem

Galectins influence the progress of pregnancy by regulating key processes associated with embryo-maternal cross talk, including angiogenesis and placentation. Galectin family members exert multiple roles in the context of hemochorial and epitheliochorial placentation; however, the galectin profile in endotheliochorial placenta remains to be investigated.

Method of study

Here, we used immunohistochemistry to analyze galectin (gal)-1, gal-3 and gal-9 expression during early and late endotheliochorial placentation in two different species (dogs and cats).

Results

We found that during early feline gestation, all three galectin members were more strongly expressed on trophoblast and maternal vessels compared to the decidua. This was accompanied by an overall decrease of gal-1, gal-3 and gal-9 expressions in late feline gestation. In canine early pregnancy, we observed that gal-1 and gal-9 were expressed strongly in cytotrophoblast (CTB) cells compared to gal-3, and no galectin expression was observed in syncytiotrophoblast (STB) cells. Progression of canine gestation was accompanied by increased gal-1 and gal-3 expressions on STB cells, whereas gal-9 expression remained similar in CTB and STB.

Conclusion

These data suggest that both the maternal and fetal compartments are characterized by a spatiotemporal regulation of galectin expression during endotheliochorial placentation. This strongly suggests the involvement of the galectin family in important developmental processes during gestation including immunomodulation, trophoblast invasion and angiogenesis. A conserved functional role for galectins during mammalian placental development emerges from these studies.

nutrition and gas exchange for the developing fetus. As a consequence of this process, tolerance mechanisms must be employed at the maternal–fetal interface to protect the developing semiallogenic fetus

from recognition by the maternal immune system.¹ According to the extent of trophoblast invasion into the uterus, placentation can be classified as hemochorial, endotheliochorial or epitheliochorial. Humans and mice exhibit hemochorial placentation; the most invasive placental type in which trophoblast cells penetrate the maternal blood vessels and come into direct contact with maternal blood.² In contrast, epitheliochorial placentation, characteristic of large farm animals such as cows, pig and sheep, is non-invasive, and trophoblast cells simply attach to the uterine epithelium.³ The third type of placentation, endotheliochorial, is found in dogs and cats; here trophoblast cells invade the uterus until they border the maternal blood vessels.^{4–6} Regardless of the type of placentation, this organ ensures normal growth and development of the embryo and fetus while supporting pregnancy-related changes in maternal physiological systems.

Galectins are a family of highly conserved proteins that mediate numerous biological processes by cross-linking glycans on cell membranes.⁷ All members of this family are composed of a distinct carbohydrate recognition domain (CRD) with conserved consensus regions and beta-galactoside specific lectin activity. While some galectins contain one CRD and are biologically active as either homodimers (e.g., galectin-1) or oligomers that aggregate through their non-lectin domain (galectin-3); other galectins contain two CRDs connected by a short linker peptide (e.g., galectin-9).⁸ Several members of this family play an active role in pregnancy maintenance by regulating trophoblast cell growth and differentiation, angiogenesis and inflammation throughout gestation.^{9–14} Within this family, galectin-1 (gal-1), gal-3 and gal-9 are of particular interest as they are highly expressed at the maternal–fetal interface.¹⁵ Their expression is spatiotemporally regulated during pregnancy, and the biological function of a particular galectin may vary according to the site of expression and availability of particular ligands.⁸ For instance, gal-1 appears to play specific roles during implantation, decidualization, vascularization and later on, during placentation.^{14,16–18} During early gestational stages, gal-1 exhibits proangiogenic functions by promoting decidual vascular expansion through vascular endothelial growth factor (VEGF) receptor 2 signaling.¹² Putative roles played by gal-1 during hemochorial placentation include organization of the extracellular matrix, regulation of trophoblast differentiation and cell motility^{17,19,20} and modulation of class Ib human leukocyte antigen

(HLA) expression;¹¹ the later representing a potent immunosuppressive mechanism triggered by trophoblast cells.^{21–23} In addition, gal-1 is also strongly expressed in immune cells such as decidual natural killer (NK) cells²⁴ and T regulatory cells (Tregs)²⁵ and we have demonstrated that this lectin plays a critical role in immune tolerance by inducing tolerogenic dendritic cells and the subsequent expansion of IL-10 producing regulatory T cells.⁹ During epitheliochorial placentation gal-1 is present in decidual stromal cells and early gestational trophoblast giant cells.²⁶

Regarding the function of gal-3 and gal-9 during pregnancy, specific roles for these proteins remain to be elucidated. Human reports demonstrate that gal-3 expression at the fetal–maternal interface partially overlaps that of gal-1.^{17,27} In mice, uterine gal-3 expression is selectively upregulated within the decidua during early stages of pregnancy, whereas at later stages it is predominantly expressed in the placenta.²⁸ Furthermore, decreased implantation rates were observed upon tissue-specific gal-3 knockdown in the murine uterus,²⁹ suggesting a role for this lectin during the embryo–maternal cross talk driving implantation. Little information is available on epitheliochorial placentation where gal-3 is confined to uterine epithelial cells.²⁶ As for gal-9, this lectin is substantially upregulated in human decidua compared to proliferative endometrium.³⁰ We recently showed the expression of at least six gal-9 isoforms in decidual tissue, and that the *Lgals9* D5 variant can selectively suppress IFN- γ production by uterine NK cells.¹³ Finally, expression of gal-9 has also been reported in the epitheliochorial placenta, though at lower levels than in the decidual compartment.²⁶

Although the role that galectins play in placentation is presently a topic of much interest, the information concentrates mainly on the hemochorial placental type. Information on the involvement of galectins in endotheliochorial and epitheliochorial placental types is minimal, and limited mainly to farm relevant animals such as cows,²⁶ sheep^{31,32} and goats.³³ To our knowledge there are presently no studies investigating galectin expression in endotheliochorial placental mammals such as dogs and cats. As a first step in illuminating this subject, the objective of this research was to examine gal-1, gal-3 and gal-9 expression patterns at the feto–maternal interface in cats and dogs, and to identify changes in expression between early and late gestation. This new knowledge will provide a strong foundation for galectin research and pregnancy in veterinary relevant species.

Materials and methods

Tissue Collection

The cat (*Felis catus*) and dog (*Canis lupus familiaris*) specimens for this study were obtained with owners' approval, during hysterectomy or Cesarean section at the University of La Plata. All procedures were approved by the Universidad Nacional de La Plata ethical committee. Gestational ages were estimated by embryo or fetal length. Average gestation length for dogs and cats is 63–65 days. Accordingly, samples were collected at estimated period of gestation: early (<30 days) or late (>45 days). For each estimated period of gestation, more than one visible embryonic vesicle was used in the study: early ($n = 7$ dogs and 4 cats), late ($n = 5$ dogs and 4 cats). We collected samples of different fetal sacs from each mother. The samples included central areas of placental girdle and excluded marginal hematoma. Maternal and embryo or fetal placental structures were collected and fixed in buffered formalin for 24 h following our standard protocol.¹²

Immunohistochemistry

After deparaffinization and rehydration, serial dog and cat paraffin-embedded tissue sections (4 μm) were washed in TBS, followed by blocking of endogenous peroxidase (PO) through incubation with 3% H_2O_2 in methanol for 30 min at room temperature. After incubation with 2% normal serum for 20 min, primary antibodies rabbit against gal-1 (1:500, sc-28248), gal-3 (1:200, sc-20157) and goat against gal-9 (1:100, sc-19292, all purchased from Santa Cruz Biotechnology) were incubated overnight at 4°C. The slides were then washed and incubated for 1 hr at room temperature with goat anti-rabbit PO-conjugated secondary Ab (1:200, cat. #111-035-003, ImmunoResearch) for gal-1 and gal-3 determination or donkey anti-goat PO-conjugated secondary Ab (1:200, cat. #705-035-147 Jackson ImmunoResearch) for gal-9. The signal was detected using a liquid DAB + Substrate Chromogen System (cat. #K3467, DAKO) at room temperature. After washing, nuclei were counterstained with 0.1% Mayer's hematoxylin followed by a standard dehydration procedure and mounting in Vitro-Clud medium (R. Langenbrinck, Germany). Negative controls were run in parallel for each galectin primary antibody. Briefly, we pre-absorbed the rabbit anti-gal-1, gal-3

or gal-9 primary antibody with the respective blocking peptide (overnight at 4°C), and then proceeded with the above described protocol. Negative controls showed no specific immuno-reactivity. For overview, image tissue sections were stained with hematoxylin and eosin (H&E).

Analysis

Analysis of galectin-stained slides was performed using a Keyence BZ9000 microscope and by three independent observers. Images were taken at different magnifications for each staining. In both species, galectin expression of labyrinthine structures was described, taking into account: maternal vessels (mv), syncytiotrophoblast (STB) and cytotrophoblast (CTB), fetal mesenchyma (fm) and fetal vessels (fv). Galectin expression in the decidual cells of the cat placenta was also assessed. Each observer quantified staining of the respective galectins in the mentioned placental structures with the following score: strong (+++), moderate (++) , weak (+) staining and negative (-). Furthermore, positive cells were quantified by counting within the different areas in each tissue sample including mv, STB, CTB, fm, fv and decidua (Dec). For each sample, at least four different regions (i.e., fields) were recorded. Percentage of positive cells was expressed as a mean count for each of the analyzed group.

Results

Differential Expression of gal-1, gal-3 and gal-9 Between Early and Late Healthy Feline Gestation

Early feline gestation samples showed expression of gal-1, gal-3 and gal-9 within the trophoblast cytoplasm, maternal vessels and decidual cytoplasm (Fig. 1a–d; Table I, Figure S1a). In all galectin stainings, the cyto- and syncytiotrophoblasts and maternal vessels stained more strongly than decidual tissue (Fig. 1b–d). Interestingly, comparing between the different galectin stainings, gal-3 and gal-9 were more strongly expressed than gal-1 in early gestation. The fetal mesenchyma only expressed low levels of gal-1 (Fig. 1b). The fetal vessels were negative for all three galectins tested. Also noteworthy, strong cytoplasmic expression of gal-9 was observed in approximately 25% of decidual stromal cells (Figs 1d and S1a). No nuclear staining was evident in any of the cell types observed.




Fig. 1 Galectins expression during early feline gestation. (a) Left panel: Representative overview image of hematoxylin and eosin (H&E) staining in the labyrinth of endotheliochorial early (<30 days) feline placenta. Scale bar = 100 μm; magnification 100x. Right panel: Scale bar = 25 μm; magnification 400x. (b–d) Representative photomicrographs of gal-1 (b), gal-3 (c) and gal-9 (d) staining at the maternal–fetal interface of early (<30 days) healthy feline gestations. Bar represents 25 μm; magnification 400x. Dec, decidua; mv, maternal vessel; CTB, cytotrophoblast; STB, syncytiotrophoblast; fv, fetal vessels; fm, fetal mesenchyme.

Table I Quantification of Galectin Distribution in Early and Late Feline Placenta Samples

Placental structure	Gestation	gal-1	gal-3	gal-9
Maternal vessels	Early	++	++	++(+)
	Late	++	+++	+(+)
Syncytiotrophoblast	Early	++	++	++
	Late	++	+++	++
Cytotrophoblast	Early	+	++	+++
	Late	++	+++	++
Fetal mesenchyma	Early	–	–	–
	Late	+	+	++
Fetal vessels	Early	+	–	–
	Late	–	–	–
Decidua	Early	+	+	++(+)
	Late	++	+	++(+)

Quantification of staining: strong +++, moderate ++, weak +, negative –. Brackets are indicative of heterogeneity of labeling intensities [i.e., ++(+), weak to moderate staining].

Analysis of mature feline placenta revealed that gal-1 was moderately expressed at similar levels in maternal vessels, the trophoblast and the cytoplasm of decidual cells (Fig. 2b; Table I, Figure S1b). Interestingly, the nuclei of decidual cells were also stained. However, gal-1 expression was almost absent from the fetal vessels during late gestation, while the fetal mesenchyma started to express gal-1 at low levels (Fig. 2b). Gal-3 expression in mature placenta was also increased overall when compared with early pregnancy samples; however, the staining pattern remained the same with the trophoblast

cell cytoplasm and maternal blood vessels staining more strongly than decidual cell cytoplasm. The fetal mesenchyma was weakly stained for gal-3 (Figs 2c and S1b). As for gal-9, a differential pattern of expression was observed between early and late stages of feline gestations. Maternal vessel and trophoblast expression of gal-9 decreased in the mature placenta but decidual cells maintained strong expression. Additionally, the decidual cells presented a nuclear staining. The syncytio- and cytotrophoblasts displayed a vesicular rather than uniform staining in the cytoplasm. The fetal mesenchyma exhibited moderate gal-9 expression in all areas of the mature placenta (Figs 2d and S1b), whereas the fetal vessels were negative for all galectins during late gestation.

Spatiotemporal Expression of gal-1, gal-3 and gal-9 During Normal Canine Gestation

Staining of early canine gestation placental samples revealed that the CTB expresses all three galectins: gal-1 and gal-9 were strongly stained while gal-3 was only weakly expressed (Fig. 3a–d, Table II, Figure S1c). No galectin expression was observed in the STB in early gestation. Maternal blood vessels also expressed gal-3 and gal-9 and no staining of fetal vessels was observed (Fig. 3c,d). When analyzing late canine gestation samples, gal-1 expression was evident in maternal vessels and surrounding STB cells (Fig. 4a–d, Table II, Figure S1d). This lectin was also expressed in CTB, albeit more weakly than in

Fig. 2 Galectins expression pattern in late feline endotheliochorial placentation. (a) Left panel: Representative overview image of hematoxylin & eosin (H&E) staining in the labyrinth of endotheliochorial late (>35 days) feline placenta. Scale bar = 100 μ m; magnification 100 \times . Right panel: Scale bar = 25 μ m; magnification 400 \times . (b–d) Representative examples of gal-1 (b), gal-3 (c) and gal-9 (d) staining at the maternal–fetal interface of late (>45 days) healthy feline gestations. Insert on the bottom left of each panel show negative controls for gal-1 (b), gal-3 (c) and gal-9 (d) staining. Bar represents 25 μ m; magnification 400 \times . Dec, decidua; mv, maternal vessel; CTB, cytotrophoblast; STB, syncytiotrophoblast; fv, fetal vessels; fm, fetal mesenchyme.





Fig. 3 Characterization of galectin expression patterns during early canine endotheliochorial placentation. (a) Top panel: Representative H&E staining shows the maternal zone (Mz), Junctional zone (Jz) and labyrinth (Lab) of the endotheliochorial early canine placenta (<30 days). Scale bar = 250 μ m; magnification 50 \times . Bottom panels: Details of the maternal zone (Mz), Junctional zone (Jz) and labyrinth (Lab) of the endotheliochorial canine placenta during early are shown. Bars represent 25 μ m; magnification 400 \times . (b–d) Representative examples of gal-1 (b), gal-3 (c) and gal-9 (d) staining showing the maternal zone (left), junctional zone (middle) and labyrinth (right) panels of the maternal–fetal interface during early stage (<30 days) healthy canine gestation. Scale bar represent 25 μ m; magnification 400 \times . CTB, cytotrophoblast; STB, syncytiotrophoblast; fv, fetal vessels; fm, fetal mesenchyme; mv, maternal vessel.

early gestation. Regarding gal-3 expression in late canine gestation, we found immunoreactivity only in STB cytoplasm, whereas no CTB staining was observed. In addition, maternal vessels and fetal mesenchyme also weakly expressed gal-3 (Fig. 4c).

Finally, gal-9 expression was detected in maternal vessels and all trophoblast cell populations, with most strong labeling in the CTB and STB cytoplasm. However, the CTB gal-9 expression was heterogeneous (Fig. 4d).

Table II Quantification of Immunohistochemical Detection of Galectin Expression in Early and Late Canine Gestation

Placental structure	Gestation	gal-1	gal-3	gal-9
Maternal vessels	Early	–	++	++
	Late	++	+	+
Syncytiotrophoblast	Early	–	–	–
	Late	++	++	+++
Cytotrophoblast	Early	+++	+	+++
	Late	+	–	++(+)
Fetal mesenchyma	Early	–	–	–
	Late	–	–	–
Fetal vessels	Early	–	–	–
	Late	–	+	–

Quantification of staining: strong +++, moderate ++, weak +, negative –. Brackets are indicative of heterogeneity of labeling intensities [i.e., ++(+), weak to moderate staining].

Discussion

Studies in reproductive tissues from different species have provided important insights into the role played by galectins in pregnancy. As a delicate interplay between maternal and fetal galectin expression is critical for healthy gestation, identification of galectin profiles in placental types with different invasive capacities can help to understand the roles of these important molecules. In this study, we provide the first evidence that endotheliochorial placentas (both canine and feline) have a temporal and cell specific expression of three galectin members from early to late gestation.

Galectins display a unique combination of biological functions, with immunomodulation at the fetomaternal interface being one of the best-described mechanisms of this lectin family.³⁴ Here, we show that gal-1, gal-3 and gal-9 are expressed during early gestation in endotheliochorial placentation, implying that these galectins could also contribute to the protection of canine and feline embryos from attack by the maternal immune system. In this context, canine embryos do not express major histocompatibility complex (MHC) I and II but express FasL, preventing their recognition as foreign antigens and eliminating Fas-bearing cytotoxic maternal T cells, respectively.³⁵ Interestingly, Fas/Apo-1 has been identified as a target for gal-1 recognition in Jurkat cells, sensitizing them to caspase-8 mediated apoptosis.³⁶ Thus, gal-1 expression in early gestation endotheliochorial placentas could constitute a complementary mechanism contributing to fetomaternal

tolerance, which is consistent with its described role facilitating the apoptosis of activated maternal T cells and inducing Treg cells in hemochorial placentation.^{9,10} Similarly to gal-1, gal-9 also functions as an anti-inflammatory signal that influences T-cell survival by promoting Treg and suppressing Th17 cell activity,³⁷ and therefore, its expression during early endotheliochorial placentation could be involved in promoting fetomaternal tolerance. Additionally, our study showed that gal-3 expression was abundant during early endotheliochorial gestation, whereas its expression decreased at later stages. This distribution could imply a role of this lectin in the embryo-maternal cross talk driving implantation, as a similar pattern of gal-3 expression has been found in hemochorial placentation during early gestation.^{18,27,28}

Besides their roles in immunomodulation, galectins also modulate angiogenic responses *in vitro* and *in vivo* by influencing endothelial cell activation, proliferation, migration and tube formation.³⁸ During early endotheliochorial placentation, localization of gal-1, gal-3 and gal-9 to the endothelia of maternal blood vessels within the labyrinth could represent a marker of endothelial cell activation. This is in agreement with studies that show increased galectin expression in the maternal associated vasculature during hemochorial placentation³⁹ and tumorigenesis.^{40–42} As VEGF signaling is increased during the early stages of endotheliochorial placentation,³⁵ it is tempting to speculate that the expression of these three galectins in the maternal vessel endothelia could contribute to the angiogenesis process. Supporting this notion, gal-1 interacts with neuropilin to promote signaling via the VEGF receptor 2,⁴³ enhances H-Ras signaling in endothelial cells⁴⁴ and is associated with fetal growth delay in *Lgals-1* deficient mice due to hampered placental vascularization.¹²

Similarly to gal-1, gal-3 also exhibits pro-angiogenic functions. Gal-3 binding to $\alpha v \beta 3$ integrin modulates VEGF and basic fibroblast growth factor (bFGF)-mediated angiogenesis,⁴⁵ and can also activate VEGFR2 by regulating receptor internalization.⁴⁶ Notably, studies profiling the expression of the VEGF system in canine and mink endotheliochorial placentas have demonstrated that the VEGFR2 is the predominant receptor expressed on microvascular endothelial cells during early pregnancy.^{47,48} This expression largely correlates with the pattern described for several matrix metalloproteases (MMP) in canine and feline pregnancies,^{49,50} particularly with MMP-2 and MMP-9. MMP-2 and MMP-9 target




Fig. 4 Distribution of galectins at the maternal–fetal interface during late (>35 days) normal canine pregnancies. Top panel: Representative H&E staining shows the maternal zone (Mz), Junctional zone (Jz) and labyrinth (Lab) of the endotheliochorial late canine placenta during late (>45 days). Scale bar = 250 μ m; magnification 25 \times . Bottom panels: Details of the Junctional zone (Jz) and labyrinth (Lab) of the endotheliochorial canine placenta during late are shown. Bars represent 25 μ m; magnification 400 \times . (b–d) Representative images of gal-1 (b), gal-3 (c) and gal-9 (d) immunohistochemical staining in the junctional zone (left) and labyrinth (right) of the endotheliochorial canine placenta are shown. Inserts show negative controls. CTB, cytotrophoblast; STB, syncytiotrophoblast; eCTB, extra villous CTB; fv, fetal vessels; fm, fetal mesenchyme; mv, maternal vessel.

the non-galectin domain of gal-3, enhancing the chemotactic properties of this lectin toward endothelial cells and promoting migration and onset of angiogenesis.⁵¹ While pro-angiogenic functions of gal-9 are still elusive, Heusschen et al.⁴² demonstrated that the endogenous Lgals9 Δ 5 is able to influence human endothelial cell proliferation *in vitro*, suggesting that this splice variant of gal-9 can modulate angiogenesis *in vivo*. Together, these findings suggest that galectin expression on maternal blood vessel endothelia in the labyrinth is likely associated with angiogenesis in the canine and feline placenta. These results encourage further studies on galectin interaction with the VEGF system during early endotheliochorial placentalation.

Within endotheliochorial placentalation, comparison of galectin expression from early to late gestation in

feline and canine placentas revealed some striking differences. In the feline placenta, gal-1, gal-3 and gal-9 were expressed in both CTB and STB, whereas at later stages these galectins were more weakly expressed in the trophoblast. This suggests that progression of endotheliochorial placentalation leads to a decrease in galectin expression. Contrasting this, analysis of canine early gestation trophoblast cells revealed that only the CTB expressed gal-1, gal-3 and gal-9. In late canine gestation, galectin expression decreased in the CTB and appeared in the STB, suggesting a gain of galectin expression in canines as CTB differentiates into STB. These results appear to contradict findings in human hemochorial placentalation, where gal-1 expression is lost during the differentiation of CTB to STB.¹¹ These contradictory results may be due to the different degree of

trophoblast invasiveness in hemochorial and endotheliochorial placentation and may also account for the finding that decidual galectin expression (-1, -3 and 9) appears to be lower in canine and feline specimens than in species with deeper invasion such as mice and humans.

The spatiotemporal expression of gal-1, gal-3 and gal-9 in the endotheliochorial placenta provides clues into the roles of these molecules in pregnancy. Research into species with alternative placental strategies can help to unravel the fine details of galectin function, specifically with regard to immune tolerance and angiogenesis at the maternal–fetal interface.

Acknowledgements

This work was supported by Deutsche Forschungsgemeinschaft (DFG) grant BL1115/2-1 to S.M.B and grants from Universidad Nacional de la Plata to C.B. We thank P. Moschansky and Evelyn Hagen (Blois's laboratory) for their excellent technical assistance in generating this work. N.F. received a doctoral fellowship from Charité and G.B. was granted a Re-invitation fellowship from the Deutscher Akademischer Austausch Dienst (DAAD). M.L.C. was awarded with a Rachel Hirsch Habilitation fellowship.

Conflict of interest

The authors declare that no conflict of interest exists.

References

- Arck PC, Hecher K: Fetomaternal immune cross-talk and its consequences for maternal and offspring's health. *Nat Med* 2013; 19:548–556.
- Moffett A, Loke C: Immunology of placentation in eutherian mammals. *Nat Rev Immunol* 2006; 6:584–594.
- Bazer FW, Johnson GA: Pig blastocyst-uterine interactions. *Differentiation* 2014; 87: 52–65.
- Enders AC, Carter AM: Review: The evolving placenta: different developmental paths to a hemochorial relationship. *Placenta* 2012; 33(Suppl.):S92–S98.
- Fernandez P, Diessler M, Pachame A, Ortega H, Gimeno E, Portiansky E, Barbeito C: Intermediate filament proteins expression and carbohydrate moieties in trophoblast and decidual cells of mature cat placenta. *Reprod Domest Anim* 2014; 49: 263–269.
- Fernandez PE, Barbeito CG, Portiansky EL, Gimeno EJ: Intermediate filament protein expression and sugar moieties in normal canine placenta. *Histol Histopathol* 2000; 15:1–6.
- Than NG, Romero R, Goodman M, Weckle A, Xing J, Dong Z, Xu Y, Tarquini F, Szilagyi A, Gal P, Hou Z, Tarca AL, Kim CJ, Kim JS, Haidarian S, Uddin M, Bohn H, Benirschke K, Santolaya-Forgas J, Grossman LI, Erez O, Hassan SS, Zavodszky P, Papp Z, Wildman DE: A primate subfamily of galectins expressed at the maternal-fetal interface that promote immune cell death. *Proc Natl Acad Sci USA* 2009; 106:9731–9736.
- Vasta GR: Galectins as pattern recognition receptors: structure, function, and evolution. *Adv Exp Med Biol* 2012; 946:21–36.
- Blois SM, Ilarregui JM, Tometten M, Garcia M, Orsal AS, Cordo-Russo R, Toscano MA, Bianco GA, Kobelt P, Handjiski B, Tirado I, Markert UR, Klapp BF, Poirier F, Szekeres-Bartho J, Rabinovich GA, Arck PC: A pivotal role for galectin-1 in fetomaternal tolerance. *Nat Med* 2007; 13:1450–1457.
- Kopcow HD, Rosetti F, Leung Y, Allan DS, Kutok JL, Strominger JL: T cell apoptosis at the maternal-fetal interface in early human pregnancy, involvement of galectin-1. *Proc Natl Acad Sci USA* 2008; 105:18472–18477.
- Tirado-Gonzalez I, Freitag N, Barrientos G, Shaikly V, Nagaeva O, Strand M, Kjellberg L, Klapp BF, Mincheva-Nilsson L, Cohen M, Blois SM: Galectin-1 influences trophoblast immune evasion and emerges as a predictive factor for the outcome of pregnancy. *Mol Hum Reprod* 2013; 19:43–53.
- Freitag N, Tirado-Gonzalez I, Barrientos G, Herse F, Thijssen VL, Weedon-Fekjaer SM, Schulz H, Wallukat G, Klapp BF, Nevers T, Sharma S, Staff AC, Dechend R, Blois SM: Interfering with Gal-1-mediated angiogenesis contributes to the pathogenesis of preeclampsia. *Proc Natl Acad Sci USA* 2013; 110:11451–11456.
- Heusschen R, Freitag N, Tirado-Gonzalez I, Barrientos G, Moschansky P, Munoz-Fernandez R, Leno-Duran E, Klapp BF, Thijssen VL, Blois SM: Profiling Lgals9 splice variant expression at the fetal-maternal interface: implications in normal and pathological human pregnancy. *Biol Reprod* 2013; 88:22.
- Barrientos G, Freitag N, Tirado-Gonzalez I, Unverdorben L, Jeschke U, Thijssen VL, Blois SM: Involvement of galectin-1 in reproduction: past, present and future. *Hum Reprod Update* 2014; 20: 175–193.
- Than NG, Romero R, Kim CJ, McGowen MR, Papp Z, Wildman DE: Galectins: guardians of eutherian pregnancy at the maternal-fetal interface. *Trends Endocrinol Metab* 2012; 23:23–31.
- Jeschke U, Hutter S, Heublein S, Vrekoussis T, Andergassen U, Unverdorben L, Papadakis G, Makrigiannakis A: Expression and function of galectins in the endometrium and at the human fetomaternal interface. *Placenta* 2013; 34:863–872.
- Vicovac L, Jankovic M, Cuperlovic M: Galectin-1 and -3 in cells of the first trimester placental bed. *Hum Reprod* 1998; 13:730–735.
- Maquoi E, van den Brule FA, Castronovo V, Foidart JM: Changes in the distribution pattern of galectin-1 and galectin-3 in human placenta correlates with the differentiation pathways of trophoblasts. *Placenta* 1997; 18:433–439.
- Jeschke U, Toth B, Scholz C, Friese K, Makrigiannakis A: Glycoprotein and carbohydrate binding protein expression in the placenta in early pregnancy loss. *J Reprod Immunol* 2010; 85:99–105.
- Kolundzic N, Bojic-Trbojevic Z, Kovacevic T, Stefanoska I, Kadoya T, Vicovac L: Galectin-1 is part of human trophoblast invasion machinery—a functional study in vitro. *PLoS ONE* 2011; 6:e28514.
- Apps R, Gardner L, Sharkey AM, Holmes N, Moffett A: A homodimeric complex of HLA-G on normal trophoblast cells modulates antigen-presenting cells via LILRB1. *Eur J Immunol* 2007; 37:1924–1937.
- Chen LJ, Han ZQ, Zhou H, Zou L, Zou P: Inhibition of HLA-G expression via RNAi abolishes resistance of extravillous trophoblast cell line TEV-1 to NK lysis. *Placenta* 2010; 31:519–527.

- 23 Solier C, Aguerre-Girr M, Lenfant F, Campan A, Berrebi A, Rebmann V, Grosse-Wilde H, Le Bouteiller P: Secretion of proapoptotic intron 4-retaining soluble HLA-G1 by human villous trophoblast. *Eur J Immunol* 2002; 32:3576–3586.
- 24 Koopman LA, Kopcow HD, Rybalov B, Boyson JE, Orange JS, Schatz F, Masch R, Lockwood CJ, Schachter AD, Park PJ, Strominger JL: Human decidual natural killer cells are a unique NK cell subset with immunomodulatory potential. *J Exp Med* 2003; 198:1201–1212.
- 25 Garin MI, Chu CC, Golshayan D, Cernuda-Morollon E, Wait R, Lechler RI: Galectin-1: a key effector of regulation mediated by CD4 + CD25 + T cells. *Blood* 2007; 109:2058–2065.
- 26 Froehlich R, Hambuch N, Haeger JD, Dilly M, Kaltner H, Gabius HJ, Pfarrer C: Galectin fingerprinting detects differences in expression profiles between bovine endometrium and placentomes as well as early and late gestational stages. *Placenta* 2012; 33:195–201.
- 27 von Wolff M, Wang X, Gabius HJ, Strowitzki T: Galectin fingerprinting in human endometrium and decidua during the menstrual cycle and in early gestation. *Mol Hum Reprod* 2005; 11:189–194.
- 28 Phillips B, Knisley K, Weitlauf KD, Dorsett J, Lee V, Weitlauf H: Differential expression of two beta-galactoside-binding lectins in the reproductive tracts of pregnant mice. *Biol Reprod* 1996; 55:548–558.
- 29 Yang H, Lei C, Zhang W: Expression of galectin-3 in mouse endometrium and its effect during embryo implantation. *Reprod Biomed Online* 2012; 24:116–122.
- 30 Popovici RM, Krause MS, Germeyer A, Strowitzki T, von Wolff M: Galectin-9: a new endometrial epithelial marker for the mid- and late-secretory and decidual phases in humans. *J Clin Endocrinol Metab* 2005; 90:6170–6176.
- 31 Gray CA, Dunlap KA, Burghardt RC, Spencer TE: Galectin-15 in ovine uteroplacental tissues. *Reproduction* 2005; 130:231–240.
- 32 Iglesias MM, Rabinovich GA, Ambrosio AL, Castagna LF, Sotomayor CE, Wolfenstein-Todel C: Purification of galectin-3 from ovine placenta: developmentally regulated expression and immunological relevance. *Glycobiology* 1998; 8:59–65.
- 33 Woldeesenbet S, Garcia R, Igbo N, Leake J, Lewis SK, Newton GR: Lectin receptors for endometrial H-type 1 antigen on goat conceptuses. *Am J Reprod Immunol* 2004; 52:74–80.
- 34 Blois SM, Barrientos G: Galectin signature in normal pregnancy and preeclampsia. *J Reprod Immunol* 2014; 101–102:127–134.
- 35 Schafer-Somi S: Early canine pregnancy—a battle for successful growth and angiogenesis. *Reprod Domest Anim* 2012; 47(Suppl. 6):165–168.
- 36 Brandt B, Buchse T, Abou-Eladab EF, Tiedge M, Krause E, Jeschke U, Walzel H: Galectin-1 induced activation of the apoptotic death-receptor pathway in human Jurkat T lymphocytes. *Histochem Cell Biol* 2008; 129:599–609.
- 37 Oomizu S, Arikawa T, Niki T, Kadowaki T, Ueno M, Nishi N, Yamauchi A, Hirashima M: Galectin-9 suppresses Th17 cell development in an IL-2-dependent but Tim-3-independent manner. *Clin Immunol* 2012; 143:51–58.
- 38 Thijssen VL, Rabinovich GA, Griffioen AW: Vascular galectins: regulators of tumor progression and targets for cancer therapy. *Cytokine Growth Factor Rev* 2013; 24:547–558.
- 39 Thijssen VL, Hulsmans S, Griffioen AW: The galectin profile of the endothelium: altered expression and localization in activated and tumor endothelial cells. *Am J Pathol* 2008; 172:545–553.
- 40 Thijssen VL, Postel R, Brandwijk RJ, Dings RP, Nesmelova I, Satijn S, Verhofstad N, Nakabeppu Y, Baum LG, Bakkers J, Mayo KH, Poirier F, Griffioen AW: Galectin-1 is essential in tumor angiogenesis and is a target for antiangiogenesis therapy. *Proc Natl Acad Sci USA* 2006; 103:15975–15980.
- 41 D'Haene N, Catteau X, Maris C, Martin B, Salmon I, Decaestecker C: Endothelial hyperplasia and endothelial galectin-3 expression are prognostic factors in primary central nervous system lymphomas. *Br J Haematol* 2008; 140:402–410.
- 42 Heusschen R, Schulkens IA, van Beijnum J, Griffioen AW, Thijssen VL: Endothelial LGALS9 splice variant expression in endothelial cell biology and angiogenesis. *Biochim Biophys Acta* 2014; 1842:284–292.
- 43 Hsieh SH, Ying NW, Wu MH, Chiang WF, Hsu CL, Wong TY, Jin YT, Hong TM, Chen YL: Galectin-1, a novel ligand of neuropilin-1, activates VEGFR-2 signaling and modulates the migration of vascular endothelial cells. *Oncogene* 2008; 27:3746–3753.
- 44 Thijssen VL, Barkan B, Shoji H, Aries IM, Mathieu V, Deltour L, Hackeng TM, Kiss R, Kloog Y, Poirier F, Griffioen AW: Tumor cells secrete galectin-1 to enhance endothelial cell activity. *Cancer Res* 2010; 70:6216–6224.
- 45 Markowska AI, Jefferies KC, Panjwani N: Galectin-3 protein modulates cell surface expression and activation of vascular endothelial growth factor receptor 2 in human endothelial cells. *J Biol Chem* 2011; 286:29913–29921.
- 46 Markowska AI, Liu FT, Panjwani N: Galectin-3 is an important mediator of VEGF- and bFGF-mediated angiogenic response. *J Exp Med* 2010; 207:1981–1993.
- 47 Winther H, Dantzer V: Co-localization of vascular endothelial growth factor and its two receptors flt-1 and kdr in the mink placenta. *Placenta* 2001; 22:457–465.
- 48 Schafer-Somi S, Sabitzer S, Klein D, Reinbacher E, Kanca H, Beceriklisoy HB, Aksoy OA, Kucukaslan I, Macun HC, Aslan S: Vascular endothelial (VEGF) and epithelial growth factor (EGF) as well as platelet-activating factor (PAF) and receptors are expressed in the early pregnant canine uterus. *Reprod Domest Anim* 2012; 48:20–26.
- 49 Walter I, Schonkypf S: Extracellular matrix components and matrix degrading enzymes in the feline placenta during gestation. *Placenta* 2006; 27:291–306.
- 50 Beceriklisoy HB, Walter I, Schafer-Somi S, Miller I, Kanca H, Izgur H, Aslan S: Matrix metalloproteinase (MMP)-2 and MMP-9 activity in the canine uterus before and during placentation. *Reprod Domest Anim* 2007; 42:654–659.
- 51 Nangia-Makker P, Wang Y, Raz T, Tait L, Balan V, Hogan V, Raz A: Cleavage of galectin-3 by matrix metalloproteases induces angiogenesis in breast cancer. *Int J Cancer* 2010; 127:2530–2541.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Figure S1. Quantification of the galectins immunohistochemical detection during endotheliochorial placentation.