



Planning in entropy-regularized Markov decision processes and games

Jean-Bastien Grill, Omar Domingues, Pierre Ménard, Rémi Munos, Michal Valko

► To cite this version:

Jean-Bastien Grill, Omar Domingues, Pierre Ménard, Rémi Munos, Michal Valko. Planning in entropy-regularized Markov decision processes and games. Neural Information Processing Systems, 2019, Vancouver, Canada. hal-02387515

HAL Id: hal-02387515

<https://hal.inria.fr/hal-02387515>

Submitted on 29 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Planning in entropy-regularized Markov decision processes and games

Jean-Bastien Grill*
DeepMind Paris
jbgrill@google.com

Omar D. Domingues*
SequeL team, Inria Lille
omar.darwiche-domingues@inria.fr

Pierre Ménard
SequeL team, Inria Lille
pierre.menard@inria.fr

Rémi Munos
DeepMind Paris
munos@google.com

Michal Valko
DeepMind Paris
valkom@deepmind.com

Abstract

We propose SmoothCruiser, a new planning algorithm for estimating the value function in entropy-regularized Markov decision processes and two-player games, given a generative model of the environment. SmoothCruiser makes use of the smoothness of the Bellman operator promoted by the regularization to achieve *problem-independent sample complexity* of order $\tilde{O}(1/\varepsilon^4)$ for a desired accuracy ε , whereas for non-regularized settings there are no known algorithms with guaranteed polynomial sample complexity in the worst case.

1 Introduction

Planning with a generative model is *thinking before acting*. An agent thinks using a world model that it has built from prior experience [Sutton, 1991, Sutton and Barto, 2018]. In the present paper, we study planning in two types of environments, *Markov decision processes* (MDPs) and *two-player turn-based zero-sum games*. In both settings, agents interact with an environment by taking actions and receiving rewards. Each action changes the state of the environment and the agent aims to choose actions to maximize the sum of rewards. We assume that we are given a generative model of the environment, that takes as input a state and an action and returns a reward and a next state as output. Such generative models, called *oracles*, are typically built from known data and involve simulations, for example, a physics simulation. In many cases, simulations are costly. For example, simulations may require the computation of approximate solutions of differential equations or the discretization of continuous state spaces. Therefore, a smart algorithm *makes only a small the number of oracles calls* required to estimate the value of a state. The total number of oracle calls made by an algorithm is referred to as *sample complexity*.

The value of a state s , denoted by $V(s)$, the maximum of the sum of discounted rewards that can be obtained from that state. We want an algorithm that returns an estimate of precision ε of the $V(s)$ for any fixed s and has a low sample complexity, which should naturally be a function of ε . An agent can then use this algorithm to predict the value of the possible actions at any given state and choose the best one. The main advantage in estimating the value of a *single* given state s at a time instead of the complete value function² $s \mapsto V(s)$ is that we can have algorithms whose sample complexity does not depend on the size of the state space, which is important when our state space is very large or continuous. On the other hand, the disadvantage is that the algorithm must be run each time a new state is encountered.

*equal contribution

²as done by approximate dynamic programming

Our main contribution is an algorithm that *estimates the value function in a given state* in planning problems that satisfy specific smoothness conditions, which is the case when the rewards are regularized by adding an entropy term. We exploit this smoothness property to obtain a polynomial sample complexity of order $\tilde{O}(1/\varepsilon^4)$ that is *problem independent*.

Related work Kearns et al. [1999] came up with a sparse sampling algorithm (SSA) for planning in MDPs with finite actions and arbitrary state spaces. SSA estimates the value of a state s by building a sparse look-ahead tree starting from s . However, SSA achieves a sample complexity of $\mathcal{O}((1/\varepsilon)^{\log(1/\varepsilon)})$, which is non-polynomial in $1/\varepsilon$. SSA is slow since its search is *uniform*, i.e., it does not select actions adaptively. Walsh et al. [2010] gave an improved version of SSA with adaptive action selection, but its sample complexity is still non-polynomial. The UCT algorithm [Kocsis and Szepesvári, 2006], used for planning in MDPs and games, selects actions based on optimistic estimates of their values and has good empirical performance in several applications. However, the sample complexity of UCT can be worse than exponential in $1/\varepsilon$ for some environments, which is mainly due to exploration issues [Coquelin and Munos, 2007]. Algorithms with sample complexities of order $\mathcal{O}(1/\varepsilon^d)$, where d is a problem-dependent quantity, have been proposed for deterministic dynamics [Hren and Munos, 2008], and in an open-loop³ setting [Bubeck and Munos, 2010, Leurent and Maillard, 2019, Bartlett et al., 2019], for bounded number of next states and a full MDP model is known [Buşoniu and Munos, 2012], for bounded number of next states in a *finite-horizon* setting [Feldman and Domshlak, 2014], for bounded number of next states [Szörényi et al., 2014], and for general MDPs [Grill et al., 2016]. In general, when the state space is infinite and the transitions are stochastic, the problem-dependent quantity d can make the sample complexity guarantees exponential. For a related setting, when rewards are only obtained in the leaves of a fixed tree, Kaufmann and Koolen [2017] and Huang et al. [2017] present algorithms to identify the optimal action in a game based on best-arm identification tools.

Entropy regularization in MDPs and reinforcement learning have been employed in several commonly used algorithms. In the context of policy gradient algorithms, common examples are the TRPO algorithm [Schulman et al., 2015] which uses the Kullback-Leibler divergence between the current and the updated policy to constrain the gradient step sizes, the A3C algorithm [Mnih et al., 2016] that penalizes policies with low entropy to improve exploration, and the work of Neu et al. [2017] presenting a theoretical framework for entropy regularization using the joint state-action distribution. Formulations with entropy-augmented rewards, which is the case in our work, have been used to learn multi-modal policies to improve exploration and robustness [Haarnoja et al., 2017, 2018] and can also be related to policy gradient methods [Schulman et al., 2017]. Furthermore, Geist et al. [2019] propose a theory of regularized MDPs which includes entropy as a special case. Summing up, reinforcement learning knows *how* to employ entropy regularization. In this work, we tasked ourselves to give insights on *why*.

2 Setting and motivation

Both MDPs and two-player games can be formalized as a tuple $(\mathcal{S}, \mathcal{A}, P, R, \gamma)$, where \mathcal{S} is the set of states, \mathcal{A} is the set of actions, $P \triangleq \{P(\cdot|s, a)\}_{s, a \in \mathcal{S} \times \mathcal{A}}$ is a set of probability distributions over \mathcal{S} , $R : \mathcal{S} \times \mathcal{A} \rightarrow [0, 1]$ is a (possibly random) reward function and $\gamma \in [0, 1[$ is the known discount factor. In the MDP case, at each round t , an agent is at state s , chooses action a and observes a reward $R(s, a)$ and a transition to a next state $z \sim P(\cdot|s, a)$. In the case of turn-based two-player games, there are two agents and, at each round t , an agent chooses an action, observes a reward and a transition; at round $t + 1$ it's the other player's turn. This is equivalent to an MDP with an augmented state space $\mathcal{S}^+ \triangleq \mathcal{S} \times \{1, 2\}$ and transition probabilities such that $P((z, j)|(s, i), a) = 0$ if $i = j$. We assume that the action space \mathcal{A} is a finite with cardinality K and the state space \mathcal{S} has arbitrary (possibly infinite) cardinality.

Our objective is to find an algorithm that outputs a good estimate of the value $V(s)$ for any given state s as quickly as possible. An agent can then use this algorithm to choose the best action in an MDP or a game. More precisely, for a state $s \in \mathcal{S}$ and given $\varepsilon > 0$ and $\delta > 0$, our goal is to compute an estimate $\hat{V}(s)$ of $V(s)$ such that $\mathbb{P}\left[|\hat{V}(s) - V(s)| > \varepsilon\right] \leq \delta$ with small number of oracle calls

³This means that the policy is seen as a function of time, not the states. The open-loop setting is particularly adapted to environments with deterministic transitions.

required to compute this estimate. In our setting, we consider the case of *entropy-regularized* MDPs and games, where the objective is augmented with an entropy term.

2.1 Value functions

Markov decision process The policy π of an agent is a function from \mathcal{S} to $\mathcal{P}(\mathcal{A})$, the set of probability distributions over \mathcal{A} . We denote by $\pi(a|s)$ the probability of the agent choosing action a at state s . In MDPs, the value function at a state s , $V(s)$, is defined as the supremum over all possible policies of the expected sum of discounted rewards obtained starting from s , which satisfies the Bellman equations [Puterman, 1994],

$$\forall s \in \mathcal{S}, V(s) = \max_{\pi(\cdot|s) \in \mathcal{P}(\mathcal{A})} \mathbb{E}[R(s, a) + \gamma V(z)], \quad a \sim \pi(\cdot|s), \quad z \sim P(\cdot|s, a). \quad (1)$$

Two-player turn-based zero-sum games In this case, there are two agents (1 and 2), each one with its own policy and different goals. If the policy of Agent 2 is fixed, Agent 1 aims to find a policy that *maximizes* the sum of discounted rewards. Conversely, if the policy of Agent 1 is fixed, Agent 2 aims to find a policy that *minimizes* this sum. Optimal strategies for both agents can be shown to exist and for any $(s, i) \in \mathcal{S}^+ \triangleq \mathcal{S} \times \{1, 2\}$, the value function is defined as [Hansen et al., 2013]

$$V(s, i) \triangleq \begin{cases} \max_{\pi(\cdot|s) \in \mathcal{P}(\mathcal{A})} \mathbb{E}[R((s, i), a) + \gamma V(z, j)], & \text{if } i = 1, \\ \min_{\pi(\cdot|s) \in \mathcal{P}(\mathcal{A})} \mathbb{E}[R((s, i), a) + \gamma V(z, j)], & \text{if } i = 2, \end{cases} \quad (2)$$

with $a \sim \pi(\cdot|s)$ and $(z, j) \sim P(\cdot|(s, i), a)$. In this case, the function $s \mapsto V(s, i)$ is the optimal value function for Agent i when the other agent follows its optimal strategy.

Entropy-regularized value functions Consider a regularization factor $\lambda > 0$. In the case of MDPs, when rewards are augmented by an entropy term, the value function at state s is given by [Haarnoja et al., 2017, Dai et al., 2018, Geist et al., 2019]

$$\begin{aligned} V(s) &\triangleq \max_{\pi(\cdot|s) \in \mathcal{P}(\mathcal{A})} \left\{ \mathbb{E}[R(s, a) + \gamma V(z)] + \lambda \mathcal{H}(\pi(\cdot|s)) \right\}, \quad a \sim \pi(\cdot|s), \quad z \sim P(\cdot|s, a) \\ &= \lambda \log \sum_{a \in \mathcal{A}} \exp\left(\frac{1}{\lambda} \mathbb{E}[R(s, a) + \gamma V(z)]\right), \quad z \sim P(\cdot|s, a), \end{aligned} \quad (3)$$

where $\mathcal{H}(\pi(\cdot|s))$ is the entropy of the probability distribution $\pi(\cdot|s) \in \mathcal{P}(\mathcal{A})$.

The function $\text{LogSumExp}_\lambda : \mathbb{R}^K \rightarrow \mathbb{R}$, defined as $\text{LogSumExp}_\lambda(x) \triangleq \lambda \log \sum_{i=1}^K \exp(x_i/\lambda)$, is a smooth approximation of the max function, since $\|\max - \text{LogSumExp}_\lambda\|_\infty \leq \lambda \log K$. Similarly, the function $-\text{LogSumExp}_{-\lambda}$ is a smooth approximation of the min function. This allows us to define the regularized version of the value function for turn-based two player games, in which both players have regularized rewards, by replacing the max and the min in Equation 2 by their smooth approximations.

For any state s , let $F_s \triangleq \text{LogSumExp}_\lambda$ or $F_s \triangleq -\text{LogSumExp}_{-\lambda}$ depending on s . Both for MDPs and games, we can write the entropy-regularized value functions as

$$V(s) = F_s(Q_s), \quad \text{with } Q_s(a) \triangleq \mathbb{E}[R(s, a) + \gamma V(z)], \quad z \sim P(\cdot|s, a), \quad (4)$$

where $Q_s \triangleq (Q_s(a))_{a \in \mathcal{A}}$, the Q function at state s , is a vector in \mathbb{R}^K representing the value of each action. The function F_s is the *Bellman operator* at state s , which becomes smooth due to the entropy regularization.

Useful properties Our algorithm exploits the smoothness property of F_s defined above. In particular, these functions are L -smooth, that is, for any $Q, Q' \in \mathbb{R}^K$, we have

$$|F_s(Q) - F_s(Q') - (Q - Q')^\top \nabla F_s(Q')| \leq L \|Q - Q'\|_2^2, \quad \text{with } L = 1/\lambda. \quad (5)$$

Furthermore, the functions F_s have two important properties: $\nabla F_s(Q) \succeq 0$ and $\|\nabla F_s(Q)\|_1 = 1$ for all $Q \in \mathbb{R}^K$. This implies that the gradient $\nabla F_s(Q)$ defines a probability distribution.⁵

⁴ $\nabla F_s(Q)$ is the gradient of $F_s(Q)$ with respect to Q .

⁵It is a Boltzmann distribution with temperature λ .

Assumptions We assume that \mathcal{S} , \mathcal{A} , λ , and γ are given to the learner. Moreover, we assume that we can access a generative model, the *oracle*, from which we can get reward and transition samples from arbitrary state-action pairs. Formally, when called with parameter $(s, a) \in \mathcal{S} \times \mathcal{A}$, the oracle outputs a new random variable (R, Z) independent from any other outputs received from the generative model so far such that $Z \sim P(\cdot | s, a)$ and R has same distribution as $R(s, a)$. We denote a call to the oracle as $R, Z \leftarrow \text{oracle}(s, a)$.

2.2 Using regularization for the polynomial sample complexity

To pave the road for `SmoothCruiser`, we consider two extreme cases, based on the strength of the regularization:

1. **Strong regularization** In this case, $\lambda \rightarrow \infty$ and $L = 0$, that is, F_s is linear for all s : $F_s(x) = w_s^\top x$, with $\|w_s\|_1 = 1$, $w_s \in \mathbb{R}^k$ and $w_s \succeq 0$,
2. **No regularization** In this case, $\lambda = 0$ and $L \rightarrow \infty$, that is, F_s cannot be well approximated by a linear function.⁶

In the strongly regularized case, we can approximate the value $V(s)$ with $\tilde{\mathcal{O}}(1/\varepsilon^2)$ oracle calls. This the linearity of F_s , since the value function can be written as $V(s) = \mathbb{E}[\sum_{t=0}^{\infty} \gamma^t R(S_t, A_t) | S_0 = s]$ where A_t is distributed according to the probability vector w_{S_t} . As a result, $V(s)$ can be estimated by Monte-Carlo sampling of trajectories.

With no regularization, we can apply a simple adaptation of the sparse sampling algorithm of [Kearns et al. \[1999\]](#) that we briefly describe. Assume that we have an subroutine that provides an approximation of the value function with precision $\varepsilon/\sqrt{\gamma}$, denoted by $\hat{V}_{\varepsilon/\sqrt{\gamma}}(s)$, for any s . We can call this subroutine several times as well as the oracle to get improved estimate \hat{V} defined as

$$\hat{V}(s) = F_s(\hat{Q}_s) \quad \text{with} \quad \hat{Q}_s(a) \leftarrow \frac{1}{N} \sum_{i=1}^N [r_i(s, a) + \gamma \hat{V}_{\varepsilon/\sqrt{\gamma}}(z_i)],$$

where $r_i(s, a)$ and z_i are rewards and next states sampled by calling the oracle with parameters (s, a) . By Hoeffding's inequality, we can choose $N = \mathcal{O}(1/\varepsilon^2)$ such that $\hat{V}(s)$ is an approximation of $V(s)$ with precision ε with high probability. By applying this idea recursively, we start with $\hat{V} = 0$, which is an approximation of the value function with precision $1/(1 - \gamma)$, and progressively improve the estimates towards a desired precision ε , which can be reached at a recursion depth of $H = \mathcal{O}(\log(1/\varepsilon))$. Following the same reasoning as [Kearns et al. \[1999\]](#), this approach has a sample complexity of $\mathcal{O}((1/\varepsilon)^{\log(1/\varepsilon)})$: to estimate the value at a given recursion depth, we make $\mathcal{O}(1/\varepsilon^2)$ recursive calls and stop once we reach the maximum depth, resulting in a sample complexity of

$$\underbrace{\frac{1}{\varepsilon^2} \times \dots \times \frac{1}{\varepsilon^2}}_{\mathcal{O}(\log(1/\varepsilon)) \text{ times}} = \left(\frac{1}{\varepsilon}\right)^{\mathcal{O}(\log(\frac{1}{\varepsilon}))}.$$

In the next section, we provide `SmoothCruiser` (Algorithm 1), that uses the assumption that the functions F_s are L -smooth with $0 < L < \infty$ to interpolate between the two cases above and obtain a sample complexity of $\tilde{\mathcal{O}}(1/\varepsilon^4)$.

3 SmoothCruiser

We now describe our planning algorithm. Its building blocks are two procedures, `sampleV` (Algorithm 2) and `estimateQ` (Algorithm 3) that recursively call each other. The procedure `sampleV` returns a noisy estimate of $V(s)$ with a bias bounded by ε . The procedure `estimateQ` averages the outputs of several calls to `sampleV` to obtain an estimate \hat{Q}_s that is an approximation of Q_s with precision ε with high probability. Finally, `SmoothCruiser` calls `estimateQ(s, \varepsilon)` to obtain \hat{Q}_s and outputs $\hat{V}(s) = F_s(\hat{Q}_s)$. Using the assumption that F_s is 1-Lipschitz, we can show that $\hat{V}(s)$ is an approximation of $V(s)$ with precision ε . Figure 1 illustrates a call to `SmoothCruiser`.

⁶This is the case of the max and min functions.




Figure 1: Visualization of a call to $\text{SmoothCruiser}(s_0, \varepsilon_0, \delta')$.

3.1 Smooth sailing

The most important part of the algorithm is the procedure `sampleV`, that returns a low-bias estimate of the value function. Having the estimate of the value function, the procedure `estimateQ` averages the outputs of `sampleV` to obtain a good estimate of the Q function with high probability. The main idea of `sampleV` is to first compute an estimate of precision $\mathcal{O}(\sqrt{\varepsilon})$ of the value of each action $\{\hat{Q}_s(a)\}_{a \in \mathcal{A}}$ to linearly approximate the function F_s around \hat{Q}_s .

The local approximation of F_s around \hat{Q}_s is subsequently used to estimate the value of s with a better precision, of order $\mathcal{O}(\varepsilon)$, which is possible due to the smoothness of F_s .

Algorithm 1 SmoothCruiser

Input: $(s, \varepsilon, \delta') \in \mathcal{S} \times \mathbb{R}_+ \times \mathbb{R}_+$
 $M_\lambda \leftarrow \sup_{s \in \mathcal{S}} |F_s(0)| = \lambda \log K$
 $\kappa \leftarrow (1 - \sqrt{\gamma}) / (KL)$
Set δ' , κ , and M_λ as global parameters
 $\hat{Q}_s \leftarrow \text{estimateQ}(s, \varepsilon)$
Output: $F_s(\hat{Q}_s)$

Algorithm 2 sampleV

- 1: **Input:** $(s, \varepsilon) \in \mathcal{S} \times \mathbb{R}_+$
- 2: **if** $\varepsilon \geq (1 + M_\lambda) / (1 - \gamma)$ **then**
- 3: **Output:** 0
- 4: **else if** $\varepsilon \geq \kappa$ **then**
- 5: $\hat{Q}_s \leftarrow \text{estimateQ}(s, \varepsilon)$
- 6: **Output:** $F_s(\hat{Q}_s)$
- 7: **else if** $\varepsilon < \kappa$ **then**
- 8: $\hat{Q}_s \leftarrow \text{estimateQ}(s, \sqrt{\kappa\varepsilon})$
- 9: $A \leftarrow$ action drawn from $\nabla F_s(\hat{Q}_s)$
- 10: $(R, Z) \leftarrow \text{oracle}(s, A)$
- 11: $\hat{V} \leftarrow \text{sampleV}(Z, \varepsilon / \sqrt{\gamma})$
- 12: **Output:**
- 13: $F_s(\hat{Q}_s) - \hat{Q}_s^\top \nabla F_s(\hat{Q}_s) + (R + \gamma \hat{V})$
- 14: **end if**

Algorithm 3 estimateQ

- 1: **Input:** (s, ε)
- 2: $N(\varepsilon) \leftarrow \left\lceil \frac{18(1+M_\lambda)^2 \log(2K/\delta')}{(1-\gamma)^4 (1-\sqrt{\gamma})^2 \varepsilon^2} \right\rceil$
- 3: **for** $a \in \mathcal{A}$ **do**
- 4: $q_i \leftarrow 0$ for $i = 1, \dots, N(\varepsilon)$
- 5: **for** $i = 1, \dots, N(\varepsilon)$ **do**
- 6: $(R, Z) \leftarrow \text{oracle}(s, a)$.
- 7: $\hat{V} \leftarrow \text{sampleV}(Z, \varepsilon / \sqrt{\gamma})$
- 8: $q_i \leftarrow R + \gamma \hat{V}$
- 9: **end for**
- 10: $\hat{Q}_s(a) \leftarrow \text{mean}(q_1, \dots, q_N)$
- 11: clip $\hat{Q}_s(a)$ to $[0, (1 + M_\lambda) / (1 - \gamma)]$
- 12: $\hat{Q}_s(a) \leftarrow \max(0, \hat{Q}_s(a))$
- 13: $\hat{Q}_s(a) \leftarrow \min((1 + M_\lambda) / (1 - \gamma), \hat{Q}_s(a))$
- 14: **end for**
- 15: **Output:** $\hat{Q}_s(a)$

For a target accuracy ε at state s , `sampleV` distinguishes three cases, based on a reference threshold $\kappa \triangleq (1 - \sqrt{\gamma}) / (KL)$, which is the maximum value of ε for which we can compute a good estimate of the value function using linear approximations of F_s .

- **First**, if $\varepsilon \geq (1 + \lambda \log K) / (1 - \gamma)$, then 0 is a valid output, since $V(s)$ is bounded by $(1 + \lambda \log K) / (1 - \gamma)$. This case furthermore ensures that our algorithm terminates, since the recursive calls are made with increasing values of ε .
- **Second**, if $\kappa \leq \varepsilon \leq (1 + \lambda \log K) / (1 - \gamma)$, we run $F_s(\text{estimateQ}(s, \varepsilon))$ in which for each action, both the oracle and `sampleV` are called $\mathcal{O}(1/\varepsilon^2)$ times in order to return $\hat{V}(s)$ which is with high probability an ε -approximation of $V(s)$.

- **Finally**, if $\varepsilon < \kappa$, we take advantage of the smoothness of F_s to compute an ε -approximation of $V(s)$ in a more efficient way than calling the oracle and `sampleV` $\mathcal{O}(1/\varepsilon^2)$ times. We achieve it by calling `estimateQ` with a precision $\sqrt{\kappa\varepsilon}$ instead of ε , which requires $\mathcal{O}(1/\varepsilon)$ calls instead.

3.2 Smoothness guarantee an improved sample complexity

In this part, we describe the key ideas that allows us to exploit the smoothness of the Bellman operator to obtain a better sample complexity. Notice that when $\varepsilon < \kappa$, the procedure `estimateQ` is called to obtain an estimate \widehat{Q}_s such that

$$\|\widehat{Q}_s - Q_s\|_2 = \mathcal{O}\left(\sqrt{\varepsilon/L}\right).$$

The procedure `sampleV` then continues with computing a linear approximation of $F_s(Q_s)$ around \widehat{Q}_s . Using the L -smoothness of F_s , we guarantee the ε -approximation,

$$|F_s(Q_s) - \{F_s(\widehat{Q}_s) + (Q_s - \widehat{Q}_s)^\top \nabla F_s(\widehat{Q}_s)\}| \leq L\|\widehat{Q}_s - Q_s\|_2^2 = \mathcal{O}(\varepsilon).$$

We wish to output this linear approximation, but we need to handle the fact that the vector Q_s (the true Q -function at s) is unknown. Notice that the vector $\nabla F_s(\widehat{Q}_s)$ represents a probability distribution. The term $Q_s^\top \nabla F_s(\widehat{Q}_s)$ in the linear approximation of $F_s(Q_s)$ above can be expressed as

$$Q_s^\top \nabla F_s(\widehat{Q}_s) = \mathbb{E}\left[Q_s(A) \Big| \widehat{Q}_s\right], \text{ with } A \sim \nabla F_s(\widehat{Q}_s).$$

Therefore, we can build a low-bias estimate of $Q_s^\top \nabla F_s(\widehat{Q}_s)$ from estimating only $Q_s(A)$:

- sample action $A \sim \nabla F_s(\widehat{Q}_s)$
- call the generative model to sample a reward and a next state $R_{s,A}, Z_{s,A} \leftarrow \text{oracle}(s, A)$
- obtain an $\mathcal{O}(\varepsilon)$ -approximation of $Q_s(A)$: $\widetilde{Q}(A) = R_{s,A} + \gamma \text{sampleV}(Z_{s,A}, \varepsilon/\sqrt{\gamma})$
- output $\widehat{V}(s) = F_s(\widehat{Q}_s) - \widehat{Q}_s^\top \nabla F_s(\widehat{Q}_s) + \widetilde{Q}(A)$

We show that $\widehat{V}(s)$ is an ε -approximation of the true value function $V(s)$. The benefit of such approach is that we can call `estimateQ` with a precision $\mathcal{O}(\sqrt{\varepsilon})$ instead of $\mathcal{O}(\varepsilon)$, which thanks to the smoothness of F_s , reduces the sample complexity. In particular, one call to `sampleV(s, ε)` will make $\mathcal{O}(1/\varepsilon)$ recursive calls to `sampleV(s, $\mathcal{O}(\sqrt{\varepsilon})$)`, and the total number of calls to `sampleV` behaves as

$$\frac{1}{\varepsilon} \times \frac{1}{\varepsilon^{1/2}} \times \frac{1}{\varepsilon^{1/4}} \times \dots \leq \frac{1}{\varepsilon^2}.$$

Therefore, the number of `sampleV` calls made by `SmoothCruiser` is of order $\mathcal{O}(1/\varepsilon^2)$, which implies that the total sample complexity is of $\mathcal{O}(1/\varepsilon^4)$.

3.3 Comparison to Monte-Carlo tree search

Several planning algorithms are based on Monte-Carlo tree search (MCTS, [Coulom, 2007](#), [Kocsis and Szepesvári, 2006](#)). Algorithm 4 gives a template for MCTS, which uses the procedure `search` that calls `selectAction` and `evaluateLeaf`. Algorithm 5, `search`, returns an estimate of the value function; `selectAction` selects the action to be executed (also called *tree policy*); and `evaluateLeaf` returns an estimate of the value of a leaf. We now provide the analogies that make it possible to see `SmoothCruiser` as an MCTS algorithm:

Algorithm 4 genericMCTS

Input: state s
repeat `search(s, 0)`
until timeout
Output: estimate of best action or value.

- `sampleV` corresponds to the function `search`

- `selectAction` is implemented by calling `estimateQ` to compute \widehat{Q}_s , followed by sampling an action with probability proportional to $\nabla F_s(\widehat{Q}_s)$
- `evaluateLeaf` is implemented using the sparse sampling strategy of Kearns et al. [1999], if we see leaves as the nodes reached when $\varepsilon \geq \kappa$

4 Theoretical guarantees

In Theorem 1 we bound the sample complexity. Note that `SmoothCruiser` is non-adaptive, hence its sample complexity is *deterministic* and *problem independent*. Indeed, since our algorithm is agnostic to the output of the oracle, it performs the same number of oracle calls for any given ε and δ' , regardless of the *random* outcome of these calls.

Theorem 1. *Let $n(\varepsilon, \delta')$ be the number of oracle calls before `SmoothCruiser` terminates. For any state $s \in \mathcal{S}$ and $\varepsilon, \delta' > 0$,*

$$n(\varepsilon, \delta') \leq \frac{c_1}{\varepsilon^4} \log\left(\frac{c_2}{\delta'}\right) \left[c_3 \log\left(\frac{c_4}{\varepsilon}\right) \right]^{\log_2(c_5(\log(\frac{c_2}{\delta'})))} = \tilde{\mathcal{O}}\left(\frac{1}{\varepsilon^4}\right),$$

where c_1, c_2, c_3, c_4 , and c_5 are constants that depend only on K, L , and γ .

The proof of Theorem 1 with the exact constants is in the appendix. In Theorem 2, we provide our consistency result, stating that the output of `SmoothCruiser` applied to a state $s \in \mathcal{S}$ is a good approximation of $V(s)$ with high probability.

Theorem 2. *For any $s \in \mathcal{S}$, $\varepsilon > 0$, and $\delta > 0$, there exists a δ' that depends on ε and δ such that the output $\widehat{V}(s)$ of `SmoothCruiser`(s, ε, δ') satisfies*

$$\mathbb{P}\left[|\widehat{V}(s) - V(s)| > \varepsilon\right] \leq \delta.$$

and such that $n(\varepsilon, \delta') = \mathcal{O}(1/\varepsilon^{4+c})$ for any $c > 0$.

More precisely, in the proof of Theorem 2, we establish that

$$\mathbb{P}\left[|\widehat{V}(s) - V(s)| > \varepsilon\right] \leq \delta' n(\varepsilon, \delta').$$

Therefore, for any parameter δ' satisfying $\delta' n(\varepsilon, \delta') \leq \delta$, `SmoothCruiser` with parameters ε and δ' provides an approximation of $V(s)$ which is (ε, δ) correct.

Impact of regularization constant For a regularization constant λ , the smoothness constant is $L = 1/\lambda$. In Theorem 1 we did not make the dependence on L explicit to preserve simplicity. However, it is easy to analyze the sample complexity in the two limits:

strong regularization $L \rightarrow 0$ and F_s is linear

no regularization $L \rightarrow \infty$ and F_s is not smooth

As $L \rightarrow 0$, the condition $\kappa \leq \varepsilon \leq (1 + \lambda \log K)/(1 - \gamma)$ will be met less and eventually the algorithm will sample $N = \mathcal{O}(1/\varepsilon^2)$ trajectories, which implies a sample complexity of order $\mathcal{O}(1/\varepsilon^2)$. On the other hand, as L goes to ∞ , the condition $\varepsilon < \kappa$ will be met less and the algorithm eventually runs a uniform sampling strategy of Kearns et al. [1999], which results in a sample complexity of order $\mathcal{O}((1/\varepsilon)^{\log(1/\varepsilon)})$, which is non-polynomial in $1/\varepsilon$.

Let $V_\lambda(s)$ be the entropy regularized value function and $V_0(s)$ be its non-regularized version. Since F_s is 1-Lipschitz and $\|\text{LogSumExp}_\lambda - \max\|_\infty \leq \lambda \log K$, we can prove that $\sup_s |V_\lambda(s) - V_0(s)| \leq \lambda \log K/(1 - \gamma)$. Thus, we can interpret $V_\lambda(s)$ as an approximate value function which we can estimate faster.

Comparison to lower bound For non-regularized problems, Kearns et al. [1999] prove a sample complexity lower bound of $\Omega((1/\varepsilon)^{1/\log(1/\gamma)})$, which is polynomial in $1/\varepsilon$, but its exponent grows as γ approaches 1. For regularized problems, Theorem 1 shows that the sample complexity is polynomial with an exponent that is *independent* of γ . Hence, when γ is close to 1, regularization gives us a better asymptotic behavior with respect to $1/\varepsilon$ than the lower bound for the non-regularized case, although we are not estimating the same value.

5 Generalization of SmoothCruiser

Consider the general definition of value functions in Equation 4. Although we focused on the case where F_s is the LogSumExp function, which arises as a consequence of entropy regularization, our theoretical results hold for any set of functions $\{F_s\}_{s \in \mathcal{S}}$ that for any s satisfy the following conditions:

1. F_s is differentiable
2. $\forall Q \in \mathbb{R}^K, 0 < \|\nabla F_s(Q)\|_1 \leq 1$
3. (nonnegative gradient) $\forall Q \in \mathbb{R}^K, \nabla F_s(Q) \succeq 0$
4. (L -smooth) there exists $L \geq 0$ such that for any $Q, Q' \in \mathbb{R}^K$

$$|F_s(Q) - F_s(Q') - (Q - Q')^\top \nabla F_s(Q')| \leq L \|Q - Q'\|_2^2$$

For the more general definition above, we need to make two simple modifications of the procedure `sampleV`. When $\varepsilon < \kappa$, the action A in `sampleV` is sampled according to

$$A \sim \frac{\nabla F_s(\hat{Q}_s)}{\|\nabla F_s(\hat{Q}_s)\|_1}$$

and its output is modified to

$$F_s(\hat{Q}_s) - \hat{Q}_s^\top \nabla F_s(\hat{Q}_s) + (R + \gamma \hat{v}) \|\nabla F_s(\hat{Q}_s)\|_1.$$

In particular, SmoothCruiser can be used for more general regularization schemes, as long as the Bellman operators satisfy the assumptions above. One such example is presented in Appendix E.

6 Conclusion

We provided SmoothCruiser that estimates the value function of MDPs and discounted games defined through smooth approximations of the optimal Bellman operator, which is the case in entropy-regularized value functions. More generally, our algorithm can also be used when value functions are defined through *any* smooth Bellman operator with nonnegative gradients. We showed that our algorithm has a polynomial sample complexity of $\tilde{\mathcal{O}}(1/\varepsilon^4)$, where ε is the desired precision. This guarantee is problem independent and holds for *state spaces of arbitrary cardinality*.

One interesting interpretation of our results is that computing entropy-regularized value functions, which are commonly employed for reinforcement learning, can be seen as a smooth relaxation of a planning problem for which we can obtain a much better sample complexity in terms of the required precision ε . Unsurprisingly, when the regularization tends to zero, we recover the well-known non-polynomial bound $\mathcal{O}((1/\varepsilon)^{\log(1/\varepsilon)})$ of Kearns et al. [1999]. Hence, an interesting direction for future work is to study adaptive regularization schemes in order to accelerate planning algorithms. Although SmoothCruiser makes large amount of recursive calls, which makes it impractical in most situations, we believe it might help us to understand how regularization speeds planning and inspire more practical algorithms. This might be possible by exploiting its similarities to Monte-Carlo tree search that we have outlined above.

Acknowledgments The research presented was supported by European CHIST-ERA project DELTA, French Ministry of Higher Education and Research, Nord-Pas-de-Calais Regional Council, Inria and Otto-von-Guericke-Universität Magdeburg associated-team north-European project Allocate, and French National Research Agency project BoB (grant n.ANR-16-CE23-0003), FMJH Program PGMO with the support of this program from Criteo.

References

- Peter L Bartlett, Victor Gabillon, Jennifer Healey, and Michal Valko. [Scale-free adaptive planning for deterministic dynamics & discounted rewards](#). In *International Conference on Machine Learning*, 2019.
- Sébastien Bubeck and Rémi Munos. [Open-loop optimistic planning](#). In *Conference on Learning Theory*, 2010.
- Lucian Buşoniu and Rémi Munos. [Optimistic planning for Markov decision processes](#). In *International Conference on Artificial Intelligence and Statistics*, 2012.
- Pierre-Arnaud Coquelin and Rémi Munos. [Bandit algorithms for tree search](#). In *Uncertainty in Artificial Intelligence*, 2007.
- Rémi Coulom. [Efficient selectivity and backup operators in Monte-Carlo tree search](#). *Computers and games*, 4630:72–83, 2007.
- Bo Dai, Albert Shaw, Lihong Li, Lin Xiao, Niao He, Zhen Liu, Jianshu Chen, and Le Song. [SBEED: Convergent reinforcement learning with nonlinear function approximation](#). In *International Conference on Machine Learning*, 2018.
- Zohar Feldman and Carmel Domshlak. [Simple regret optimization in online planning for Markov decision processes](#). *Journal of Artificial Intelligence Research*, 2014.
- Matthieu Geist, Bruno Scherrer, and Olivier Pietquin. [A Theory of regularized Markov decision processes](#). In *International Conference on Machine Learning*, pages 2160–2169, 2019.
- Jean-Bastien Grill, Michal Valko, and Rémi Munos. [Blazing the trails before beating the path: Sample-efficient Monte-Carlo planning](#). In *Neural Information Processing Systems*, 2016.
- Tuomas Haarnoja, Haoran Tang, Pieter Abbeel, and Sergey Levine. [Reinforcement learning with deep energy-based policies](#). In *International Conference on Machine Learning*, 2017.
- Tuomas Haarnoja, Aurick Zhou, Pieter Abbeel, and Sergey Levine. [Soft actor-critic: Off-policy maximum entropy deep reinforcement learning with a stochastic actor](#). In *International Conference on Machine Learning*, 2018.
- Thomas Dueholm Hansen, Peter Bro Miltersen, and Uri Zwick. [Strategy iteration is strongly polynomial for 2-player turn-based stochastic games with a constant discount factor](#). *Journal of the ACM*, 60, 2013.
- Jean-Francois Hren and Rémi Munos. [Optimistic planning of deterministic systems](#). In *European Workshop on Reinforcement Learning*, 2008.
- Ruitong Huang, Mohammad M. Ajallooeian, Csaba Szepesvári, and Martin Müller. [Structured best-arm identification with fixed confidence](#). In *Algorithmic Learning Theory*, 2017.
- Emilie Kaufmann and Wouter M Koolen. [Monte-carlo tree search by best-arm identification](#). In *Neural Information Processing Systems*, 2017.
- Michael Kearns, Yishay Mansour, and Andrew Y. Ng. [A sparse sampling algorithm for near-optimal planning in large Markov decision processes](#). In *International Conference on Artificial Intelligence and Statistics*, 1999.
- Levente Kocsis and Csaba Szepesvári. [Bandit-based Monte-Carlo planning](#). In *European Conference on Machine Learning*, 2006.
- Edouard Leurent and Odalric-Ambrym Maillard. [Practical open-loop optimistic planning](#). In *European Conference on Machine Learning*, 2019.
- Volodymyr Mnih, Adria Puigdomenech Badia, Mehdi Mirza, Alex Graves, Timothy Lillicrap, Tim Harley, David Silver, and Koray Kavukcuoglu. [Asynchronous methods for deep reinforcement learning](#). In *International Conference on Machine Learning*, 2016.

- Gergely Neu, Anders Jonsson, and Vicenç Gómez. **A unified view of entropy-regularized Markov decision processes**. In *arXiv:1705.07798*, 2017.
- Martin L Puterman. *Markov Decision Processes: Discrete Stochastic Dynamic Programming*. John Wiley & Sons, New York, NY, 1994.
- John Schulman, Sergey Levine, Pieter Abbeel, Michael Jordan, and Philipp Moritz. **Trust region policy optimization**. In *International Conference on Machine Learning*, 2015.
- John Schulman, Xi Chen, and Pieter Abbeel. **Equivalence between policy gradients and soft Q-learning**. In *arXiv:1704.06440*, 2017.
- Richard S. Sutton. **Dyna, an integrated architecture for learning, planning, and reacting**. *SIGART Bulletin*, 2(4):160–163, 1991.
- Richard S. Sutton and Andrew G. Barto. *Reinforcement learning: An introduction*. The MIT Press, second edition, 2018.
- Balázs Szörényi, Gunnar Kedenburg, and Rémi Munos. **Optimistic planning in Markov decision processes using a generative model**. In *Neural Information Processing Systems*, 2014.
- Thomas J Walsh, Sergiu Goschin, and Michael L Littman. **Integrating sample-based planning and model-based reinforcement learning**. *AAAI Conference on Artificial Intelligence*, 2010.

A Preliminaries

A.1 General definition of value functions

We consider the general definition of value functions in Equation 4 and we assume that all the functions F_s satisfy

1. F_s is differentiable,
2. $\forall x \in \mathbb{R}^K, 0 < \|\nabla F_s(x)\|_1 \leq 1$,
3. (nonnegative gradient) $\forall x \in \mathbb{R}^K, \nabla F_s(x) \succeq 0$,
4. (L -smooth) There exists $L \geq 0$ such that for any $x_0, x \in \mathbb{R}^K$,

$$|F_s(x) - F_s(x_0) - (x - x_0)^\top \nabla F_s(x_0)| \leq L \|x - x_0\|_2^2,$$

which is the case for the functions LogSumExp_λ and $-\text{LogSumExp}_{-\lambda}$ that we study in the present paper. In particular, the second requirement implies that F_s is 1-Lipschitz,

$$\forall x, y \in \mathbb{R}^K, |F_s(x) - F_s(y)| \leq \|x - y\|_\infty.$$

For this more general definition, we modify the output of `sampleV` when $\varepsilon < \kappa$ to

$$\mathbf{output} = F_s(\widehat{Q}_s) - (\widehat{Q}_s)^\top \nabla F_s(\widehat{Q}_s) + (R + \gamma \widehat{v}) \|\nabla F_s(\widehat{Q}_s)\|_1$$

and the action sampled in `sampleV` is sampled according to

$$A \sim \frac{\nabla F_s(\widehat{Q}_s)}{\|\nabla F_s(\widehat{Q}_s)\|_1} \quad \text{instead of} \quad A \sim \nabla F_s(\widehat{Q}_s).$$

A.2 Other definitions

The constant M_λ is defined as

$$M_\lambda \triangleq \sup_{s \in \mathcal{S}} |F_s(0)|.$$

For any $c \in \mathbb{R}$, the function $\text{clip}_c : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is defined component-wise as

$$\text{clip}_c(x)_i = \begin{cases} 0 & \text{if } x_i \leq 0, \\ x_i & \text{if } -c < x_i < c, \\ c & \text{if } x_i \geq c. \end{cases}$$

B Sample complexity

Theorem 1. *Let $n(\varepsilon, \delta')$ be the number of oracle calls before SmoothCruiser terminates. For any state $s \in \mathcal{S}$ and $\varepsilon, \delta' > 0$,*

$$n(\varepsilon, \delta') \leq \frac{c_1}{\varepsilon^4} \log\left(\frac{c_2}{\delta'}\right) \left[c_3 \log\left(\frac{c_4}{\varepsilon}\right) \right]^{\log_2(c_5(\log(\frac{c_2}{\delta'})))} = \tilde{\mathcal{O}}\left(\frac{1}{\varepsilon^4}\right),$$

where c_1, c_2, c_3, c_4 , and c_5 are constants that depend only on K, L , and γ .

To bound the sample complexity, we make the following steps.

- Proposition 1 bounds the number of recursive calls of `sampleV` in the uniform sampling phase ($\varepsilon \geq \kappa$) and is similar to the results of Kearns et al. [1999].
- Lemma 1 bounds the number of recursive calls of `sampleV` when $\varepsilon < \kappa$.
- By noticing that the number of recursive calls of `sampleV` is equal to the number of oracle calls, we bound the sample complexity of SmoothCruiser in Theorem 1.

Let $n_{\text{sampleV}}(s, \varepsilon, \delta')$ be the total number of recursive calls to `sampleV` after an initial call with parameters (s, ε) , and including the initial call. Since this number does not depend on the state s , we denote it by $n_{\text{sampleV}}(\varepsilon, \delta')$.

Proposition 1. *Let $\varepsilon \geq \kappa$. For all $h \in \mathbb{N}$, $\forall \varepsilon$ such that $\frac{(1+M_\lambda)\sqrt{\gamma}^h}{1-\gamma} \leq \varepsilon \leq \frac{1+M_\lambda}{1-\gamma}$, we have*

$$\begin{aligned} n_{\text{sampleV}}(\varepsilon, \delta') &\leq \gamma^{\frac{1}{2}H(\varepsilon)(H(\varepsilon)-1)} \left(\frac{2\alpha(\delta')}{\varepsilon^2} \right)^{H(\varepsilon)} \\ &\leq \gamma^{\frac{1}{2}H(\kappa)(H(\kappa)-1)} \left(\frac{2\alpha(\delta')}{\kappa^2} \right)^{H(\kappa)} \end{aligned}$$

where

$$H(\varepsilon) = \left\lceil 2 \log_\gamma \left(\frac{\varepsilon(1-\gamma)}{1+M_\lambda} \right) \right\rceil$$

and

$$\alpha(\delta') = \frac{18(1+M_\lambda)^2 K}{(1-\gamma)^4 (1-\sqrt{\gamma})^2} \log \left(\frac{2K}{\delta'} \right)$$

Proof. We want to prove that $n_{\text{sampleV}}(\varepsilon, \delta') \leq G(\varepsilon)$, where

$$G(\varepsilon) = \gamma^{\frac{1}{2}H(\varepsilon)(H(\varepsilon)-1)} \left(\frac{2\alpha(\delta')}{\varepsilon^2} \right)^{H(\varepsilon)}$$

We proceed by induction on h .

Base case Let $h = 0$. We have $\varepsilon = \frac{1+M_\lambda}{1-\gamma}$, which implies $n_{\text{sampleV}}(\varepsilon, \delta') = 1$ and $G(\varepsilon) = 1$ (since $H(\varepsilon) = 0$). Hence, the proposition is true for $h = 0$.

Induction hypothesis Assume true for h .

Induction step Let $\varepsilon \geq \frac{(1+M_\lambda)\sqrt{\gamma}^{h+1}}{1-\gamma}$. Since $\frac{\varepsilon}{\sqrt{\gamma}} \geq \frac{(1+M_\lambda)\sqrt{\gamma}^h}{1-\gamma}$, we use the induction hypothesis to obtain

$$\begin{aligned} n_{\text{sampleV}}(\varepsilon, \delta') &= \underbrace{1}_{\text{current call}} + \underbrace{KN(\varepsilon)n_{\text{sampleV}}\left(\frac{\varepsilon}{\sqrt{\gamma}}, \delta'\right)}_{\text{calls in estimateQ}} \\ &\leq \frac{2\alpha(\delta')}{\varepsilon^2} n_{\text{sampleV}}\left(\frac{\varepsilon}{\sqrt{\gamma}}, \delta'\right) \\ &\leq \frac{2\alpha(\delta')}{\varepsilon^2} \gamma^{\frac{1}{2}(H(\varepsilon)-1)(H(\varepsilon)-2)} \left(\frac{\gamma 2\alpha(\delta')}{\varepsilon^2} \right)^{H(\varepsilon)-1}, \quad \text{since } H\left(\frac{\varepsilon}{\sqrt{\gamma}}\right) = H(\varepsilon) - 1 \\ &= \gamma^{\frac{1}{2}H(\varepsilon)(H(\varepsilon)-1)} \left(\frac{2\alpha(\delta')}{\varepsilon^2} \right)^{H(\varepsilon)}, \end{aligned}$$

which completes the proof. □

Lemma 1. *Let $\varepsilon \leq \kappa$. For all $h \in \mathbb{N}$, $\forall \varepsilon \geq \kappa\sqrt{\gamma}^h$, we have*

$$n_{\text{sampleV}}(\varepsilon, \delta') \leq \eta_1 \left[\log_{\frac{1}{\gamma}} \left(\frac{\kappa/\gamma}{\varepsilon} \right) \right]^{\eta_2(\delta')} \frac{1}{\varepsilon^2}$$

where

$$\begin{aligned}\kappa &= \frac{1 - \sqrt{\gamma}}{KL} \\ \eta_1 &= \kappa^2 n_{\text{sampleV}}(\kappa, \delta') \\ \eta_2(\delta') &= \log_2 \left(\frac{\gamma}{1 - \gamma} \frac{2\beta(\delta')}{\kappa} \right) \\ \beta(\delta') &= \frac{18(1 + M_\lambda)^2 K^2 L}{(1 - \gamma)^4 (1 - \sqrt{\gamma})^3} \log \left(\frac{2K}{\delta'} \right)\end{aligned}$$

under the condition that

$$\log_2 \left(\frac{\gamma}{1 - \gamma} \frac{2\beta(\delta')}{\kappa} \right) \geq 0, \quad \text{i.e.,} \quad \beta(\delta') \geq \frac{(1 - \gamma)(1 - \sqrt{\gamma})}{2\gamma KL} \quad (6)$$

which is satisfied by choosing δ' small enough.

Proof. First, let us define some auxiliary quantities,

$$B_1(\varepsilon) \triangleq \left[\log_{\frac{1}{\gamma}} \left(\frac{\kappa/\gamma}{\varepsilon} \right) \right]^{\eta_2(\delta')}, \quad (7)$$

$$B_2(\varepsilon) \triangleq \frac{\eta_1}{\varepsilon^2} \quad \text{and} \quad (8)$$

$$B(\varepsilon) \triangleq B_1(\varepsilon) B_2(\varepsilon) \quad (9)$$

We want to prove that $n_{\text{sampleV}}(\varepsilon, \delta') \leq B(\varepsilon)$ and we proceed by induction on h .

Base case For $h = 0$, we have $\varepsilon \geq \kappa$ and, by assumption, $\varepsilon \leq \kappa$. Therefore, $\varepsilon = \kappa$. It can be easily verified that $B(\kappa) = n_{\text{sampleV}}(\kappa, \delta')$, hence the lemma is true for $h = 0$.

Induction hypothesis Assume that the lemma is true for h .

Induction step Let $\varepsilon \geq \kappa \sqrt{\gamma}^{h+1}$. We have that

$$\begin{aligned}n_{\text{sampleV}}(\varepsilon, \delta') &= \underbrace{1}_{\text{current call}} + \underbrace{n_{\text{sampleV}} \left(\frac{\varepsilon}{\sqrt{\gamma}}, \delta' \right)}_{\text{call in line 11 of sampleV}} + \underbrace{KN(\sqrt{\kappa\varepsilon}) n_{\text{sampleV}} \left(\sqrt{\frac{\kappa\varepsilon}{\gamma}}, \delta' \right)}_{\text{calls in estimateQ}} \\ &= 1 + n_{\text{sampleV}} \left(\frac{\varepsilon}{\sqrt{\gamma}}, \delta' \right) + \frac{\beta(\delta')}{\varepsilon} n_{\text{sampleV}} \left(\sqrt{\frac{\kappa\varepsilon}{\gamma}}, \delta' \right) \\ &\leq n_{\text{sampleV}} \left(\frac{\varepsilon}{\sqrt{\gamma}}, \delta' \right) + \frac{2\beta(\delta')}{\varepsilon} n_{\text{sampleV}} \left(\sqrt{\frac{\kappa\varepsilon}{\gamma}}, \delta' \right)\end{aligned}$$

Since $\varepsilon \geq \kappa \sqrt{\gamma}^{h+1}$ and $\varepsilon \leq \kappa$, we have $\sqrt{\frac{\kappa\varepsilon}{\gamma}} \geq \frac{\varepsilon}{\sqrt{\gamma}} \geq \kappa \sqrt{\gamma}^h$. This allows us to use our induction hypothesis to get

$$n_{\text{sampleV}}(\varepsilon, \delta') \leq B \left(\frac{\varepsilon}{\sqrt{\gamma}} \right) + \frac{2\beta(\delta')}{\varepsilon} B \left(\sqrt{\frac{\kappa\varepsilon}{\gamma}} \right).$$

We will need the equation below, which is easily verified as

$$\log \left(\frac{\kappa}{\sqrt{\frac{\kappa\varepsilon}{\gamma}}} \right) = \frac{1}{2} \log \left(\frac{\kappa/\gamma}{\varepsilon} \right) \quad (10)$$

We have that

$$\begin{aligned} \frac{B\left(\frac{\varepsilon}{\sqrt{\gamma}}\right)}{B(\varepsilon)} &= \frac{B_1\left(\frac{\varepsilon}{\sqrt{\gamma}}\right)}{B_1(\varepsilon)} \frac{B_2\left(\frac{\varepsilon}{\sqrt{\gamma}}\right)}{B_2(\varepsilon)} \\ &= \gamma \underbrace{\left[\frac{\log\left(\frac{\kappa/\gamma}{\varepsilon}\right) - \frac{1}{2} \log \frac{1}{\gamma}}{\log\left(\frac{\kappa/\gamma}{\varepsilon}\right)} \right]^{\eta_2(\delta')}}_{<1} \\ &\leq \gamma, \end{aligned}$$

where we used the assumption that $\eta_2(\delta') \geq 0$.

Also we get that

$$\begin{aligned} \frac{B\left(\sqrt{\frac{\kappa\varepsilon}{\gamma}}\right)}{B(\varepsilon)} &= \frac{\varepsilon\gamma}{\kappa} \frac{B_1\left(\sqrt{\frac{\kappa\varepsilon}{\gamma}}\right)}{B_1(\varepsilon)} \\ &= \frac{\varepsilon\gamma}{\kappa} \left[\frac{\log_{\frac{1}{\gamma}}\left(\frac{\kappa}{\sqrt{\frac{\kappa\varepsilon}{\gamma}}\gamma}\right)}{\log_{\frac{1}{\gamma}}\left(\frac{\kappa/\gamma}{\varepsilon}\right)} \right]^{\eta_2(\delta')} \\ &= \frac{\varepsilon\gamma}{\kappa} \left[\frac{\frac{1}{2} \log_{\frac{1}{\gamma}}\left(\frac{\kappa/\gamma}{\varepsilon}\right)}{\log_{\frac{1}{\gamma}}\left(\frac{\kappa/\gamma}{\varepsilon}\right)} \right]^{\eta_2(\delta')} \\ &= \frac{\varepsilon\gamma}{\kappa} \left(\frac{1}{2}\right)^{\eta_2(\delta')} = \frac{\varepsilon\gamma(1-\gamma)}{\kappa} \frac{\kappa}{\gamma} \frac{1}{2\beta(\delta')} = \frac{(1-\gamma)\varepsilon}{2\beta(\delta')} \end{aligned}$$

Finally, we obtain

$$\begin{aligned} n_{\text{sampleV}}(\varepsilon, \delta') &\leq B\left(\frac{\varepsilon}{\sqrt{\gamma}}\right) + \frac{2\beta(\delta')}{\varepsilon} B\left(\sqrt{\frac{\kappa\varepsilon}{\gamma}}\right) \\ &\leq \gamma B(\varepsilon) + \frac{2\beta(\delta')}{\varepsilon} \frac{(1-\gamma)\varepsilon}{2\beta(\delta')} B(\varepsilon) \\ &= B(\varepsilon), \end{aligned}$$

which proves the lemma. \square

Now we can prove Theorem 1, which is restated below.

Theorem. *Let $n(\varepsilon, \delta')$ be the number of calls to the generative model (oracle) before the algorithm terminates. For any state $s \in \mathcal{S}$ and $\varepsilon, \delta' > 0$,*

$$n(\varepsilon, \delta') \leq \frac{c_1}{\varepsilon^4} \log\left(\frac{c_2}{\delta'}\right) \left[c_3 \log\left(\frac{c_4}{\varepsilon}\right) \right]^{\log_2(c_5(\log(\frac{c_2}{\delta'})))} = \tilde{\mathcal{O}}\left(\frac{1}{\varepsilon^4}\right)$$

where c_1, c_2, c_3, c_4 and c_5 are constants that depend only on K, L and γ .

Proof. First, notice that the number of calls to the generative model is smaller than the total number of calls to `sampleV`. `SmoothCruiser` makes one call to `estimateQ`, which makes $N(\varepsilon)$ calls to `sampleV`. If $\varepsilon \geq \kappa$, Proposition 1 shows that the sample complexity is bounded by a constant. Lemma 1 bounds the sample complexity for $\varepsilon \leq \kappa$, and we use it to bound $n(\varepsilon, \delta')$:

$$\begin{aligned} n(\varepsilon, \delta') &= N(\varepsilon) n_{\text{sampleV}}(\varepsilon, \delta') \\ &\leq N(\varepsilon) \eta_1 \left[\log_{\frac{1}{\gamma}}\left(\frac{\kappa/\gamma}{\varepsilon}\right) \right]^{\eta_2(\delta')} \frac{1}{\varepsilon^2} \\ &\leq \frac{c_1}{\varepsilon^4} \log\left(\frac{c_2}{\delta'}\right) \left[c_3 \log\left(\frac{c_4}{\varepsilon}\right) \right]^{\log_2(c_5(\log(\frac{c_2}{\delta'})))} = \tilde{\mathcal{O}}\left(\frac{1}{\varepsilon^4}\right) \end{aligned}$$

by using the definition of $N(\varepsilon)$ for $\varepsilon \leq \kappa$ and the definition of $\eta_2(\delta')$ in Lemma 1.

The constants are given by:

- $c_1 = \frac{18(1+M_\lambda)^2 n_{\text{sampleV}}(\kappa, \delta')}{K^2 L^2 (1-\gamma)^4}$;
- $c_2 = 2K$;
- $c_3 = [\log(1/\gamma)]^{-1}$;
- $c_4 = (1 - \sqrt{\gamma})/(\gamma KL)$;
- $c_5 = \frac{36(1+M_\lambda)^2 \gamma K^3 L^2}{(1-\gamma)^5 (1-\sqrt{\gamma})^4}$.

□

C Consistency

Theorem 2. *For any $s \in \mathcal{S}$, $\varepsilon > 0$, and $\delta > 0$, there exists a δ' that depends on ε and δ such that the output $\widehat{V}(s)$ of `SmoothCruiser`(s, ε, δ') satisfies*

$$\mathbb{P}\left[|\widehat{V}(s) - V(s)| > \varepsilon\right] \leq \delta.$$

and such that $n(\varepsilon, \delta') = \mathcal{O}(1/\varepsilon^{4+c})$ for any $c > 0$.

To prove that our algorithm outputs a good estimate of the value function with high probability, we proceed as follows:

- In Lemma 2, we prove that the output of `sampleV`, conditioned on an event \mathcal{A} , is a low-bias estimate of the true value function, and that \mathcal{A} happens with high probability;
- Given Lemma 2, the proof of Theorem 2 is straightforward.

Throughout the proof, we will make distinctions between two cases:

- **Case 1:** $\kappa \leq \varepsilon < \frac{1+M_\lambda}{1-\gamma}$
- **Case 2:** $\varepsilon < \kappa$

C.1 Definitions

We define the function $\zeta(\varepsilon)$ as

$$\zeta(\varepsilon) = \begin{cases} \varepsilon, & \text{if } \kappa \leq \varepsilon < \frac{1+M_\lambda}{1-\gamma}, \\ \sqrt{\kappa\varepsilon}, & \text{if } \varepsilon < \kappa, \\ \infty, & \text{otherwise.} \end{cases} \quad (11)$$

Define `params`(s, ε) as the (random) set of parameters used to call `sampleV` after a call to `sampleV`(s, ε), that is

$$\text{params}(s, \varepsilon) = \left\{ \left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right) \text{ for } k = 1, \dots, N(\varepsilon); a \in \mathcal{A} \right\} \quad (12)$$

in case 1 and

$$\text{params}(s, \varepsilon) = \left\{ \left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right) \text{ for } k = 1, \dots, N(\varepsilon); a \in \mathcal{A} \right\} \cup \left\{ \left(Z_{s,A}, \frac{\varepsilon}{\sqrt{\gamma}} \right) \right\} \quad (13)$$

in case 2, where $Z_{s,a}^{(k)}$ are the next states sampled in `estimateQ` and $Z_{s,A}$ is the next state sampled `sampleV`(s, ε).

A call to `sampleV`(s, ε) makes one call to `estimateQ`. Denote the output of this call to `estimateQ` by $\widehat{Q}_s^\varepsilon$. We define the event $\mathcal{A}(s, \varepsilon)$ as follows:

$$\mathcal{A}(s, \varepsilon) = \begin{cases} \left\{ \|\widehat{Q}_s^\varepsilon - Q_s\|_\infty \leq \zeta(\varepsilon) \right\} \cap \mathcal{B}(s, \varepsilon), & \text{if } 0 < \varepsilon < \frac{1+M_\lambda}{1-\gamma}, \\ \Omega, & \text{if } \varepsilon \geq \frac{1+M_\lambda}{1-\gamma}. \end{cases} \quad (14)$$

where Ω is the whole sample space and

$$\mathcal{B}(s, \varepsilon) = \bigcap_{(z,e) \in \text{params}(s,\varepsilon)} \mathcal{A}(z, e) \quad (15)$$

Define C_γ as:

$$C_\gamma = \frac{3(1+M_\lambda)}{(1-\gamma)^2} \quad (16)$$

C.2 Proofs

Lemma 2. Let $\widehat{V}_\varepsilon(s) = \text{sampleV}(s, \varepsilon)$. For all $h \in \mathbb{N}$, $s \in \mathcal{S}$, $\varepsilon \geq \frac{(1+M_\lambda)\sqrt{\gamma}^h}{1-\gamma}$, we have:

- (i) $|\mathbb{E}[\widehat{V}_\varepsilon(s) | \mathcal{A}(s, \varepsilon)] - V(s)| \leq \varepsilon$, and
- (ii) $\mathbb{P}[|\widehat{V}_\varepsilon(s)| \leq C_\gamma | \mathcal{A}(s, \varepsilon)] = 1$
- (iii) $\mathbb{P}[\mathcal{A}(s, \varepsilon)] \geq 1 - \delta' n_{\text{sampleV}}(\varepsilon, \delta')$

where

$$n_{\text{sampleV}}(\varepsilon, \delta') = 1 + \sum_{(z,e) \in \text{params}(s,\varepsilon)} n_{\text{sampleV}}(e, \delta') \quad (17)$$

is the total number of recursive calls to `sampleV` after an initial call with parameters (s, ε) .

Proof. We proceed by induction over h .

(1) Base case. If $h = 0$, $\varepsilon \geq \frac{1+M_\lambda}{1-\gamma}$ and $\mathcal{A}(s, \varepsilon) = \Omega$. The output is then $\widehat{V}_\varepsilon(s) = 0$. Point (i) is verified by using the fact that $|V(s)| \leq \frac{1+M_\lambda}{1-\gamma} \leq \varepsilon$; points (ii) and (iii) are trivially verified.

(2) Induction hypothesis. Assume that (i), (ii) and (iii) are true for h .

(3) Induction step. Let $\varepsilon \geq \frac{(1+M_\lambda)\sqrt{\gamma}^{h+1}}{1-\gamma}$. This implies that $\varepsilon/\sqrt{\gamma}$ and $\zeta(\varepsilon)/\sqrt{\gamma}$ are both greater than $\frac{(1+M_\lambda)\sqrt{\gamma}^h}{1-\gamma}$, which will allow us to use our induction hypothesis.

We start by proving (iii).

Let $\widehat{Q}_s^\varepsilon = \text{estimateQ}(s, \zeta(\varepsilon))$. Let the reward $R_{s,a}^{(k)}$ and state $Z_{s,a}^{(k)}$ be the random variables associated to the k -th call to the generative model used to compute $\widehat{Q}_s^\varepsilon$ in `estimateQ`, for $k \in \{1, \dots, N(\varepsilon)\}$. Let

$$q_s^k(a) := R_{s,a}^{(k)} + \gamma \text{sampleV}\left(Z_{s,a}^{(k)}, \zeta(\varepsilon)/\sqrt{\gamma}\right) \quad (18)$$

and let

$$\bar{Q}_s^\varepsilon(a) = \frac{1}{N(\varepsilon)} \sum_{k=1}^{N(\varepsilon)} q_s^k(a) \quad (19)$$

so that:

$$\widehat{Q}_s^\varepsilon = \mathbf{clip}_{(1+M_\lambda)(1-\gamma)^{-1}} \left(\bar{Q}_s^\varepsilon(a) \right) \quad (20)$$

Using Fact 2, we have:

$$|\widehat{Q}_s^\varepsilon(a) - Q_s(a)| \leq |\bar{Q}_s^\varepsilon(a) - Q_s(a)| \quad (21)$$

$$\leq \underbrace{|\bar{Q}_s^\varepsilon(a) - \mathbb{E}[\bar{Q}_s^\varepsilon(a) | \mathcal{B}(s, \varepsilon)]|}_{\text{(I)}} + \underbrace{|\mathbb{E}[\bar{Q}_s^\varepsilon(a) | \mathcal{B}(s, \varepsilon)] - Q_s(a)|}_{\text{(II)}} \quad (22)$$

We'd like to use Hoeffding's inequality to bound (I) in probability. For that, we need to verify that the random variables $\{q_s^k(a)\}_{k=1}^{N(\varepsilon)}$ are bounded and independent conditionally on $\mathcal{B}(s, \varepsilon)$.

Boundedness. By induction hypothesis (ii) In the event $\mathcal{B}(s, \varepsilon)$, the random variables $\text{sampleV}\left(Z_{s,a}^{(k)}, \zeta(\varepsilon)/\sqrt{\gamma}\right)$, for all k , are bounded by C_γ . Using the fact that the rewards are in $[0, 1]$ and that $C_\gamma \geq 1/(1-\gamma)$, we obtain $q_s^k(a)$ is also bounded by C_γ .

Independence. Let $E_k = \mathcal{A}(Z_{s,a}^k, \zeta(\varepsilon)/\sqrt{\gamma})$. For any $t \in \mathbb{R}^{N(\varepsilon)}$, the characteristic function of $\{q_s^k(a)\}_{k=1}^{N(\varepsilon)}$ conditionally on $\mathcal{B}(s, \varepsilon)$ is given by

$$\begin{aligned} \mathbb{E} \left[\exp \left(i \sum_k t_k q_s^k(a) \right) \middle| \mathcal{B}(s, \varepsilon) \right] &\stackrel{\text{(a)}}{=} \mathbb{E} \left[\exp \left(i \sum_k t_k q_s^k(a) \right) \middle| \bigcap_k E_k \right] \\ &= \frac{\mathbb{E} [\exp(i \sum_k t_k q_s^k(a)) \prod_k \mathbb{I}_{\{E_k\}}]}{\mathbb{E} [\prod_k \mathbb{I}_{\{E_k\}}]} \\ &= \frac{\mathbb{E} [\prod_k \exp(it_k q_s^k(a)) \mathbb{I}_{\{E_k\}}]}{\mathbb{E} [\prod_k \mathbb{I}_{\{E_k\}}]} \\ &\stackrel{\text{(b)}}{=} \frac{\prod_k \mathbb{E} [\exp(it_k q_s^k(a)) \mathbb{I}_{\{E_k\}}]}{\prod_k \mathbb{E} [\mathbb{I}_{\{E_k\}}]} \\ &= \prod_k \mathbb{E} [\exp(it_k q_s^k(a)) | E_k] \\ &\stackrel{\text{(c)}}{=} \prod_k \mathbb{E} [\exp(it_k q_s^k(a)) | \mathcal{B}(s, \varepsilon)] \end{aligned}$$

which is justified by

- (a) Definition of $\mathcal{B}(s, \varepsilon)$ and the fact that $\{q_s^k(a)\}_{k=1}^{N(\varepsilon)}$ are independent of $\mathcal{A}\left(Z_{s,A}, \frac{\varepsilon}{\sqrt{\gamma}}\right)$;
- (b) The random variables $\{q_s^k(a)\}_{k=1}^{N(\varepsilon)}$ are independent and the events $\{E_k\}_{i=1}^{N(\varepsilon)}$ are also independent;
- (c) The random variable $q_s^k(a)$ is independent of every E_j for $j \neq k$.

Since the characteristic function of $\{q_s^k(a)\}_{k=1}^{N(\varepsilon)}$ is the product of their characteristic functions, these random variables are independent given $\mathcal{B}(s, \varepsilon)$.

Now we can use Hoeffding's inequality:

$$\begin{aligned}
& \mathbb{P} \left[\left| \overline{Q}_s^\varepsilon(a) - \mathbb{E} \left[\overline{Q}_s^\varepsilon(a) \mid \mathcal{B}(s, \varepsilon) \right] \right| \geq (1 - \sqrt{\gamma}) \zeta(\varepsilon) \mid \mathcal{B}(s, \varepsilon) \right] \\
&= \mathbb{P} \left[\left| \frac{1}{N(\varepsilon)} \sum_{k=1}^{N(\varepsilon)} q_s^k(a) - \mathbb{E} \left[q_s^k(a) \mid \mathcal{B}(s, \varepsilon) \right] \right| \geq (1 - \sqrt{\gamma}) \zeta(\varepsilon) \mid \mathcal{B}(s, \varepsilon) \right] \\
&\leq 2 \exp \left(- \frac{N(\varepsilon) (1 - \sqrt{\gamma})^2 \zeta(\varepsilon)^2}{2C_\gamma^2} \right) \\
&\leq \frac{\delta'}{K}
\end{aligned}$$

And **(II)** is bounded by using the induction hypothesis **(i)**:

$$\begin{aligned}
& \left| \mathbb{E} \left[q_s^k(a) \mid \mathcal{B}(s, \varepsilon) \right] - Q_s(a) \right| \\
&\stackrel{\text{(a)}}{=} \gamma \left| \mathbb{E} \left[\text{sampleV} \left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right) \mid \mathcal{B}(s, \varepsilon) \right] - \mathbb{E} \left[V(Z_{s,a}^{(k)}) \mid \mathcal{B}(s, \varepsilon) \right] \right| \\
&\stackrel{\text{(b)}}{=} \gamma \left| \mathbb{E} \left[\text{sampleV} \left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right) \mid \mathcal{A} \left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right) \right] - \mathbb{E} \left[V(Z_{s,a}^{(k)}) \mid \mathcal{A} \left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right) \right] \right| \\
&\stackrel{\text{(c)}}{=} \gamma \left| \mathbb{E} \left[\mathbb{E} \left[\text{sampleV} \left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right) \mid Z_{s,a}^{(k)}, \mathcal{A} \left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right) \right] - V(Z_{s,a}^{(k)}) \mid \mathcal{A} \left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right) \right] \right| \\
&\stackrel{\text{(d)}}{\leq} \gamma \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \\
&= \sqrt{\gamma} \zeta(\varepsilon)
\end{aligned}$$

which is justified by the following:

- (a) $\mathbb{E} \left[R_{s,a}^{(k)} \mid \mathcal{B}(s, \varepsilon) \right] = \mathbb{E} \left[R_{s,a}^{(k)} \right]$, since the reward depends only on s, a ;
- (b) The term $\left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right)$ depends on $\mathcal{B}(s, \varepsilon)$ only through $\mathcal{A} \left(Z_{s,a}^{(k)}, \frac{\zeta(\varepsilon)}{\sqrt{\gamma}} \right)$;
- (c) Law of total expectation;
- (d) Consequence of induction hypothesis **(i)**.

Putting together the bounds for **(I)** and **(II)** and doing an union bound over all actions, we obtain:

$$\mathbb{P} \left[\left\| \widehat{Q}_s^\varepsilon - Q_s \right\|_\infty \geq \zeta(\varepsilon) \mid \mathcal{B}(s, \varepsilon) \right] \leq \delta'$$

We can now give a lower bound to the probability of the event $\mathcal{A}(s, \varepsilon)$. Let

$$\mathcal{E} = \left\{ \left\| \widehat{Q}_s^\varepsilon - Q_s \right\|_\infty < \zeta(\varepsilon) \right\} \tag{23}$$

We have:

$$\begin{aligned}
\mathbb{P}[\mathcal{A}(s, \varepsilon)] &\geq \mathbb{P}[\mathcal{E} \cap \mathcal{B}(s, \varepsilon)] \\
&= \mathbb{P}[\mathcal{E} \mid \mathcal{B}(s, \varepsilon)] \mathbb{P}[\mathcal{B}(s, \varepsilon)] \\
&= \left(1 - \mathbb{P}[\mathcal{E}^c \mid \mathcal{B}(s, \varepsilon)]\right) \mathbb{P}[\mathcal{B}(s, \varepsilon)] \\
&\geq \mathbb{P}[\mathcal{B}(s, \varepsilon)] - \delta' \\
&\geq 1 - \delta' n_{\text{sampleV}}(\varepsilon, \delta')
\end{aligned}$$

since

$$\begin{aligned}
\mathbb{P}[\mathcal{B}(s, \varepsilon)] &= 1 - \mathbb{P}[\mathcal{B}(s, \varepsilon)^c] \\
&= 1 - \mathbb{P}\left[\bigcup_{(z, e) \in \text{params}(s, \varepsilon)} \mathcal{A}(z, e)^c\right] \\
&\geq 1 - \sum_{(z, e) \in \text{params}(s, \varepsilon)} \mathbb{P}[\mathcal{A}(z, e)^c] \\
&\geq 1 - \delta' \sum_{(z, e) \in \text{params}(s, \varepsilon)} n_{\text{sampleV}}(e, \delta') \quad \text{by induction hypothesis (iii)} \\
&= 1 - \delta' (n_{\text{sampleV}}(\varepsilon, \delta') - 1)
\end{aligned}$$

This proves **(iii)**. Now, let's prove **(i)**.

For any event \mathcal{E} , we write

$$\mathbb{E}_{\mathcal{E}}[\cdot] = \mathbb{E}[\cdot \mid \mathcal{E}]$$

Case 1. We start with case 1, $\kappa \leq \varepsilon < \frac{1+M\lambda}{1-\gamma}$, where $\zeta(\varepsilon) = \varepsilon$ and

$$\widehat{V}_{\varepsilon}(s) = F_s(\widehat{Q}_s^{\varepsilon}) \tag{24}$$

We have:

$$\begin{aligned}
|\mathbb{E}_{\mathcal{A}(s, \varepsilon)}[\widehat{V}_{\varepsilon}(s)] - V(s)| &= |\mathbb{E}_{\mathcal{A}(s, \varepsilon)}[F_s(\widehat{Q}_s^{\varepsilon}) - F_s(Q_s)]| \\
&\leq \mathbb{E}_{\mathcal{A}(s, \varepsilon)}[|F_s(\widehat{Q}_s^{\varepsilon}) - F_s(Q_s)|] \\
&\leq \mathbb{E}_{\mathcal{A}(s, \varepsilon)}[\|\widehat{Q}_s^{\varepsilon}(a) - Q_s(a)\|_{\infty}] \\
&\leq \zeta(\varepsilon) = \varepsilon
\end{aligned}$$

and **(i)** is verified for case 1.

Case 2. Consider now the case 2, $\varepsilon < \kappa$, where $\zeta(\varepsilon) = \sqrt{\kappa\varepsilon}$.

Let A be the action following the distribution $\frac{\nabla F_s(\widehat{Q}_s^{\varepsilon})}{\|\nabla F_s(\widehat{Q}_s^{\varepsilon})\|_1}$, and let the reward $R_{s,A}$ and the state $Z_{s,A}$ be the random variables associated to the call to the generative model with parameters (s, A) . Let $\widehat{v} = \text{sampleV}(Z_{s,A}, \varepsilon/\sqrt{\gamma})$. The output in this case is given by

$$\widehat{V}_{\varepsilon}(s) = F_s(\widehat{Q}_s^{\varepsilon}) - (\widehat{Q}_s^{\varepsilon})^{\top} \nabla F_s(\widehat{Q}_s^{\varepsilon}) + (R + \gamma \widehat{v}) \|\nabla F_s(\widehat{Q}_s^{\varepsilon})\|_1 \tag{25}$$

Let

$$\begin{aligned} Q_s(A) &= \mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[R_{s,A} + \gamma V(Z_{s,A}) \mid A, \widehat{Q}_s^\varepsilon \right] \\ &= \mathbb{E}_{\mathcal{A}(s,\varepsilon)} [R_{s,A} + \gamma V(Z_{s,A}) \mid A] \end{aligned}$$

and let

$$\widetilde{V}(s) = \mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[F_s(\widehat{Q}_s^\varepsilon) - (\widehat{Q}_s^\varepsilon)^\top \nabla F_s(\widehat{Q}_s^\varepsilon) + Q_s(A) \|\nabla F_s(\widehat{Q}_s^\varepsilon)\|_1 \right] \quad (26)$$

We have

$$\begin{aligned} & |\mathbb{E}_{\mathcal{A}(s,\varepsilon)} [\widehat{V}_\varepsilon(s)] - \widetilde{V}(s)| \\ & \stackrel{\text{(a)}}{=} \gamma |\mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[\mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[\text{sampleV} \left(Z_{s,A}, \frac{\varepsilon}{\sqrt{\gamma}} \right) - V(Z_{s,A}) \mid A, \widehat{Q}_s^\varepsilon, Z_{s,A} \right] \|\nabla F_s(\widehat{Q}_s^\varepsilon)\|_1 \right]| \\ & \stackrel{\text{(b)}}{=} \gamma |\mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[\left(\mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[\text{sampleV} \left(Z_{s,A}, \frac{\varepsilon}{\sqrt{\gamma}} \right) \mid A, \widehat{Q}_s^\varepsilon, Z_{s,A} \right] - V(Z_{s,A}) \right) \|\nabla F_s(\widehat{Q}_s^\varepsilon)\|_1 \right]| \\ & \stackrel{\text{(c)}}{\leq} \gamma \mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[\left| \mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[\text{sampleV} \left(Z_{s,A}, \frac{\varepsilon}{\sqrt{\gamma}} \right) \mid A, \widehat{Q}_s^\varepsilon, Z_{s,A} \right] - V(Z_{s,A}) \right| \right] \\ & \stackrel{\text{(d)}}{=} \gamma \mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[\left| \mathbb{E}_{\mathcal{A}(Z_{s,A}, \varepsilon/\sqrt{\gamma})} \left[\text{sampleV} \left(Z_{s,A}, \frac{\varepsilon}{\sqrt{\gamma}} \right) \mid Z_{s,A} \right] - V(Z_{s,A}) \right| \right] \\ & \stackrel{\text{(e)}}{\leq} \gamma \frac{\varepsilon}{\sqrt{\gamma}} = \sqrt{\gamma} \varepsilon \end{aligned}$$

which is justified by the following points:

- (a) The reward depend only on s, a and law of total expectation;
- (b) $V(Z_{s,A})$ is a function of $Z_{s,A}$ and no other random variable;
- (c) Jensen's inequality and the fact that $\|\nabla F_s(\widehat{Q}_s^\varepsilon)\|_1 \leq 1$;
- (d) Given $Z_{s,A}$, the term $\text{sampleV} \left(Z_{s,A}, \frac{\varepsilon}{\sqrt{\gamma}} \right)$ depends on $\mathcal{A}(s,\varepsilon)$ only through $\mathcal{A}(Z_{s,A}, \varepsilon/\sqrt{\gamma})$;
- (e) Induction hypothesis **(i)**.

Now, $\mathbb{E}_{\mathcal{A}(s,\varepsilon)} [Q_s(A) \|\nabla F_s(\widehat{Q}_s^\varepsilon)\|_1]$ can be written as

$$\begin{aligned} & \mathbb{E}_{\mathcal{A}(s,\varepsilon)} [Q_s(A) \|\nabla F_s(\widehat{Q}_s^\varepsilon)\|_1] \\ & = \mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[\mathbb{E}_{\mathcal{A}(s,\varepsilon)} [Q_s(A) \mid \widehat{Q}_s^\varepsilon] \|\nabla F_s(\widehat{Q}_s^\varepsilon)\|_1 \right] \\ & = \mathbb{E}_{\mathcal{A}(s,\varepsilon)} [Q_s^\top \nabla F_s(\widehat{Q}_s^\varepsilon)] \end{aligned}$$

so that $\widetilde{V}(s)$ is given by

$$\widetilde{V}(s) = \mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[F_s(\widehat{Q}_s^\varepsilon) + (Q_s - \widehat{Q}_s^\varepsilon)^\top \nabla F_s(\widehat{Q}_s^\varepsilon) \right] \quad (27)$$

Finally, we bound the difference between $\widetilde{V}(s)$ and $V(s)$:

$$\begin{aligned}
|\tilde{V}(s) - V(s)| &\leq \mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[|F_s(\hat{Q}_s^\varepsilon) + (Q_s - \hat{Q}_s^\varepsilon)^\top \nabla F_s(\hat{Q}_s^\varepsilon) - V(s)| \right] \\
&\leq L \mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[\|Q_s - \hat{Q}_s^\varepsilon\|_2^2 \right] \\
&\stackrel{\text{(a)}}{\leq} KL \mathbb{E}_{\mathcal{A}(s,\varepsilon)} \left[\|Q_s - \hat{Q}_s^\varepsilon\|_\infty^2 \right] \\
&\leq KL\zeta(\varepsilon)^2 \\
&= KL\kappa\varepsilon \\
&= (1 - \sqrt{\gamma})\varepsilon
\end{aligned}$$

by using the fact that we are on $\mathcal{A}(s, \varepsilon)$ and (a) uses the fact that for all $x \in \mathbb{R}^K$, $\|x\|_2^2 \leq K\|x\|_\infty^2$.

We can now prove (i) for case 2:

$$|\mathbb{E}_{\mathcal{A}(s,\varepsilon)} [\hat{V}_\varepsilon(s)] - V(s)| \leq |\mathbb{E}_{\mathcal{A}(s,\varepsilon)} [\hat{V}_\varepsilon(s)] - \tilde{V}(s)| + |\tilde{V}(s) - V(s)| \quad (28)$$

$$\leq \sqrt{\gamma}\varepsilon + (1 - \sqrt{\gamma})\varepsilon = \varepsilon \quad (29)$$

Finally, let's prove (ii).

Case 1. In this case, $\hat{V}_\varepsilon(s) = F_s(\hat{Q}_s^\varepsilon)$ with $\|\hat{Q}_s^\varepsilon\|_\infty \leq (1 + M_\lambda)/(1 - \gamma)$, since each component of \hat{Q}_s^ε is clipped and lie in the interval $\left[0, \frac{1+M_\lambda}{1-\gamma}\right]$. The assumptions on F_s imply that $|\hat{V}_\varepsilon(s)| \leq \frac{1+M_\lambda}{1-\gamma} \leq C_\gamma$.

Case 2. In this case, we have:

$$\begin{aligned}
|\hat{V}_\varepsilon(s)| &\leq |F_s(\hat{Q}_s^\varepsilon) - (\hat{Q}_s^\varepsilon)^\top \nabla F_s(\hat{Q}_s^\varepsilon)| + |R + \gamma\hat{v}| \|\nabla F_s(\hat{Q}_s^\varepsilon)\|_1 \\
&\leq 2\|\hat{Q}_s^\varepsilon\|_\infty + M_\lambda + 1 + \gamma C_\gamma \\
&\leq \frac{2(1 + M_\lambda)}{1 - \gamma} + M_\lambda + 1 + \gamma C_\gamma \\
&\leq C_\gamma
\end{aligned}$$

since $|\hat{v}| \leq C_\gamma$ by induction hypothesis (ii).

This proves (ii) for case 2:

$$\mathbb{P} \left[|\hat{V}(s)| \leq C_\gamma \mid \mathcal{A}(s, \varepsilon) \right] = 1 \quad (30)$$

□

Now, we can prove Theorem 2, which is restated as follows:

Theorem. Let $\hat{V}(s)$ be the output of `SmoothCruiser`(s, ε, δ'). For any state $s \in \mathcal{S}$ and $\varepsilon, \delta' > 0$,

$$\mathbb{P} \left[|\hat{V}(s) - V(s)| > \varepsilon \right] \leq \delta' n(\varepsilon, \delta').$$

Proof. Let $\hat{Q}_s = \text{estimateQ}(s, \varepsilon)$. We have $\hat{V}(s) = F_s(\hat{Q}_s)$. As in the proof of Lemma 2, let the reward $R_{s,a}^{(k)}$ and state $Z_{s,a}^{(k)}$ be the random variables associated to the k -th call to the generative model used to compute $\hat{Q}_s(a)$ in `estimateQ`, for $k \in \{1, \dots, N(\varepsilon)\}$.

We have:

$$\widehat{Q}_s(a) = \frac{1}{N(\varepsilon)} \sum_{k=1}^{N(\varepsilon)} R_{s,a}^{(k)} + \gamma \text{sampleV}\left(Z_{s,a}^{(k)}, \varepsilon/\sqrt{\gamma}\right) \quad (31)$$

Consider the event \mathcal{E} defined by:

$$\mathcal{E} = \bigcap_{k=1}^{N(\varepsilon)} \mathcal{A}\left(Z_{s,a}^{(k)}, \frac{\varepsilon}{\sqrt{\gamma}}\right) \quad (32)$$

By the same arguments as in the proof of Lemma 2, we have:

- In \mathcal{E} , we have $\|\widehat{Q}_s - Q_s\|_\infty \leq \varepsilon$;
- $\mathbb{P}[\mathcal{E}] \geq 1 - \delta' N(\varepsilon) n_{\text{sampleV}}(\varepsilon, \delta') = 1 - \delta' n(\varepsilon, \delta')$.

This implies the result, since $|\widehat{V}(s) - V(s)| \leq \|\widehat{Q}_s - Q_s\|_\infty$.

Now, for every $\varepsilon > 0$ and every $\delta > 0$, we need to be able to find a value of δ' such that $\delta' n(\varepsilon, \delta') \leq \delta$. That is, given ε and δ , we need to find δ' such that

$$\delta' \frac{c_1}{\varepsilon^4} \log\left(\frac{c_2}{\delta'}\right) \left[c_3 \log\left(\frac{c_4}{\varepsilon}\right) \right]^{\log_2(c_5(\log(\frac{c_2}{\delta'})))} \leq \delta. \quad (33)$$

Such value exists, since the term on the LHS tends to 0 as $\delta' \rightarrow 0$, and it depends on ε . We will show that this dependence is polynomial when $\varepsilon \rightarrow 0$.

Let $\delta' = \varepsilon^5$. There exists a value $\tilde{\varepsilon}$ that depends on δ such that:

$$\forall \varepsilon \leq \tilde{\varepsilon}, \quad \varepsilon^5 \frac{c_1}{\varepsilon^4} \log\left(\frac{c_2}{\varepsilon^5}\right) \left[c_3 \log\left(\frac{c_4}{\varepsilon}\right) \right]^{\log_2(c_5(\log(\frac{c_2}{\varepsilon^5})))} \leq \delta. \quad (34)$$

since the term on the LHS tends to 0 as $\varepsilon \rightarrow 0$, as a consequence of Proposition 2.

Putting it all together, we can choose δ' as follows:

$$\delta' = \begin{cases} \tilde{\delta} \text{ such that } \tilde{\delta} \frac{c_1}{\varepsilon^4} \log\left(\frac{c_2}{\tilde{\delta}}\right) \left[c_3 \log\left(\frac{c_4}{\varepsilon}\right) \right]^{\log_2(c_5(\log(\frac{c_2}{\tilde{\delta}})))} \leq \delta, & \text{if } \varepsilon > \tilde{\varepsilon}, \\ \varepsilon^5, & \text{if } \varepsilon \leq \tilde{\varepsilon} \end{cases} \quad (35)$$

which is $\mathcal{O}(\varepsilon^5)$.

Proposition 3 implies that, for this choice of δ' , the sample complexity is still of order $\mathcal{O}(1/\varepsilon^{4+c})$ for any $c > 0$. □

D Auxiliary results

Fact 1. For all $s \in \mathcal{S}$ and all $x \in \mathbb{R}^K$, we have $F_s(x) \leq \|x\|_\infty + \sup_s |F_s(0)|$.

Proof. By the assumptions on F_s , we have:

$$|F_s(x)| = |F_s(x) - F_s(0) + F_s(0)| \leq |F_s(x) - F_s(0)| + |F_s(0)| \quad (36)$$

$$\leq \|x - 0\|_\infty + |F_s(0)| \leq \|x\|_\infty + \sup_s |F_s(0)|. \quad (37)$$

□

Fact 2. Let $x, q \in \mathbb{R}^d$ be such that $0 \leq q_i \leq c$ for all i . Let $\tilde{x} = \mathbf{clip}_c(x)$. Then, $\|\tilde{x} - q\|_\infty \leq \|x - q\|_\infty$.

Proof. For any $i \in \{1, \dots, d\}$, we have $|\tilde{x}_i - q_i| \leq |x_i - q_i|$, since $0 \leq q_i \leq c$. The result follows. \square

Proposition 2. $\forall a, b, c > 0$

$$\lim_{x \rightarrow \infty} \frac{1}{x^c} \exp(a[\log \log(x^b)]^2) = 0$$

Proof. We have

$$\begin{aligned} \frac{1}{x^c} \exp(a[\log \log(x^b)]^2) &= \exp(a[\log \log(x^b)]^2 - c \log x) \\ &= \exp\left(a[\log u]^2 - \frac{c}{b}u\right), \quad \text{by setting } u = \log(x^b) \end{aligned}$$

And, for any $k > 0$, we have

$$\lim_{u \rightarrow \infty} \log^2 u - ku = -\infty. \quad (38)$$

which allows us to conclude. \square

Proposition 3. If we set $\delta' = \delta'(\varepsilon) = \varepsilon^5$, we have:

$$n(\varepsilon, \delta'(\varepsilon)) = \mathcal{O}\left(\frac{1}{\varepsilon^{4+c}}\right), \quad \forall c > 0$$

Proof. We have:

$$\begin{aligned} n_{\text{sampleV}}(\varepsilon, \delta'(\varepsilon)) &\leq \eta_1 \left[\log_{\frac{1}{\gamma}} \left(\frac{\bar{\varepsilon}/\gamma}{\varepsilon} \right) \right]^{\eta_2(\varepsilon^3)} \frac{1}{\varepsilon^2} \\ &= \underbrace{\left[\log_{\frac{1}{\gamma}} \left(\frac{\bar{\varepsilon}/\gamma}{\varepsilon} \right) \right]^{\log_2(k \log(\frac{2K}{\varepsilon^3}))}}_{(A)} \frac{1}{\varepsilon^2} \end{aligned}$$

where k is a constant that does not depend on ε . The term (A) can be rewritten as:

$$\begin{aligned} \left[\log_{\frac{1}{\gamma}} \left(\frac{\bar{\varepsilon}/\gamma}{\varepsilon} \right) \right]^{\log_2(k \log(\frac{2K}{\varepsilon^3}))} &= \left[c_1 \log \left(\frac{c_2}{\varepsilon} \right) \right]^{c_3 \log[k \log(\frac{c_4}{\varepsilon^3})]} \\ &= \exp \left\{ c_3 \log \left[k \log \left(\frac{c_4}{\varepsilon^3} \right) \right] \log \left(c_1 \log \left(\frac{c_2}{\varepsilon} \right) \right) \right\} \end{aligned}$$

which can be shown to be $\mathcal{O}(\frac{1}{\varepsilon^c})$ for any $c > 0$ by applying proposition 2 after some algebraic manipulations.

Hence,

$$n_{\text{sampleV}}(\varepsilon, \delta'(\varepsilon)) = \frac{1}{\varepsilon^2} \mathcal{O}\left(\frac{1}{\varepsilon^c}\right) = \mathcal{O}\left(\frac{1}{\varepsilon^{2+c}}\right), \quad \forall c > 0.$$

Since we have

$$n(\varepsilon, \delta') = N(\varepsilon) n_{\text{sampleV}}(\varepsilon, \delta')$$

with $N(\varepsilon) = \tilde{\mathcal{O}}(1/\varepsilon^2)$, this proves the result. \square

Corollary 1. *If we set $\delta' = \delta'(\varepsilon) = \varepsilon^5$, we have:*

$$\lim_{\varepsilon \rightarrow 0} \delta'(\varepsilon) n(\varepsilon, \delta'(\varepsilon)) = 0$$

Proof. It is an immediate consequence of proposition 3 by taking $c \in]0, 1[$. □

E On other smooth approximations of the max

In this paper we focus on the LogSumExp_λ function as a smooth approximation to the maximum function. Yet our proof is more general and can handle any approximation of the max function which verifies the properties listed in Section 5. For instance let's consider the following regularization of the Bellman equation:

$$F(Q) = \max_{(\pi_a)_{a \in \mathcal{A}}} \sum_{a \in \mathcal{A}} (Q_a \cdot \pi_a + \lambda \sqrt{\pi_a}) \quad (39)$$

This smooth function is particularly interesting because it approximates the distribution of pulled arms of the UCB algorithm by taking $\lambda = 2c \cdot \sqrt{\frac{\ln(n)}{n}}$ (see 41 and notice that $\pi_a^* \cdot n$ approximates n_a). We show that this smooth approximation of the maximum verifies the assumptions made in Section 5. We have

$$F(Q) = \sum_{a \in \mathcal{A}} (Q_a \cdot \pi_a^* + \lambda \sqrt{\pi_a^*}) \quad (40)$$

and we can show that $\nabla_Q F(Q) = \pi^*$. Therefore point 1, 2 and 3 of Section 5 are verified. Now by differentiating with respect to π this time:

$$\forall a \in \mathcal{A} \quad Q_a + \frac{\lambda}{2\sqrt{\pi_a^*}} = U \quad (41)$$

where U is the Lagrange multiplier. Using the fact that $\sum_{a \in \mathcal{A}} \pi_a^* = 1$, we get

$$\sum_{a \in \mathcal{A}} \left(\frac{\lambda/2}{U - Q_a} \right)^2 = 1 \quad (42)$$

Because $U > \max_a \pi_a^*$ the derivative of the left side with respect to U is positive for all $Q_a \in [0, (1 + M_\lambda)/(1 - \gamma)]^{|\mathcal{A}|}$. Using the inverse function theorem we get that U is differentiable with respect to Q and that $\pi_a^* = \left(\frac{\lambda/2}{U - Q_a} \right)^2$ is also differentiable with respect to Q . Finally because $[0, (1 + M_\lambda)/(1 - \gamma)]^{|\mathcal{A}|}$ is compact we can conclude that F is L -smooth for some $L \geq 0$ verifying point 4 of Section 5.

F Experimental validation of the theoretical results

In this section, we present the experiments we made to verify the correctness of our sample complexity bounds (Theorem 1) and of our consistency results (Theorem 2).

F.1 Checking the sample complexity guarantee

The key step for proving Theorem 1 is using Lemma 1, that bounds the number of calls to the generative model made by a call to $\text{sampleV}(s, \varepsilon)$.

Figure 2 shows the simulated number of calls to the generative model made by $n_{\text{sampleV}}(\varepsilon, \delta')$ as a function of $1/\varepsilon$ and compares it to our theoretical bound in Lemma 1 and to the number of calls that would be required by a Sparse Sampling strategy, which corresponds to the bound in Proposition 1

extrapolated to all values of ε . The simulated values were obtained by computing the following recurrence for several values of ε :

$$n_{\text{sampleV}}^{\text{sim}}(\varepsilon, \delta') = \begin{cases} 1 + n_{\text{sampleV}}^{\text{sim}}\left(\frac{\varepsilon}{\sqrt{\gamma}}, \delta'\right) + KN(\sqrt{\kappa\varepsilon})n_{\text{sampleV}}^{\text{sim}}\left(\sqrt{\frac{\kappa\varepsilon}{\gamma}}, \delta'\right), & \text{if } \varepsilon < \kappa, \\ \gamma^{\frac{1}{2}H(\varepsilon)(H(\varepsilon)-1)}\left(\frac{2\alpha(\delta')}{\varepsilon^2}\right)^{H(\varepsilon)}, & \text{otherwise.} \end{cases}$$

Figure 3 shows the number of calls to the generative model made by n_{sampleV} as a function of the regularization parameter λ in order to achieve a relative error of 0.01^7 and its ratio with respect to the number of calls that would be required by Sparse Sampling in the same setting. We see that fewer samples are required as the regularization increases. We also see that, for small λ , there is no advantage with respect to Sparse Sampling, but SmoothCruiser has a very large advantage when the regularization λ grows.




Figure 2: Simulated number of calls to the generative model made by $n_{\text{sampleV}}(\varepsilon, \delta')$ as a function of $1/\varepsilon$ compared to our theoretical bound (Lemma 1) and to the number of calls that would be required by a Sparse Sampling strategy. The parameters used were: $\gamma = 0.2$, $\delta' = 0.1$, $K = 2$ and $\lambda = 0.1$.

F.2 Checking the consistency guarantee

Using our MCTS analogy in Section 3.3, the two most computationally costly operations of SmoothCruiser are the `selectAction` and the `evaluateLeaf` functions. They both rely on estimates of the Q function with some required accuracy. Hence, for a *sanity-check*, we implemented the function `sampleV` by replacing its calls to `estimateQ(s, accuracy)` by the true Q function at state s plus some accuracy-dependent noise, and we denote this simplified version of `sampleV` by `sampleVcheck`. This allowed us to verify that our bounds for the bias of the `sampleV` outputs (Lemma 2) are correct. After N_{sim} calls to `sampleVcheck`(ε, δ'), we compute the error

$$\widehat{\Delta}(s, \varepsilon) = \frac{1}{N_{\text{sim}}} \sum_{i=1}^{N_{\text{sim}}} (\widehat{V}_i(s, \varepsilon) - V(s)) \quad (43)$$

⁷ We set $\varepsilon = 0.01V_{\lambda}^{\text{max}}$, where $V_{\lambda}^{\text{max}} = (1 + \lambda \log K)/(1 - \gamma)$ is an upper bound on the regularized value function.




Figure 3: Number of calls to the generative model made by n_{sampleV} as a function of the regularization parameter λ in order to achieve a relative error of 0.01 (left) and its ratio with respect to the number of calls that would be required by Sparse Sampling in the same setting (right). The parameters used were: $\gamma = 0.2$, $\delta' = 0.1$ and $K = 2$.

where s is a reference state and $\widehat{V}_i(s, \varepsilon)$ is the output of the i -th call to $\text{sampleV}_{\text{check}}(s, \varepsilon)$. Lemma 2 states that, for some high probability event B , we have $-\varepsilon \leq \mathbb{E}[\widehat{\Delta}(s, \varepsilon)|B] \leq \varepsilon$. Hence, for large N_{sim} , we should have $-\varepsilon \leq \widehat{\Delta}(s, \varepsilon) \leq \varepsilon$ approximately.

Table 1 shows simulated values of $\widehat{\Delta}(s, \varepsilon)$ and their standard deviations for different environments. The value of N_{sim} was chosen so that $\widehat{\Delta}(s, \varepsilon)$ is close to its mean, by using Hoeffding's inequality and assuming that $\widehat{V}_i(s, \varepsilon)$ is bounded by C_γ (which holds with high probability, by Lemma 2).

Environment	$\widehat{\Delta}(s, \varepsilon)$
5-Chain	$(-1.21 \pm 1.65) \times 10^{-2}$
10-Chain	$(-1.20 \pm 1.63) \times 10^{-2}$
5x5-GridWorld	$(-0.71 \pm 2.04) \times 10^{-2}$
10x10-GridWorld	$(-0.71 \pm 2.03) \times 10^{-2}$

Table 1: Simulated values of $\widehat{\Delta}(s, \varepsilon)$ and its standard deviation for different environments, for $\varepsilon = 0.35$. The value of ε was chosen such that $\varepsilon \leq \kappa/4$ in all environments. The parameters used were: $N_{\text{sim}} = 32723$, $\gamma = 0.2$ and $\lambda = 10$. The n -Chain environments have $K = 2$ and n states and the $n \times n$ -GridWorld environments have $K = 4$ and n^2 states.

The code for the experiments is at <https://github.com/omardrwh/smoothcruiser-check>.