



# MarLIN

## Marine Information Network

Information on the species and habitats around the coasts and sea of the British Isles

# *Saccharina latissima* and/or *Saccorhiza polyschides* on exposed infralittoral rock

MarLIN – Marine Life Information Network  
Marine Evidence-based Sensitivity Assessment (MarESA) Review

Dr Keith Hiscock

2002-05-31

A report from:

The Marine Life Information Network, Marine Biological Association of the United Kingdom.

**Please note.** This MarESA report is a dated version of the online review. Please refer to the website for the most up-to-date version [<https://www.marlin.ac.uk/habitats/detail/237>]. All terms and the MarESA methodology are outlined on the website (<https://www.marlin.ac.uk>)


This review can be cited as:

Hiscock, K. 2002. [*Saccharina latissima*] and/or [*Saccorhiza polyschides*] on exposed infralittoral rock. In Tyler-Walters H. and Hiscock K. (eds) *Marine Life Information Network: Biology and Sensitivity Key Information Reviews*, [on-line]. Plymouth: Marine Biological Association of the United Kingdom. DOI <https://dx.doi.org/10.17031/marlinhab.237.1>



The information (TEXT ONLY) provided by the Marine Life Information Network (MarLIN) is licensed under a Creative Commons Attribution-Non-Commercial-Share Alike 2.0 UK: England & Wales License. Note that images and other media featured on this page are each governed by their own terms and conditions and they may or may not be available for reuse. Permissions beyond the scope of this license are available [here](http://www.marlin.ac.uk). Based on a work at [www.marlin.ac.uk](http://www.marlin.ac.uk)


(page left blank)



*Saccharina latissima* and/or *Saccorhiza polyschides* on exposed infralittoral rock.

Photographer: Paul Brazier

Copyright: Joint Nature Conservation Committee (JNCC)



Researched by Dr Keith Hiscock

Refereed by This information is not refereed.

## Summary

### ☰ UK and Ireland classification

EUNIS 2008	A3.122	<i>Laminaria saccharina</i> and/or <i>Saccorhiza polyschides</i> on exposed infralittoral rock
JNCC 2015	IR.HIR.KSed.SlatSac	<i>Saccharina latissima</i> and/or <i>Saccorhiza polyschides</i> on exposed infralittoral rock
JNCC 2004	IR.HIR.KSed.LsacSac	<i>Laminaria saccharina</i> and/or <i>Saccorhiza polyschides</i> on exposed infralittoral rock
1997 Biotope	IR.EIR.KFaR.LsacSac	<i>Laminaria saccharina</i> and/or <i>Saccorhiza polyschides</i> on exposed infralittoral rock

### 🔍 Description

A forest or park of the fast-growing, opportunistic kelps *Saccharina latissima* and/or *Saccorhiza polyschides* often occurs on seasonally unstable or scoured infralittoral rock. The substratum varies from large boulders in exposed areas to smaller boulders and cobbles in areas of moderate wave

exposure or nearby bedrock. In these cases, movement of the substratum during winter storms prevents a longer-lived forest of *Laminaria hyperborea* from becoming established. This biotope may also develop on bedrock where it is affected by its close proximity to unstable substrata. Other fast-growing algae such as *Ulva* spp., *Alaria esculenta*, *Desmarestia* spp. and *Chorda filum* are often present. This biotope can be found below the *Laminaria hyperborea* zone (EIR.LhypFa or EIR.LhypR), especially where close to a rock/ sand interface (subjected to sand scour in winter?). Some *Laminaria hyperborea* plants may occur in this biotope, but they are typically small since the plants do not survive many years. In St Kilda, this biotope is present on steep/vertical rock between the sublittoral fringe of *Alaria esculenta* and the *Laminaria hyperborea* forest below. In such places this biotope occurs because intense wave action in winter storms is too severe to allow *Laminaria hyperborea* to develop and remain in shallow water. (Information taken from the Marine Biotope Classification for Britain and Ireland, Version 97.06: Connor *et al.*, 1997a, b). NB. *Saccharina latissima* is now considered to be a synonym of *Saccharina latissima*.

### ↓ Depth range

0-5 m, 5-10 m, 10-20 m, 20-30 m

### Additional information

None.

### ✓ Listed By

- none -

### Further information sources

Search on:



## Habitat review

### Ecology

#### Ecological and functional relationships

The species in this biotope are either short lived opportunist species or are mobile species that migrate into the area. Kelps provide a habitat for a variety of encrusting bryozoans and the holdfasts may be colonized by mobile species including polychaete worms, crustaceans and prosobranch molluscs. The limpet *Patella pellucida* may feed on the kelps especially *Saccorhiza polyschides*. Grazing species such as the sea urchin *Echinus esculentus* are likely to keep rocks clear of a dense growth of erect algae. Kelps may provide local shelter for small fish such as the two-spotted goby *Gobiusculus flavescens*.

#### Seasonal and longer term change

This biotope will be highly changeable in relation especially to the onset of winter storms when most of the algae can be expected to be removed by scour. However, some parts of algae are likely to remain and, in less stormy winters, the biotope might remain largely intact and begin to show growth of longer lived kelps especially *Laminaria hyperborea*.

#### Habitat structure and complexity

Complexity in the biotope is brought about particularly by the structures in the kelp forest where fronds stipes and holdfasts provide different habitats for occupancy by associated species. For instance, *Saccorhiza polyschides* are known to shelter large animals such as large polychaetes, squat lobsters and fish inside the bulbous holdfast, while amphipods, brittle stars and polychaetes occur in the space between the base of the bulb and the rock surface to which it is attached (McKenzie & Moore, 1981). Similar species are associated with *Saccharina latissima* although holdfast faunas are of smaller species that live in the interstices between haptera. Fronds of kelps are grazed by urchins such as *Echinus esculentus* and *Paracentrotus lividus*, and the blue-rayed limpet *Patella pellucida*. There will also be significant shelter afforded by foliose algae. Rocks may have fissures and crevices providing further refuges for animal species.

#### Productivity

This biotope would appear to be productive of organic matter from seaweeds especially (primary production).

#### Recruitment processes

The dominant and characteristic species are recruited from planktonic larvae and spores. Other species such as fish, sea urchins and crustaceans are mainly migratory.

#### Time for community to reach maturity

Providing that sources of larvae, spores and mobile animals are nearby, the biotope would develop rapidly on new substrata so that, in appearance based on visually dominant species, it would be likely to be established in less than a year.

## Additional information

None.

## Preferences & Distribution

### Habitat preferences

<b>Depth Range</b>	0-5 m, 5-10 m, 10-20 m, 20-30 m
<b>Water clarity preferences</b>	
<b>Limiting Nutrients</b>	No information found
<b>Salinity preferences</b>	Full (30-40 psu)
<b>Physiographic preferences</b>	Open coast
<b>Biological zone preferences</b>	Infralittoral
<b>Substratum/habitat preferences</b>	Bedrock, Large to very large boulders, Small boulders, Cobbles
<b>Tidal strength preferences</b>	Moderately Strong 1 to 3 knots (0.5-1.5 m/sec.), Weak < 1 knot (<0.5 m/sec.)
<b>Wave exposure preferences</b>	Exposed, Moderately exposed, Very exposed
<b>Other preferences</b>	Areas subject to scour.

### Additional Information

The biotope is characteristic of seasonally unstable or scoured infralittoral rock. The biotopes classification (Connor *et al.* 1997a and JNCC, 1999) suggest that the biotope is restricted to Scotland but suitable habitats most likely occur elsewhere where it might have been identified as MIR.SedK.Sac (*Saccorhiza polyschides* and other opportunistic kelps on disturbed sublittoral fringe rock), which occurs mainly in southwest Britain. However, restriction of the mapped distribution to Scotland suggests that the biotope is one that occurs in cold waters.

## Species composition

### Species found especially in this biotope

- None.

### Rare or scarce species associated with this biotope

-

### Additional information

None.

## Sensitivity review

### Sensitivity characteristics of the habitat and relevant characteristic species

IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, and IR.HIR.KSed.XKScrR are within the sediment-affected or disturbed kelp and seaweed communities (IR.HIR.KSed) habitat complex. As a result of nearby sediment scouring or seasonally unstable infralittoral rock, opportunistic brown seaweeds; *Desmarestia* spp., *Saccharina latissima* (formerly *Laminaria saccharina*) and *Saccorhiza polyschides* can proliferate. *Laminaria hyperborea* can be present within the community, however, due to the disturbed nature of IR.HIR.KSed biotopes, does not become fully established and sporophytes do not typically survive beyond a couple of seasons (Connor *et al.*, 2004).

Due to the disturbed nature of IR.HIR.KSed biotopes the understory community can vary locally and is characterized by scour tolerant or ephemeral red seaweeds, such as *Corallina officinalis*, *Plocamium cartilagineum*, *Chondrus crispus*, *Dilsea carnosa*, and encrusting coralline algae. Faunal diversity and abundance are also generally low and typically limited to encrusting bryozoans and/or sponges. In areas sheltered from sediment scour or sediment, the stability biological diversity increases and *Laminaria hyperborea* becomes more dominant (Connor *et al.*, 2004).

In undertaking this assessment of sensitivity, an account is taken of knowledge of the biology of all characterizing species/taxa in the biotope. However, 'indicative species' are particularly important in undertaking the assessment as they structure and characterize the biotope. For this sensitivity assessment the opportunistic brown seaweeds; *Desmarestia* spp., *Saccharina latissima* and *Saccorhiza polyschides* are the primary foci of research. Examples of important species groups are mentioned where appropriate.

### Resilience and recovery rates of habitat

*Desmarestia* spp., *Saccorhiza polyschides* and *Saccharina latissima* (formerly *Laminaria saccharina*) are opportunistic seaweeds which have relatively fast growth rates when compared to other perennial species, and dominate areas subject to recurrent disturbance or in areas where environmental conditions limit competition from *Laminaria hyperborea*.

*Desmarestia* spp. and *Saccharina latissima* are widely distributed in the north Atlantic from Svalbard to Portugal (Birkett *et al.*, 1998b; Connor *et al.*, 2004; Bekby & Moy 2011; Moy & Christie 2012), *Saccorhiza polyschides* from mid-Norway to Ghana, and present in parts of the Mediterranean (Lüning, 1990). *Desmarestia* spp. are a genus of annual seaweeds with a life expectancy of ca 8 months (Gagnon *et al.*, 2013). *Saccorhiza polyschides* is also termed as an annual and can reach maturity in 8 months, although sporophytes that do not reach maturity within the first growth season can overwinter and have a life expectancy of 16 months (Birkett *et al.*, 1998; Fernández, 2011), during which time fronds can reach a length of 3-4m (D. Birkett, pers. obs in Birkett *et al.*, 1998). *Saccharina latissima* is a perennial kelp which can reach maturity in 15-20 months and has a life expectancy of 2-4 years.

*Desmarestiales* and *Laminariales* have heteromorphic life strategies (Edwards, 1998). Mature sporophytes broadcast zoospores which settle onto rock and develop into gametophytes, following fertilization these germinate into juvenile sporophytes. Kelp zoospores are expected to have a large dispersal range, but zoospore density and the rate of successful fertilization decreases exponentially with distance from the parental source (Fredriksen *et al.*, 1995). Hence, recruitment following disturbance can be influenced by the proximity of mature kelp beds producing viable

zoospores to the disturbed area (Kain, 1979; Fredriksen *et al.*, 1995). The exact mechanism of zoospore release in *Desmarestia* spp. is unknown but it may occur during a period of senescence (mentioned below)(Gagnon *et al.*, 2013). *Saccorhiza polyschides* & *Saccharina latissima* both release zoospores from reproductive structures known as sori, located centrally on the blade (*Saccharina latissima* & *Saccorhiza polyschides*), stipe and holdfast/bulb (*Saccorhiza polyschides*).

*Desmarestia* spp. and *Saccorhiza polyschides* sporophytes appear from March-April, beyond which is a period of rapid growth. *Desmarestia* spp. reach their maximum size by September (ca 60 cm).

Sporophytes then begin to decay (known as the senescence period) and typically die off by late October (Edwards, 1998; Gagnon *et al.*, 2013). *Saccorhiza polyschides* sporophytes are capable of growing  $\leq 6.2$  cm per week (Norton, 1970; Fernández, 2011) and reaching a maximum length of 3-4 m (Birket *et al.*, 1998; Fernández, 2011). The onset of maturity triggers a phase of senescence in which growth ceases and the frond erodes, resulting in the blade becoming progressively smaller and by winter the entire sporophyte can disappear (Birket *et al.*, 1998; Fernández, 2011).

*Saccharina latissima* recruits appear in late winter early spring beyond which is a period of rapid growth, in late summer and autumn growth rates slow and spores are released from autumn to winter (Parke, 1948; Lüning, 1979; Birket *et al.*, 1998). The overall length of the sporophyte may not change during the growing season due to marginal erosion but the growth of the blade has been measured at 1.1 cm/day, with a total length addition of  $\geq 2.25$  m per year (Birket *et al.*, 1998).

Light intensity and temperature are key development triggers for *Desmarestia* spp. (Edwards, 1998). However, other factors, such as nutrient availability and the abundance of coralline algae may also influence recruitment (Edwards, 1998). *Desmarestia* spp. sporophytes are typically rare in areas with established kelp canopies but have rapid growth in response to increases in light intensity and changes from red-blue wave lengths, indicating an opportunistic life history when kelp canopies are thinned/cleared (Chapman & Burrows, 1970; Müller & Lüthe, 1981; Edwards, 1998). Edwards (1998) found *Desmarestia ligulata* recruitment was cued by seasonal changes in day length but the recruitment was increased in areas where kelp canopies were cleared. In kelp clearances, *Desmarestia ligulata* was capable of rapidly achieving ca 50-90% coverage whereas abundance remained low under kelp canopies at ca <10% coverage. Field and experimental observations of *Desmarestia aculeata* in Port Erin, UK have found that light intensity is a principal factor in the development of gametophyte and sporophyte development, and hence recruitment processes (Kain, 1966; Chapman & Burrows, 1970). In winter, a season in which *Desmarestia aculeata* sporophytes are absent from marine habitats, Kain (1966) collected visually bare stones from Port Erin, Isle of Man. When the stones were exposed to high illumination (2,780 lux) for 18 hours a day and maintained at 5°C, *Desmarestia aculeata* sporophytes grew successfully. Demonstrating that increases in light intensity are an important trigger for *Desmarestia* spp. growth and recruitment.

*Saccharina latissima* can be quite ephemeral in nature and appear early in algal succession. For example, Lienaas & Christie (1996) removed *Strongylocentrotus droebachiensis* from "Urchin Barrens" and observed a succession effect. The substratum was colonized initially by filamentous algae, but after a couple of weeks, these were out-competed and the habitat dominated by *Saccharina latissima* which themselves were subsequently out-competed by *Laminaria hyperborea*. In the Isle of Man, Kain (1975) cleared sublittoral blocks of *Laminaria hyperborea* at different times of the year for several years. The first colonizers and succession community differed between blocks and at what time of year the blocks were cleared. However, *Saccharina latissima* was an early colonizer, but within 2 years of clearance the blocks were dominated by *Laminaria hyperborea*.

**Resilience assessment.** All three canopy forming seaweeds that characterize IR.HIR.KSed.Sac,



IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, and IR.HIR.KSed.XKScrR are opportunistic species with rapid colonization and growth rates. Both *Desmarestia spp.* & *Saccorhiza polyschides* are capable of reaching maturity within a year. *Saccharina latissima* has been shown to be an early colonizer within macroalgal succession, appearing within 2 weeks of clearance. Therefore, resilience has been assessed as 'High'.

## Hydrological Pressures

	Resistance	Resilience	Sensitivity
<b>Temperature increase (local)</b>	<b>None</b>	<b>High</b>	<b>Medium</b>
	Q: High A: High C: High	Q: High A: High C: High	Q: High A: High C: High

*Saccorhiza polyschides* has a wide geographic distribution and can tolerate a wide range of temperatures. Sporophyte growth can occur from 3-24°C and gametophyte development from 5-25°C (Norton, 1977). Fernández (2011) however suggested that summer temperatures of >20°C sustained for longer than a period of 30 days may inhibit development and recruitment.

Desmarestiales are unusual in that they produce and accumulate sulphuric acid within intracellular vacuoles (McClintock *et al.*, 1982; Connor *et al.*, 2004; Gagnon *et al.*, 2013). Seasonal increases in temperatures limit the ability of the storage vacuoles to contain the acid and release it into the surrounding environment. The continued release of acid results in progressive decolourisation, tissue degradation and mortality of *Desmarestia* sporophytes (Gagnon *et al.*, 2013). Gagnon *et al.* (2013) exposed *Desmarestia viridis* samples during 30 hour salinity and temperature treatments. At 29 and 32psu (MNCR: Full Salinity scale) *Desmarestia viridis* was able to tolerate changes in temperature from 5-12°C, but exposure to 18°C was lethal to sporophytes.

The geographic distribution of *Saccharina latissima* is determined by the 19-20°C isotherm (Müller *et al.*, 2009). Gametophytes can develop in ≤23°C (Lüning, 1990), but the optimal temperature range for sporophyte growth is 10-15°C (Bolton & Lüning, 1982). Bolton & Lüning (1982) experimentally observed that sporophyte growth was inhibited by 50-70% at 20°C and following 7 days at 23°C all specimens completely disintegrated. In the field *Saccharina latissima* has significant regional variation in its acclimation to temperature change. For example Gerard & Dubois (1988) observed sporophytes of *Saccharina latissima* which were regularly exposed to ≥20°C could tolerate these temperatures, whereas sporophytes from other populations which rarely experience ≥17°C showed 100% mortality after 3 weeks of exposure to 20°C. Therefore, the response of *Saccharina latissima* to a change in temperatures is likely to be locally variable.

Anderson *et al.* (2011) transplanted *Saccharina latissima* from the Skagerrak region, Norway an area which has experienced a 50.7-83% *Saccharina latissima* decline since 2002 (Moy & Christie, 2012). Since 1960-2009 sea surface temperatures in the region have regularly exceeded 20°C (the temperature at which *Saccharina latissima* growth is severely inhibited) and so has the number of days which remain above 20°C. Anderson *et al.* (2011) hypothesised that high sea temperatures were indirectly linked to *Saccharina latissima* deforestation in the region, causing high ephytic loading of sporophyte fronds (estimated to cover 80 & 100% of transplanted sporophytes). High sea temperatures have been linked to the slow growth of *Saccharina latissima* which is likely to decrease the photosynthetic ability of, and increase the vulnerability of *Saccharina latissima* to ephytic loading, bacterial and viral attacks (Anderson *et al.*, 2011).

Desmarestiales are sensitive to high temperatures and low salinities (Gagnon *et al.*, 2013).

Desmarestiales are unusual in that they produce and accumulate sulphuric acid ( $H_2SO_4$ ) within intracellular vacuoles (McClintock *et al.*, 1982; Connor *et al.*, 2004; Gagnon *et al.*, 2013). Increases in temperatures and low salinities limit the ability of the storage vacuoles to contain the acid and release it into the surrounding environment. The continued release of acid results in progressive decolourisation, tissue degradation and mortality of *Desmarestia* sporophytes (Gagnon *et al.*, 2013). Gagnon *et al.* (2013) exposed *Desmarestia viridis* samples during 30 hour salinity and temperature treatments, observing at 29 and 32 psu (MNCR: Full Salinity scale) *Desmarestia viridis* was tolerant to changes in temperature from 5-12°C, exposure to 18°C was lethal to sporophytes. Furthermore, sporophytes that had already begun the senescence phase were exposed to  $10.8 \pm 0.3^\circ C$  and completely shed all tissue from the stipe and laterals within ca15 days, whereas those exposed to lower temperatures of  $2.5 \pm 0.1^\circ C$  lasted ca30 days. Gagnon *et al.* (2013) also observed *Desmarestia spp.* degraded progressively in low salinity treatments of 26, 23 & 20psu (<20psu was not tested). Therefore indicating *Desmarestia spp.* are highly sensitive to both high temperature (12-18°C) and low salinities (<26 psu).

IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, and IR.HIR.KSed.XKScrR are distributed throughout the UK (Connor *et al.*, 2004). Northern to southern Sea Surface Temperature (SST) ranges from 8-16°C in summer and 6-13°C in winter (Beszczynska-Möller & Dye, 2013)

**Sensitivity assessment.** Acute 5°C increases in temperature for a period of 1 month combined with high summer temperatures may exceed the threshold temperature of 18-20°C in biotopes within the south of the UK, which would likely cause mortality of *Desmarestia spp.*, and severely limit *Saccorhiza polyschides* recruitment & *Saccharina latissima* sporophyte growth. A 2°C increase in temperature for a period of 1 year would likely result in the exceeding an 18°C temperature threshold in the south of the UK. This temperature threshold would likely result in high mortality of *Desmarestia spp.* *Saccharina latissima* which are not acclimated to similar temperatures may also experience high and rapid mortality. Resistance has been assessed as '**None**', Resilience as '**High**'. Sensitivity has been assessed as '**Medium**'.

#### Temperature decrease (local)

**Low**

Q: High A: High C: High

**High**

Q: High A: High C: High

**Low**

Q: High A: High C: High

*Desmarestia spp.* are a cold water adapted genus with a polar distribution, which can grow abundantly in water temperatures of ca 0.5°C (Gagnon *et al.*, 2013). *Saccharina latissima* is also widely spread throughout the Arctic and has a lower temperature threshold for sporophyte growth at 0°C (Lüning, 1990). Subtidal red algae can survive at -2°C (Lüning, 1990; Kain & Norton, 1990). These temperatures are well below that considered within this pressure benchmark. *Desmarestia spp.* *Saccharina latissima* are therefore unlikely to be adversely affected by a decrease in temperature at the benchmark level.

*Saccorhiza polyschides* sporophyte growth can occur within a range from 3-24°C and gametophyte development within a range of 5-25°C (Norton, 1977). Norton (1977) experimentally observed that at 3-5°C gametophytes failed to develop into viable sporophytes. This temperature range also corresponds with *Saccorhiza polyschides* northern range edge (ca 65° 35'N, mid-Norway), above which the average winter temperature is  $\leq 4^\circ C$  (U.S. Navy, 1958).

IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, and IR.HIR.KSed.XKScrR are distributed throughout the UK (Connor *et al.*, 2004). Northern to southern Sea Surface

Temperature (SST) ranges from 8-16°C in summer and 6-13°C in winter (Beszczynska-Möller & Dye, 2013)

**Sensitivity assessment.** Both long-term and acute temperature decrease 2-5°C combined with low winter temperatures could negatively affect *Saccorhiza polyschides* recruitment in biotopes located in the north of the UK. Resistance has been assessed as 'Low', resilience as 'High'. Sensitivity has been assessed as 'Low'.

### Salinity increase (local)

Low

Q: Medium A: High C: High

High

Q: High A: Low C: High

Low

Q: Medium A: Low C: High

Lüning (1990) suggest that kelps are stenohaline, their general tolerance to salinity as a phenotypic group covering 16-50 psu over a 24 hr period. Optimal growth probably occurring between 30-35 psu (MNCR category-Full Salinity) and growth rates are likely to be affected by periodic salinity stress. Birkett *et al.* (1998b) suggested that long-term increases in salinity may affect kelp growth and may result in loss of affected kelp, and, therefore, loss of the biotope.

Karsten (2007) tested the photosynthetic ability of *Saccharina latissima* under acute 2 & 5 day exposure to salinity treatments ranging from 5-60 psu. A control experiment was also carried at 34 psu (Salinity under normal marine conditions). Between 25-55 psu *Saccharina latissima* had a high photosynthetic ability at >80% of the control levels.

**Sensitivity assessment.** The evidence suggests that *Saccharina latissima* can tolerate exposure to hypersaline conditions of  $\geq 40\text{‰}$  (MNCR full salinity range=30-40‰). Optimal salinities for other kelps are assumed to be 30-35 psu. Hypersaline tolerances for *Desmarestia spp.* are unknown. Resistance has been assessed as 'Low', resilience as 'High'. The sensitivity of this biotope to an increase in salinity has been assessed as 'Low'.

### Salinity decrease (local)

Low

Q: High A: High C: High

High

Q: High A: High C: High

Low

Q: High A: High C: High

*Saccorhiza polyschides* is affected by low salinities. Norton & South (1969) observed at  $\leq 9\text{‰}$  zoospores often burst due to internal osmotic pressure and none developed. At  $\leq 25\text{‰}$ , only 25% of gametophytes germinated and at  $\leq 20\text{‰}$  sporophyte growth was often retarded. At  $\leq 35\text{‰}$ , 76% of gametophytes germinated. These results demonstrate that at  $\leq 25\text{‰}$  recruitment may be inhibited and sporophyte growth retarded.

Karsten (2007) tested the photosynthetic ability of *Saccharina latissima* under acute 2 & 5 day exposure to salinity treatments ranging from 5-60 psu. A control experiment was also carried at 34 psu (Salinity under normal marine conditions). Between 25-55 psu *Saccharina latissima* had a high photosynthetic ability at >80% of the control levels. Hyposaline treatment of 10-20 psu led to a gradual decline of photosynthetic ability. After 2 days at 5 psu, *Saccharina latissima* photosynthetic ability was ~30% of control. After 5 days at 5 psu, *Saccharina latissima* specimens became bleached and showed signs of severe damage. The experiment was conducted on *Saccharina latissima* from the Arctic, and the authors suggest that at extremely low water temperatures (1-5°C) macroalgae acclimation to rapid salinity changes could be slower than at temperate latitudes. It is, therefore, possible that *Saccharina latissima* of the UK may be able to

acclimate to salinity changes more effectively and quicker.

Gagnon *et al.* (2013) observed *Desmarestia spp.* sporophytes degraded progressively in low salinity treatments of 26, 23 & 20 psu (<20 psu was not tested). *Desmarestia spp.* accumulate sulphuric acid throughout the growth season which is released when the sporophyte becomes stressed under high temperatures or low salinities. Acid release causes progressive degradation of the sporophyte and mortality. Gagnon *et al.* (2013) exposed *Desmarestia viridis* samples during 30 hour salinity and temperature treatments, observing *Desmarestia spp.* degraded progressively in low salinity treatments of 26, 23 & 20psu (<20psu was not tested).

**Sensitivity assessment.** A decrease in one MNCR salinity scale from “Full Salinity” (30-40psu) to “Reduced Salinity” (18-30 psu) may result in mortality of *Desmarestia spp.* inhibit *Saccorhiza polyschides* recruitment and inhibit *Saccharina latissima* photosynthesis. Resistance has been assessed as ‘**Low**’ resilience as ‘**High**’. The sensitivity of this biotope to a decrease in salinity has been assessed as ‘**Low**’.

#### Water flow (tidal current) changes (local)

**High**

Q: Medium A: High C: High

**High**

Q: High A: High C: High

**Not sensitive**

Q: Medium A: High C: High

Peteiro & Freire (2013) measured *Saccharina latissima* growth from 2 sites, the 1<sup>st</sup> had maximal water velocities of 0.3 m/sec and the 2<sup>nd</sup> 0.1 m/sec. At site 1 *Saccharina latissima* had significantly larger biomass than at site 2 (16kg /m to 12 kg /m respectively). Peteiro & Freire (2013) suggested that faster water velocities were beneficial to *Saccharina latissima* growth. However, Gerard & Mann (1979) measured *Saccharina latissima* productivity at greater water velocities and found *Saccharina latissima* productivity is reduced in moderately strong tidal streams ( $\leq 1\text{m/sec}$ ) when compared to weak tidal streams ( $<0.5\text{m/sec}$ ).

IR.HIR.KSed.SlatSac can be found from very strong ( $>3\text{m/sec}$ ) to very weak tidal streams. IR.HIR.KSed.Sac, IR.HIR.KSed.DesFilR, IR.HIR.KSed.XKScrR can be found from moderately strong (0.5-1.5m/sec) to weak tidal streams (0.5m/sec). An increase in tidal flow may increase local sediment mobility and scour, potentially increasing dislodgment of kelps (Birket *et al.*, 1998) and *Desmarestia spp.*

**Sensitivity assessment.** Due to the range of tidal streams which these biotopes can be found a change of 0.1m/s to 0.2m/s, is not likely to dramatically affect biotope structure. Resistance has been assessed as ‘**High**’, resilience as ‘**High**’. Sensitivity has been assessed as ‘**Not Sensitive**’.

#### Emergence regime changes

**Low**

Q: Low A: NR C: NR

**High**

Q: High A: Low C: High

**Low**

Q: Low A: NR C: NR

IR.HIR.KSed.Sac and IR.HIR.KSed.XKScr can be found on sublittoral fringe rock so that the characterizing species would be exposed during low spring tides. IR.HIR.KSed.SlatSac and IR.HIR.KSed.DesFilR are found in the infralittoral and as such would only be exposed on extreme low tides (Connor *et al.*, 2004).

An increase in emergence will result in an increased risk of desiccation and mortality of the dominant seaweeds (*Desmarestia spp.*, *Saccorhiza polyschides*, *Saccharina latissima*). Removal of canopy forming seaweeds has also been shown to increase desiccation and mortality of the understory macroalgae (Hawkins & Harkin, 1985). Providing that suitable substrata are present,

the biotope is likely to re-establish further down the shore within a similar emergence regime to that which existed previously. A decrease in emergence could, however, result in an extension of the biotope further up the shore, although its lower limit is still probably controlled by light penetration, competition, and grazing so that entire extent of the biotope may move.

**Sensitivity assessment.** Resistance has been assessed as 'Low'. Resilience as 'High'. The sensitivity of this biotope to a change in emergence is considered as 'Low'.

### Wave exposure changes (local)

High

Q: Low A: NR C: NR

High

Q: High A: Low C: High

Not sensitive

Q: Low A: NR C: NR

IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, IR.HIR.KSed.XKScrR are recorded from extremely exposed to wave sheltered sites and characterized by rapidly colonising macroalgae (Connor *et al.*, 2004). As a result of rapid recovery, the community is relatively resistant to disturbance when compared to other kelp biotopes (e.g. IR.HIR.Kfar.LhypR). Birkett *et al.*, (1998) suggest that *Saccharina latissima* and *Saccorhiza polyschides* are rarely present in areas of wave exposure, where they may be spatially out-competed by *Laminaria hyperborea*. However, the seasonally unstable nature of the substrata or periodic sediment scour within these biotopes is likely to inhibit the growth of long-lived species, such as *Laminaria hyperborea* and allow opportunistic species such as *Desmarestia spp.*, *Saccharina latissima* and *Saccorhiza polyschides* to proliferate. An increase in local wave height may increase local sediment mobility and scour, potentially increasing dislodgment of kelps (Birket *et al.*, 1998) and *Desmarestia spp.* The biotopes may appear sparse after winter storms but the biotope recovers again due to rapid colonization and growth by the dominant kelps and *Desmarestia spp.*

**Sensitivity assessment.** The biotope is dominated by rapid colonizing species that tolerate or rapidly recover from scour, siltation and burial. They occur across a broad wave exposure range, and, therefore, a change in nearshore significant wave height of 3-5% is not likely to have a significant effect on biotope structure. Resistance has been assessed as 'High', resilience as 'High'. Therefore, sensitivity has been assessed as 'Not Sensitive' at the benchmark level.

## Chemical Pressures

### Transition elements & organo-metal contamination

Resistance

Not Assessed (NA)

Q: NR A: NR C: NR

Resilience

Not assessed (NA)

Q: NR A: NR C: NR

Sensitivity

Not assessed (NA)

Q: NR A: NR C: NR

This pressure is **Not assessed** but evidence is presented where available.

Bryan (1984) suggested that the general order for heavy metal toxicity in seaweeds is: Organic Hg > inorganic Hg > Cu > Ag > Zn > Cd > Pb. Cole *et al.*, (1999) reported that Hg was very toxic to macrophytes. Similarly, Hopkin & Kain (1978) demonstrated sub-lethal effects of heavy metals on kelp gametophytes and sporophytes, including reduced growth and respiration. Sheppard *et al.* (1980) noted that increasing levels of heavy metal contamination along the west coast of Britain reduced species number and richness in holdfast fauna, except for suspension feeders which became increasingly dominant. Gastropods may be relatively tolerant of heavy metal pollution (Bryan, 1984). Although macroalgae species may not be killed, except by high levels of contamination, reduced growth rates may impair the ability of the biotope to recover from other



environmental disturbances. Thompson & Burrows (1984) observed the growth of *Saccharina latissima* sporophyte growth was significantly inhibited at 50 µg Cu/l, 1000 µg Zn/l and 50 µg Hg/l. Zoospores were found to be more intolerant and significant reductions in survival rates were observed at 25 µg Cu/l, 1000 µg Zn/l and 5 µg/l.

#### Hydrocarbon & PAH contamination

Not Assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

This pressure is **Not assessed** but evidence is presented where available.

IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, IR.HIR.KSed.XKScrR, are predominantly recorded in the sub-tidal (<0 m). These habitats are therefore not likely to come into contact with freshly released oil but only to sinking emulsified oil and oil adsorbed onto particles (Birket *et al.*, 1998). The mucilaginous slime layer coating of laminarians may protect them from smothering by oil. Hydrocarbons in solution reduce photosynthesis and may be algicidal. However, Holt *et al.* (1995) reported that oil spills in the USA and from the *Torrey Canyon* had little effect on kelp forests. Similarly, surveys of subtidal communities at a number of sites between 1-22.5m below chart datum showed no noticeable impacts of the Sea Empress oil spill and clean up (Rostron & Bunker, 1997). Laboratory studies of the effects of oil and dispersants on several red algae species, including *Delesseria sanguinea* (Grandy 1984; cited in Holt *et al.*, 1995) concluded that they were all sensitive to oil/dispersant mixtures, with little differences between adults, sporelings, diploid or haploid life stages.

#### Synthetic compound contamination

Not Assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

This pressure is **Not assessed** but evidence is presented where available.

O'Brian & Dixon (1976) suggested that red algae were the most sensitive group of macrophytes to oil and dispersant contamination (see Smith, 1968). *Saccharina latissima* has also been found to be sensitive to antifouling compounds. Johansson (2009) exposed samples of *Saccharina latissima* to several antifouling compounds, observing chlorothalonil, DCOIT, dichlofluanid and tolylfluanid inhibited photosynthesis. Exposure to Chlorothalonil and tolylfluanid, was also found to continue inhibiting oxygen evolution after exposure had finished, and may cause irreversible damage.

Smith (1968) observed that epiphytic and benthic red algae were intolerant of dispersant or oil contamination during the *Torrey Canyon* oil spill; only the epiphytes *Cryptopleura ramosa* and *Spermothamnion repens* and some tufts of *Jania rubens* survived together with *Osmundea pinnatifida*, *Gigartina pistillata* and *Phyllophora crispa* from the sublittoral fringe. *Delesseria sanguinea* was probably the most intolerant since it was damaged at depths of 6m (Smith, 1968). Holt *et al.* (1995) suggested that *Delesseria sanguinea* is probably generally sensitive of chemical contamination.

#### Radionuclide contamination

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

No evidence (NEv)

Q: NR A: NR C: NR

No evidence

**Introduction of other substances**

Not Assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

This pressure is **Not assessed**.

**De-oxygenation**

High

Q: Medium A: High C: High

High

Q: Medium A: High C: High

Not sensitive

Q: Medium A: High C: High

Reduced oxygen concentrations can inhibit both photosynthesis and respiration in macroalgae (Kinne, 1977). Despite this, macroalgae are thought to buffer the environmental conditions of low oxygen, thereby acting as a refuge for organisms in oxygen-depleted regions especially if the oxygen depletion is short-term (Frieder *et al.*, 2012). If levels do drop below 4 mg/l negative effects on these organisms can be expected with adverse effects occurring below 2mg/l (Cole *et al.*, 1999). However, in wave exposed, highly mixed, areas, the hypoxic conditions are likely to be transient.

**Sensitivity Assessment.** Reduced oxygen levels are likely to inhibit photosynthesis and respiration but not cause a loss of the macroalgae population directly. Resistance has been assessed as 'High', Resilience as 'High'. Sensitivity has been assessed as 'Not sensitive' at the benchmark level.

**Nutrient enrichment**

Medium

Q: Medium A: High C: High

High

Q: High A: Medium C: High

Low

Q: Medium A: Medium C: High

Conolly & Drew (1985) found that *Saccharina latissima* sporophytes plants at the most eutrophic site in a study on the east coast of Scotland where nutrient levels were 25% higher than ambient levels exhibited a high growth rate. However, Read *et al.* (1983) reported after removal of a major sewage pollution in the Firth of Forth, *Saccharina latissima* became abundant where previously it had been absent. Bokn *et al.* (2003) conducted a nutrient loading experiment on intertidal fucoids. Within 3 years of the experiment no significant effect was observed in the communities, however, 4-5 years into the experiment a shift occurred from perennials to ephemeral algae occurred. Although Bokn *et al.* (2003) focussed on fucoids the results could indicate that long-term (>4 years) nutrient loading can result in community shift to ephemeral algae species, and explain the disparity between the findings of Conolly & Drew (1985) & Read *et al.* (1983).

Johnston & Roberts (2009) conducted a meta-analysis, which reviewed 216 papers to assess how a variety of contaminants (including sewage and nutrient loading) affected 6 marine habitats (including subtidal reefs). A 30-50% reduction in species diversity and richness was identified from all habitats exposed to the contaminant types. Johnston & Roberts (2009) however also highlighted that macro-algal communities are relatively tolerant to contamination, but that contaminated communities can have low diversity assemblages which are dominated by opportunistic and fast growing species (Johnston & Roberts, 2009 and references therein). Nutrient enrichment may also result in phytoplankton blooms that increase turbidity and, therefore, may negatively impact photosynthesis.

**Sensitivity assessment.** However, the biotope is assessed as 'Not sensitive' at the pressure benchmark that assumes compliance with good status as defined by the WFD.

**Organic enrichment**

Medium

Q: Medium A: High C: High

High

Q: High A: Medium C: High

Low

Q: Medium A: Medium C: High

Conolly & Drew (1985) found *Saccharina latissima* sporophytes had relatively higher growth rates close to a sewage outlet in St Andrews, UK when compared to other sites along the east coast of Scotland. At St Andrews, nitrate levels were 20.22µM, which represents an approx 25% increase when compared to other comparable sites (approx 15.87 µM). Handå *et al.* (2013) also reported *Saccharina latissima* sporophytes grew approx 1% faster per day when in close proximity to Norwegian Salmon farms, where elevated ammonium can be readily absorbed. Read *et al.* (1983) reported after the installation of a new sewage treatment works, which reduced the suspended solid content of liquid effluent by 60% in the Firth of Forth, *Saccharina latissima* became abundant where previously it had been absent. Bokn *et al.* (2003) conducted a nutrient loading experiment on intertidal fucoids. Within 3 years of the experiment no significant effect was observed in the communities, however, 4-5 years into the experiment a shift occurred from perennials to ephemeral algae occurred. Although Bokn *et al.* (2003) focussed on fucoids the results could indicate that long-term (>4 years) nutrient loading can result in community shift to ephemeral algae species. Differences between the findings of the aforementioned studies are likely to be related to the level of organic enrichment, however, could also be time dependent.

Johnston & Roberts (2009) conducted a meta-analysis, which reviewed 216 papers to assess how a variety of contaminants (including sewage and nutrient loading) affected 6 marine habitats (including subtidal reefs). A 30-50% reduction in species diversity and richness was identified from all habitats exposed to the contaminant types. Johnston & Roberts (2009) however also highlighted that macroalgal communities are relatively tolerant to contamination, but that contaminated communities can have low diversity assemblages which are dominated by opportunistic and fast growing species (Johnston & Roberts, 2009 and references therein). Organic enrichment may also result in phytoplankton blooms that increase turbidity and, therefore, may negatively impact photosynthesis.

**Sensitivity assessment.** Although short-term exposure (<4 years) to organic enrichment may not affect seaweeds directly, indirect effects such as turbidity may significantly affect photosynthesis. Therefore, resistance has been assessed as '**Medium**', resilience as '**High**' and sensitivity as '**Low**'.

## A Physical Pressures

	Resistance	Resilience	Sensitivity
Physical loss (to land or freshwater habitat)	<b>None</b> Q: High A: High C: High	<b>Very Low</b> Q: High A: High C: High	<b>High</b> Q: High A: High C: High

All marine habitats and benthic species are considered to have a resistance of 'None' to this pressure and to be unable to recover from a permanent loss of habitat (resilience is '**Very Low**'). Sensitivity within the direct spatial footprint of this pressure is therefore '**High**'. Although no specific evidence is described confidence in this assessment is '**High**', due to the incontrovertible nature of this pressure.

Physical change (to another seabed type)	<b>None</b> Q: High A: High C: High	<b>Very Low</b> Q: High A: High C: High	<b>High</b> Q: High A: High C: High
--	--	--	--

If rock substrata were replaced with sedimentary substrata this would represent a fundamental change in habitat type, which *Saccharina latissima*, *Saccorhiza polyschides* and *Desmarestia spp.* would not be able to tolerate. The biotope would be lost.



**Sensitivity assessment.** Resistance to the pressure is considered '**None**', and resilience '**Very Low**'. The sensitivity of this biotope to change from sedimentary or soft rock substrata to hard rock or artificial substrata or vice-versa is assessed as '**High**'.

<b>Physical change (to another sediment type)</b>	Not relevant (NR) Q: NR A: NR C: NR	Not relevant (NR) Q: NR A: NR C: NR	Not relevant (NR) Q: NR A: NR C: NR
---	--	--	--

Not relevant

<b>Habitat structure changes - removal of substratum (extraction)</b>	Not relevant (NR) Q: NR A: NR C: NR	Not relevant (NR) Q: NR A: NR C: NR	Not relevant (NR) Q: NR A: NR C: NR
---	--	--	--

Not relevant

<b>Abrasion/disturbance of the surface of the substratum or seabed</b>	<b>None</b> Q: Low A: NR C: NR	<b>High</b> Q: High A: High C: High	<b>Medium</b> Q: Low A: NR C: NR
--	-----------------------------------	--	-------------------------------------

Abrasion of the substratum e.g. from bottom or pot fishing gear, cable laying etc. may cause localised mobility of the substrata and mortality of the resident community. The effect would be situation dependent however if bottom fishing gear were towed over a site it may mobilise a high proportion of the rock substrata and cause high mortality in the resident community. However, the characteristic species within IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFiIR, and IR.HIR.KSed.XKScrR have rapid growth rates and are distinctive of 'disturbed' areas. Information on from experimental clearances is summarised under resilience above.

**Sensitivity assessment.** Resistance has been assessed as '**None**', Resilience as '**High**'. Sensitivity has been assessed as '**Medium**'.

<b>Penetration or disturbance of the substratum subsurface</b>	Not relevant (NR) Q: NR A: NR C: NR	Not relevant (NR) Q: NR A: NR C: NR	Not relevant (NR) Q: NR A: NR C: NR
--	--	--	--

Not relevant, please refer to pressure "Abrasion/disturbance of the substratum on the surface of the seabed".

<b>Changes in suspended solids (water clarity)</b>	<b>Low</b> Q: Medium A: Medium C: Medium	<b>High</b> Q: High A: High C: High	<b>Low</b> Q: Medium A: Medium C: Medium
--	---	--	---

Suspended Particle Matter (SPM) concentration has a linear relationship with subsurface light attenuation ( $K_d$ ) (Devlin *et al.*, 1998). Light penetration influences the maximum depth at which kelp species can grow and it has been reported that laminarians grow at depths at which the light levels are reduced to 1 percent of incident light at the surface. Maximal depth distribution of laminarians, therefore, varies from 100 m in the Mediterranean to only 6-7 m in the silt-laden German Bight. In Atlantic European waters, the depth limit is typically 35 m. In very turbid waters the depth at which kelp is found may be reduced, or in some cases excluded completely (e.g. Severn

Estuary), because of the alteration in light attenuation by suspended sediment (Lüning, 1990; Birkett *et al.*, 1998b). Limited information is available on which to assess the effect of a decrease in water clarity on Laminariales show a decrease of 50% photosynthetic activity when turbidity increases by 0.1/m (light attenuation coefficient =0.1-0.2/m; Staehr & Wernberg, 2009). *Desmarestia* spp. recruitment has also been found highly affected by light attenuation and frequency, typically rare (ca <10% coverage (Edwards, 1998) beneath kelp canopies where light levels can be 1-5% of surface irradiance (Kitching, 1941).

A decrease in water clarity as a result of mobilised sediments may also increase sediment scour of biotopes within close proximity. However, the characterizing species within IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, IR.HIR.KSed.XKScrR have rapid growth and colonisation rates are as such relatively resilient to sediment scouring.

**Sensitivity assessment.** An increase in water clarity from clear to intermediate (10-100 mg/l) represent a change in light attenuation of ca 0.67-6.7 Kd/m. and is likely to result in a greater than 50% reduction in photosynthesis of *Laminaria* spp. Therefore, an increase in SPM from intermediate to moderate turbidity is likely to significantly reduce the depth at which laminarians can grow. Resistance to this pressure is defined as 'Low' as the biotope is typical of sediment affected habitats. Resilience to this pressure is defined as 'High' and this biotope is regarded as having a sensitivity of 'Low'.

#### Smothering and siltation rate changes (light)

High

Q: Low A: NR C: NR

High

Q: High A: High C: High

Not sensitive

Q: Low A: Low C: Low

Smothering by sediment e.g. 5 cm material during a discrete event, is unlikely to damage *Saccharina latissima*, *Saccorhiza polyschides* and *Desmarestia* spp. sporophytes but may provide a physical barrier to zoospore settlement and therefore negatively impact on recruitment processes (Moy & Christie, 2012). Laboratory studies showed that kelp and gametophytes can survive in darkness for between 6-16 months at 8°C and would probably survive smothering by a discrete event and once returned to normal conditions gametophytes resumed growth or maturation within 1 month (Dieck, 1993). *Saccorhiza polyschides* zoospores successfully developed after 180 days of darkness (Norton, 1977). Intolerance to this factor is likely to be higher during the peak periods of sporulation and/or spore settlement.

IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, IR.HIR.KSed.XKScrR are found from extreme wave exposed-sheltered sites (Connor *et al.*, 2004). In wave exposed biotopes deposited sediment is unlikely to remain for more than a few tidal cycles (due to water flow or wave action). In sheltered biotopes deposited sediment could remain and the effects of deposition could be longer lasting.

**Sensitivity assessment.** Resistance has been assessed as 'High', resilience as 'High' and sensitivity as 'Not Sensitive'.

#### Smothering and siltation rate changes (heavy)

Medium

Q: Low A: NR C: NR

High

Q: High A: High C: High

Low

Q: Low A: Low C: Low

Smothering by sediment e.g. 30 cm material during a discrete event, is unlikely to damage *Saccharina latissima*, *Saccorhiza polyschides* and *Desmarestia* spp. sporophytes but may provide a physical barrier to zoospore settlement and therefore negatively impact on recruitment processes

(Moy & Christie, 2012). The volume of sediment may also inundate juvenile sporophytes. Given the short life expectancy of *Saccharina latissima* (2-4 years; Parke, 1948), self sustaining populations are likely to be dependent on annual recruitment (Moy & Christie, 2012). Given the microscopic size of the gametophyte, 30 cm of sediment could be expected to significantly inhibit growth. Laboratory studies showed that kelp and gametophytes can survive in darkness for between 6-16 months at 8°C and would probably survive smothering by a discrete event and once returned to normal conditions gametophytes resumed growth or maturation within 1 month (Dieck, 1993). *Saccorhiza polyschides* zoospores, specifically, successfully developed after 180 days of darkness (Norton, 1977). Intolerance to this factor is likely to be higher during the peak periods of sporulation and/or spore settlement.

IR.HIR.KSed.XKScrR is subject to periodic burial from surrounding sediments, however IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, IR.HIR.KSed.XKScrR are recorded from extreme wave exposed-sheltered sites (Connor *et al.*, 2004) and therefore the effects of burial are likely to be mediated. In highly wave exposed biotopes deposited sediment is unlikely to remain for more than a few tidal cycles (due to water flow or wave action). In sheltered biotopes the high volume of deposited sediment could remain and the effects could be longer lasting. However, these biotopes are periodically disturbed by winter storms or sediment scour therefore deposited sediments are unlikely to remain for a full season.

Sensitivity assessment. Resistance has been assessed as '**Medium**', resilience as '**High**'. Sensitivity has been assessed as '**Low**'.

## Litter

Not Assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

Not assessed.

## Electromagnetic changes

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

No evidence (NEv)

Q: NR A: NR C: NR

No evidence

## Underwater noise changes

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant

## Introduction of light or shading

Low

Q: Low A: NR C: NR

Medium

Q: Low A: NR C: NR

Medium

Q: Low A: NR C: NR

There is no evidence to suggest that anthropogenic light sources would affect *Saccharina latissima*, *Saccorhiza polyschides* and *Desmarestia spp.* Shading of the biotope (e.g. by the construction of a pontoon, pier etc) could adversely affect the biotope in areas where the water clarity is also low, and tip the balance to shade tolerant species, resulting in the loss of the biotope directly within the shaded area, or a reduction in seaweed abundance.

**Sensitivity assessment.** Resistance is probably '**Low**', with a '**Medium**' resilience and a sensitivity

of 'Medium', albeit with 'low' confidence due to the lack of direct evidence.

<b>Barrier to species movement</b>	Not relevant (NR)	Not relevant (NR)	Not relevant (NR)
	Q: NR A: NR C: NR	Q: NR A: NR C: NR	Q: NR A: NR C: NR

**Not relevant.** This pressure is considered applicable to mobile species, e.g. fish and marine mammals rather than seabed habitats. Physical and hydrographic barriers may limit the dispersal of spores. But spore dispersal is not considered under the pressure definition and benchmark.

<b>Death or injury by collision</b>	Not relevant (NR)	Not relevant (NR)	Not relevant (NR)
	Q: NR A: NR C: NR	Q: NR A: NR C: NR	Q: NR A: NR C: NR

**Not relevant.** Collision from grounding vessels is addressed under abrasion above.

<b>Visual disturbance</b>	Not relevant (NR)	Not relevant (NR)	Not relevant (NR)
	Q: NR A: NR C: NR	Q: NR A: NR C: NR	Q: NR A: NR C: NR

Not relevant

## Biological Pressures

	Resistance	Resilience	Sensitivity
<b>Genetic modification &amp; translocation of indigenous species</b>	No evidence (NEv)	Not relevant (NR)	No evidence (NEv)
	Q: NR A: NR C: NR	Q: NR A: NR C: NR	Q: NR A: NR C: NR

**No evidence** regarding the genetic modification or effects of translocation of native kelp populations was found.

<b>Introduction or spread of invasive non-indigenous species</b>	Low	Very Low	High
	Q: High A: High C: High	Q: High A: High C: High	Q: High A: High C: High

*Undaria pinnatifida* has received a large amount of research attention as a major Invasive Non-Indigenous Species (INIS) which could out-compete native UK kelp habitats (see Farrell & Fletcher, 2006; Thompson & Schiel, 2012, Brodie *et al.*, 2014; Hieser *et al.*, 2014). *Undaria pinnatifida* was first recorded in the UK, Hamble Estuary, in June 1994 (Fletcher & Manfredi, 1995) and has since spread to a number of British ports. *Undaria pinnatifida* is an annual species, sporophytes appear in Autumn and grow rapidly throughout winter and spring during which they can reach a length of 1.65m (Birket *et al.*, 1998). Farrell & Fletcher (2006) suggested that native short lived species that occupy similar ecological niches to *Undaria pinnatifida*, such as *Saccharina latissima*, *Saccorhiza polyschides* or *Desmarestia spp.*, are likely to be worst affected and out-competed by *Undaria pinnatifida*. Where present, an abundance of *Undaria pinnatifida* has corresponded to a decline in *Saccharina latissima* (Farrel & Fletcher, 2006) and *Laminaria hyperborea* (Hieser *et al.*, 2014). Furthermore, disturbance has been found a significant pathway for the establishment of *Undaria pinnatifida* (Valentine & Johnson, 2003). Therefore, periodic disturbance within this biotope could facilitate the establishment of *Undaria pinnatifida*.

In new Zealand, Thompson & Schiel (2012) observed that native fucoids could out-compete *U.pinnatifida* and re-dominate the substratum. However, Thompson & Schiel (2012) suggested the fucoid recovery was partially due to an annual *Undaria pinnatifida* die back, which as noted by Heiser *et al.*, (2014) does not occur in Plymouth sound, UK. *Undaria pinnatifida* was successfully eradicated on a sunken ship in Clatham Islands, New Zealand, by applying a heat treatment of 70°C (Wotton *et al.*, 2004) however numerous other eradication attempts have failed, and as noted by Fletcher & Farrell, (1999) once established *Undaria pinnatifida* resists most attempts at long-term removal. The biotope is unlikely to fully recover until *Undaria pinnatifida* is fully removed from the habitat, which as stated above is unlikely to occur.

**Sensitivity assessment.** Resistance to the pressure is considered ‘**Low**’, and resilience ‘**Very low**’. The sensitivity of this biotope to the introduction of microbial pathogens is assessed as ‘**High**’.

<b>Introduction of microbial pathogens</b>	<b>No evidence (NEv)</b>	<b>No evidence (NEv)</b>	<b>No evidence (NEv)</b>
	Q: NR A: NR C: NR	Q: NR A: NR C: NR	Q: NR A: NR C: NR

Laminarians may be infected by the microscopic brown alga *Streblonema acidioides*. Infected algae show symptoms of Streblonema disease, i.e. alterations of the blade and stipe ranging from dark spots to heavy deformations and completely crippled thalli. Infection can reduce growth rates of host algae (Peters & Scaffelke, 1996). The marine fungi *Eurychasma spp* can also infect early life stages of laminarians and *Desmarestia viridis*, however, the effects of infection are unknown (Müller *et al.*, 1999).

**Sensitivity assessment.** Due to a lack of conclusive evidence the sensitivity of this biotope cannot be assessed against this pressure.

<b>Removal of target species</b>	<b>None</b>	<b>High</b>	<b>Medium</b>
	Q: Low A: NR C: NR	Q: High A: High C: High	Q: Low A: NR C: NR

There has been recent commercial interest in *Saccharina latissima* as a consumable called “sea vegetables” (Birket *et al.*, 1998). However, *Saccharina latissima* sporophytes are typically matured on ropes (Handå *et al.* 2013) and not directly extracted from the seabed, as with *Laminaria hyperborea* (Christie *et al.*, 1998). No evidence has been found for commercial extraction of *Saccorhiza polyschides* or *Desmarestia spp*. However, if the biotopes were subject to harvest, then a large proportion of the resident kelp population could be removed. . Thus evidence to assess the resistance of IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, IR.HIR.KSed.XKScrR to direct harvesting is limited. It has been assumed that if targeted harvesting were in operation it would remove >75% of sporophytes.

**Sensitivity assessment.** Resistance has been assessed as ‘**None**’, Resilience as ‘**High**’. Sensitivity has been assessed as ‘**Medium**’.

<b>Removal of non-target species</b>	<b>None</b>	<b>High</b>	<b>Medium</b>
	Q: Low A: NR C: NR	Q: High A: High C: High	Q: Low A: NR C: NR

IR.HIR.KSed.Sac, IR.HIR.KSed.SlatSac, IR.HIR.KSed.DesFilR, IR.HIR.KSed.XKScrR Are characterised by a canopy of *Saccorhiza polyschides*, *Saccharina latissima*, and *Desmarestia spp*. If the canopy were removed the red seaweeds understory community may become bleached, and/or

perish (Hawkins & Harkin, 1985), leading to further reductions in biodiversity. The biotope is however naturally periodically disturbed and as such would recover rapidly.

**Sensitivity assessment.** Resistance has been assessed as '**None**', Resilience as '**High**'. Sensitivity has been assessed as '**Medium**'.



## Bibliography

- Baardseth, E., 1956. The growth rings in *Alaria stipes*. In *Proceedings of the International Seaweed Symposium, 2, Trondheim* (eds T. Braarud & N.A. Sorensen) pp. 153-157. London: Pergamon Press.
- Bégin, C., Johnson, L.E. & Himmelman, J.H., 2004. Macroalgal canopies: distribution and diversity of associated invertebrates and effects on the recruitment and growth of mussels. *Marine Ecology Progress Series*, **271** (1), 121-132.
- Bekkby, T. & Moy, F.E., 2011. Developing spatial models of sugar kelp (*Saccharina latissima*) potential distribution under natural conditions and areas of its disappearance in Skagerrak. *Estuarine Coastal and Shelf Science*, **95** (4), 477-483.
- Beszczynska-Möller, A., & Dye, S.R., 2013. ICES Report on Ocean Climate 2012. In *ICES Cooperative Research Report*, vol. 321 pp. 73.
- Birkett, D.A., Maggs, C.A., Dring, M.J. & Boaden, P.J.S., 1998b. Infralittoral reef biotopes with kelp species: an overview of dynamic and sensitivity characteristics for conservation management of marine SACs. *Natura 2000 report prepared by Scottish Association of Marine Science (SAMS) for the UK Marine SACs Project.*, Scottish Association for Marine Science. (UK Marine SACs Project, vol V.). Available from: <http://www.ukmarinesac.org.uk/publications.htm>
- Bishop, G.M., 1985. *Aspects of the reproductive ecology of the sea urchin Echinus esculentus L.* Ph.D. thesis, University of Exeter, UK.
- Booolootian, R.A., 1966. *Physiology of Echinodermata.* (Ed. R.A. Booolootian), pp. 822-822. New York: John Wiley & Sons.
- Bower, S.M., 1996. *Synopsis of Infectious Diseases and Parasites of Commercially Exploited Shellfish: Bald-sea-urchin Disease.* [On-line]. Fisheries and Oceans Canada. [cited 26/01/16]. Available from: <http://www.dfo-mpo.gc.ca/science/aah-saa/diseases-maladies/bsudsu-eng.html>
- Bradshaw, C., Veale, L.O., Hill, A.S. & Brand, A.R., 2000. The effects of scallop dredging on gravelly seabed communities. In: *Effects of fishing on non-target species and habitats* (ed. M.J. Kaiser & de S.J. Groot), pp. 83-104. Oxford: Blackwell Science.
- Brodie J., Williamson, C.J., Smale, D.A., Kamenos, N.A., Mieszkowska, N., Santos, R., Cunliffe, M., Steinke, M., Yesson, C. & Anderson, K.M., 2014. The future of the northeast Atlantic benthic flora in a high CO<sub>2</sub> world. *Ecology and Evolution*, **4** (13), 2787-2798.
- Bryan, G.W., 1984. Pollution due to heavy metals and their compounds. In *Marine Ecology: A Comprehensive, Integrated Treatise on Life in the Oceans and Coastal Waters*, vol. 5. *Ocean Management*, part 3, (ed. O. Kinne), pp.1289-1431. New York: John Wiley & Sons.
- Burrows, M.T., Smale, D., O'Connor, N., Rein, H.V. & Moore, P., 2014. Marine Strategy Framework Directive Indicators for UK Kelp Habitats Part 1: Developing proposals for potential indicators. *Joint Nature Conservation Committee*, Peterborough. Report no. 525.
- Casas, G., Scrosati, R. & Piriz, M.L., 2004. The invasive kelp *Undaria pinnatifida* (Phaeophyceae, Laminariales) reduces native seaweed diversity in Nuevo Gulf (Patagonia, Argentina). *Biological Invasions*, **6** (4), 411-416.
- Christie, H., Fredriksen, S. & Rinde, E., 1998. Regrowth of kelp and colonization of epiphyte and fauna community after kelp trawling at the coast of Norway. *Hydrobiologia*, **375/376**, 49-58.
- Cole, S., Codling, I.D., Parr, W., Zabel, T., 1999. Guidelines for managing water quality impacts within UK European marine sites [On-line]. *UK Marine SACs Project.* [Cited 26/01/16]. Available from: [http://www.ukmarinesac.org.uk/pdfs/water\\_quality.pdf](http://www.ukmarinesac.org.uk/pdfs/water_quality.pdf)
- Connor, D.W., Allen, J.H., Golding, N., Howell, K.L., Lieberknecht, L.M., Northen, K.O. & Reker, J.B., 2004. The Marine Habitat Classification for Britain and Ireland. Version 04.05. ISBN 1 861 07561 8. In *JNCC (2015), The Marine Habitat Classification for Britain and Ireland Version 15.03.* [2019-07-24]. Joint Nature Conservation Committee, Peterborough. Available from <https://mhc.jncc.gov.uk/>
- Connor, D.W., Dalkin, M.J., Hill, T.O., Holt, R.H.F. & Sanderson, W.G., 1997a. Marine biotope classification for Britain and Ireland. Vol. 2. Sublittoral biotopes. *Joint Nature Conservation Committee, Peterborough, JNCC Report no. 230, Version 97.06.*, *Joint Nature Conservation Committee, Peterborough, JNCC Report no. 230, Version 97.06.*
- Conolly N.J. & Drew, E.A., 1985. Physiology of *Laminaria*. III. Effect of a coastal eutrophication on seasonal patterns of growth and tissue composition in *Laminaria digitata* and *L. saccharina*. *Marine Ecology, Pubblicazioni della Stazione Zoologica di Napoli I*, **6**, 181-195.
- Cosse, A., Potin, P. & Leblanc, C., 2009. Patterns of gene expression induced by oligoguluronates reveal conserved and environment-specific molecular defence responses in the brown alga *Laminaria digitata*. *New Phytologist*, **182** (1), 239-250.
- Crump, R.G., Morley, H.S., & Williams, A.D., 1999. West Angle Bay, a case study. Littoral monitoring of permanent quadrats before and after the *Sea Empress* oil spill. *Field Studies*, **9**, 497-511.
- Dauvin, J.C., Bellan, G., Bellan-Santini, D., Castric, A., Francour, P., Gentil, F., Girard, A., Gofas, S., Mahe, C., Noel, P., & Reviers, B. de., 1994. Typologie des ZNIEFF-Mer. Liste des parametres et des biocoenoses des cotes francaises metropolitaines. 2nd ed. *Secretariat Faune-Flore, Museum National d'Histoire Naturelle, Paris (Collection Patrimoines Naturels, Serie Patrimoine Ecologique, No. 12).* Coll. Patrimoines Naturels, vol. 12, Secretariat Faune-Flore, Paris.
- Davies, C.E. & Moss, D., 1998. European Union Nature Information System (EUNIS) Habitat Classification. *Report to European Topic Centre on Nature Conservation from the Institute of Terrestrial Ecology, Monks Wood, Cambridgeshire.* [Final draft with further revisions to marine habitats.], Brussels: European Environment Agency.
- Dayton, P.K., Tegner, M.J., Parnell, P.E. & Edwards, P.B., 1992. Temporal and spatial patterns of disturbance and recovery in a kelp

forest community. *Ecological Monographs*, **62**, 421-445.

Devlin, M.J., Barry, J., Mills, D.K., Gowen, R.J., Foden, J., Sivyer, D. & Tett, P., 2008. Relationships between suspended particulate material, light attenuation and Secchi depth in UK marine waters. *Estuarine, Coastal and Shelf Science*, **79** (3), 429-439.

Dieck, T.I., 1992. North Pacific and North Atlantic digitate *Laminaria* species (Phaeophyta): hybridization experiments and temperature responses. *Phycologia*, **31**, 147-163.

Dieck, T.I., 1993. Temperature tolerance and survival in darkness of kelp gametophytes (Laminariales: Phaeophyta) - ecological and biogeographical implications. *Marine Ecology Progress Series*, **100**, 253-264.

Dinesen G., 1999. *Modiolus modiolus* and the associated fauna. In Bruntse, G., Lein, T.E., Nielsen, R. (eds), *Marine benthic algae and invertebrate communities from the shallow waters of the Faroe Islands: a base line study*. Kaldbak Marine Biological Laboratory, pp. 66-71.

Dinnel, P.A., Pagano, G.G., & Oshido, P.S., 1988. A sea urchin test system for marine environmental monitoring. In *Echinoderm Biology. Proceedings of the Sixth International Echinoderm Conference, Victoria, 23-28 August 1987*, (R.D. Burke, P.V. Mladenov, P. Lambert, Parsley, R.L. ed.), pp 611-619. Rotterdam: A.A. Balkema.

Drew, E.A., Forster, G.R., Gage, J., Harwood, G., Larkum, A.W.D., Lythgoe, J.M & Potts, G.W. 1967. "Torrey Canyon" report. Underwater Association Report for 1966-67, pp. 53-60.

Eckman, J.E., 1983. Hydrodynamic processes affecting benthic recruitment 1. *Limnology and Oceanography*, **28** (2), 241-257.

Edwards, A., 1980. Ecological studies of the kelp *Laminaria hyperborea* and its associated fauna in south-west Ireland. *Ophelia*, **9**, 47-60.

Elnor, R.W. & Vadas, R.L., 1990. Inference in ecology: the sea urchin phenomenon in the northwest Atlantic. *American Naturalist*, **136**, 108-125.

Elsäßer, B., Fariñas-Franco, J.M., Wilson, C.D., Kregting, L. & Roberts, D., 2013. Identifying optimal sites for natural recovery and restoration of impacted biogenic habitats in a special area of conservation using hydrodynamic and habitat suitability modelling. *Journal of Sea Research*, **77**, 11-21.

Erwin, D.G., Picton, B.E., Connor, D.W., Howson, C.M., Gilleece, P. & Bogues, M.J., 1990. Inshore Marine Life of Northern Ireland. *Report of a survey carried out by the diving team of the Botany and Zoology Department of the Ulster Museum in fulfilment of a contract with Conservation Branch of the Department of the Environment (N.I.)*, Ulster Museum, Belfast: HMSO.

Farrell, P. & Fletcher, R., 2006. An investigation of dispersal of the introduced brown alga *Undaria pinnatifida* (Harvey) Suringar and its competition with some species on the man-made structures of Torquay Marina (Devon, UK). *Journal of Experimental Marine Biology and Ecology*, **334** (2), 236-243.

Fletcher, R.L., 1996. The occurrence of 'green tides' - a review. In *Marine Benthic Vegetation. Recent changes and the Effects of Eutrophication* (ed. W. Schramm & P.H. Nienhuis). Berlin Heidelberg: Springer-Verlag. [Ecological Studies, vol. 123].

Fredriksen, S., Sjøtun, K., Lein, T.E. & Rueness, J., 1995. Spore dispersal in *Laminaria hyperborea* (Laminariales, Phaeophyceae). *Sarsia*, **80** (1), 47-53.

Frieder, C., Nam, S., Martz, T. & Levin, L., 2012. High temporal and spatial variability of dissolved oxygen and pH in a nearshore California kelp forest. *Biogeosciences*, **9** (10), 3917-3930.

Gainey, L.F., 1994. Volume regulation in three species of marine mussels. *Journal of Experimental Marine Biology and Ecology*, **181** (2), 201-211.

Gomez, J.L.C. & Miguez-Rodriguez, L.J., 1999. Effects of oil pollution on skeleton and tissues of *Echinus esculentus* L. 1758 (Echinodermata, Echinoidea) in a population of A Coruna Bay, Galicia, Spain. In *Echinoderm Research 1998. Proceedings of the Fifth European Conference on Echinoderms, Milan, 7-12 September 1998*, (ed. M.D.C. Carnevali & F. Bonasoro) pp. 439-447. Rotterdam: A.A. Balkema.

Gorman, D., Bajjouk, T., Populus, J., Vasquez, M. & Ehrhold, A., 2013. Modeling kelp forest distribution and biomass along temperate rocky coastlines. *Marine Biology*, **160** (2), 309-325.

Grandy, N., 1984. *The effects of oil and dispersants on subtidal red algae*. Ph.D. Thesis. University of Liverpool.

Griffiths, A.B., Dennis, R. & Potts, G.W., 1979. Mortality associated with a phytoplankton bloom off Penzance in Mount's Bay. *Journal of the Marine Biological Association of the United Kingdom*, **59**, 515-528.

Hammer, L., 1972. Anaerobiosis in marine algae and marine phanerogams. In *Proceedings of the Seventh International Seaweed Symposium, Sapporo, Japan, August 8-12, 1971* (ed. K. Nisizawa, S. Arasaki, Chihara, M., Hirose, H., Nakamura V., Tsuchiya, Y.), pp. 414-419. Tokyo: Tokyo University Press.

Harkin, E., 1981. Fluctuations in epiphyte biomass following *Laminaria hyperborea* canopy removal. In *Proceedings of the X<sup>th</sup> International Seaweed Symposium, Gøteborg, 11-15 August 1980* (ed. T. Levring), pp.303-308. Berlin: Walter de Gruyter.

Hawkins, S.J. & Harkin, E., 1985. Preliminary canopy removal experiments in algal dominated communities low on the shore and in the shallow subtidal on the Isle of Man. *Botanica Marina*, **28**, 223-30.

Hawkins, S.J. & Hartnoll, R.G., 1985. Factors determining the upper limits of intertidal canopy-forming algae. *Marine Ecology Progress Series*, **20**, 265-271.

Hayward, P.J. 1988. *Animals on seaweed*. Richmond, Surrey: Richmond Publishing Co. Ltd. [Naturalists Handbooks 9].

Heiser, S., Hall-Spencer, J.M. & Hiscock, K., 2014. Assessing the extent of establishment of *Undaria pinnatifida* in Plymouth Sound



- Special Area of Conservation, UK. *Marine Biodiversity Records*, **7**, e93.
- Hiscock, K. & Mitchell, R., 1980. *The Description and Classification of Sublittoral Epibenthic Ecosystems*. In *The Shore Environment*, Vol. 2, *Ecosystems*, (ed. J.H. Price, D.E.G. Irvine, & W.F. Farnham), 323-370. London and New York: Academic Press. [Systematics Association Special Volume no. 17(b)].
- Hoare, R. & Hiscock, K., 1974. An ecological survey of the rocky coast adjacent to the effluent of a bromine extraction plant. *Estuarine and Coastal Marine Science*, **2** (4), 329-348.
- Holt, T.J., Jones, D.R., Hawkins, S.J. & Hartnoll, R.G., 1995. The sensitivity of marine communities to man induced change - a scoping report. *Countryside Council for Wales, Bangor, Contract Science Report*, no. 65.
- Hopkin, R. & Kain, J.M., 1978. The effects of some pollutants on the survival, growth and respiration of *Laminaria hyperborea*. *Estuarine and Coastal Marine Science*, **7**, 531-553.
- Irvine, L. M. & Chamberlain, Y. M., 1994. *Seaweeds of the British Isles*, vol. 1. *Rhodophyta*, Part 2B *Corallinales, Hildenbrandiales*. London: Her Majesty's Stationery Office.
- JNCC, 2015. The Marine Habitat Classification for Britain and Ireland Version 15.03. (20/05/2015). Available from <https://mhc.jncc.gov.uk/>
- JNCC, 2015. The Marine Habitat Classification for Britain and Ireland Version 15.03. (20/05/2015). Available from <https://mhc.jncc.gov.uk/>
- JNCC (Joint Nature Conservation Committee), 1999. *Marine Environment Resource Mapping And Information Database (MERMAID): Marine Nature Conservation Review Survey Database*. [on-line] <http://www.jncc.gov.uk/mermaid>
- Jones, C.G., Lawton, J.H. & Shackak, M., 1994. Organisms as ecosystem engineers. *Oikos*, **69**, 373-386.
- Jones, D.J., 1971. Ecological studies on macro-invertebrate communities associated with polluted kelp forest in the North Sea. *Helgolander Wissenschaftliche Meeresuntersuchungen*, **22**, 417-431.
- Jones, L.A., Hiscock, K. & Connor, D.W., 2000. Marine habitat reviews. A summary of ecological requirements and sensitivity characteristics for the conservation and management of marine SACs. *Joint Nature Conservation Committee, Peterborough. (UK Marine SACs Project report)*. Available from: <http://www.ukmarinesac.org.uk/pdfs/marine-habitats-review.pdf>
- Jones, N.S. & Kain, J.M., 1967. Subtidal algal recolonisation following removal of *Echinus*. *Helgolander Wissenschaftliche Meeresuntersuchungen*, **15**, 460-466.
- Kain, J.M., 1964. Aspects of the biology of *Laminaria hyperborea* III. Survival and growth of gametophytes. *Journal of the Marine Biological Association of the United Kingdom*, **44** (2), 415-433.
- Kain, J.M. & Svendsen, P., 1969. A note on the behaviour of *Patina pellucida* in Britain and Norway. *Sarsia*, **38**, 25-30.
- Kain, J.M., 1967. Populations of *Laminaria hyperborea* at various latitudes. *Helgolander Wissenschaftliche Meeresuntersuchungen*, **15**, 489-499.
- Kain, J.M., 1971a. Synopsis of biological data on *Laminaria hyperborea*. *FAO Fisheries Synopsis*, no. 87.
- Kain, J.M., 1975a. Algal recolonization of some cleared subtidal areas. *Journal of Ecology*, **63**, 739-765.
- Kain, J.M., 1979. A view of the genus *Laminaria*. *Oceanography and Marine Biology: an Annual Review*, **17**, 101-161.
- Kain, J.M., 1987. Photoperiod and temperature as triggers in the seasonality of *Delesseria sanguinea*. *Helgolander Meeresuntersuchungen*, **41**, 355-370.
- Kain, J.M., & Norton, T.A., 1990. Marine Ecology. In *Biology of the Red Algae*, (ed. K.M. Cole & Sheath, R.G.). Cambridge: Cambridge University Press.
- Kain, J.M., Drew, E.A. & Jupp, B.P., 1975. Light and the ecology of *Laminaria hyperborea* II. In *Proceedings of the Sixteenth Symposium of the British Ecological Society, 26-28 March 1974. Light as an Ecological Factor: II* (ed. G.C. Evans, R. Bainbridge & O. Rackham), pp. 63-92. Oxford: Blackwell Scientific Publications.
- Karsten, U., 2007. Research note: salinity tolerance of Arctic kelps from Spitsbergen. *Phycological Research*, **55** (4), 257-262.
- Kelly, M.S., 2000. The reproductive cycle of the sea urchin *Psammechinus miliaris* (Echinodermata: Echinoidea) in a Scottish sea loch. *Journal of the Marine Biological Association of the United Kingdom*, **80**, 909-919.
- Kinne, O. (ed.), 1984. *Marine Ecology: A Comprehensive, Integrated Treatise on Life in Oceans and Coastal Waters*. Vol. V. *Ocean Management Part 3: Pollution and Protection of the Seas - Radioactive Materials, Heavy Metals and Oil*. Chichester: John Wiley & Sons.
- Kinne, O., 1977. International Helgoland Symposium "Ecosystem research": summary, conclusions and closing. *Helgoländer Wissenschaftliche Meeresuntersuchungen*, **30**(1-4), 709-727.
- Kitching, J., 1941. Studies in sublittoral ecology III. *Laminaria* forest on the west coast of Scotland; a study of zonation in relation to wave action and illumination. *The Biological Bulletin*, **80** (3), 324-337
- Kregting, L., Blight, A., Elsässer, B. & Savidge, G., 2013. The influence of water motion on the growth rate of the kelp *Laminaria hyperborea*. *Journal of Experimental Marine Biology and Ecology*, **448**, 337-345.
- Kruuk, H., Wansink, D. & Moorhouse, A., 1990. Feeding patches and diving success of otters, *Lutra lutra*, in Shetland. *Oikos*, **57**, 68-72.
- Lang, C. & Mann, K., 1976. Changes in sea urchin populations after the destruction of kelp beds. *Marine Biology*, **36** (4), 321-326.
- Lein, T.E., Sjutun, K. & Wakili, S., 1991. Mass - occurrence of a brown filamentous endophyte in the lamina of the kelp *Laminaria*

- hyperborea* (Gunnerus) Foslie along the south western coast of Norway *Sarsia*, **76**, 187-193.
- Leinaas, H.P. & Christie, H., 1996. Effects of removing sea urchins (*Strongylocentrotus droebachiensis*): stability of the barren state and succession of kelp forest recovery in the east Atlantic. *Oecologia*, **105**(4), 524-536.
- Lobban, C.S. & Harrison, P.J., 1997. *Seaweed ecology and physiology*. Cambridge: Cambridge University Press.
- Lüning, K., 1990. *Seaweeds: their environment, biogeography, and ecophysiology*: John Wiley & Sons.
- Lyngby, J.E. & Mortensen, S.M., 1996. Effects of dredging activities on growth of *Laminaria saccharina*. *Marine Ecology, Pubblicazioni della Stazione Zoologica di Napoli I*, **17**, 345-354.
- Mann, K.H., 1982. Kelp, sea urchins, and predators: a review of strong interactions in rocky subtidal systems of eastern Canada, 1970-1980. *Netherlands Journal of Sea Research*, **16**, 414-423.
- MarLIN, 2015. MarLIN (Marine Life Network). (13-10-2015). <http://www.marlin.ac.uk>
- McKenzie, J.D. & Moore, P.G., 1981. The microdistribution of animals associated with the bulbous holdfasts of *Saccorhiza polyschides* (Phaeophyta). *Ophelia*, **20**, 201-213.
- Miller III, H.L., Neale, P.J. & Dunton, K.H., 2009. Biological weighting functions for UV inhibition of photosynthesis in the kelp *Laminaria hyperborea* (Phaeophyceae) 1. *Journal of Phycology*, **45** (3), 571-584.
- Moore, P.G., 1973a. The kelp fauna of north east Britain I. Function of the physical environment. *Journal of Experimental Marine Biology and Ecology*, **13**, 97-125.
- Moore, P.G., 1973b. The kelp fauna of north east Britain. II. Multivariate classification: turbidity as an ecological factor. *Journal of Experimental Marine Biology and Ecology*, **13**, 127-163.
- Moore, P.G., 1978. Turbidity and kelp holdfast Amphipoda. I. Wales and S.W. England. *Journal of Experimental Marine Biology and Ecology*, **32**, 53-96.
- Moore, P.G., 1985. Levels of heterogeneity and the amphipod fauna of kelp holdfasts. In *The Ecology of Rocky Coasts: essays presented to J.R. Lewis, D.Sc.* (ed. P.G. Moore & R. Seed), 274-289. London: Hodder & Stoughton Ltd.
- NBN, 2015. National Biodiversity Network 2015(20/05/2015). <https://data.nbn.org.uk/>
- Nichols, D., 1981. The Cornish Sea-urchin Fishery. *Cornish Studies*, **9**, 5-18.
- Nichols, D., 1984. An investigation of the population dynamics of the common edible sea urchin (*Echinus esculentus* L.) in relation to species conservation management. *Report to Department of the Environment and Nature Conservancy Council from the Department of Biological Sciences, University of Exeter*.
- Norderhaug, K., 2004. Use of red algae as hosts by kelp-associated amphipods. *Marine Biology*, **144** (2), 225-230.
- Norderhaug, K.M. & Christie, H.C., 2009. Sea urchin grazing and kelp re-vegetation in the NE Atlantic. *Marine Biology Research*, **5** (6), 515-528.
- Norderhaug, K.M., Christie, H. & Fredriksen, S., 2007. Is habitat size an important factor for faunal abundances on kelp (*Laminaria hyperborea*)? *Journal of Sea Research*, **58** (2), 120-124.
- Nordheim, van, H., Andersen, O.N. & Thissen, J., 1996. Red lists of Biotopes, Flora and Fauna of the Trilateral Wadden Sea area, 1995. *Helgolander Meeresuntersuchungen*, **50** (Suppl.), 1-136.
- Norton, T.A. & South, G.R., 1969. Influence of reduced salinity on the distribution of two laminarian algae. *Oikos*, **20**, 320-326
- Norton, T.A., 1970. Synopsis of biological data on *Saccorhiza polyschides*. *FAO Fisheries Synopsis*, No. 83, 1-35.
- Norton, T.A., 1978. The factors influencing the distribution of *Saccorhiza polyschides* in the region of Lough Ine. *Journal of the Marine Biological Association of the United Kingdom*, **58**, 527-536.
- Norton, T.A., 1992. Dispersal by macroalgae. *British Phycological Journal*, **27**, 293-301.
- Norton, T.A., Hiscock, K. & Kitching, J.A., 1977. The Ecology of Lough Ine XX. The *Laminaria* forest at Carrigathorna. *Journal of Ecology*, **65**, 919-941.
- O'Brien, P.J. & Dixon, P.S., 1976. Effects of oils and oil components on algae: a review. *British Phycological Journal*, **11**, 115-142.
- Pedersen, M.F., Nejrup, L.B., Fredriksen, S., Christie, H. & Norderhaug, K.M., 2012. Effects of wave exposure on population structure, demography, biomass and productivity of the kelp *Laminaria hyperborea*. *Marine Ecology Progress Series*, **451**, 45-60.
- Penfold, R., Hughson, S., & Boyle, N., 1996. *The potential for a sea urchin fishery in Shetland*. <http://www.nafc.ac.uk/publish/note5/note5.htm>, 2000-04-14
- Peters, A.F. & Schaffelke, B., 1996. *Streblonema* (Ectocarpales, Phaeophyceae) infection in the kelp *Laminaria saccharina* in the western Baltic. *Hydrobiologia*, **326/327**, 111-116.
- Philippart, C.J., Anadón, R., Danovaro, R., Dippner, J.W., Drinkwater, K.F., Hawkins, S.J., Oguz, T., O'Sullivan, G. & Reid, P.C., 2011. Impacts of climate change on European marine ecosystems: observations, expectations and indicators. *Journal of Experimental Marine Biology and Ecology*, **400** (1), 52-69.
- Raffaelli, D. & Hawkins, S., 1999. *Intertidal Ecology* 2nd edn.. London: Kluwer Academic Publishers.
- Read, P.A., Anderson, K.J., Matthews, J.E., Watson, P.G., Halliday, M.C. & Shiells, G.M., 1983. Effects of pollution on the benthos of the Firth of Forth. *Marine Pollution Bulletin*, **14**, 12-16.
- Rinde, E. & Sjøtun, K., 2005. Demographic variation in the kelp *Laminaria hyperborea* along a latitudinal gradient. *Marine Biology*,

146 (6), 1051-1062.

- Rostron, D.M. & Bunker, F. St P.D., 1997. An assessment of sublittoral epibenthic communities and species following the Sea Empress oil spill. A report to the Countryside Council for Wales from Marine Seen & Sub-Sea Survey., Countryside Council for Wales, Bangor, CCW Sea Empress Contact Science, no. 177.
- Santos, R., 1993. A multivariate study of biotic and abiotic relationships in a subtidal algal stand. *Marine Ecology Progress Series*, **94**, 181-190.
- Schiel, D.R. & Foster, M.S., 1986. The structure of subtidal algal stands in temperate waters. *Oceanography and Marine Biology: an Annual Review*, **24**, 265-307.
- Sheppard, C.R.C., Bellamy, D.J. & Sheppard, A.L.S., 1980. Study of the fauna inhabiting the holdfasts of *Laminaria hyperborea* (Gunn.) Fosl. along some environmental and geographical gradients. *Marine Environmental Research*, **4**, 25-51.
- Sivertsen, K., 1997. Geographic and environmental factors affecting the distribution of kelp beds and barren grounds and changes in biota associated with kelp reduction at sites along the Norwegian coast. *Canadian Journal of Fisheries and Aquatic Sciences*, **54**, 2872-2887.
- Sjøtun, K., Christie, H. & Helge Fosså, J., 2006. The combined effect of canopy shading and sea urchin grazing on recruitment in kelp forest (*Laminaria hyperborea*). *Marine Biology Research*, **2** (1), 24-32.
- Sjøtun, K. & Schoschina, E.V., 2002. Gametophytic development of *Laminaria* spp. (Laminariales, Phaeophyta) at low temperatures. *Phycologia*, **41**, 147-152.
- Smale, D.A., Burrows, M.T., Moore, P., O'Connor, N. & Hawkins, S.J., 2013. Threats and knowledge gaps for ecosystem services provided by kelp forests: a northeast Atlantic perspective. *Ecology and evolution*, **3** (11), 4016-4038.
- Smale, D.A., Wernberg, T., Yunnice, A.L. & Vance, T., 2014. The rise of *Laminaria ochroleuca* in the Western English Channel (UK) and comparisons with its competitor and assemblage dominant *Laminaria hyperborea*. *Marine ecology*.
- Smith, J.E. (ed.), 1968. 'Torrey Canyon'. *Pollution and marine life*. Cambridge: Cambridge University Press.
- Somerfield, P.J. & Warwick, R.M., 1999. Appraisal of environmental impact and recovery using *Laminaria* holdfast faunas. *Sea Empress, Environmental Evaluation Committee., Countryside Council for Wales, Bangor, CCW Sea Empress Contract Science, Report no. 321*.
- Staehr, P.A. & Wernberg, T., 2009. Physiological responses of *Ecklonia radiata* (Laminariales) to a latitudinal gradient in ocean temperature. *Journal of Phycology*, **45**, 91-99.
- Steneck, R.S., Graham, M.H., Bourque, B.J., Corbett, D., Erlandson, J.M., Estes, J.A. & Tegner, M.J., 2002. Kelp forest ecosystems: biodiversity, stability, resilience and future. *Environmental conservation*, **29** (04), 436-459.
- Steneck, R.S., Vavrinc, J. & Leland, A.V., 2004. Accelerating trophic-level dysfunction in kelp forest ecosystems of the western North Atlantic. *Ecosystems*, **7** (4), 323-332.
- Svendsen, P. & Kain, J.M., 1971. The taxonomic status, distribution, and morphology of *Laminaria cucullata* sensu Jorde and Klavestad. *Sarsia*, **46** (1), 1-22.
- Thompson, G.A. & Schiel, D.R., 2012. Resistance and facilitation by native algal communities in the invasion success of *Undaria pinnatifida*. *Marine Ecology, Progress Series*, **468**, 95-105.
- Thompson, R.S. & Burrows, E.M., 1984. The toxicity of copper, zinc and mercury to the brown macroalga *Laminaria saccharina*. In *Ecotoxicological testing for the marine environment* (ed. G. Persoone, E. Jaspers, & C. Claus), Vol. 2, pp. 259-269. Ghent: Laboratory for biological research in aquatic pollution, State University of Ghent.
- Ursin, E., 1960. A quantitative investigation of the echinoderm fauna of the central North Sea. *Meddelelser fra Danmark Fiskeri-og-Havundersogelser*, **2** (24), pp. 204.
- Vadas, R.L. & Elner, R.W., 1992. *Plant-animal interactions in the north-west Atlantic*. In *Plant-animal interactions in the marine benthos*, (ed. D.M. John, S.J. Hawkins & J.H. Price), 33-60. Oxford: Clarendon Press. [Systematics Association Special Volume, no. 46].
- Vadas, R.L., Johnson, S. & Norton, T.A., 1992. Recruitment and mortality of early post-settlement stages of benthic algae. *British Phycological Journal*, **27**, 331-351.
- Van den Hoek, C., 1982. The distribution of benthic marine algae in relation to the temperature regulation of their life histories. *Biological Journal of the Linnean Society*, **18**, 81-144.
- Vost, L.M., 1983. The influence of *Echinus esculentus* grazing on subtidal algal communities. *British Phycological Journal*, **18**, 211.
- Werner, A. & Kraan, S., 2004. Review of the potential mechanisation of kelp harvesting in Ireland. *Marine Environment and Health Series*, (No. 17).
- Whittick, A., 1983. Spatial and temporal distributions of dominant epiphytes on the stipes of *Laminaria hyperborea* (Gunn.) Fosl. (Phaeophyta: Laminariales) in S.E. Scotland. *Journal of Experimental Marine Biology and Ecology*, **73**, 1-10.
- Wotton, D.M., O'Brien, C., Stuart, M.D. & Fergus, D.J., 2004. Eradication success down under: heat treatment of a sunken trawler to kill the invasive seaweed *Undaria pinnatifida*. *Marine Pollution Bulletin*, **49** (9), 844-849.