



MarLIN

Marine Information Network

Information on the species and habitats around the coasts and sea of the British Isles

A bristleworm (*Cirratulus cirratus*)

MarLIN – Marine Life Information Network
Biology and Sensitivity Key Information Review

Ken Neal & Susie Ballerstedt

2006-05-02

A report from:

The Marine Life Information Network, Marine Biological Association of the United Kingdom.

Please note. This MarESA report is a dated version of the online review. Please refer to the website for the most up-to-date version [<https://www.marlin.ac.uk/species/detail/1786>]. All terms and the MarESA methodology are outlined on the website (<https://www.marlin.ac.uk>)

This review can be cited as:

Neal, K.J. & Ballerstedt, S. 2006. *Cirratulus cirratus* A bristleworm. In Tyler-Walters H. and Hiscock K. (eds) *Marine Life Information Network: Biology and Sensitivity Key Information Reviews*, [on-line]. Plymouth: Marine Biological Association of the United Kingdom.
DOI <https://dx.doi.org/10.17031/marlin.sp.1786.2>



The information (TEXT ONLY) provided by the Marine Life Information Network (MarLIN) is licensed under a Creative Commons Attribution-Non-Commercial-Share Alike 2.0 UK: England & Wales License. Note that images and other media featured on this page are each governed by their own terms and conditions and they may or may not be available for reuse. Permissions beyond the scope of this license are available [here](#). Based on a work at www.marlin.ac.uk

(page left blank)



Cirratulus spp.

Photographer: Teresa Darbyshire

Copyright: National Museum Wales (NMW)

See online review for
distribution map

Distribution data supplied by the Ocean Biogeographic Information System (OBIS). To interrogate UK data visit the NBN Atlas.

Researched by	Ken Neal & Susie Ballerstedt	Refereed by	This information is not refereed.
Authority	(O. F. Müller, 1776)		
Other common names	-	Synonyms	-

Summary

Description

Cirratulus cirratus has a long, slender, orange, pinkish or brownish-red body, with 75 to 130 segments, and can reach 12 cm in length. The head is a blunt cone with a row of 4 to 8 large black eyes either side that may meet on top of the head. There are two groups of up to 8 feeding tentacles on the first segment. Pairs of long slender gills arise at intervals from the whole length of the body and these appear as a mass of reddish threads when the worm is buried.

Recorded distribution in Britain and Ireland

Found in suitable habitats all round the coast of Britain and Ireland.

Global distribution

Most north west European coasts and also in the south Atlantic.

Habitat

Occurs on the lower shore in mud or muddy sand beneath or between rocks.

Depth range

Mainly intertidal but may be circalittoral.

🔍 Identifying features

- Long, slender body with 75 to 130 segments.
- Up to 12 cm in length.
- Blunt head with 4 to 8 large, black eyes.
- Body is orange, pinkish or brownish red.
- Two pairs of up to 8 feeding tentacles near the head.
- Paired, thread-like gills present along most of the body.

🏛️ Additional information

Cirratulus cirratus is usually found in aggregations of up to 200 individuals. During the breeding season their colour changes, the females become bright yellow and the males white.

✓ Listed by

🔗 Further information sources

Search on:



Biology review

Taxonomy

Phylum	Annelida	Segmented worms e.g. ragworms, tubeworms, fanworms and spoon worms
Class	Polychaeta	Bristleworms, e.g. ragworms, scaleworms, paddleworms, fanworms, tubeworms and spoon worms
Order	Terebellida	
Family	Cirratulidae	
Genus	Cirratulus	
Authority	(O. F. Müller, 1776)	
Recent Synonyms	-	

Biology

Typical abundance	Moderate density
Male size range	5 - 130mm
Male size at maturity	20mm
Female size range	Medium(11-20 cm)
Female size at maturity	
Growth form	Cylindrical
Growth rate	See additional information.
Body flexibility	High (greater than 45 degrees)
Mobility	
Characteristic feeding method	Non-feeding, Surface deposit feeder, Surface deposit feeder
Diet/food source	
Typically feeds on	Diatoms and algal detritus.
Sociability	
Environmental position	Infaunal
Dependency	No information found.
Supports	No information
Is the species harmful?	No

Biology information

Little information on the general biology or life history characteristics of this species was found. *Cirratulus cirratus* is regarded as a generally tolerant species and can be found in moderate densities in areas of high environmental disturbance (e.g. 120 per m² 500 m away from an oil platform) (Levell *et al.*, 1989). Once larvae and juveniles settle, they remain in their burrow and adults do not move. It can grow up to 2 cm between reproductive episodes, which occur every 1-2 years (Olive, 1970).

Habitat preferences

Physiographic preferences	Open coast, Offshore seabed, Strait / sound
----------------------------------	---

Biological zone preferences	Lower circalittoral, Lower eulittoral, Lower infralittoral, Mid eulittoral, Sublittoral fringe, Upper circalittoral, Upper eulittoral
Substratum / habitat preferences	Mud, Muddy gravel, Under boulders
Tidal strength preferences	Weak < 1 knot (<0.5 m/sec.)
Wave exposure preferences	Extremely sheltered, Sheltered, Very sheltered
Salinity preferences	Full (30-40 psu), Reduced (18-30 psu), Variable (18-40 psu)
Depth range	Mainly intertidal but may be circalittoral.
Other preferences	No text entered
Migration Pattern	Non-migratory / resident

Habitat Information

In Northumberland it is the dominant crevice organism on rocky shores between low and high water neaps (Olive, 1970). *Cirratulus cirratus* has been described as an opportunistic deposit feeder that is characteristic of areas of organic enrichment (Penry & Jumars, 1990). *Cirratulus cirratus* is mostly intertidal but is sometimes found subtidally (up to 50 m depth off the Devon coast) (Garwood, 1982; Olive, 1970).

Life history

Adult characteristics

Reproductive type	Gonochoristic (dioecious)
Reproductive frequency	Biannual episodic
Fecundity (number of eggs)	No information
Generation time	1-2 years
Age at maturity	1-2 years
Season	See additional text
Life span	5-10 years

Larval characteristics

Larval/propagule type	-
Larval/juvenile development	Lecithotrophic
Duration of larval stage	< 1 day
Larval dispersal potential	No information
Larval settlement period	Insufficient information

Life history information

Reproduction in *Cirratulus cirratus* is asynchronous i.e. it is not entrained to any of the seasons and members of the population are at different stages of reproductive development at any one time (Garwood, 1982; Gibbs, 1971). Oocytes are 150 µm in diameter and once fertilized are deposited in a jelly mass on the surface of rocks (Petersen, 1999). The eggs hatch as a ciliated post-trochophore after 6 days. The larvae are entirely benthic for the duration of their development,

living off yolk for around 24 days after hatching and then commence adult style deposit feeding (Olive, 1970). Females can spawn 2-3 times in their lifetime and it takes 1-2 years after each spawning to mature a new clutch of oocytes (Olive, 1970). There are separate sexes, the males are white, females are lemon-yellow due to the colour of coelomic oocytes (Gibbs, 1971). Sex ratios vary and have been recorded as 1:1 (Olive, 1970) 1:1.7 and 1:2.8 (Gibbs, 1971). Asexual reproduction by epitoky (clones growing from the posterior end of the worm) may occur in *Cirratulus cirratus*. However, the taxonomic status of *Cirratulus* is in constant review and epitokes may be formed by another species that has been erroneously identified as *Cirratulus cirratus* (Petersen, 1999).

Sensitivity review

This MarLIN sensitivity assessment has been superseded by the MarESA approach to sensitivity assessment. MarLIN assessments used an approach that has now been modified to reflect the most recent conservation imperatives and terminology and are due to be updated by 2016/17.

A Physical Pressures

	Intolerance	Recoverability	Sensitivity	Confidence
Substratum Loss	High	Low	High	Very low
<p><i>Cirratulus cirratus</i> needs stones to live under in a muddy environment and if these were to be removed, mortality is likely to be very high due to desiccation and predation and an intolerance of high has been recorded. For recoverability see additional information.</p>				
Smothering	High	Low	High	Very low
<p><i>Cirratulus cirratus</i> lives in mud under stones with its feeding tentacles spread out on the mud surface. A sudden influx of sediment would probably interfere with feeding and gas exchange and cause high mortality. Therefore, an intolerance of high has been recorded. For recoverability see additional information.</p>				
Increase in suspended sediment	Tolerant*	Not relevant	Not sensitive*	Very low
<p><i>Cirratulus cirratus</i> is often found in estuaries (Clay, 1967g), which are areas of high suspended sediment and it is likely that it is tolerant to an increase in suspended sediment. <i>Cirratulus cirratus</i> feeds on precipitating particles and is likely to benefit from an increase in suspended organic matter. Therefore tolerant* has been recorded.</p>				
Decrease in suspended sediment	Low	Very high	Very Low	
<p>A decrease in suspended sediment is unlikely to make <i>Cirratulus cirratus</i> more vulnerable to predation. <i>Cirratulus cirratus</i> relies on particulate organic matter precipitating onto the substratum for food, so that a decrease in suspended particulates may reduce its food supply. However, the reduced turbidity may increase benthic primary productivity, which would be of benefit to <i>Cirratulus cirratus</i>. Overall, its food supply may be reduced and an intolerance of low has been recorded.</p>				
Desiccation	Not relevant	Not relevant	Not relevant	Not relevant
<p><i>Cirratulus cirratus</i> lives in mud under stones and is therefore unlikely to be subject to desiccation. Not relevant has been recorded.</p>				
Increase in emergence regime	Intermediate	High	Low	Very low
<p><i>Cirratulus cirratus</i> is found from the subtidal up to the high water of neap tide level and an increase in emergence time is likely to exclude worms from the upper limit of their range. There would probably be some mortality near high water neap tide level and a shift of the population down to the new high water neaps level. Therefore, an intolerance of intermediate has been recorded.</p>				
Decrease in emergence regime	Tolerant*	Not relevant	Not sensitive*	
<p>A decrease in emergence is likely to have the opposite effect of an increase and extend the range of the population up the shore to the new high water neaps level. Therefore, tolerant* has been recorded.</p>				

Increase in water flow rate Intermediate High Low Very low

Cirratulus cirratus feeds by laying its tentacles out on the surface of the mud and passing food particles to the mouth by ciliary movements. An increase in water flow rate may affect the ability of *Cirratulus cirratus* to collect food particles. Also, an increase in water flow rate may change sediment characteristics and remove the preferred sediment type of *Cirratulus cirratus*. In increase in water flow, e.g. from weak to strong (see benchmark) is likely to significantly affect the substratum, removing fine muddy deposits, and reducing the area of suitable habitat for *Cirratulus cirratus*. Therefore, an intolerance of intermediate has been recorded.

Decrease in water flow rate Not relevant Not relevant Not relevant

Cirratulus cirratus is found in estuaries and on muddy shores, where the water flow rate is slow so a decrease in water flow rate is unlikely to affect this species and this factor has been assessed as not relevant.

Increase in temperature Tolerant Not relevant Not sensitive Moderate

Cirratulus cirratus probably has wide temperature tolerances (7-25°C (Gibbs, 1971). *Cirratulus cirratus* is probably tolerant of temperature changes at the benchmark level.

Decrease in temperature Low High Low High

Cirratulus cirratus probably has wide temperature tolerances (7-25°C (Gibbs, 1971). During the extremely cold winter in the United Kingdom in 1962-63, the distribution of *Cirratulus cirratus* did not change (George, 1968). *Cirratulus cirratus* survived temperatures as low as -2°C for long periods but was killed after a few hours in -4°C due to ice crystals forming in its tissues (George, 1968). At the benchmark level, *Cirratulus cirratus* will probably not be adversely affected by a decrease in temperature and an intolerance of low has been recorded.

Increase in turbidity Tolerant Not relevant Not sensitive Very low

Reduced illumination due to turbidity may reduce the productivity of the microalgae that *Cirratulus cirratus* feeds upon. However, it also feeds on particulate organic matter and, therefore, is not likely to be adversely affected.

Decrease in turbidity Tolerant Not relevant Not sensitive Not relevant

A decrease in turbidity is likely to increase benthic microalgal productivity, which could potentially benefit *Cirratulus cirratus*. However, the relative contribution of benthic microalgae and organic matter to its diet is unknown. Therefore, tolerant has been recorded.

Increase in wave exposure Intermediate High Low Low

Increasing wave exposure increases the incidence of turnover of rocks on the shore and will also increase the size of rocks disturbed (Osman, 1977). If the rocks on the shore are turned over, anything underneath is likely to be washed out of the sediment by subsequent waves and suffer increased predation. *Cirratulus cirratus* is not a very motile species and if it were revealed by loss of a protecting rock, mortality is likely to be high and an intolerance of intermediate has been recorded to account for this.

Decrease in wave exposure Not relevant Not relevant Not relevant

Cirratulus cirratus is found from sheltered to extremely sheltered shores and therefore a decrease in wave exposure is not relevant.

Noise Tolerant Not relevant Not sensitive High

At most, *Cirratulus cirratus* will have only a limited ability to detect sound or vibration and therefore is unlikely to be sensitive to noise.

Visual Presence Tolerant Not relevant Not sensitive High

Cirratulus cirratus does have eyes but it lives under rocks with only the deposit feeding tentacles exposed and so is probably tolerant to visual presence at the benchmark level.

Abrasion & physical disturbance Intermediate High Low Very low

At the benchmark level, abrasion is likely to cause some mortality by moving stones and unearthing the worms. Disturbing rocks may also affect the survivorship of embryos in eggs attached to the rock surface. Therefore an intolerance of intermediate has been recorded.

Displacement Not relevant

Cirratulus cirratus establishes under rocks in mud as a larva and there was no information on whether adults reburrow if disturbed. There is insufficient information to assess the intolerance of *Cirratulus cirratus* to displacement.

Chemical Pressures

Synthetic compound contamination Intolerance High Recoverability Low Sensitivity High Confidence High

After a spill of fuel oil in Southampton Water, dispersants were used to clean certain areas and high mortalities of *Cirratulus cirratus* were observed (George, 1971). Three dispersants were tested for their effects on *Cirratulus cirratus* survivorship, BP 1002, Essolvne (which was used in Southampton Water) and Corexit 7664. BP 1002 and Essolvne become toxic at 100 ppm. BP 1002 caused 50% mortality at 129 ppm and 100% mortality at 144 ppm. Essolvne was slightly less toxic, causing 50% mortality at 162 ppm and 100% mortality at 200 ppm. Corexit was far less toxic than either BP 1002 or Essolvne. It took a concentration of 100,000 ppm of Corexit 7664 to cause 50% to *Cirratulus cirratus*. Longer exposure to sublethal concentrations of BP 1002 or Essolvne completely prevented the maturation of oocytes (George, 1971). The evidence presented above suggests an intolerance of high to synthetic chemicals. For recoverability, see additional information below.

Heavy metal contamination Not relevant

Insufficient information.

Hydrocarbon contamination Low High Low High

A spill of fuel oil in Southampton water lead to widespread oiling of intertidal mud but this had very little effect on *Cirratulus cirratus* abundance. The thickness of the oil was not sufficient to prevent oxygen reaching the sediment and at high tide, the oil refloated so that *Cirratulus cirratus* could feed as normal. Embryo development was also unaffected by oil (George, 1971) and an intolerance of low has been recorded.

Radionuclide contamination Not relevant

Insufficient information.

Changes in nutrient levels Tolerant* Not relevant Not sensitive* Moderate

Cirratulus cirratus is characteristic of areas of organic enrichment (Penry & Jumars, 1990) and therefore is probably tolerant* of an increase in nutrient levels.

Increase in salinity Not relevant

No information on hypersaline conditions was found.

Decrease in salinity Low High Low

Cirratulus cirratus can tolerate salinities down to 17 psu (Gibbs, 1971) and so is likely to survive the benchmark chronic change as it is normally found intertidally at full salinity. An acute change will probably stress a population of *Cirratulus cirratus* but not cause high mortality and an intolerance of low has been recorded.

Changes in oxygenation Low High Low Very low

Cirratulus cirratus is characteristic of areas of organic enrichment (Penry & Jumars, 1990), although no further information was found. Therefore, an intolerance of low has been recorded, albeit with very low confidence.

Biological Pressures

Intolerance Recoverability Sensitivity Confidence

Introduction of microbial pathogens/parasites

Insufficient information

Introduction of non-native species

Insufficient information

Extraction of this species

Not relevant Not relevant Not relevant Not relevant

Cirratulus cirratus is not targeted for extraction.

Extraction of other species

Not relevant Not relevant Not relevant Not relevant

No co-occurring species to *Cirratulus cirratus* are known to be extracted.

Additional information

Recoverability

The fecundity of *Cirratulus cirratus* is unknown but the larvae are entirely benthic throughout their development (Olive, 1970) and, if an area is completely defaunated, recolonization by *Cirratulus cirratus* may be slow (George, 1968). Recovery by populations that suffer partial mortality could take up to 2 years. However, populations that are completely wiped out may not recover at all due to the limited dispersal capability of the larvae of *Cirratulus cirratus* (George, 1968).

Importance review

Policy/legislation

- no data -

Status

National (GB)
importance -

Global red list
(IUCN) category -

Non-native

Native -

Origin -

Date Arrived -

Importance information

-none-

Bibliography

- Clay, E., 1967g. *Literature survey of the common fauna of estuaries*, 7. *Cirratulus cirratus* O.F. Müller. I.C.I. Research Laboratory, Brixham. PVM45/B/380.
- Fauchald, K., 1977. *The polychaete worms. Definitions and keys to the orders, families and genera*. USA: Natural History Museum of Los Angeles County.
- Fish, J.D. & Fish, S., 1996. *A student's guide to the seashore*. Cambridge: Cambridge University Press.
- Garwood, P.R., 1982. Polychaeta - Sedentaria incl. Archiannelida. *Report of the Dove Marine Laboratory Third Series*, **23**, 273p.
- George, J.D., 1968. The effect of the 1962-63 winter on the distribution of the cirratulid polychaetes, *Cirratulus cirratus* (Müller) and *Cirriformia tentaculata* (Montagu) in the British Isles. *Journal of Animal Ecology*, **37**, 321-31.
- George, J.D., 1971. The effects of pollution by oil and oil dispersants on the common intertidal polychaetes, *Cirriformia tentaculata* and *Cirratulus cirratus*. *Journal of Applied Ecology*, **8**, 411-420.
- Gibbs, P.E., 1971. Reproductive cycles in four polychaete species belonging to the family Cirratulidae. *Journal of the Marine Biological Association of the United Kingdom*, **51**, 745-769.
- Hayward, P., Nelson-Smith, T. & Shields, C. 1996. *Collins pocket guide. Sea shore of Britain and northern Europe*. London: HarperCollins.
- Hayward, P.J. & Ryland, J.S. (ed.) 1995b. *Handbook of the marine fauna of North-West Europe*. Oxford: Oxford University Press.
- Howson, C.M. & Picton, B.E., 1997. *The species directory of the marine fauna and flora of the British Isles and surrounding seas*. Belfast: Ulster Museum. [Ulster Museum publication, no. 276.]
- JNCC (Joint Nature Conservation Committee), 1999. *Marine Environment Resource Mapping And Information Database (MERMAID): Marine Nature Conservation Review Survey Database*. [on-line] <http://www.jncc.gov.uk/mermaid>
- Levell, D., Rostron, D. & Dixon, I.M.T., 1989. Sediment macrobenthic communities from oil ports to offshore oilfields. In *Ecological Impacts of the Oil Industry*, Ed. B. Dicks. Chichester: John Wiley & Sons Ltd.
- Olive, P.J.W., 1970. Reproduction of a Northumberland population of the polychaete *Cirratulus cirratus*. *Marine Biology*, **5**, 259-273.
- Osman, R.W., 1977. The establishment and development of a marine epifaunal community. *Ecological Monographs*, **47**, 37-63.
- Penry, D.L. & Jumars, P.A., 1990. Gut architecture, digestive constraints and feeding ecology of deposit-feeding and carnivorous polychaetes. *Oecologia*, **82**, 1-11.
- Petersen, M.E., 1999. Reproduction and development in Cirratulidae (Annelida: Polychaeta). *Marine Biology*, **8**, 243-259.

Datasets

- Centre for Environmental Data and Recording, 2018. Ulster Museum Marine Surveys of Northern Ireland Coastal Waters. Occurrence dataset <https://www.nmni.com/CEDaR/CEDaR-Centre-for-Environmental-Data-and-Recording.aspx> accessed via NBNAtlas.org on 2018-09-25.
- Fenwick, 2018. Aphotomarine. Occurrence dataset <http://www.aphotomarine.com/index.html> Accessed via NBNAtlas.org on 2018-10-01
- Kent Wildlife Trust, 2018. Kent Wildlife Trust Shoresearch Intertidal Survey 2004 onwards. Occurrence dataset: <https://www.kentwildlifetrust.org.uk/> accessed via NBNAtlas.org on 2018-10-01.
- NBN (National Biodiversity Network) Atlas. Available from: <https://www.nbnatlas.org>.
- OBIS (Ocean Biogeographic Information System), 2019. Global map of species distribution using gridded data. Available from: Ocean Biogeographic Information System. www.iobis.org. Accessed: 2019-03-21
- South East Wales Biodiversity Records Centre, 2018. SEWBRc Worms (South East Wales). Occurrence dataset: <https://doi.org/10.15468/5vh0w8> accessed via GBIF.org on 2018-10-02.