

OPEN

JAIDS Journal of Acquired Immune Deficiency Syndromes Publish Ahead of Print

DOI: 10.1097/QAI.0000000000002263

Improving Early Infant Diagnosis Observations: Estimates of timely HIV testing and mortality among HIV-exposed infants

Karen Webb, MSc^{1,2} §, Vivian Chitiyo, MPhil¹, Nyikadzino Mahachi, MD MPH³, Solomon Huruva Mukungunugwa, MD, MBChB, MPH³, Angela Mushavi MBChB MMed³, Simukai Zizhou, MBChB, MPH⁴, Barbara Engelsmann, MD MPH¹, Rashida Abbas Ferrand MD, MSc, PhD², Melissa Neuman, ScD MS², Wendy Hartogensis, PhD MPH⁵, Elvin Geng, MD MPH⁵

¹Organization for Public Health Interventions and Development, Harare, Zimbabwe;

²London School of Hygiene and Tropical Medicine, London, UK;

³Ministry of Health and Child Care, National PMTCT Program, Harare, Zimbabwe;

⁴Ministry of Health and Child Care, Mashonaland East Province, Zimbabwe;

⁵University of California, San Francisco, USA

Corresponding Author:

Karen Webb,

Organization for Public Health Interventions and Development

20 Cork Road, Belgravia

Harare, Zimbabwe

Tel: +263772141699/+263242252772; Email: kwebb@ophid.co.zw

Conflicts of interest: None

Word Count:

Abstract: 249

Manuscript: 1, 982

Number of Tables: 0

Number of Figures: 2

Meetings at which parts of data were presented:

1. 8th IAS Conference on HIV Pathogenesis, Treatment and Prevention (IAS 2015), Vancouver Canada, 19-22 July 2015; Abstract: MOPED747.
2. 18th International Conference on AIDS and STIs in Africa (ICASA 2015), Harare, Zimbabwe, 29 November – 4 December 2015; Abstract: THUAC1105.
3. 21st International AIDS Conference (AIDS 2016), 18-22 July 2016, Durban, South Africa; Abstract: LBPE036.

Funding: Supported by the President's Emergency Plan for AIDS Relief (PEPFAR) through USAID and Families and Communities for Elimination of HIV in Zimbabwe (AID-613-A-12-00003, FACE HIV) (OPHID); K24 AI 134413 (EG); CFAR ISWG (P30 AI027763) (EG)

Running head:

Loss to follow up for Early Infant Diagnosis

This is an open access article distributed under the [Creative Commons Attribution License 4.0 \(CCBY\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract/Key Word Page

Background: Improving efforts towards elimination of mother-to-child transmission of HIV requires timely early infant diagnosis (EID) among all HIV-exposed infants, but the occurrence of timely EID and infant survival may be under-ascertained in routine, facility-bound program data.

Methods: From March-May 2015, we traced a random sample of HIV positive mother and HIV-exposed infant pairs lost to follow-up for EID in facility registers in Zimbabwe. We incorporated updated information into weighted survival analyses to estimate incidence of EID and death. Reasons for no EID were surveyed from caregivers.

Results: Among 2,651 HIV-positive women attending antenatal care, 1,823 (68.8%) infants had no documented EID by 3 months of age. Among a random sample of 643 (35.3%) HIV-exposed infants lost to follow-up for EID, vital status was ascertained among 371 (57.7%) and updated care status obtained from 256 (39.8%) mothers traced. Among all HIV-infected mother HIV-exposed infant pairs, weighted estimates found cumulative incidence of infant death by 90 days of 3.9% (95% CI: 3.4% to 4.4%). Cumulative incidence of timely EID with death as a competing risk was 60%. The most frequently cited reasons for failure to uptake EID were 'my child died' and 'I didn't know I should have my child tested'.

Conclusions: Our findings indicate uptake of timely EID among HIV-exposed infants is underestimated in routine health information systems. High, early mortality among HIV-exposed infants underscores the need to more effectively identify HIV positive mother-HIV exposed infant pairs at high risk of adverse outcomes and loss to follow-up for enhanced interventions.

Keywords: HIV-exposed infants, early infant diagnosis, loss to follow-up, prevention of mother-to-child HIV transmission, retention

INTRODUCTION

The timeliness of early infant diagnosis (EID) – HIV testing of exposed infants six to eight weeks after birth – and proportion of infants testing positive, are cardinal indicators of prevention of mother to child transmission (PMTCT) program success.¹ However, in routine program settings, completion of EID is assessed at individual facilities, from information across several paper-based registers, which may be incomplete and/or inaccurate.^{2,3}

New approaches are needed to improve confidence in estimates of EID completion and survival of HIV-exposed infants to guide on-going quality improvement and inform national modelling estimates. These become increasingly important as countries such as Zimbabwe, with an HIV prevalence of 16.0% among women,⁴ seek to validate elimination of mother to child transmission.⁵

In this study, we randomly sampled lost-to-follow up (LTFU) HIV-positive mother, HIV-exposed baby pairs (MB) pairs from routine health facility-based data. We actively traced MB pairs in the community to assess infant and maternal survival and EID uptake. Finally, we incorporated these findings to correct estimates of HIV-exposed infant mortality and timely EID in the entire clinic population of MB pairs using a probability weight. Although this sampling-based approach has been used in clinic-based cohorts of adult patients in HIV treatment programs and found to alter estimates of retention and mortality,^{6,7} our study represents the first time the approach has been used in PMTCT programs. We reflect on key lessons from our

research on program progress and persisting gaps for strengthening EID coverage on the path to elimination of MTCT in Zimbabwe.

METHODS

Study Design

We defined LTFU as no documented sample taken for EID polymerase chain reaction (PCR) testing among infants of HIV-positive antenatal care (ANC) patients within 90 days of delivery. At the time of study, Zimbabwe HIV care and treatment guidelines recommended HIV testing of all exposed children 6 weeks postnatal.⁸ Our sample included all women with HIV who accessed ANC at public sector health facilities in Mashonaland East Province (referral sites excluded), using a multi-stage clustered survey sampling approach.

Phase I: Register Tracing: We selected 5 clinics from each of the 9 Districts within Mashonaland East Province based on probability proportional to size, using program data on HIV-positive women accessing ANC over the previous year. In the absence of inter-facility-linked electronic patient monitoring systems, we used a combination of individual identifiers including clinic-allocated PMTCT number, patient name and date of birth, to trace each identified HIV positive pregnant woman in ANC and her HIV-exposed infant through five paper-based registers (ANC, Delivery, HIV-Exposed Infants, Clinic Registers, HIV Infant Diagnosis and EID Lab Request Form Booklets) to determine LTFU for EID status. We targeted a random sample of at least 10% of lost MB pairs based on practical considerations regarding limitations of completeness and accuracy of patient information in multiple paper-based registers,⁹ and available resources to intensively trace those identified as LTFU for EID over the study timeframe. Based upon known challenges in tracing defaulters from HIV care and treatment,¹⁰⁻¹² we over-sampled by 100% to ensure we met our target sample.

Phase II: Tracing of LTFU MB Pairs. In the second stage of sampling, a random sample of patients identified as lacking documented EID in Stage 1 was selected for active tracing to determine outcomes through direct patient interviews. We trained MOHCC Village Health Workers (VHWs) as ascertainers due to their familiarity with the surrounding community and existing role to support health facilities with defaulter tracing.^{13,14}

VHWs were assigned selected LTFU patients residing within their geographical catchment area for tracing. If the individual was unable to be located, mother and infant vital status was obtained from informants (friends, neighbors or relatives); however, to maintain confidentiality, informants were not asked any specific questions related to HIV services. Successfully traced and consenting mothers were interviewed using a standardized questionnaire to ascertain: survival outcomes, EID status and timing, and reasons for failure to obtain EID or for self-transfer of care to another facility. Analyses were conducted using Stata v.13.1. This study was approved by the Medical Research Council of Zimbabwe (MRCZ/A/1844) and MOHCC authorities.

Analyses

EID estimates: First, we estimated EID incidence based on clinic registry data only, using facility weights inverse to the probability of clinic selection to yield an estimate of EID completion based purely on paper registers kept at the facilities. Facility and individual factors on documented completion of EID were explored using Poisson regression with robust standard errors adjusted for confounders to estimate risk ratios.^{15,16}

Second, we generated corrected estimates of EID completion using data from clinic registries as well as data ascertained through interviews with a random sample of MB pairs missing EID in facility-based registers.⁷ Deaths before EID (including fetal deaths) were considered as competing risks. To obtain an estimate corrected for outcomes not captured in the facility registers, we used additional sampling weights inverse to the probability of being successfully sought to represent all LTFU HIV-infected pregnant women at each clinic.

Among women with no documented EID in facility registers who were traced and interviewed, reasons for silent transfer to a different clinic from antenatal care for EID and for not bringing HIV-exposed child in for HIV testing (no EID) were grouped into four categories informed by a socio-ecological framework^{17,18}: structural; clinic-based; psychosocial or patient-related; or medical factors.

Mortality estimates: For the weighted mortality estimate, the traced patients contributed time from the delivery date to the date of the patient interview or the date of the death of the infant at any time prior (i.e., all deaths including those later than 90 days were included in the mortality estimate); patients with documented EID contributed time from the date of delivery to the date of the 6-week postnatal visit, which is the latest date at which we have confirmed vital status for those infants. Kaplan-Meier methods were used to estimate mortality.

RESULTS

Phase I: Facility Register Data

Among 18,065 women registered for ANC between April 2012 and May 2013, 2,651 (14.7%) were HIV-positive; 31.2% (95%CI: 29.5%-33.0%) had documented uptake of EID for their infant within three months of delivery in clinic registers. After adjustment for register

information and site characteristics, factors associated with documented EID completion included earlier gestational age at presentation (Risk Ratio (RR): 0.97 per two weeks; 95%CI:0.95-0.99; p=0.013), later calendar time of ANC presentation (RR: 1.04 per 30 days; 95%CI:1.02-1.06, p= 0.011) and smaller site volume (RR: 1.85 1-200 ANC patient volume, 1001-1500 volume referent; 95%CI: 1.44-2.38, p<0.001).

Phase II: Community Tracing LTFU

Among 1,652 mother-baby pairs identified as LTFU with any documented locator information, a random sample of 643 (38.9%) were selected for community tracing between March-May 2015. In 371/643 (57.7%), updated vital or EID status information was obtained (22.5%; 371/1,652 of the total LTFU sample). The primary reason for failure to locate clients was insufficient location information. Among 371 successfully traced patients, 256 (69.0% of located) mothers were interviewed directly on infant vital status and EID uptake and for the remaining 115 (31.0%), informants were interviewed regarding MB pair survival outcomes (not HIV-related) (Figure 1). Among the 371 mother-baby pairs for whom vital status outcomes were determined, 66 infants (17.8%; 95% CI: 14.0 – 22.1%) and 18 mothers (4.9%; 95% CI: 2.9%-7.6%) were found to be deceased.

Most mothers interviewed (190/256; 74.2%; 95% CI: 68.4% - 79.5%) reported their infant had received HIV testing; though fewer than half received EID testing prior to 3 months of age (92/190; 48.4%; 95%CI: 41.1% - 55.8%). Our corrected estimate following tracing resulted in a cumulative incidence of EID with death as a competing risk of 60.0% (95%CI: 58.7% - 61.3%). We estimated a cumulative incidence of mortality among HIV-exposed infants at three months of 3.9% (95%CI: 3.4% - 4.4%) and at one-year of 7.7% (95%CI: 4.7% - 13.5%). Among the 66

infants with no EID at any time, the most frequently cited reason for failure to have EID was ‘my child died’ (36.6%; 95% CI: 25.8%-49.0%), Figure 2A. Among infants with timing of death ascertained, most did not survive to the age of recommended EID testing (6 weeks) (26/42; 61.9%). Among mothers of living infants, ‘I didn’t know I should have my child tested’ was the most frequently cited reason for no EID (16/45; 35.6%). Relocation to a different area (21.8%; 95% CI: 13.2%-32.6%) and transport being easier/cheaper at new clinic (20.5%; 95% CI: 12.2%-31.1%) were the most commonly reported reasons for silent transfer (Figure 2B).

DISCUSSION

In a representative sample of Mashonaland East Province of Zimbabwe, after tracing a sample of MB pairs identified as LTFU for EID, our corrected estimate of EID almost doubled (from 31.2%-60.0%). These findings underscore the risk of equating LTFU of MB pairs in health information systems with disengagement from care where such systems do not facilitate electronic, longitudinal tracing of service uptake within and between health facilities.¹⁹ MOHCC is currently expanding efforts to strengthen facility-based documentation, retention monitoring and longitudinal outcome reporting through use of the Mother-Baby Pair Register to track longitudinal outcomes of MB Pairs, together with appointment diary systems.²⁰ As Zimbabwe transitions towards electronic health record systems, we indicate the value of sampling-based methods for improving accuracy of EID coverage estimates and informing current progress towards EMTCT validation.²¹

We found high mortality among HIV-exposed infants, with ‘my child died’ being the most frequently cited reason for no EID testing. Passive facility-based monitoring may underestimate true mortality by up to 80%²² and under-reporting of infant mortality in high HIV burden settings

is acknowledged to bias child mortality estimates downward.^{23,24} Our findings reinforce the need to act early among HIV-positive mother-exposed infant pairs at high risk of defaulting or adverse clinical outcomes with enhanced PMTCT interventions.^{25,26} Subsequent MOHCC guidelines recommending prioritisation of viral load monitoring for pregnant and lactating mothers, birth testing for high-risk infants^{20,27}, and case-based surveillance of infants testing HIV positive^{20,28} are intended strengthen program evidence and action among high risk MB pairs. Our findings emphasize that monitoring of facility-level implementation fidelity, data quality, and robust analysis of resulting data will be central to realising the benefit of such efforts for improved PMTCT program strategies and impact.

Finally, we not only improved our understanding of ‘true EID’ rates, but also developed evidence on factors influencing timely infant HIV testing for informing quality improvement. At patient-level, the predominance of psychosocial reasons (‘I didn’t know’) for failure to uptake EID among mothers of living infants emphasizes need to provide information about the importance of timely infant HIV testing during initial engagement in antenatal care and continued emphasis at every subsequent visit. Reasons for silent transfer are consistent with studies of adult ART retention and highlight the role of structural²⁹ and clinic-based factors³⁰ for optimizing retention in care along the PMTCT cascade. Findings have guided PMTCT program planning and actions including strengthening of patient education and problem solving, appointment monitoring, active follow-up and outcome documentation in Zimbabwe and other settings.^{20,31-33}

Limitations

While oversampling enabled achievement of our targeted sample of at least 10% of MB pairs LTFU for EID (we traced 22%), our findings highlight need to strengthen routine documentation and other forms of observational data to ascertain uptake of services and mortality.³⁴ We acknowledge non-response due to incomplete tracing details may introduce some uncertainty in the resulting estimates. However, our findings of lower LTFU and higher mortality following tracing are concordant with larger, more robust LTFU studies.³⁵ In the absence of effective inter-facility electronic patient monitoring systems, we demonstrate the value of leveraging routine observational data in ‘real-life’ program settings for improving accuracy of estimates, identifying bottlenecks and guiding programmatic actions at local level.^{34,36,37}

Conclusions

Our findings suggest that routinely reported EID completion in public health information systems may be substantially underestimated. Accurate determination of mother-infant pair outcomes in PMTCT programs is complicated by resource-constrained health information systems that involve multiple paper-based registers, lack of unique identifiers and challenges with completeness and accuracy of information recorded. We demonstrate the value of sampling-based approaches in pediatric HIV research for providing important, context-based evidence for policy and programs.

ACKNOWLEDGEMENTS

We are grateful to the Zimbabwe Ministry of Health and Child Care, participating health facilities, village health workers and study participants. This research was made possible through support from the President’s Emergency Plan for AIDS Relief (PEPFAR) through USAID and

Families and Communities for Elimination of HIV in Zimbabwe (AID-613-A-12-00003, FACE HIV) (OPHID); K24 AI 134413 (EG); CFAR ISWG (P30 AI027763) (EG). The opinions expressed in this article do not necessarily reflect the views or policies of PEPFAR, USAID, NIH or CFAR.

REFERENCES

1. Ghadrshenas A, Ben Amor Y, Chang J, et al. Improved access to early infant diagnosis is a critical part of a child-centric prevention of mother-to-child transmission agenda. *AIDS*. Nov 2013;27 Suppl 2:S197-205.
2. Mate KS, Bennett B, Mphatswe W, Barker P, Rollins N. Challenges for routine health system data management in a large public programme to prevent mother-to-child HIV transmission in South Africa. *PloS one*. 2009;4(5):e5483.
3. Gourlay A, Wringe A, Todd J, et al. Challenges with routine data sources for PMTCT programme monitoring in East Africa: insights from Tanzania. *Global health action*. 2015;8:29987.
4. Ministry of Health and Child Care (MOHCC). *Zimbabwe Population-Based HIV Impact Assessment (ZIMPHIA) 2015-16: First Report*. Harare: MOHCC; July 2017.
5. WHO. *Global guidance on criteria and processes for validation: elimination of mother-to-child transmission of HIV and syphilis*. 2nd Edition. Geneva: WHO; 2017.
6. Geng EH, Bangsberg DR, Musinguzi N, et al. Understanding reasons for and outcomes of patients lost to follow-up in antiretroviral therapy programs in Africa through a sampling-based approach. *J Acquir Immune Defic Syndr*. Mar 2010;53(3):405-411.

7. Geng EH, Odeny TA, Lyamuya RE, et al. Estimation of mortality among HIV-infected people on antiretroviral treatment in East Africa: a sampling based approach in an observational, multisite, cohort study. *The lancet. HIV*. Mar 2015;2(3):e107-116.
8. MOHCC. Guidelines for Antiretroviral Therapy for the Prevention and Treatment of HIV in Zimbabwe, NMTPAC and MOHCC. Harare, 2013.
9. Matyanga P, Webb K, Patel, D, et al. Drowning in registers: Task-oriented job analysis of rural health nurses in Zimbabwe in the era of Option B+ [WEPE451]. Presented at: *20th International AIDS Conference (AIDS 2014)*. Melbourne, Australia, 2014.
10. Vogt F, Ferreyra C, Bernasconi A, et al. Tracing defaulters in HIV prevention of mother-to-child transmission programmes through community health workers: results from a rural setting in Zimbabwe. *Journal of the International AIDS Society*. 2015;18:20022.
11. Tweya H, Gugsu S, Hosseinipour M, et al. Understanding factors, outcomes and reasons for loss to follow-up among women in Option B+ PMTCT programme in Lilongwe, Malawi. *Tropical medicine & international health : TM & IH*. Nov 2014;19(11):1360-1366.
12. Tweya H, Feldacker C, Estill J, et al. Are they really lost? "true" status and reasons for treatment discontinuation among HIV infected patients on antiretroviral therapy considered lost to follow up in Urban Malawi. *PloS one*. 2013;8(9):e75761.
13. Schmitz K, Basera TJ, Egbujie B, et al. Impact of lay health worker programmes on the health outcomes of mother-child pairs of HIV exposed children in Africa: A scoping review. *PloS one*. 2019;14(1):e0211439.

14. MOHCC. The Village Health Worker Handbook. Harare, Zimbabwe.: MOHCC; 2010.
15. Geng EH, Hare CB, Kahn JO, et al. The effect of a "universal antiretroviral therapy" recommendation on HIV RNA levels among HIV-infected patients entering care with a CD4 count greater than 500/muL in a public health setting. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. Dec 2012;55(12):1690-1697.
16. Zou G. A modified poisson regression approach to prospective studies with binary data. *American journal of epidemiology*. Apr 1 2004;159(7):702-706.
17. Geng EH, Odeny TA, Lyamuya R, et al. Retention in Care and Patient-Reported Reasons for Undocumented Transfer or Stopping Care Among HIV-Infected Patients on Antiretroviral Therapy in Eastern Africa: Application of a Sampling-Based Approach. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. Apr 01 2016;62(7):935-944.
18. Rachlis B, Naanyu V, Wachira J, et al. Identifying common barriers and facilitators to linkage and retention in chronic disease care in western Kenya. *BMC public health*. Aug 08 2016;16:741.
19. Myer L, Phillips TK. Beyond "Option B+": Understanding Antiretroviral Therapy (ART) Adherence, Retention in Care and Engagement in ART Services Among Pregnant and Postpartum Women Initiating Therapy in Sub-Saharan Africa. *J Acquir Immune Defic Syndr*. Jun 01 2017;75 Suppl 2:S115-S122.

20. MOHCC. *National strategic plan for eliminating new HIV infections in children and keeping mothers and families alive (2014-2018)*. Harare: Zimbabwe Ministry of Health and Child Care; 2014.
21. WHO. Validation of EMTCT of HIV and/or syphilis: Tools and checklists for in-country evaluation of four required components. Geneva: WHO; July 2017.
22. Yiannoutsos CT, An MW, Frangakis CE, et al. Sampling-based approaches to improve estimation of mortality among patient dropouts: experience from a large PEPFAR-funded program in Western Kenya. *PloS one*. 2008;3(12):e3843.
23. Hug L, Alexander M, You D, Alkema L. National, regional, and global levels and trends in neonatal mortality between 1990 and 2017, with scenario-based projections to 2030: a systematic analysis. *The Lancet. Global health*. Jun 2019;7(6):e710-e720.
24. Hallett TB, Gregson S, Kurwa F, et al. Measuring and correcting biased child mortality statistics in countries with generalized epidemics of HIV infection. *Bulletin of the World Health Organization*. Oct 1 2010;88(10):761-768.
25. Bhardwaj S, Carter B, Aarons GA, Chi BH. Implementation Research for the Prevention of Mother-to-Child HIV Transmission in Sub-Saharan Africa: Existing Evidence, Current Gaps, and New Opportunities. *Current HIV/AIDS reports*. Jun 2015;12(2):246-255.
26. Myer L, Essajee S, Broyles LN, et al. Pregnant and breastfeeding women: A priority population for HIV viral load monitoring. *PLoS medicine*. Aug 2017;14(8):e1002375.
27. MOHCC. *Guidelines for Antiretroviral Therapy for the Prevention and Treatment of HIV in Zimbabwe, 2016*. Harare: NMTPAC and MOHCC; 2016.

28. Webb K, Chitiyo V, Page-Mtongwiza S, Nesara P, Mbetu P, Patel D, Chinyanga T, Mukungunugwa S, Mushavi A. Too little, too late: Early lessons on MTCT risk from HIV positive infant case investigation in Zimbabwe [30]. Paper presented at: International Workshop on HIV Pediatrics; July 19-20, 2019; Mexico City, Mexico.
29. McCoy SI, Buzdugan R, Mushavi A, et al. Food insecurity is a barrier to prevention of mother-to-child HIV transmission services in Zimbabwe: a cross-sectional study. *BMC public health*. 2015;15:420.
30. Zanolini A, Sikombe K, Sikazwe I, et al. Understanding preferences for HIV care and treatment in Zambia: Evidence from a discrete choice experiment among patients who have been lost to follow-up. *PLoS medicine*. Aug 2018;15(8):e1002636.
31. Nance N, Pendo P, Masanja J, et al. Short-term effectiveness of a community health worker intervention for HIV-infected pregnant women in Tanzania to improve treatment adherence and retention in care: A cluster-randomized trial. *PloS one*. 2017;12(8):e0181919.
32. Webb K, Chitiyo V, Patel D, et al. The Action Birth Card: Evaluation of an innovative goal-setting tool to increase demand and uptake of underutilized services along the PMTCT cascade [MOPEC466]. *8th IAS Conference on HIV Pathogenesis, Treatment and Prevention (IAS 2015)*, 2015; Vancouver, Canada.
33. MOHCC. *National strategic plan for eliminating new HIV infections in children and keeping mothers and families alive (2014-2018)*. Harare: Zimbabwe Ministry of Health and Child Care; 2014.

34. Sohn AH, Judd A, Mofenson L, et al. Using Observational Data to Inform HIV Policy Change for Children and Youth. *J Acquir Immune Defic Syndr*. 2018;15(78):S22-S26.
35. McMahon JH, Elliott JH, Hong SY, Bertagnolio S, Jordan MR. Effects of physical tracing on estimates of loss to follow-up, mortality and retention in low and middle income country antiretroviral therapy programs: a systematic review. *PloS one*. 2013;8(2):e56047.
36. Rice B, Boule A, Baral S, et al. Strengthening Routine Data Systems to Track the HIV Epidemic and Guide the Response in Sub-Saharan Africa. *JMIR public health and surveillance*. 2018;4(2):e36-e36.
37. Mark D, Geng E, Vorkoper S, et al. Making Implementation Science Work for Children and Adolescents Living With HIV. *J Acquir Immune Defic Syndr*. Aug 15 2018;78 Suppl 1:S58-S62.

FIGURE LEGENDS

Figure 1

Title: Flowchart of outcomes in the study population (N=18,065)

Caption:

Abbreviations: ANC, antenatal care; HIV, human immunodeficiency virus; EID, early infant diagnosis.


Figure 2

Title: Prevalence of patient-reported reasons for no EID (N=71) (A) and switching site of EID from ANC care (N=78) (B)


Caption:

Prevalence of patient-reported reasons for no Early Infant Diagnosis (EID) (A) and switching site of care (B). Structural barriers stem from material conditions of life in resource-limited settings (e.g., transportation cost and availability, family conflict, not enough money) and are shown in red bars. Psychosocial barriers are related to knowledge, beliefs, or attitudes of the patients in the given social setting (e.g., ‘didn’t know’ child should be tested, fear of disclosure, preference for spiritual healing) and are shown in yellow bars. Clinic-based barriers are related to delivery processes at a clinic site (e.g., long waiting times, healthcare worker friendliness, quality of care) and are shown in green bars. Medical barriers related to health status (such as infant death or mother too sick to bring child to clinic) are shown in blue bars.

Abbreviations: ANC, antenatal care; HIV, human immunodeficiency virus; EID, early infant diagnosis.



A



B

