

OPEN

Rising CO₂ enhances hypoxia tolerance in a marine fish

Daniel W. Montgomery^{1*}, Stephen D. Simpson¹, Georg H. Engelhard^{2,3},
Silvana N. R. Birchenough² & Rod W. Wilson^{1*}

Global environmental change is increasing hypoxia in aquatic ecosystems. During hypoxic events, bacterial respiration causes an increase in carbon dioxide (CO₂) while oxygen (O₂) declines. This is rarely accounted for when assessing hypoxia tolerances of aquatic organisms. We investigated the impact of environmentally realistic increases in CO₂ on responses to hypoxia in European sea bass (*Dicentrarchus labrax*). We conducted a critical oxygen (O_{2crit}) test, a common measure of hypoxia tolerance, using two treatments in which O₂ levels were reduced with constant ambient CO₂ levels (~530 μatm), or with reciprocal increases in CO₂ (rising to ~2,500 μatm). We also assessed blood acid-base chemistry and haemoglobin-O₂ binding affinity of sea bass in hypoxic conditions with ambient (~650 μatm) or raised CO₂ (~1770 μatm) levels. Sea bass exhibited greater hypoxia tolerance (~20% reduced O_{2crit}), associated with increased haemoglobin-O₂ affinity (~32% fall in P₅₀) of red blood cells, when exposed to reciprocal changes in O₂ and CO₂. This indicates that rising CO₂ which accompanies environmental hypoxia facilitates increased O₂ uptake by the blood in low O₂ conditions, enhancing hypoxia tolerance. We recommend that when impacts of hypoxia on aquatic organisms are assessed, due consideration is given to associated environmental increases in CO₂.

A lack of O₂ is one of the greatest challenges that most life can face. In terrestrial ecosystems conditions of low O₂ are rare. In contrast low O₂, referred to as hypoxia, is much more common in freshwater and marine ecosystems¹⁻⁴. Hypoxia occurs because high biological demand for O₂ can exceed the rate of O₂ supply to the ecosystem, leading to a reduction in environmental O₂ levels^{5,6}. However, the challenges of hypoxia are not solely a result of reduced O₂. Organisms must also contend with simultaneous but reciprocal changes in the other respiratory gas, CO₂.

When O₂ decreases in aquatic systems there is a corresponding increase in CO₂^{7,8}. This is a by-product of respiration, the same process that causes depletion of O₂. As such high CO₂ during hypoxia is ubiquitous and unavoidable. This coupling of O₂ and CO₂ has been highlighted numerous times in oceanographic sciences, most recently by Robinson⁵. Yet unaccountably, despite the known link between decreasing O₂ and increasing CO₂ during hypoxia^{8,9}, the issue of increased environmental CO₂ during periods of low O₂ has been relatively overlooked by biologists.

Implications of rising CO₂ during hypoxia on aquatic organisms are particularly important to address in the face of human driven climate change. Hypoxic areas are predicted to become more common and more severe, particularly in marine systems, with the de-oxygenation of the world's oceans recently highlighted as a major component of climate change^{2,3,10-12}. In addition, there will be an increase in ambient CO₂ as rising atmospheric CO₂ is absorbed by the world's oceans¹³. Non-linear interactive effects between higher atmospheric CO₂ and CO₂ accumulation during hypoxia will lead to increased CO₂ levels during hypoxia in future oceans¹⁴. This means that effects of rising CO₂ during hypoxia in marine systems will be amplified by climate change.

Typically, experiments which test responses to hypoxia or impacts from hypoxia on aquatic organisms create hypoxic conditions by off-gassing oxygen from water by gassing with pure nitrogen or a mix of nitrogen (N₂) and O₂ (for examples see¹⁵⁻¹⁸). This creates low O₂ conditions without the concurrent CO₂ increase that would be expected in the environment. The lack of studies in which an environmentally realistic simultaneous decrease in O₂ and increase of CO₂ have been conducted may lead to mismeasurement of responses to hypoxia. Recently, several studies on marine fish and invertebrates have demonstrated interactive effects of low oxygen and increased CO₂^{9,19,20}, with some species exhibiting loss of equilibrium (LoE) and death at higher O₂ concentrations when

¹Bioscience Department, College of Life and Environmental Sciences, University of Exeter, Exeter, UK. ²Centre for Environment, Fisheries & Aquaculture Science (Cefas), Pakefield Road, Lowestoft, NR33 0HT, UK. ³School of Environmental Sciences, University of East Anglia, Norwich, NR4 7TJ, UK. *email: dm513@exeter.ac.uk; R.W.Wilson@exeter.ac.uk




Figure 1. Calculated critical oxygen level (O_{2crit}) of European sea bass, *Dicentrarchus labrax*, when O_{2crit} tests include a constant ambient CO_2 level ($\sim 500 \mu\text{atm}$, $N = 8$) or an ecologically realistic rise in CO_2 (~ 500 – $2500 \mu\text{atm}$, $N = 7$) during the test. *Indicates significant difference between CO_2 regimes ($p < 0.05$). Boxes represent median value and inter-quartile range, whiskers represent minimum and maximum values.

CO_2 is simultaneously elevated^{19,21}. However, these experiments do not give insight into the physiological mechanisms underlying the influence of CO_2 on hypoxic responses of fish.

Previously observed impacts of hypoxia-associated rises in CO_2 on hypoxia tolerance of fish could be a result of changes in O_2 uptake, as CO_2 has been shown to impact upon several aspects of organismal biology that are involved in O_2 uptake and transport^{22,23}. We aimed to assess whether concurrent increases in CO_2 during decreases in O_2 affect O_2 uptake in a marine fish, the European sea bass (*Dicentrarchus labrax*) by conducting a standard critical O_2 level test (O_{2crit}). Under normal O_2 (normoxic) conditions fish maintain a minimum level of O_2 consumption rate ($\dot{M}O_2$), referred to as the standard metabolic rate (SMR), in order to meet maintenance energetic demands of essential processes through aerobic respiration²⁴. As the level of O_2 in water drops fish deploy a number of responses (i.e. increased ventilatory water flow and cardiac output, increased haematocrit, functional changes in gill morphology, changes in Hb- O_2 affinity) in order to maintain and regulate this minimum level of $\dot{M}O_2$ ^{25–27}. If environmental O_2 continues to drop there comes a point at which fish are unable to regulate $\dot{M}O_2$ to meet minimal energy demands, referred to as the critical O_2 level (O_{2crit}). At O_2 levels below O_{2crit} fish become oxy-conformers (where $\dot{M}O_2$ is directly proportional to environmental O_2 availability) and fish become increasingly reliant on anaerobic metabolism which is unsustainable in the medium to long term. In the past the measure of O_{2crit} has been used as a proxy for overall hypoxia tolerance but recently this approach has been questioned^{28,29}. Nevertheless O_{2crit} does provide information related to the ability of fish to maintain O_2 uptake and supply during hypoxia, and its prevalence in the literature allows comparison of responses between species³⁰. Furthermore, we investigated whether any changes in O_{2crit} could be linked to changes in blood acid-base chemistry and blood gas transport via alteration of Hb- O_2 binding caused by rising environmental CO_2 . Our hypothesis was that the simultaneous increase in CO_2 during a progressive decrease in O_2 would decrease hypoxia tolerance (increase O_{2crit}) and that this response may be a result of blood acid-base disturbance decreasing Hb- O_2 affinity and O_2 transport.

Results

O_{2crit} tests. There was evidence of enhanced tolerance to hypoxia in sea bass exposed to rising compared to constant CO_2 conditions. This was indicated by measurements of O_{2crit} in European sea bass being significantly different between fish exposed to either constant, or rising CO_2 levels during O_{2crit} tests when accounting for variation in SMR (Fig. 1; ANCOVA, $F_{1,12} = 7.525$, $p = 0.0178$). A CO_2 increase during O_{2crit} tests resulted in a 20% reduction of O_{2crit} ($3.88 \pm 0.19 \text{ kPa } O_2$, $18.7 \pm 0.9\%$ air saturation, mean \pm S.E.) when compared to tests in which CO_2 levels were maintained at ambient levels ($4.87 \pm 0.22 \text{ kPa } O_2$, $23.4 \pm 1.1\%$ air saturation, mean \pm S.E.).

Blood chemistry analysis. A comparison of blood chemistry parameters between the two treatment groups indicated that sea bass fully compensated for the rise in CO_2 during hypoxia within 5 hours (the period of exposure prior to blood sampling). Blood pH (pH_b) was not different between fish exposed to constant ambient CO_2 (7.87 ± 0.03) and fish exposed to progressively rising CO_2 (7.88 ± 0.02) (Fig. 2A, GLM, $F_{1,14} = 0.23$, $p = 0.64$). The acidifying effect of the $\sim 79\%$ rise in blood pCO_2 levels in the rising CO_2 regime ($0.272 \pm 0.019 \text{ kPa } CO_2$) compared to the constant ambient CO_2 regime ($0.152 \pm 0.015 \text{ kPa } CO_2$) (Fig. 2C, GLM, $F_{1,13} = 23.9$, $p < 0.001$) was fully compensated by elevating blood HCO_3^- (Fig. 2D, GLM, $F_{1,13} = 40$, $p < 0.001$). Plasma HCO_3^- was 88% higher under the rising CO_2 regime ($6.76 \pm 0.29 \text{ mM}$) when compared to the constant ambient CO_2 ($3.60 \pm 0.43 \text{ mM}$). There were no differences in haematocrit (Fig. 2B, general linear model, $F_{1,13} = 0.69$, $p = 0.42$) or plasma lactate (Fig. 2E, general linear model, $F_{1,14} = 1.48$, $p = 0.24$) between fish sampled under a constant ambient CO_2 regime (haematocrit = $39.4 \pm 0.8\%$, lactate = $0.88 \pm 0.26 \text{ mM}$) or rising CO_2 regime (haematocrit = $38.2 \pm 1.1\%$, lactate = $0.53 \pm 0.12 \text{ mM}$). Blood glucose levels were $\sim 26\%$ lower in fish exposed to a progressively rising CO_2 regime ($4.50 \pm 0.53 \text{ mM}$) when compared to a constant ambient CO_2 regime ($6.09 \pm 0.31 \text{ mM}$) (Fig. 2F, GLM, $F_{1,14} = 6.74$, $p = 0.021$).




Figure 2. Blood chemistry characteristics of European sea bass sampled at ~8.4 kPa O₂ (~40% air saturation) following a progressive O₂ decline accompanied by either constant CO₂ (~650 μatm CO₂, N = 8) or a progressive increase in CO₂ (sampled at ~1770 μatm CO₂, N = 8). Blood pH (A), haematocrit (B), plasma lactate (E) and plasma glucose (F) were directly measured whilst blood pCO₂ (C) and plasma HCO₃⁻ (D) were calculated (see Methods for details). *Indicates statistical significance with $p < 0.05$ and ***indicates statistical significance with $p < 0.001$. Boxes represent median value and inter-quartile range, whiskers represent minimum and maximum values.

Haemoglobin affinity for oxygen. Oxygen affinity of haemoglobin was increased in fish sampled under the progressively rising CO₂ regime (Fig. 3A, GLM, $F_{1,12} = 10.42$, $p = 0.0073$). Haemoglobin P₅₀ was decreased by ~32.5% in fish under the progressively rising CO₂ regime (1.64 ± 0.15 kPa O₂, $7.9 \pm 0.7\%$ air saturation) compared to fish sampled with a constant ambient CO₂ (2.43 ± 0.20 kPa O₂, $11.7 \pm 1\%$ air saturation). There was no significant change in Hills number between treatments (Fig. 3B, GLM, $F_{1,12} = 0.50$, $p = 0.494$).

Discussion

Our results highlight the biological importance of simultaneously rising CO₂ under conditions where O₂ levels in water are depleted – a scenario that reflects the natural conditions during hypoxia which will be exacerbated by climate change – by demonstrating that ecologically relevant changes of CO₂ impact physiological performance of a marine fish at both the molecular and whole organism level. We hypothesised that rising CO₂ during progressive O₂ decreases would lead to an increase in O_{2crit} as a result of increased blood CO₂, decreased blood pH and the associated Bohr/Root effect of fish haemoglobin (Hb), in which Hb-O₂ affinity (Bohr effect) and the total capacity of Hb for O₂ (Root effect) are reduced when pH falls. In contrast, we show that increasing CO₂ as O₂ declined led to enhanced hypoxia tolerance of sea bass with a 20% lower critical oxygen level (Fig. 1). This change in whole organism hypoxic response was accompanied by an increase in Hb-O₂ affinity of blood cells in fish exposed to concurrent CO₂ rises (Fig. 3). The change in Hb-O₂ affinity was not accompanied by a change in blood pH (although the one hour acclimation to treatment pCO₂ prior to blood sampling may have contributed




Figure 3. Haemoglobin P_{50} (A) and Hills number (B) for fish sampled at ~ 8.4 kPa O_2 ($\sim 40\%$ air saturation) following a progressive O_2 decline accompanied by either constant CO_2 (~ 650 $\mu\text{atm } CO_2$, $N = 7$) or a progressive increase in CO_2 (sampled at ~ 1770 $\mu\text{atm } CO_2$, $N = 7$). Measurements were made using a gas mix which matched the calculated blood pCO_2 of each individual fish blood sample. **Indicates statistical significance with $p < 0.01$. Boxes represent median value and inter-quartile range, whiskers represent minimum and maximum values.

to this result). This provides a potential mechanistic basis to explain improved O_{2crit} , enabling sea bass to enhance O_2 uptake during hypoxia and thus maintain normal aerobic metabolism to lower environmental O_2 .

The driver of increased Hb- O_2 affinity in sea bass exposed to concurrent O_2 decline and CO_2 rise is not clear from our results. Several allosteric factors that can modulate the affinity of haemoglobin for O_2 could be involved, including pH, organic phosphates and inorganic ions. Fish haemoglobin is highly sensitive to pH, which modulates Hb- O_2 affinity and carrying capacity via the Bohr and Root effects³¹, but we found no differences in blood pH of sea bass between treatment groups. In addition, the *in-vivo* increase in pCO_2 in fish exposed to concurrent CO_2 rises during hypoxia led to an opposite response of Hb- O_2 affinity than would be expected by an *in-vitro* rise in pCO_2 which would result in pH induced Bohr/Root effects. Increased Hb- O_2 affinity could result from increased intracellular pH of erythrocytes³², as acute hypoxic exposure has been shown to stimulate a β -adrenergic stimulated increase in intracellular erythrocyte pH in rainbow trout³³. Alternatively, increased Hb- O_2 affinity could be due to decreased red cell nucleoside triphosphates (NTPs)^{31,34}, a known hypoxia adaptation, but this can take more than 6 days to complete³⁵. Sea bass may also have a particularly strong β -adrenergic response and/or fast NTP response, although there is little evidence to suggest this because P_{50} measurements from fish exposed to hypoxia at ambient CO_2 levels do not differ from P_{50} measurements in normoxic fish from the same population (Montgomery *et al.*, Unpublished data). It is possible that rising CO_2 during hypoxia may modulate the β -adrenergic response and/or the red cell NTP response within the time frame (~ 4 – 6 hours) of our treatments. However, direct measurements of red cell pH_i and NTP content of sea bass in each treatment group would be needed to confirm this.

A third possible driver of increased Hb- O_2 affinity in sea bass exposed to concurrent CO_2 rise during hypoxia could be decreased erythrocyte chloride (Cl^-)³⁶. Although erythrocyte Cl^- was not directly measured in our study, plasma HCO_3^- was approximately 3 mM higher in fish exposed to concurrent CO_2 rises during hypoxia than fish which experienced constant ambient CO_2 during hypoxia (Fig. 2D). The higher HCO_3^- in fish exposed to rising ambient CO_2 during hypoxia is likely a result of rapid compensation for a respiratory acidosis due to rising blood pCO_2 . This change in plasma HCO_3^- is typically mirrored by a reciprocal change in plasma Cl^- ^{37,38} which is likely to be followed by a similar decline in erythrocyte Cl^- .

Analysis of O_{2crit} is a common measure of hypoxia tolerance in fish but concurrent CO_2 increases during hypoxia have been generally unaccounted for. A recent meta-analysis by Rogers *et al.*³⁹ constructed a database of O_{2crit} research of fish (both freshwater and marine). This analysis identified two broad methods employed in O_{2crit} measurements:

1. Closed respirometry where O_2 is usually reduced by the O_2 consumption of the fish (52 identified studies) or;
2. Intermittent or flow-through respirometry in which O_2 is usually reduced via gassing with pure N_2 or combined N_2 and O_2 mixes (32 identified studies).

The use of closed respirometry in the majority of studies would result in concurrent CO_2 rises as O_2 is depleted by fish O_2 consumption. The increase in ambient CO_2 during closed respirometry is well known and often used as a criticism of this respirometry technique⁴⁰. In contrast, use of intermittent-flow respirometry in O_{2crit} trials normally necessitates the reduction of O_2 in the water by aeration with N_2 or a mix of N_2 & O_2 . As a result CO_2 would likely decrease during the time course of hypoxia induction (as the gas mixture would contain zero CO_2 , rather than ~ 400 μatm present in atmospheric air). Such a change in CO_2 during the O_{2crit} trial would be the opposite of that seen in nature. Therefore it may be considered that closed respirometry provides conditions which give a more environmentally relevant measure of O_{2crit} ²⁸.

As our results indicate that rising CO_2 during hypoxia directly affects the ability of sea bass to maintain O_2 uptake, it could be expected that the use of closed respirometry methods would result in lower measurements

of O_{2crit} than intermittent-flow methods for the same species. This effect has not been documented for species in which a direct comparison has been made – with either there being no effect of respirometry method on O_{2crit} ^{39,41} or higher O_{2crit} measurements when closed respirometry is used⁴¹. However, such comparisons are complicated by differences in the rate of hypoxia induction (RHI) by different studies, which in turn will influence how much time fish have to regulate blood pH when ambient CO_2 is rising. For example, Regan and Richards⁴¹ have demonstrated that the faster rates of hypoxia induction (RHI) typical of closed respirometry O_{2crit} trials lead to higher values of O_{2crit} (i.e. lower hypoxia tolerance) when compared with longer trials using slower RHI's typical of the intermittent-flow method. The effect of RHI on O_{2crit} was proposed by Regan and Richards⁴⁰ as a potential explanation of the results of Snyder *et al.* (i.e. higher O_{2crit} in closed respirometry compared to intermittent-flow respirometry)⁴². The speed of RHI during closed respirometry will also effect the speed of CO_2 rise. Almost all studies using closed respirometry to measure O_{2crit} do not report changes in CO_2 over the course of measurement period. When accumulation of CO_2 during a closed respirometry O_{2crit} trial was measured by Regan and Richards, CO_2 levels were $\sim 8,000 \mu atm$ after ~ 90 minutes⁴¹. However, we should note that this pCO_2 level was measured after the O_{2crit} point, when anaerobic metabolism continues to produce CO_2 in the absence of O_2 consumption, but also metabolic acid production and excretion further drives up water pCO_2 in the respirometer due to excess H^+ ions titrating ambient HCO_3^- to CO_2 . Regardless, at O_2 levels above O_{2crit} the rate of CO_2 onset will be faster than those used in our current study as a result of the faster RHI. Increased speed of CO_2 onset in closed respirometry trials may 'outstrip' the ability of fish to acid-base regulate, causing an uncompensated respiratory acidosis during the time of the trial, which in turn would decrease Hb- O_2 affinity via the Bohr & Root effects and potentially increase O_{2crit} . Similarly, fish species which have reduced ability to acid-base regulate may have an increased O_{2crit} when rising CO_2 is included in trials.

Our results indicate improved hypoxia tolerance during rising CO_2 in European sea bass. This contrasts with previous research investigating interactive effects of CO_2 and hypoxia on O_{2crit} of fish. Woolly sculpin, *Clinocottus analis* (an intertidal species that can breathe air), exposed to $\sim 1100 \mu atm$ CO_2 showed no impact on O_{2crit} after 7 days acclimation but after 28 days had O_{2crit} measurements $\sim 34\%$ higher than fish held in ambient ($\sim 400 \mu atm$) conditions⁴³. Higher O_{2crit} after 28 days corresponded with higher RMR and Na^+ , K^+ , ATPase activity. This contrast in results could indicate that the beneficial effect of acute rises in CO_2 associated with natural hypoxia documented in our study are potentially reversed when fish are exposed to long term constantly high CO_2 associated with anthropogenic climate change. Moreover, acute changes in CO_2 had no effect on O_{2crit} of the estuarine fish species mummichog, *Fundulus heteroclitus*, and Norfolk spot, *Leiostomus xanthurus* when they were exposed to $\sim 8,000$ – $10,000 \mu atm$ CO_2 immediately prior to an O_{2crit} trial⁴⁴. As such the effect of CO_2 on O_{2crit} will likely depend on differences in physiological responses to CO_2 and O_2 between species.

Simultaneously rising CO_2 also shows variable impacts on non-metabolic responses to hypoxia of several species. Cycling CO_2 had no effect on aquatic surface respiration (ASR), the use of the thin surface layer of water for aquatic respiration⁴⁵, or survival in juvenile *Menidia menidia*, *Fundulus majalis*, *Fundulus heteroclitus* or *Morone saxatilis* exposed to short term cycles of O_2 ⁴⁶. In contrast, combined hypoxia and acidification resulted in an increase in the O_2 level at which *Menidia menidia* and *Menidia beryllina* first performed ASR, consistently performed ASR, exhibited LoE, and finally died²¹. Additionally, combined high CO_2 ($\sim 2,000 \mu atm$) and hypoxia had no effect on survival of larval *Cyprinodon variegatus*, an additive negative effect on larval *Menidia beryllina*, and a synergistic negative effect on larval *Menidia menidia*⁴⁷. This variation in effect of CO_2 on hypoxia responses could be a result of methodological differences (e.g. constant high CO_2 in Dixon *et al.*⁴⁶, cycling DO/pH in DePasquale *et al.*⁴⁷, and concurrent CO_2 rise/ O_2 decrease in Miller *et al.*²¹), the level of CO_2 used in studies (e.g. CO_2 levels used by Miller *et al.*²¹ were $\sim 23,000 \mu atm$ which is much higher than levels likely to be commonly found in the environment during hypoxia and may have contributed to the negative effects of rising CO_2 noted in the study), differences in species and life stages used (changes in physiological tolerance across life stages have been noted for thermal tolerance by Komoroske *et al.*⁴⁸), or possibly variability in response as a result of differences in previous environmental experience⁴⁹. The role of environmental variability in species sensitivities to CO_2 has recently been outlined in the proposed Ocean Variability Hypothesis (OVH)⁵⁰ and warrants testing on various species in the future.

Overall our results indicate that the environmentally realistic, simultaneous rises in CO_2 during a hypoxic event increased the hypoxia tolerance (i.e. reduces O_{2crit}) of European sea bass which is at least partly explained by an enhanced ability of fish to uptake O_2 via increased Hb- O_2 affinity. Miller *et al.*²¹ also demonstrated impacts of concurrent CO_2 rise on measurements of hypoxia tolerance, although in an opposite direction to that noted in our study. As concurrent CO_2 rises during hypoxia are the norm in nature, evidence that this affects physiology of organisms exposed to hypoxia highlights an important shortcoming of research to predict tolerances to hypoxia of fish. More research on this issue is needed to clarify how common this modifying effect of CO_2 on the response to hypoxia is and whether such measurements in the lab are ecologically relevant. A greater understanding of this issue may allow more accurate assessments of the impacts of hypoxic events on marine fish in nature, aiding management and conservation of fish species. With specific regard to measurements of O_{2crit} we believe future studies should include concurrent rising CO_2 in the following ways:

1. Intermittent-flow respirometer studies should include increases of CO_2 relevant to hypoxic events that organisms may experience,
2. Closed respirometry studies should report the start and end CO_2 levels in the respirometer.

In addition, both methods should aim to create an environmentally relevant rate of hypoxia induction/ CO_2 increase for the species studied, and consistently report CO_2 levels measured. By incorporating these recommendations we believe that future studies of O_{2crit} will give more representative estimations of species hypoxia tolerance.

Time held in system (days)	Temperature (°C)	pH	Salinity	Total Alkalinity (mM/kgSW)	pCO ₂ (µatm)
318	18.01 ± 0.03	8.04 ± 0.04	33.25 ± 0.70	2064.6 ± 136.0	516.9 ± 41.1

Table 1. Water chemistry parameters of the recirculating aquaculture system in which sea bass were held prior to experimental work (means ± S.E. shown).

Materials and Methods

Fish collection and husbandry. We collected juvenile sea bass from estuaries and coastal lagoons on the south Dorset coast and Isle of Wight in June 2017 (Marine Management Organisation permit #030/17 & Natural England permit #OLD1002654). Prior to experimentation, these fish were held in the University of Exeter's Aquatic Resource Centre in an aerated recirculating aquaculture system and fed a commercial pellet at a ration of ~1–2% body weight per day three times a week (for system water chemistry see Table 1). All fish were starved for a minimum of 72 hours prior to the start of all measurements to ensure their metabolism was not affected by digestion (i.e. specific dynamic action²⁴). All experimental procedures were carried out under home office licence P88687E07 and approved by the University of Exeter's Animal Welfare and Ethical Review Board.

Measuring hypoxia tolerance. We determined oxygen consumption rates ($\dot{M}O_2$) of sea bass using an intermittent-flow respirometer system. The respirometer system set up followed recommendations set out by Svendsen *et al.*⁵¹. Briefly, the system comprised of a sealed 4.515 L respirometer chamber connected to a recirculating loop, including an in-line recirculating pump (Eheim universal 600, Deizisau, Germany), and a measurement chamber into which a temperature-compensated fibre optic oxygen optode (Firesting O₂ oxygen meter, Pyroscience GmbH, Germany) was placed. Oxygen optodes were calibrated in water at the start of experiments at 100% air saturation and 0% air saturation according to manual instructions. Respirometry was conducted in a semi-closed system consisting of three 100 L experimental tanks fed by a 100 L sump, with overflowing water from the experimental tanks recirculating back to the sump. A second pump was used to periodically flush the respirometer system with water from the surrounding tank. This pump was controlled by an automated computer program (AquaResp 3, AquaResp[®]) to intermittently flush the respirometer. Five respirometer chambers were distributed between the three experimental tanks (maximum of 2 chambers per experimental 100 L tank). The sump was temperature controlled (18.27 ± 0.02 °C, mean ± S.E.) using a heater/chiller unit (Grant TX150 R2, Grant Instruments, Cambridge, UK) attached to a temperature exchange coil. Together these tanks formed a 400 L system with the same temperature, oxygen and water chemistry parameters for all respirometers.

Individual sea bass (average mass = 131.2 ± 7.5 g), chosen at random, were placed inside the respirometers and allowed an overnight recovery period, for a minimum of 13 hours, before O_{2crit} tests began. While sea bass were in the respirometers measurements of $\dot{M}O_2$ were conducted every 10 minutes, including a flush period of 300 s, a wait period of 60 s and a measurement period of 240 s. During the wait and measurement period the chamber was sealed by switching off the flush pump and the decline in dissolved O₂ within the chamber was continuously measured by the fibre optic O₂ electrode.

Following the overnight recovery period a standard O_{2crit} test was conducted. Oxygen levels in the respirometer system were reduced from ~100% air saturation to ~15% air saturation over the course of 6 hours (decline in O₂ was ~20% air saturation per hour between ~100% air saturation and ~40% air saturation and ~10% air saturation per hour from ~40% air saturation until the end of trials). Oxygen was regulated by gassing the sump and experimental tanks with a mix of N₂ and O₂ (G400 Gas mixing system, Qubit Biology Inc.) at a rate of 10 L min⁻¹ following a pre-set automated protocol (Flowvision, Alicat software). Levels of CO₂ within the system were controlled under one of two treatments (with 8 fish exposed to Treatment 1 and 8 separate fish exposed Treatment 2):

- Treatment 1 (Constant CO₂) - ambient levels of CO₂ were maintained by including 0.04% CO₂ as part of the gas mix delivered to the respirometer system.
- Treatment 2 (Rising CO₂) - the proportion of CO₂ in the gas mix was gradually increased as O₂ was decreased. This increase in CO₂ was designed to reflect environmentally realistic increases in CO₂ predicted as a result of depletion of O₂ by bacterial respiration (assumed respiratory quotient of 1)⁵², using the seawater carbonate chemistry calculator CO2sys (see supplementary material for predictions of increased CO₂ during hypoxia).

Water chemistry of treatment 2 was monitored once per hour by measuring pH_{NBS}, temperature and salinity as well as taking a 12 mL water sample to measure total CO₂ (TCO₂)/Dissolved Inorganic Carbon (DIC). Water chemistry of treatment 1 was monitored at the start and end of the treatment to ensure no change in water pCO₂ occurred over the time course of the O_{2crit} trial. Seawater DIC analysis was conducted using a custom built system described in detail by Lewis *et al.*⁵³. These four parameters were then input into the seawater carbon calculator programme, CO2SYS to calculate pCO₂ based on the NBS pH scale, equilibration constants from Mehrbach *et al.* refitted by Dickson and Millero, and KSO₄ dissociation constants from Dickson. The reciprocal changes in O₂ and CO₂ during O_{2crit} tests for each treatment are illustrated in Fig. 4. O_{2crit} tests were stopped once a minimum of 3 $\dot{M}O_2$ measurements showed a transition from an oxy-regulating to oxy-conforming state for each fish or fish showed a large drop in $\dot{M}O_2$ and signs of distress in the respirometer. No fish exhibited LoE during trials. Following completion of O_{2crit} trials experimental tanks were aerated with ambient air to swiftly restore O₂ and CO₂ levels.

Sea bass were left to recover in respirometers, for a minimum of 1 hour post-trial, until O₂ levels reached ~21 kPa O₂ (~100% air saturation). Fish were then removed from respirometers and background respiration was measured for a minimum of 1 hour (6 measurement cycles) for all respirometers immediately post trial.




Figure 4. Changes in partial pressure of O₂ (expressed as kPa O₂) and CO₂ (µatm) during (A) two O_{2crit} trials representing treatment 1 in which O₂ was reduced with no change in CO₂; and (B) two trials representing treatment 2 where O₂ was reduced with a corresponding rise in CO₂. Data presented are means ± S.D.

Oxygen consumption rate ($\dot{M}O_2$) analysis. Following each 240 s measurement period $\dot{M}O_2$ was automatically calculated by the AquaResp3 software. A linear regression was fitted to the O₂ versus time data for each measurement period. The slope of this regression (s, kPa O₂ h⁻¹) was then used to calculate $\dot{M}O_2$ (mg O₂ kg⁻¹ h⁻¹) using the equation outlined by Svendsen *et al.*⁵¹:

$$\dot{M}O_2 = sV_{resp}\alpha m^{-1}$$

where V_{resp} is the respirometer volume minus the volume of the fish (L), α is the solubility of O₂ in water (mgO₂ L⁻¹ kPa⁻¹) for the relevant salinity and temperature, and m is the mass of the fish (kg). Calculations of $\dot{M}O_2$ where s had a R² of <0.98 were removed from subsequent analysis. For the purpose of establishing O_{2crit} values from a plot of $\dot{M}O_2$ versus ambient O₂ level, the oxygen saturation of each measurement period was defined as the average of the dissolved O₂ measurement over the measurement period. The average background respiration, over the 1 hour post-trial measuring period, for each respirometer (average background respiration was <2% of fish $\dot{M}O_2$) was subtracted from $\dot{M}O_2$ measurements to correct fish $\dot{M}O_2$ for background respiration.

We calculated the standard metabolic rate (SMR) in R v3.5.3⁵⁴ using function calcSMR in package fish $\dot{M}O_2$ ²⁴. All $\dot{M}O_2$ values from the overnight recovery period and beginning of the O_{2crit} trial in which average dissolved O₂ saturation was >80% air saturation were included for SMR calculations. This resulted in approximately 14–16 hours of $\dot{M}O_2$ data to calculate SMR from for each fish. We estimated SMR for all fish using the mean of the lowest 10 $\dot{M}O_2$ measurements during the overnight period. Although Chabot *et al.*²⁴ recommend use of the mean of the lowest normal distribution (MLND) or quantile (q = 0.2) methods to calculate SMR we chose the mean of the 10 lowest values as it produced a value of SMR that most accurately matched the consistent low $\dot{M}O_2$ measurements of oxy-regulating fish at values of pO_2 above the O_{2crit} point. Additionally, for each fish the coefficient of variation in the mean lowest normal distribution (MLND) was assessed using a ROUT test (Q = 1%) to check whether variation in low $\dot{M}O_2$ measurements was consistent between fish (to account for potential differences in activity in the respirometer prior to O_{2crit} trials). The ROUT test removed one fish from the rising CO₂ treatment which displayed a CoV of 34.6. Mean CoV of the remaining 15 fish was 11.85 ± 1.26 (±S.E.M).

We then used function calcO2crit from the package to calculate the O_{2crit} for each individual fish, using the estimated SMR of each individual, as detailed in the supplementary material of Claireaux & Chabot⁵⁵. This function identifies the portion of the O_{2crit} test where metabolic rate data follows an oxygen conforming relationship and fits a linear regression line through this data, O_{2crit} is then calculated as the oxygen level at which this regression line crosses the calculated SMR of the individual fish. Plots of $\dot{M}O_2$ against O₂ during O_{2crit} trials showing calculated O_{2crit} for each individual can be seen in the supplementary material (Supplementary Material Figs 1–16). Calculations were conducted using a gap limit of 0.83 kPa O₂ (4% air saturation) and a maximum number of 7 $\dot{M}O_2$ points to fit the regression line through the oxygen conforming component of the data used to estimate O_{2crit}.

Measuring blood chemistry. Following O_{2crit} trials sea bass were moved to individual 7 L chambers, which were aerated and supplied with seawater from the aquarium re-circulating system at a rate of about 4 L min⁻¹.

After an overnight acclimation period we then exposed the fish to a decrease in O₂ levels to ~6.4 kPa O₂ (30% air saturation) over a period of 4 hours (equivalent to the rate of decrease in O₂ used in the previous O_{2crit} tests). We chose to blood sample fish at an O₂ level above O_{2crit} to ensure that anaerobic metabolism did not influence blood chemistry. This was combined with the same CO₂ regime each fish experienced during the O_{2crit} test (i.e. either constant or reciprocally rising CO₂). Once an O₂ level of ~6.4 kPa O₂ (30% air saturation) was achieved the fish were allowed to acclimate for 1 hour before being individually anaesthetised *in situ* using a dose of 100 mgL⁻¹ of benzocaine. Once fish were judged to be sufficiently anaesthetised (cessation of gill ventilation and lack of response to a pinch of the anal fin) they were immediately transferred within 5 seconds to a gill irrigation table (with the same pO₂ and pCO₂ levels of the respective treatment), where anaesthesia was maintained with a dose of 37.5 mgL⁻¹ of benzocaine. Gill ventilation was artificially maintained by a micro-pump, so that the operculum were just open and exhalant water flow could just be visualised. Once a stable gill water flow was established blood was sampled by caudal vessel puncture using a 1 ml heparinised syringe. This method has been demonstrated to obtain accurate measurements of blood chemistry parameters comparable to those achieved using cannulation (Davison & Wilson, University of Exeter, personal communication). At the time of blood sampling water pCO₂ was 656 ± 44 µatm (mean ± S.E.) for fish in the constant ambient CO₂ regime and 1763 ± 43 µatm (mean ± S.E.) for fish in the progressively rising CO₂ regime. Water pO₂ was 8.1 ± 0.2 kPa (mean ± S.E., ~38% air saturation) for fish in the constant ambient CO₂ regime and 8.7 ± 0.2 kPa (mean ± S.E., ~41% air saturation) for fish in the progressively rising CO₂ regime. Following blood sampling fish were transferred to seawater isolation tanks containing ~20.8 kPa O₂ (100% air-saturated) to recover from the anaesthetic. They were then monitored over a period of 24 hours before we returned them to their original holding tanks.

Immediately after sampling, whole blood pO₂ was measured at 18 °C in a temperature-controlled system (Strathkelvin 1302 electrode and 781 meter; Strathkelvin Instruments Ltd, Glasgow, UK). We measured extracellular pH on 30 µL of whole blood using an Accumet Micro pH electrode and Hanna HI8314 pH meter at 18 °C calibrated to pH_{NBS} 7.04 and 9.21 specific buffers. Three 75 µL micro capillary tubes were then filled with whole blood and sealed with Critoseal capillary tube sealant (Fisher) and paraffin oil and centrifuged for 2 minutes at 10,000 rpm. Haematocrit was measured using a Hawksley micro-haematocrit reader. Plasma was extracted from these tubes for analysis of TCO₂ using a Mettler Toledo 965D carbon dioxide analyser. Plasma pCO₂ and HCO₃⁻ were then calculated from TCO₂, temperature and blood pH using the Henderson-Hasselbalch equation with values for solubility and pK_{app}¹ based on Boutilier *et al.*^{56,57}. Haemoglobin content of the blood was also assessed by the cyanmethemoglobin method (using Drabkin's reagent, Sigma). Half the remaining whole blood was then centrifuged at 10,000 rpm for 2 minutes at 4 °C. The resulting plasma was separated and snap frozen in liquid nitrogen and stored at -80 °C before later being used to measure plasma glucose and lactate using a YSI 2900D Biochemistry Analyzer (Xylem Analytics, UK). All measurements or storage of blood occurred within 10 minutes of blood sampling.

Measuring Hb-oxygen binding. We measured Hb-O₂ affinity using a Blood Oxygen Binding System (BOBS, Loligo systems), detailed in Oellermann *et al.*⁵⁸. A sample of the same whole blood used for blood chemistry measurements was diluted at a ratio of 1:4 in its own plasma. 1 µL of this blood was then used for measurements. The BOBS exposed this blood sample to gas mixes with a progressive increase in O₂ whilst measuring absorbance of light across a spectrum ranging from 200 to 800 nm. For each individual fish the gas mix that blood was exposed to matched the calculated pCO₂ of the blood sample. The change in absorption of light at a wavelength of 435 nm was used to assess changes in oxygenation of Hb, as previously used by Verhille & Farrell⁵⁹. Background changes in absorption of the blood sample were corrected using the isosbestic wavelength of 390 nm⁵⁹. Following measurements the BOBS calculated the oxygen equilibrium curve of the sample using Hill's formula before estimating P₅₀⁵⁸.

Statistical analysis. We conducted all statistical analysis in R v3.5.3⁵⁴. There was no significant difference in mass of fish between the treatment groups (One-Way ANOVA, F_{1,13} = 2.821, p = 0.117) or of estimated SMR of fish between treatment groups (Unpaired t-test, t = 0.8455, d.f. = 13, p = 0.413). There was, however, a significant correlation between estimated SMR and calculated O_{2crit} for all fish (Pearson's correlation, df = 13, t = 3.32, R = 0.68, p = 0.0056). As such the effect of CO₂ treatment on O_{2crit} was assessed using an ANCOVA with SMR as a covariate. All other analyses were conducted using general linear modelling (GLM). All values in the text are reported as mean ± standard error (S.E.).

Data availability

Data is available via the University of Exeter's online repository at: <https://doi.org/10.24378/exe.1523>.

Received: 3 July 2019; Accepted: 25 September 2019;

Published online: 22 October 2019

References

1. Diaz, R. J. Overview of Hypoxia around the World. *J. Environ. Qual.* **30**, 275–281 (2001).
2. Breitburg, D. *et al.* Declining oxygen in the global ocean and coastal waters. *Science* (80-). **359** (2018).
3. Diaz, R. J. & Rosenberg, R. Spreading Dead Zones and Consequences for Marine Ecosystems. *Science* (80-). **321**, 926 (2008).
4. Townhill, B. L. *et al.* Consequences of climate-induced low oxygen conditions for commercially important fish. *Mar. Ecol. Prog. Ser.* **580**, 191–204 (2017).
5. Robinson, C. Microbial Respiration, the Engine of Ocean Deoxygenation. *Frontiers in Marine Science.* **5**, 533 (2019).
6. Rabalais, N. N., Turner, R. E., Díaz, R. J. & Justic, D. Global change and eutrophication of coastal waters. *ICES J. Mar. Sci.* **66**, 1528–1537 (2009).

7. Sunda, W. G. & Cai, W.-J. Eutrophication Induced CO₂-Acidification of Subsurface Coastal Waters: Interactive Effects of Temperature, Salinity, and Atmospheric PCO₂. *Environ. Sci. Technol.* **46**, 10651–10659 (2012).
8. Wallace, R. B., Baumann, H., Grear, J. S., Aller, R. C. & Gobler, C. J. Coastal ocean acidification: The other eutrophication problem. *Estuar. Coast. Shelf Sci.* **148**, 1–13 (2014).
9. Gobler, C. J. & Baumann, H. Hypoxia and acidification in ocean ecosystems: coupled dynamics and effects on marine life. *Biol. Lett.* **12** (2016).
10. Altieri, A. H. & Gedan, K. B. Climate change and dead zones. *Glob. Chang. Biol.* **21**, 1395–1406 (2015).
11. Oschlies, A., Brandt, P., Stramma, L. & Schmidtko, S. Drivers and mechanisms of ocean deoxygenation. *Nat. Geosci.* **11**, 467–473 (2018).
12. Oschlies, A., Schulz, K. G., Riebesell, U. & Schmittner, A. Simulated 21st century's increase in oceanic suboxia by CO₂-enhanced biotic carbon export. *Global Biogeochem. Cycles* **22** (2008).
13. Doney, S. C., Fabry, V. J., Feely, R. A. & Kleypas, J. A. Ocean Acidification: The Other CO₂ Problem. *Ann. Rev. Mar. Sci.* **1**, 169–192 (2009).
14. Melzner, F. *et al.* Future ocean acidification will be amplified by hypoxia in coastal habitats. *Mar. Biol.* **160**, 1875–1888 (2013).
15. Brandt, S. B., Gerken, M., Hartman, K. J. & Demers, E. Effects of hypoxia on food consumption and growth of juvenile striped bass (*Morone saxatilis*). *J. Exp. Mar. Bio. Ecol.* **381**(Suppl), S143–S149 (2009).
16. Almeida, L. Z., Guffey, S. C., Sepúlveda, M. S. & Höök, T. O. Behavioral and physiological responses of yellow perch (*Perca flavescens*) to moderate hypoxia. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* **209**, 47–55 (2017).
17. Chabot, D. & Dutil, J. D. Reduced growth of Atlantic cod in non-lethal hypoxic conditions. *J. Fish Biol.* **55**, 472–491 (1999).
18. Domenici, P., Steffensen, J. F. & Batty, R. S. The effect of progressive hypoxia on swimming activity and schooling in Atlantic herring. *J. Fish Biol.* **57**, 1526–1538 (2000).
19. Gobler, C. J., DePasquale, E. L., Griffith, A. W. & Baumann, H. Hypoxia and Acidification Have Additive and Synergistic Negative Effects on the Growth, Survival, and Metamorphosis of Early Life Stage Bivalves. *PLoS One* **9**, e83648 (2014).
20. Steckbauer, A. *et al.* Synergistic effects of hypoxia and increasing CO₂ on benthic invertebrates of the central Chilean coast. *Front. Mar. Sci.* **2**, 49 (2015).
21. Miller, S. H., Breitburg, D. L., Burrell, R. B. & Keppel, A. G. Acidification increases sensitivity to hypoxia in important forage fishes. *Mar. Ecol. Prog. Ser.* **549**, 1–8 (2016).
22. Nikinmaa, M. Oxygen and carbon dioxide transport in vertebrate erythrocytes: an evolutionary change in the role of membrane transport. *J. Exp. Biol.* **200**, 369–380 (1997).
23. Rummer, J. L. & Brauner, C. J. Root effect haemoglobins in fish may greatly enhance general oxygen delivery relative to other vertebrates. *PLoS One* **10**, e0139477 (2015).
24. Chabot, D., Steffensen, J. F. & Farrell, A. P. The determination of standard metabolic rate in fishes. *J. Fish Biol.* **88**, 81–121 (2016).
25. Perry, S. F., Jonz, M. G. & Gilmour, K. M. Chapter 5 Oxygen Sensing And The Hypoxic Ventilatory Response. In *Fish Physiology* (eds J. G. Richards, A. P. F. and C. J. B.) Volume 27, 193–253 (Academic Press, 2009).
26. Wells, R. M. G. Chapter 6 Blood-Gas Transport and Hemoglobin Function: Adaptations for Functional and Environmental Hypoxia. In *Fish Physiology* (eds Jeffrey G. Richards, A. P. F. and C. J. B.) Volume 27, 255–299 (Academic Press, 2009).
27. Gamperl, A. K. & Driedzic, W. R. Chapter 7 Cardiovascular Function and Cardiac Metabolism. In *Hypoxia* (eds Richards, J. G., Farrell, A. P. & Brauner, C. J. B. T.-F. P.) **27**, 301–360 (Academic Press, 2009).
28. Wood, C. M. The fallacy of the P_{crit} – are there more useful alternatives? *J. Exp. Biol.* **221**, jeb163717 (2018).
29. Speers-Roesch, B., Mandic, M., Groom, D. J. E. & Richards, J. G. Critical oxygen tensions as predictors of hypoxia tolerance and tissue metabolic responses during hypoxia exposure in fishes. *J. Exp. Mar. Bio. Ecol.* **449**, 239–249 (2013).
30. Regan, M. D. *et al.* Don't throw the fish out with the respirometry water. *J. Exp. Biol.* **222**, jeb200253 (2019).
31. Pelster, B. & Weber, R. E. The Physiology of the Root Effect. In *Advances in Comparative and Environmental Physiology* 51–77, https://doi.org/10.1007/978-3-642-75900-0_2LB-Pelster1991 (Springer Berlin Heidelberg, 1991).
32. Nikinmaa, M. Adrenergic regulation of haemoglobin oxygen affinity in rainbow trout red cells. *J. Comp. Physiol.* **152**, 67–72 (1983).
33. Tetens, V. & Christensen, N. J. Beta-adrenergic control of blood oxygen affinity in acutely hypoxia exposed rainbow trout. *J. Comp. Physiol. B* **157**, 667–675 (1987).
34. Weber, R. E. & Lykkeboe, G. Respiratory adaptations in carp blood influences of hypoxia, red cell organic phosphates, divalent cations and CO₂ on hemoglobin-oxygen affinity. *J. Comp. Physiol.* **128**, 127–137 (1978).
35. Soivio, A., Nikinmaa, M. & Westman, K. The blood oxygen binding properties of hypoxic *Salmo gairdneri*. *J. Comp. Physiol.* **136**, 83–87 (1980).
36. Mairböurl, H. & Weber, R. E. Oxygen Transport by Hemoglobin. *Comprehensive Physiology*, <https://doi.org/10.1002/cphy.c080113> (2012).
37. Esbaugh, A. J., Heuer, R. & Grosell, M. Impacts of ocean acidification on respiratory gas exchange and acid-base balance in a marine teleost, *Opsanus beta*. *J. Comp. Physiol. B* **182**, 921–934 (2012).
38. Perry, S. F., Goss, G. G. & Laurent, P. The interrelationships between gill chloride cell morphology and ionic uptake in four freshwater teleosts. *Can. J. Zool.* **70**, 1775–1786 (1992).
39. Rogers, N. J., Urbina, M. A., Reardon, E. E., McKenzie, D. J. & Wilson, R. W. A new analysis of hypoxia tolerance in fishes using a database of critical oxygen level (P_{crit}). *Conserv. Physiol.* **4** (2016).
40. Steffensen, J. F. Some errors in respirometry of aquatic breathers: How to avoid and correct for them. *Fish Physiol. Biochem.* **6**, 49–59 (1989).
41. Regan, M. D. & Richards, J. G. Rates of hypoxia induction alter mechanisms of O₂ tension of goldfish. *J. Exp. Biol.* **220**, 2536 LP–2544 (2017).
42. Snyder, S. *et al.* Effect of closed v. intermittent-flow respirometry on hypoxia tolerance in the shiner perch *Cymatogaster aggregata*. *J. Fish Biol.* **88**, 252–264 (2016).
43. Hancock, J. R. & Place, S. P. Impact of ocean acidification on the hypoxia tolerance of the woolly sculpin, *Clinocottus analis*. *Conserv. Physiol.* **4**, cow040–cow040 (2016).
44. Cochran, R. E. & Burnett, L. E. Respiratory responses of the salt marsh animals, *Fundulus heteroclitus*, *Leiostomus xanthurus*, and *Palaemonetes pugio* to environmental hypoxia and hypercapnia and to the organophosphate pesticide, azinphosmethyl. *J. Exp. Mar. Bio. Ecol.* **195**, 125–144 (1996).
45. Kramer, D. L. & McClure, M. Aquatic surface respiration, a widespread adaptation to hypoxia in tropical freshwater fishes. *Environ. Biol. Fishes* **7**, 47–55 (1982).
46. Dixon, R. L., Greco, P. A. & Targett, T. E. Responses of juvenile Atlantic silverside, striped killifish, mummichog, and striped bass to acute hypoxia and acidification: Aquatic surface respiration and survival. *J. Exp. Mar. Bio. Ecol.* **493**, 20–30 (2017).
47. DePasquale, E., Baumann, H. & Gobler, C. J. Vulnerability of early life stage Northwest Atlantic forage fish to ocean acidification and low oxygen. *Mar. Ecol. Prog. Ser.* **523**, 145–156 (2015).
48. Komoroske, L. M. *et al.* Ontogeny influences sensitivity to climate change stressors in an endangered fish. *Conserv. Physiol.* **2** (2014).
49. Murray, C. S., Malvezzi, A., Gobler, C. J. & Baumann, H. Offspring sensitivity to ocean acidification changes seasonally in a coastal marine fish. *Mar. Ecol. Prog. Ser.* **504**, 1–11 (2014).
50. Baumann, H. Experimental assessments of marine species sensitivities to ocean acidification and co-stressors: how far have we come? *Can. J. Zool.* 399–408, <https://doi.org/10.1139/cjz-2018-0198> (2019).

51. Svendsen, M. B. S., Bushnell, P. G. & Steffensen, J. F. Design and setup of intermittent-flow respirometry system for aquatic organisms. *J. Fish Biol.* **88**, 26–50 (2016).
52. Del Giorgio, P. A. & Duarte, C. M. Respiration in the open ocean. *Nature* **420**, 379–384 (2002).
53. Lewis, C., Clemow, K. & Holt, W. V. Metal contamination increases the sensitivity of larvae but not gametes to ocean acidification in the polychaete *Pomatoscerus lamarckii* (Quatrefages). *Mar. Biol.* **160**, 2089–2101 (2013).
54. R Core Team. R: A language and environment for statistical computing (2019).
55. Claireaux, G. & Chabot, D. Responses by fishes to environmental hypoxia: integration through Fry's concept of aerobic metabolic scope. *J. Fish Biol.* **88**, 232–251 (2016).
56. Boutilier, R. G., Heming, T. A. & Iwama, G. K. Appendix-physicochemical parameters for use in fish respiratory physiology. *Fish Physiol.* **10**, 403–430 (1984).
57. Boutilier, R. G., Iwama, G. K., Heming, T. A. & Randall, D. J. The apparent pK of carbonic acid in rainbow trout blood plasma between 5 and 15 °C. *Respir. Physiol.* **61**, 237–254 (2016).
58. Oellermann, M., Pörtner, H.-O. & Mark, F. C. Simultaneous high-resolution pH and spectrophotometric recordings of oxygen binding in blood microvolumes. *J. Exp. Biol.* **217**, 1430 LP–1436 (2014).
59. Verhille, C. & Farrell, A. P. The *in vitro* blood-O₂ affinity of triploid rainbow trout *Oncorhynchus mykiss* at different temperatures and CO₂ tensions. *J. Fish Biol.* **81**, 1124–1132 (2012).

Acknowledgements

This work was supported by a NERC GW4+ Doctoral Training Partnership studentship from the Natural Environment Research Council [NE/L002434/1] with additional funding from CASE partner, The Centre of Fisheries and Aquaculture Science (Cefas) to D.W.M., and from the Biotechnology and Biological Sciences Research Council (BB/D005108/1 and BB/J00913X/1) and NERC (NE/H017402/1) to R.W.W. The authors would like to thank the Marine Management Organisation and Natural England for granting permits to collect wild sea bass for use in this study, Simon Pengelly and the Southern Inshore Fisheries and Conservation Authority for assistance with fish collections. We would also like to thank the aquarium staff, particularly Steven Cooper, Alice Walpole and Rebecca Turner, of the Aquatic Resource Centre at Exeter University for assistance with fish husbandry and system water chemistry, and finally Will Davison, Dr. Cosima Porteus and Harriet Goodrich for assistance with blood chemistry measurements.

Author contributions

D.W.M. contributed to study concept and design, conducted all data collection, analysed all data and drafted the manuscript. S.D.S. contributed to study concept and design, and revisions of the manuscript. G.H.E. and S.N.R.B. contributed to study concept and design, data analysis and revisions of the manuscript. R.W.W. was responsible for study supervision, contributed to study concept and design, assisted with data collection, helped with interpretation of data, and contributed to revisions of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-019-51572-4>.

Correspondence and requests for materials should be addressed to D.W.M. or R.W.W.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019