

THE CUSPIDALISATION OF SECTIONS OF ARITHMETIC FUNDAMENTAL GROUPS II

MOHAMED SAÏDI

ABSTRACT. In this paper, which is a sequel to [Saïdi], we investigate the theory of cuspidalisation of sections of arithmetic fundamental groups of hyperbolic curves to cuspidally i -th and $2/p$ -th step prosolvable arithmetic fundamental groups. As a consequence we exhibit two, necessary and sufficient, conditions for sections of arithmetic fundamental groups of hyperbolic curves over p -adic local fields to arise from rational points. We also exhibit a class of sections of arithmetic fundamental groups of p -adic curves which are orthogonal to Pic^\wedge , and which satisfy (unconditionally) one of the above conditions.

CONTENTS

- §0. Introduction
- §1. Cuspidally i -th and i/p -th step prosolvable geometric fundamental groups
- §2. Cuspidalisation of sections of cuspidally i -th step prosolvable arithmetic fundamental groups
- §3. Lifting of sections to cuspidally $2/p$ -th step prosolvable arithmetic fundamental groups
- §4. Geometric sections of arithmetic fundamental groups of p -adic curves
- §5. Local sections of arithmetic fundamental groups of p -adic curves

§0. Introduction. Let k be a characteristic zero field, X a proper, smooth, and geometrically connected hyperbolic (i.e., $\text{genus}(X) \geq 2$) algebraic curve over k . Let K_X be the function field of X , K_X^{sep} a separable closure of K_X , and $\bar{k} \subset K_X^{\text{sep}}$ the algebraic closure of k . Let $\pi_1(X)$ be the étale fundamental group of X which sits in the following exact sequence

$$1 \rightarrow \pi_1(\bar{X}) \rightarrow \pi_1(X) \xrightarrow{\text{pr}} G_k \rightarrow 1,$$

where $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$, and $\pi_1(\bar{X})$ is the geometric étale fundamental group of X . Let $G_X \stackrel{\text{def}}{=} \text{Gal}(K_X^{\text{sep}}/K_X)$, and $\bar{G}_X \stackrel{\text{def}}{=} \text{Gal}(K_X^{\text{sep}}/K_X \cdot \bar{k})$. Thus, we have exact sequences

$$1 \rightarrow \bar{G}_X \rightarrow G_X \rightarrow G_k \rightarrow 1,$$

and

$$1 \rightarrow \mathcal{I}_X \rightarrow G_X \rightarrow \pi_1(X) \rightarrow 1,$$

where \mathcal{I}_X is the inertia subgroup. The theory of cuspidalisation of sections of arithmetic fundamental groups was initiated in [Saïdi], its ultimate aim is to reduce the Grothendieck anabelian section conjecture to its birational version. It can be formulated as follows (cf. loc. cit.).

The Cuspidalisation Problem for Sections of $\pi_1(X)$. Let $G_X \twoheadrightarrow H \twoheadrightarrow \pi_1(X)$ be a quotient of G_X . Given a section $s : G_k \rightarrow \pi_1(X)$ of the projection $\pi_1(X) \twoheadrightarrow G_k$, is it possible to **lift** s to a section $\tilde{s} : G_k \rightarrow H$ of the projection $H \twoheadrightarrow G_k$? i.e., is it possible to construct a section \tilde{s} such that the following diagram is commutative

$$\begin{array}{ccc} G_k & \xrightarrow{\tilde{s}} & H \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \pi_1(X) \end{array}$$

where the right vertical map is the projection $H \twoheadrightarrow \pi_1(X)$?

In [Saïdi] we investigated the cuspidalisation problem in the case $H \stackrel{\text{def}}{=} G_X^{(\text{c-ab})}$ is the maximal (geometrically) cuspidally abelian quotient of G_X . In this paper we generalise this theory to the (geometrically) *cuspidally i -th* as well as i/p -th; where p is a prime, *step prosolvable* quotient of G_X .

For $i \geq 0$, let $\mathcal{I}_X \twoheadrightarrow \mathcal{I}_{X,i}$ be the maximal i -th step prosolvable quotient of \mathcal{I}_X , and $G_X^{(i\text{-sol})} \stackrel{\text{def}}{=} G_X / \text{Ker}(\mathcal{I}_X \twoheadrightarrow \mathcal{I}_{X,i})$ the maximal (geometrically) cuspidally i -th step prosolvable quotient of G_X ($G_X^{(\text{c-ab})} \stackrel{\text{def}}{=} G_X^{(1\text{-sol})}$). For $i \geq 1$, let $s_i : G_k \rightarrow G_X^{(i\text{-sol})}$ be a section of the projection $G_X^{(i\text{-sol})} \twoheadrightarrow G_k$. In §2 we investigate the problem of lifting s_i to a section $s_{i+1} : G_k \rightarrow G_X^{(i+1\text{-sol})}$ of the projection $G_X^{(i+1\text{-sol})} \twoheadrightarrow G_k$. We say that the field k satisfies the condition **(H)** if the following holds. The Galois cohomology groups $H^1(G_k, M)$ are *finite* for every finite G_k -module M . This condition is satisfied for instance if the Galois group G_k is (topologically) finitely generated (e.g. k is a p -adic local field). One of our main results in this paper is the following (cf. Theorem 2.3.8).

Theorem 1. *Assume $i \geq 1$, and k satisfies the condition **(H)**. The section $s_i : G_k \rightarrow G_X^{(i\text{-sol})}$ **lifts** to a section $s_{i+1} : G_k \rightarrow G_X^{(i+1\text{-sol})}$ of the projection $G_X^{(i+1\text{-sol})} \twoheadrightarrow G_k$ **if and only if** for every $X' \rightarrow X$ a neighbourhood of the section s_i (i.e., corresponding to an open subgroup of $G_X^{(i\text{-sol})}$ containing $s_i(G_k)$) the class of $\text{Pic}_{X'}^1$ in $H^1(G_k, \text{Pic}_{X'}^0)$ lies in the maximal divisible subgroup of $H^1(G_k, \text{Pic}_{X'}^0)$.*

Key to the proof of Theorem 1 is the description of the G_k -module structure, induced by s_i , of $\mathcal{I}_X[i+1] \stackrel{\text{def}}{=} \text{Ker}(G_X^{(i+1\text{-sol})} \twoheadrightarrow G_X^{(i\text{-sol})})$ as the projective limit of the Tate modules of the jacobians of the neighbourhoods $\{X'\}$ as in the statement of Theorem 1 (cf. Proposition 1.1.5, and Lemma 2.3.2).

In §3 we investigate the following mod- p variant of Theorem 1, where p is a prime integer. Let $t \geq 0$, $i \geq 0$, and $\mathcal{I}_X \twoheadrightarrow \mathcal{I}_{X,i+1/p^t}$ the $i+1$ -th quotient of the $\mathbb{Z}/p^t\mathbb{Z}$ -derived series of \mathcal{I}_X (cf. 1.2). Thus, $\mathcal{I}_{X,i+1/p^t}$ is $i+1$ -step prosolvable with successive abelian quotients annihilated by p^t . Let $G_X^{(i+1/p^t\text{-sol})} \stackrel{\text{def}}{=} G_X / \text{Ker}(\mathcal{I}_X \twoheadrightarrow \mathcal{I}_{X,i+1/p^t})$ be the maximal (geometrically) cuspidally $i+1/p^t$ -th prosolvable quotient of G_X . Given a section $s : G_k \rightarrow \pi_1(X)$ of the projection $\pi_1(X) \twoheadrightarrow G_k$ we investigate

the problem of lifting s to a section $s_{i+1} : G_k \rightarrow G_X^{(i+1/p^t-\text{sol})}$ of the projection $G_X^{(i+1/p^t-\text{sol})} \twoheadrightarrow G_k$ in the **case** $\mathbf{i} = \mathbf{1}$ and $\mathbf{t} = \mathbf{1}$; the only case needed for applications in §4, and §5. For this purpose we introduce a certain quotient $G_X \twoheadrightarrow G_X^{(p,2)} \twoheadrightarrow G_X^{(2/p-\text{sol})}$, and investigate the problem of lifting the section s to a section $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$ of the projection $G_X^{(p,2)} \twoheadrightarrow G_k$ (this would give rise to a section of the projection $G_X^{(2/p-\text{sol})} \twoheadrightarrow G_k$ which lifts s). The quotient $G_X^{(p,2)}$ sits in an exact sequence $1 \rightarrow \mathcal{I}_X[p, 2] \rightarrow G_X^{(p,2)} \rightarrow G_X^{(1/p^2-\text{sol})} \rightarrow 1$, where $\mathcal{I}_X[p, 2]$ is abelian annihilated by p (cf. 3.3 for more details). In Theorem 3.4.11 we give necessary and sufficient conditions for the section s to lift to a section $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$ (cf. loc. cit. for a more precise statement).

In §4 we assume k is a p -adic local field (finite extension of \mathbb{Q}_p). We observe in this case that if $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$ is a section of the projection $G_X^{(p,2)} \twoheadrightarrow G_k$, and $s : G_k \rightarrow \pi_1(X)$ is the induced section of $\pi_1(X) \twoheadrightarrow G_k$, then s is *geometric* in the sense that it arises from a rational point $x \in X(k)$ (cf. Proposition 4.6). Further, we provide the following characterisation of sections $s : G_k \rightarrow \pi_1(X)$ which are geometric (cf. Theorem 4.5 where we prove a pro- Σ ; $p \in \Sigma$ is a set of primes, variant of Theorem 2).

Theorem 2. *Assume k is a p -adic local field. A section $s : G_k \rightarrow \pi_1(X)$ of the projection $\pi_1(X) \twoheadrightarrow G_k$ is **geometric** (cf. Definition 4.1) if and only if the following two conditions hold.*

(i) *The section s has a **cycle class uniformly orthogonal to Pic mod- p^2** (cf. Definition 3.4.1).*

(ii) *There **exists** a section $s' : G_k \rightarrow G_X^{(1/p^2-\text{sol})}$ of the projection $G_X^{(1/p^2-\text{sol})} \twoheadrightarrow G_k$ which **lifts** the section s (this holds if condition (i) is satisfied by Theorem 3.4.4) such that the following holds. For every $X' \rightarrow X$ a neighbourhood of the section s' (i.e., corresponding to an open subgroup of $G_X^{(1/p^2-\text{sol})}$ containing $s'(G_k)$) the class of $\text{Pic}_{X'}^1$ in $H^1(G_k, \text{Pic}_{X'}^0)$ is **divisible by p** .*

Condition (ii) in Theorem 2 is a necessary and sufficient condition for the section s' therein to lift to a section $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$ (cf. Theorem 3.4.10).

As an application of Theorem 2 we prove the following p -adic absolute anabelian result (cf. Theorem 4.8).

Theorem 3. *Let p_X, p_Y be prime integers, and X (resp. Y) a proper, smooth, geometrically connected hyperbolic curve over a p_X -adic local field k_X (respectively, p_Y -adic local field k_Y). Let $p_X \in \Sigma_X$ (resp. $p_Y \in \Sigma_Y$) be a non-empty set of prime integers of cardinality ≥ 2 , Π_X (resp. Π_Y) the geometrically pro- Σ_X (resp. pro- Σ_Y) arithmetic fundamental group of X (resp. Y), and $\varphi : \Pi_X \rightarrow \Pi_Y$ an isomorphism of profinite groups which fits in the following commutative diagram*

$$\begin{array}{ccc} G_X^{(p_X,2)} & \xrightarrow{\tilde{\varphi}} & G_Y^{(p_Y,2)} \\ \downarrow & & \downarrow \\ \Pi_X & \xrightarrow{\varphi} & \Pi_Y \end{array}$$

where $\tilde{\varphi}$ is an isomorphism of profinite groups. Here $G_X^{(p_X,2)}$ (resp. $G_Y^{(p_Y,2)}$) is the pro- Σ_X (resp. Σ_Y) version of the profinite group $G_X^{(p_X,2)}$ (resp. $G_Y^{(p_Y,2)}$) (cf. 3.3),

and the vertical maps are the natural projections. Then φ is geometric, i.e., arises from a uniquely determined isomorphism of schemes $X \xrightarrow{\sim} Y$.

Finally, in §5 we investigate *local* sections of arithmetic fundamental groups of p -adic curves. These are sections which arise from sections of arithmetic fundamental groups of *formal fibres* (cf. Definition 5.2). A geometric section is necessarily local in this sense. Our main result is the following (cf. Theorem 5.3).

Theorem 4. *Assume k is a p -adic local field, and $s : G_k \rightarrow \pi_1(X)$ is a local section of the projection $\pi_1(X) \twoheadrightarrow G_k$. Then s has a cycle class which is uniformly orthogonal to Pic^\wedge in the sense of [Saïdi], Definition 1.4.1(i).*

To the best of our knowledge, local sections of arithmetic fundamental groups of p -adic curves are the first non trivial (i.e., not known to be geometric a priori) examples of sections of arithmetic fundamental groups of p -adic curves which are orthogonal to Pic^\wedge . In particular, local sections satisfy condition (i) in Theorem 2.

Notations. Throughout this paper \mathfrak{Primes} denotes the set of all prime integers. For a profinite group H , we write H^{ab} for the maximal abelian quotient of H .

§1. Cuspidally i -th and i/p -th step prosolvable geometric fundamental groups. Let ℓ be an algebraically closed field of characteristic $l \geq 0$, X a proper smooth and connected hyperbolic curve (i.e., $\text{genus}(X) \geq 2$) over ℓ , and K_X its function field. Let η be a geometric point of X above its generic point; which determines a separable closure K_X^{sep} of K_X , and $\pi_1(X, \eta)$ the étale fundamental group of X with base point η .

Let $\emptyset \neq \Sigma \subseteq \mathfrak{Primes}$ be a set of prime integers. In case $\text{char}(\ell) = l > 0$ we assume that $l \notin \Sigma$. Write $\Delta_X \stackrel{\text{def}}{=} \pi_1(X, \eta)^\Sigma$ for the maximal pro- Σ quotient of $\pi_1(X, \eta)$. Let $\{x_s\}_{s=1}^n \subset X(\ell)$, $U \stackrel{\text{def}}{=} X \setminus \{x_1, \dots, x_n\}$ an open subscheme of X , $\Delta_U \stackrel{\text{def}}{=} \pi_1(U, \eta)^\Sigma$ the maximal pro- Σ quotient of the étale fundamental group $\pi_1(U, \eta)$ of U with base point η , and $I_U \stackrel{\text{def}}{=} \text{Ker}(\Delta_U \twoheadrightarrow \Delta_X)$. We shall refer to I_U as the *cuspidal subgroup* of Δ_U with respect to the natural projection $\Delta_U \twoheadrightarrow \Delta_X$ (cf. [Mochizuki], Definition 1.5); it is the subgroup of Δ_U (normally) generated by the (pro- Σ) inertia subgroups at the points $\{x_i\}_{i=1}^n$. We have the following exact sequence

$$1 \rightarrow I_U \rightarrow \Delta_U \rightarrow \Delta_X \rightarrow 1.$$

1.1 The quotient $\Delta_{U,i}$. For a profinite group H , we denote by $\overline{[H, H]}$ the closed subgroup of H topologically generated by the commutator subgroup. Consider the derived series of I_U

$$(1.1) \quad \dots \subseteq I_U(i+1) \subseteq I_U(i) \subseteq \dots \subseteq I_U(1) \subseteq I_U(0) = I_U,$$

where, for $i \geq 0$, $I_U(i+1) \stackrel{\text{def}}{=} \overline{[I_U(i), I_U(i)]}$ is the $(i+1)$ -th derived subgroup, which is a characteristic subgroup of I_U . Write

$$I_{U,i} \stackrel{\text{def}}{=} I_U / I_U(i).$$

Thus, $I_{U,i}$ is the maximal i -th step prosolvable quotient of I_U , and $I_{U,1}$ is the maximal abelian quotient of I_U . There exists a natural exact sequence

$$(1.2) \quad 1 \rightarrow I_U[i+1] \rightarrow I_{U,i+1} \rightarrow I_{U,i} \rightarrow 1$$

where $I_U[i+1]$ is the subgroup $I_U(i)/I_U(i+1)$ of $I_{U,i+1}$ and $I_U[i+1]$ is abelian. Write

$$\Delta_{U,i} \stackrel{\text{def}}{=} \Delta_U/I_U(i).$$

We shall refer to $\Delta_{U,i}$ (resp. $\Delta_{U,1}$) as the maximal **cuspidally i-th step pro-solvable** (resp. maximal **cuspidally abelian**) quotient of Δ_U (with respect to the surjection $\Delta_U \rightarrow \Delta_X$). We have the following commutative diagram of exact sequences.

$$(1.3) \quad \begin{array}{ccccccccc} & & 1 & & 1 & & & & \\ & & \downarrow & & \downarrow & & & & \\ & & I_U[i+1] & \xlongequal{\quad} & I_U[i+1] & & & & \\ & & \downarrow & & \downarrow & & & & \\ 1 & \longrightarrow & I_{U,i+1} & \longrightarrow & \Delta_{U,i+1} & \longrightarrow & \Delta_X & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \text{id}_{\Delta_X} \downarrow & & \\ 1 & \longrightarrow & I_{U,i} & \longrightarrow & \Delta_{U,i} & \longrightarrow & \Delta_X & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & & & \\ & & 1 & & 1 & & & & \end{array}$$

The profinite group $\Delta_{U,i}$; being a quotient of Δ_U , is topologically finitely generated (cf. [Grothendieck], Exposé X, Corollaire 3.10, recall $\text{char}(\ell) \notin \Sigma$). Hence there exists a sequence of characteristic open subgroups

$$\dots \subseteq \Delta_{U,i}[j+1] \subseteq \Delta_{U,i}[j] \subseteq \dots \subseteq \Delta_{U,i}[1] \stackrel{\text{def}}{=} \Delta_{U,i}$$

of $\Delta_{U,i}$ such that $\bigcap_{j \geq 1} \Delta_{U,i}[j] = \{1\}$. The open subgroup $\Delta_{U,i}[j] \subseteq \Delta_{U,i}$ corresponds to a finite (Galois) cover $X_{i,j}^U \rightarrow X$ between smooth connected and proper ℓ -curves which is étale above U . The geometric point η determines naturally a geometric point $\eta_{i,j}$ of $X_{i,j}^U$. Write $\Delta_{i,j}^U \stackrel{\text{def}}{=} \Delta_{X_{i,j}^U} \stackrel{\text{def}}{=} \pi_1(X_{i,j}^U, \eta_{i,j})^\Sigma$ for the maximal pro- Σ étale fundamental group of $X_{i,j}^U$ with base point $\eta_{i,j}$, and $(\Delta_{i,j}^U)^{\text{ab}}$ for the maximal abelian quotient of $\Delta_{i,j}^U$. The following Proposition provides a description of the structure of the profinite group $I_U[i+1]$ (cf. sequence (1.2) and diagram (1.3)) in the case $i \geq 1$. A description of the structure of $I_U[1]$ is given in [Mochizuki] Proposition 1.14 (see also [Saïdi], 2.1).

Proposition 1.1.1. *Let $i \geq 1$. There exists a natural isomorphism*

$$I_U[i+1] \xrightarrow{\sim} \varprojlim_{j \geq 1} (\Delta_{i,j}^U)^{\text{ab}}.$$

Proof of Proposition 1.1.1. Let G be a finite quotient of $\Delta_{U,i+1}$, which inserts in the following commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & I_{U,i+1} & \longrightarrow & \Delta_{U,i+1} & \longrightarrow & \Delta_X & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & I & \longrightarrow & G & \longrightarrow & G^{\text{et}} & \longrightarrow & 1 \end{array}$$

where the vertical maps are surjective. We assume, without loss of generality, that G is *not* a quotient of $\Delta_{U,i}$. The quotient G corresponds to a finite Galois cover $X_1 \rightarrow X$ with Galois group G , which factorizes as $X_1 \rightarrow X_1^{\text{et}} \rightarrow X$, where $X_1^{\text{et}} \rightarrow X$ is the maximal étale sub-cover with Galois group G^{et} , and $X_1 \rightarrow X_1^{\text{et}}$ is a (tamely) ramified Galois cover with group I . For $s \in \{1, \dots, n\}$, let $I_{x_s} \subset G$ be an inertia subgroup associated to x_s . Thus, I_{x_s} is only defined up to conjugation, and I is an $(i+1)$ -th step solvable group (normally) generated by the I_{x_s} 's. Moreover, I_{x_s} is cyclic of order $e_s \geq 1$ (coprime to $l = \text{char}(\ell)$) as the ramification is tame. The following claim follows immediately from the well-known structure of Δ_U (cf. [Grothendieck], Exposé X, Corollaire 3.10).

Claim 1.1.2. *There exists a finite quotient G' of $\Delta_{U,i}$, which inserts in the following commutative diagram (where the vertical maps are surjective)*

$$\begin{array}{ccccccc} 1 & \longrightarrow & I_{U,i} & \longrightarrow & \Delta_{U,i} & \longrightarrow & \Delta_X & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & I' & \longrightarrow & G' & \longrightarrow & G'^{\text{et}} & \longrightarrow & 1 \end{array}$$

such that the following holds. The quotient $\Delta_{U,i} \rightarrow G'$ corresponds to a finite Galois cover $X_2 \rightarrow X$ with Galois group G' , which factorizes as $X_2 \rightarrow X_2^{\text{et}} \rightarrow X$, where $X_2^{\text{et}} \rightarrow X$ is the maximal étale sub-cover with Galois group G'^{et} , and $X_2 \rightarrow X_2^{\text{et}}$ is a (tamely) ramified Galois cover with Galois group I' (an i -th step solvable group). Further, for $s \in \{1, \dots, n\}$, $I'_{x_s} \subseteq I'$ an inertia subgroup associated to x_s , then I'_{x_s} is cyclic of order $f_s = e_s h_s$ a multiple of e_s .

Next, let $K_1 \stackrel{\text{def}}{=} K_{X_1}$ (resp. $K_2 \stackrel{\text{def}}{=} K_{X_2}$) be the function field of X_1 (resp. X_2). Let $L \stackrel{\text{def}}{=} K_1.K_2$ be the compositum of K_1 and K_2 (in K_X^{sep}), and \tilde{X} the normalisation of X in L . Thus, $\tilde{X} \rightarrow X$ is a Galois cover with Galois group $H \subseteq G \times G'$ which is étale above U and factorizes as $\tilde{X} \rightarrow \tilde{X}^{\text{et}} \rightarrow X$, where $\tilde{X}^{\text{et}} \rightarrow X$ is the maximal étale sub-cover with Galois group H^{et} , and $\tilde{X} \rightarrow \tilde{X}^{\text{et}}$ is a (tamely) ramified Galois cover with group I_H : the subgroup of H (normally) generated by the inertia subgroups at the points of \tilde{X} above the $\{x_s\}_{s=1}^n$. (Thus, we have an exact sequence $1 \rightarrow I_H \rightarrow H \rightarrow H^{\text{et}} \rightarrow 1$.)

Lemma 1.1.3. *The quotient $\Delta_U \twoheadrightarrow H$ factorizes as $\Delta_U \twoheadrightarrow \Delta_{U,i+1} \twoheadrightarrow H$.*

Proof of Lemma 1.1.3. Indeed, one verifies easily that I_H is a subgroup of $I \times I'$ and $I \times I'$ is $(i+1)$ -th step solvable. \square

Next, let \tilde{I} be the maximal i -th step solvable quotient of I , which inserts in the exact sequence $1 \rightarrow I(i+1) \rightarrow I \rightarrow \tilde{I} \rightarrow 1$, with $I(i+1)$ abelian (note that $I(i+1)$ is non trivial by our assumption that G is not a quotient of $\Delta_{U,i}$). Write $\tilde{G} \stackrel{\text{def}}{=} G/I(i+1)$, which inserts in the exact sequence $1 \rightarrow \tilde{I} \rightarrow \tilde{G} \rightarrow G^{\text{et}} \rightarrow 1$. In particular, \tilde{G} is a quotient of $\Delta_{U,i}$. Let \tilde{H} be the image of H in $\tilde{G} \times G'$. We have a commutative diagram of exact sequences where the vertical maps are natural inclusions.

$$\begin{array}{ccccccc} 1 & \longrightarrow & I_H(i+1) \stackrel{\text{def}}{=} H \cap (I(i+1) \times \{1\}) & \longrightarrow & H & \longrightarrow & \tilde{H} & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & I(i+1) \times \{1\} & \longrightarrow & G \times G' & \longrightarrow & \tilde{G} \times G' & \longrightarrow & 1 \end{array}$$

Lemma 1.1.4. *The group \tilde{H} is a quotient of $\Delta_{U,i}$. Moreover, the cover $\tilde{X} \rightarrow X$ factorizes as $\tilde{X} \rightarrow \tilde{X}' \rightarrow X$, where $\tilde{X}' \rightarrow X$ is Galois (étale above U) with Galois group \tilde{H} , and $\tilde{X} \rightarrow \tilde{X}'$ is an **abelian étale** cover with Galois group $I_H(i+1)$.*

Proof of Lemma 1.1.4. The first assertion follows from the various definitions. Next, the Galois cover $X_1 \rightarrow X$ factorizes as $X_1 \rightarrow \tilde{X}_1 \rightarrow X$ where $X_1 \rightarrow \tilde{X}_1$ is Galois with Galois group $I(i+1)$, and $\tilde{X}_1 \rightarrow X$ is Galois with group \tilde{G} . Let \tilde{X}' be the normalisation of X in the compositum of the function fields of \tilde{X}_1 and X_2 . Thus, $\tilde{X}' \rightarrow X$ is a Galois cover with Galois group \tilde{H} , and we have the following commutative diagram

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & X_1 \\ \downarrow & & \downarrow \\ \tilde{X}' & \longrightarrow & \tilde{X}_1 \\ \downarrow & & \downarrow \\ X_2 & \longrightarrow & X \end{array}$$

of finite Galois covers. The ramification index in the Galois cover $X_2 \rightarrow X$ above a branched (closed) point $x_s \in X$ is divisible by the ramification index above x_s in the Galois cover $X_1 \rightarrow X$ (cf. the above condition that f_s is divisible by e_s). The fact that the morphism $\tilde{X} \rightarrow X_2$, and a fortiori $\tilde{X} \rightarrow \tilde{X}'$; which is abelian with Galois group $I_H(i+1)$, is étale follows from Abhyankar's Lemma (cf. [Grothendieck], Exposé X, Lemma 3.6). \square

Going back to the proof of Proposition 1.1.1, the above discussion shows that the finite quotients $\Delta_{U,i+1} \rightarrow H$ as in Lemma 1.1.3 form a cofinal system of finite quotients of $\Delta_{U,i+1}$. Thus, $\Delta_{U,i+1} \xrightarrow{\sim} \varprojlim_H H$. Proposition 1.1.1 then follows from the facts that the various H above fit in an exact sequence $1 \rightarrow I_H(i+1) \rightarrow H \rightarrow \tilde{H} \rightarrow 1$; $\Delta_{U,i} \xrightarrow{\sim} \varprojlim_{\tilde{H}} \tilde{H}$, and the above Galois covers $\tilde{X} \rightarrow \tilde{X}'$ with group $I_H(i+1)$ are étale abelian (cf. Lemma 1.1.4). This finishes the proof of Proposition 1.1.1. \square

Similarly, let $G_{K_X} \stackrel{\text{def}}{=} \text{Gal}(K_X^{\text{sep}}/K_X)$, and $G_X \stackrel{\text{def}}{=} G_{K_X}^{\Sigma}$ the maximal pro- Σ quotient of G_{K_X} . We have a natural exact sequence

$$1 \rightarrow \mathcal{I}_X \rightarrow G_X \rightarrow \Delta_X \rightarrow 1,$$

where $\mathcal{I}_X \stackrel{\text{def}}{=} \text{Ker}(G_X \rightarrow \Delta_X)$ is the *cuspidal subgroup* of G_X (with respect to the surjection $G_X \rightarrow \Delta_X$). Let $i \geq 0$ and write

$$\mathcal{I}_{X,i} \stackrel{\text{def}}{=} \mathcal{I}_X / \mathcal{I}_X(i).$$

Thus, $\mathcal{I}_{X,i}$ is the maximal i -th step prosolvable quotient of \mathcal{I}_X , and $\mathcal{I}_{X,1}$ is the maximal abelian quotient of \mathcal{I}_X . There exists a natural exact sequence

$$(1.4) \quad 1 \rightarrow \mathcal{I}_X[i+1] \rightarrow \mathcal{I}_{X,i+1} \rightarrow \mathcal{I}_{X,i} \rightarrow 1,$$

where $\mathcal{I}_X[i+1]$ is the subgroup $\mathcal{I}_X(i)/\mathcal{I}_X(i+1)$ of $\mathcal{I}_{X,i+1}$, and $\mathcal{I}_X[i+1]$ is abelian. Write

$$G_{X,i} \stackrel{\text{def}}{=} G_X / \mathcal{I}_X(i).$$

We shall refer to $G_{X,i}$ (resp. $G_{X,1}$) as the maximal **cuspidally i-th step pro-solvable** (resp. maximal **cuspidally abelian**) quotient of G_X (with respect to the surjection $G_X \twoheadrightarrow \Delta_X$). We have the following commutative diagram

$$\begin{array}{ccccccc}
 & & 1 & & 1 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \mathcal{I}_X[i+1] & \xlongequal{\quad} & \mathcal{I}_X[i+1] & & \\
 & & \downarrow & & \downarrow & & \\
 (1.5) & 1 & \longrightarrow & \mathcal{I}_{X,i+1} & \longrightarrow & G_{X,i+1} & \longrightarrow & \Delta_X & \longrightarrow & 1 \\
 & & & \downarrow & & \downarrow & & \downarrow & & \\
 & 1 & \longrightarrow & \mathcal{I}_{X,i} & \longrightarrow & G_{X,i} & \longrightarrow & \Delta_X & \longrightarrow & 1 \\
 & & & \downarrow & & \downarrow & & & & \\
 & & & 1 & & 1 & & & &
 \end{array}$$

of exact sequences.

Proposition 1.1.5. *There are natural isomorphisms $G_{X,i} \xrightarrow{\sim} \varprojlim_U \Delta_{U,i}$, $\mathcal{I}_{X,i} \xrightarrow{\sim} \varprojlim_U \mathcal{I}_{U,i}$, and $\mathcal{I}_X[i+1] \xrightarrow{\sim} \varprojlim_U \mathcal{I}_U[i+1]$, where the projective limit is taken over all non-empty open subschemes $U \subseteq X$. Moreover, for $i \geq 1$, we have a natural isomorphism*

$$\mathcal{I}_X[i+1] \xrightarrow{\sim} \varprojlim_U \left(\varliminf_{j \geq 1} (\Delta_{i,j}^U)^{\text{ab}} \right),$$

where $(\Delta_{i,j}^U)^{\text{ab}}$ is as in the discussion preceding Proposition 1.1.1.

Proof. Follows from the various definitions and Proposition 1.1.1. \square

1.2. The quotient $\Delta_U \twoheadrightarrow \Delta_U^{p,i+1}$. In this subsection we discuss a certain variant of the theory in 1.1, we use the same notations as in loc. cit.. For a profinite group H , a prime integer p , and an integer $t \geq 1$, write

$$\dots \subseteq H(i+1/p^t) \subseteq H(i/p^t) \subseteq \dots \subseteq H(1/p^t) \subseteq H(0/p^t) = H$$

for the $\mathbb{Z}/p^t\mathbb{Z}$ -derived series of H , where $H(i+1/p^t) \stackrel{\text{def}}{=} \overline{\langle [H(i/p^t), H(i/p^t)], H(i/p^t)^{p^t} \rangle}$ is the $i+1/p^t$ -th derived subgroup, which is a characteristic subgroup of H . Write

$$H_{i/p^t} \stackrel{\text{def}}{=} H/H(i/p^t).$$

We will refer to H_{i/p^t} as the maximal i/p^t -th step *prosolvable* quotient of H , and H_{1/p^t} as the maximal *abelian* annihilated by p^t quotient of H . There exists a natural exact sequence

$$1 \rightarrow H[i+1/p^t] \rightarrow H_{i+1/p^t} \rightarrow H_{i/p^t} \rightarrow 1$$

where $H[i + 1/p^t]$ is the subgroup $H(i/p^t)/H(i + 1/p^t)$ of H_{i+1/p^t} , and $H[i + 1/p^t]$ is abelian annihilated by p^t .

Next, let $p \in \Sigma$, and consider the $\mathbb{Z}/p\mathbb{Z}$ -derived series of I_U

$$(1.6) \quad \dots \subseteq I_U(i + 1/p) \subseteq I_U(i/p) \subseteq \dots \subseteq I_U(1/p) \subseteq I_U(0/p) = I_U$$

(cf. the above discussion in the case $t = 1$). Then $I_{U,i/p} \stackrel{\text{def}}{=} I_U/I_U(i/p)$ is the maximal i/p -th step prosolvable quotient of I_U , and $I_{U,1/p}$ is the maximal abelian annihilated by p quotient of I_U . Write $\Delta_{U,i/p} \stackrel{\text{def}}{=} \Delta_U/I_U(i/p)$, which inserts in the exact sequence

$$1 \rightarrow I_{U,i/p} \rightarrow \Delta_{U,i/p} \rightarrow \Delta_X \rightarrow 1.$$

We shall refer to $\Delta_{U,i/p}$ (resp. $\Delta_{U,1/p}$) as the maximal **cuspidally i/p -th step prosolvable** (resp. maximal **cuspidally abelian annihilated by p**) quotient of Δ_U (with respect to the surjection $\Delta_U \twoheadrightarrow \Delta_X$).

Next, we define a certain quotient $\Delta_U \twoheadrightarrow \Delta_U^{p,i+1}$ of Δ_U , which dominates $\Delta_{U,i+1/p}$. Let $i \geq 0$, and G a finite quotient of $\Delta_{U,i+1/p}$ which inserts in the following commutative diagram.

$$\begin{array}{ccccccc} 1 & \longrightarrow & I_{U,i+1/p} & \longrightarrow & \Delta_{U,i+1/p} & \longrightarrow & \Delta_X \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & I & \longrightarrow & G & \longrightarrow & G^{\text{et}} \longrightarrow 1 \end{array}$$

Thus, the quotient G corresponds to a finite Galois cover $X'_1 \rightarrow X$ with Galois group G , which factorizes as $X'_1 \rightarrow X'_1{}^{\text{et}} \rightarrow X$, where $X'_1{}^{\text{et}} \rightarrow X$ is the maximal étale sub-cover with Galois group G^{et} , and $X'_1 \rightarrow X'_1{}^{\text{et}}$ is a tamely ramified Galois cover with Galois group I . Moreover, I is an $(i + 1)$ -th step solvable group whose successive abelian quotients are annihilated by p . We will assume for the remaining discussion, without loss of generality, that G is *not* a quotient of $\Delta_{U,i/p}$.

Let $s \in \{1, \dots, n\}$, and $I_{x_s} \subset G$ an inertia subgroup associated to x_s . Then I_{x_s} is cyclic of order p^t , with $t \leq i + 1$, as follows from the structure of I . Write $\Delta_{U,1/p^{i+1}} \stackrel{\text{def}}{=} \Delta_U/I_U(1/p^{i+1})$ for the maximal cuspidally abelian annihilated by p^{i+1} quotient of Δ_U (with respect to the surjection $\Delta_U \twoheadrightarrow \Delta_X$). The following claim follows immediately from the well-known structure of Δ_U (cf. [Grothendieck], Exposé X, Corollaire 3.10).

Claim 1.2.1. *There exists a finite quotient G' of $\Delta_{U,1/p^{i+1}}$ which inserts in the following commutative diagram (where the vertical maps are surjective)*

$$\begin{array}{ccccccc} 1 & \longrightarrow & I_{U,1/p^{i+1}} & \longrightarrow & \Delta_{U,1/p^{i+1}} & \longrightarrow & \Delta_X \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & I' & \longrightarrow & G' & \longrightarrow & G'^{\text{et}} \longrightarrow 1 \end{array}$$

such that the following holds. The quotient $\Delta_{U,1/p^{i+1}} \twoheadrightarrow G'$ corresponds to a Galois cover $X'_2 \rightarrow X$ with Galois group G' which factorizes as $X'_2 \rightarrow X'_2{}^{\text{et}} \rightarrow X$, where $X'_2{}^{\text{et}} \rightarrow X$ is the maximal étale sub-cover with Galois group G'^{et} , and $X'_2 \rightarrow X'_2{}^{\text{et}}$ is a tamely ramified cover with Galois group I' (an abelian group annihilated by

p^{i+1}). Further, for $s \in \{1, \dots, n\}$, and $I'_{x_s} \subseteq I'$ an inertia subgroup associated to x_s , then I'_{x_s} is cyclic of order p^{i+1} .

Let $K'_1 \stackrel{\text{def}}{=} K_{X'_1}$ (resp. $K'_2 \stackrel{\text{def}}{=} K_{X'_2}$) be the function field of X'_1 (resp. X'_2), $L' \stackrel{\text{def}}{=} K'_1.K'_2$ the compositum of K'_1 and K'_2 , and Y the normalisation of X in L' . Thus, $Y \rightarrow X$ is a Galois cover with Galois group $H \subseteq G \times G'$ which is étale above U . Note that H maps onto G , G' , and the quotient $\Delta_U \twoheadrightarrow H$ doesn't factorize through $\Delta_U \twoheadrightarrow \Delta_{U, i+1/p}$ if $i \geq 1$.

Let $I'' \stackrel{\text{def}}{=} \text{Ker}(I \twoheadrightarrow I^{\text{ab}})$. Thus, I'' is an i -th step solvable group whose successive quotients are annihilated by p , and is a characteristic subgroup of I . Write $\tilde{G} \stackrel{\text{def}}{=} G/I''$, and let \tilde{H} be the image of H in the quotient $\tilde{G} \times G'$ of $G \times G'$. We have a commutative diagram of exact sequences

$$\begin{array}{ccccccccc} 1 & \longrightarrow & I_H \stackrel{\text{def}}{=} H \cap (I'' \times \{1\}) & \longrightarrow & H & \longrightarrow & \tilde{H} & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & I'' \times \{1\} & \longrightarrow & G \times G' & \longrightarrow & \tilde{G} \times G' & \longrightarrow & 1 \end{array}$$

where the vertical maps are natural inclusions.

Lemma 1.2.2. *The group \tilde{H} is a quotient of $\Delta_{U, 1/p^{i+1}}$. Moreover, the Galois cover $Y \rightarrow X$ factorizes as $Y \rightarrow Y' \rightarrow X$, where $Y' \rightarrow X$ is a tamely ramified Galois cover with Galois group \tilde{H} , and $Y \rightarrow Y'$ is an étale Galois cover with Galois group $I_H \subseteq I''$: an i -th step solvable group whose successive abelian quotients are annihilated by p .*

Proof. The first assertion follows from the fact that the inertia subgroup of \tilde{H} is a subgroup of $I^{\text{ab}} \times I'$. The proof of the second assertion is similar to the proof of Lemma 1.1.4 using Abhyankar's Lemma (cf. loc. cit.). \square

The profinite group $\Delta_{U, 1/p^{i+1}}$; being a quotient of Δ_U , is topologically finitely generated. Hence there exists a sequence of characteristic open subgroups

$$\dots \subseteq \Delta_{U, 1/p^{i+1}}[j+1] \subseteq \Delta_{U, 1/p^{i+1}}[j] \subseteq \dots \subseteq \Delta_{U, 1/p^{i+1}}[1] \stackrel{\text{def}}{=} \Delta_{U, 1/p^{i+1}}$$

of $\Delta_{U, 1/p^{i+1}}$ such that $\bigcap_{j \geq 1} \Delta_{U, 1/p^{i+1}}[j] = \{1\}$. The open subgroup $\Delta_{U, 1/p^{i+1}}[j] \subseteq \Delta_{U, 1/p^{i+1}}$ corresponds to a finite Galois cover $(X'_{i+1, j})^U \rightarrow X$ between smooth connected and proper ℓ -curves, with Galois group $G_{i+1, j}^U \stackrel{\text{def}}{=} \Delta_{U, 1/p^{i+1}} / \Delta_{U, 1/p^{i+1}}[j]$, and which restricts to an étale cover $(V_{i+1, j})^U \rightarrow U$. The geometric point η determines a geometric point $\eta'_{i+1, j}$ of $(X'_{i+1, j})^U$ and $(V_{i+1, j})^U$. Write $(\Delta'_{i+1, j})^U = \Delta_{(X'_{i+1, j})^U} \stackrel{\text{def}}{=} \pi_1((X'_{i+1, j})^U, \eta'_{i+1, j})^\Sigma$ for the maximal pro- Σ étale fundamental group of $(X'_{i+1, j})^U$ with base point $\eta'_{i+1, j}$, and $((\Delta'_{i+1, j})^U)_{i/p}$ for the maximal i/p -th step prosolvable quotient of $(\Delta'_{i+1, j})^U$. Consider the following push-out diagram

$$\begin{array}{ccccccccc} 1 & \longrightarrow & \pi_1((V_{i+1, j})^U, \eta'_{i+1, j})^\Sigma & \longrightarrow & \Delta_U & \longrightarrow & G_{i+1, j}^U & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & ((\Delta'_{i+1, j})^U)_{i/p} & \longrightarrow & \tilde{G}_{i+1, j}^U & \longrightarrow & G_{i+1, j}^U & \longrightarrow & 1 \end{array}$$

$(\text{ker}(\pi_1((V'_{i+1, j})^U, \eta'_{i+1, j})^\Sigma \twoheadrightarrow ((\Delta'_{i+1, j})^U)_{i/p}))$ is a normal subgroup of Δ_U .

Lemma 1.2.3. *With the above notations, G is a quotient of $\tilde{G}_{i+1,j}^U$ for some $j \geq 1$.*

Proof. Follows from Lemma 1.2.2 and the various definitions. \square

Let

$$\Delta_U^{p,i+1} \stackrel{\text{def}}{=} \varprojlim_{j \geq 1} \tilde{G}_{i+1,j}^U,$$

where $\tilde{G}_{i+1,j}^U$ is as in Lemma 1.2.3 (the $\{\tilde{G}_{i+1,j}^U\}_{j \geq 1}$ form a projective system). Thus, it follows from the various definitions that we have a natural exact sequence

$$(1.7) \quad 1 \rightarrow \varprojlim_{j \geq 1} ((\Delta'_{i+1,j})^U)_{i/p} \rightarrow \Delta_U^{p,i+1} \rightarrow \Delta_{U,1/p^{i+1}} \rightarrow 1,$$

where the $\{((\Delta'_{i+1,j})^U)_{i/p}\}_{j \geq 1}$ are defined as above.

Proposition 1.2.4. *The profinite group $\Delta_{U,i+1/p}$ is a quotient of $\Delta_U^{p,i+1}$.*

Proof. Follows from the above discussion (cf. Lemma 1.2.3). \square

Similarly, write $\mathcal{I}_{X,i+1/p} \stackrel{\text{def}}{=} \mathcal{I}_X / \mathcal{I}_X(i+1/p)$. Thus, $\mathcal{I}_{X,i+1/p}$ is the maximal $i+1/p$ -th step prosolvable quotient of \mathcal{I}_X , and $\mathcal{I}_{X,1/p}$ is the maximal abelian annihilated by p quotient of \mathcal{I}_X . Write

$$G_{X,i+1/p} \stackrel{\text{def}}{=} G_X / \mathcal{I}_X(i+1/p).$$

We shall refer to $G_{X,i+1/p}$ (resp. $G_{X,1/p}$) as the maximal **cuspidally $i+1/p$ -th step prosolvable** (resp. maximal **cuspidally abelian annihilated by p**) quotient of G_X (with respect to the surjection $G_X \twoheadrightarrow \Delta_X$). Also, write $G_{X,1/p^{i+1}} \stackrel{\text{def}}{=} G_X / \mathcal{I}_X(1/p^{i+1})$, and $G_X^{p,i+1} \stackrel{\text{def}}{=} \varprojlim_U (\Delta_U^{p,i+1})$ where the limit is taken over all open subschemes $U \subseteq X$. We have the following exact sequence

$$(1.8) \quad 1 \rightarrow \varprojlim_U (\varprojlim_{j \geq 1} ((\Delta'_{i+1,j})^U)_{i/p}) \rightarrow G_X^{p,i+1} \rightarrow G_{X,1/p^{i+1}} \rightarrow 1.$$

Lemma 1.2.5. *The profinite group $G_{X,i+1/p}$ is a quotient of $G_X^{p,i+1}$.*

Proof. Follows from the various definitions, and Proposition 1.2.4. \square

§2. Cuspidalisation of sections of cuspidally i -th step prosolvable arithmetic fundamental groups. In this section k is a field with $\text{char}(k) = l \geq 0$, X is a proper smooth and geometrically connected hyperbolic (i.e., $\text{genus}(X) \geq 2$) curve over k , and K_X its function field. Let η be a geometric point of X above its generic point; it determines an algebraic closure \bar{k} of k , and a geometric point $\bar{\eta}$ of $\bar{X} \stackrel{\text{def}}{=} X \times_k \bar{k}$.

2.1. Let $\Sigma \subseteq \mathfrak{Primes}$ be a non-empty set of prime integers. In case $\text{char}(k) = l > 0$ we assume that $l \notin \Sigma$. Write $\Delta_X \stackrel{\text{def}}{=} \pi_1(\bar{X}, \bar{\eta})^\Sigma$ for the maximal pro- Σ quotient of $\pi_1(\bar{X}, \bar{\eta})$, and $\Pi_X \stackrel{\text{def}}{=} \pi_1(X, \eta) / \text{Ker}(\pi_1(\bar{X}, \bar{\eta}) \twoheadrightarrow \pi_1(\bar{X}, \bar{\eta})^\Sigma)$. Thus, we have an exact sequence

$$(2.1) \quad 1 \rightarrow \Delta_X \rightarrow \Pi_X \xrightarrow{\text{pr}_{X,\Sigma}} G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k) \rightarrow 1.$$

We shall refer to $\pi_1(X, \eta)^{(\Sigma)} \stackrel{\text{def}}{=} \Pi_X$ as the **geometrically pro- Σ arithmetic fundamental group** of X .

2.1.1. Let $U \subseteq X$ be a nonempty open subscheme. Write $\Delta_U \stackrel{\text{def}}{=} \pi_1(\overline{U}, \overline{\eta})^\Sigma$ for the maximal pro- Σ quotient of the fundamental group $\pi_1(\overline{U}, \overline{\eta})$ of \overline{U} with base point $\overline{\eta}$, and $\Pi_U \stackrel{\text{def}}{=} \pi_1(U, \eta) / \text{Ker}(\pi_1(\overline{U}, \overline{\eta}) \twoheadrightarrow \pi_1(\overline{U}, \overline{\eta})^\Sigma)$. Thus, we have an exact sequence $1 \rightarrow \Delta_U \rightarrow \Pi_U \rightarrow G_k \rightarrow 1$. Let $I_U \stackrel{\text{def}}{=} \text{Ker}(\Pi_U \twoheadrightarrow \Pi_X) = \text{Ker}(\Delta_U \twoheadrightarrow \Delta_X)$ be the cuspidal subgroup of Π_U (with respect to the surjection $\Pi_U \twoheadrightarrow \Pi_X$). We have the following exact sequence

$$(2.2) \quad 1 \rightarrow I_U \rightarrow \Pi_U \rightarrow \Pi_X \rightarrow 1.$$

Let $i \geq 0$ be an integer, $I_{U,i}$ the maximal i -th step prosolvable quotient of I_U (cf. 1.1), and $\Pi_U^{(i\text{-sol})} \stackrel{\text{def}}{=} \Pi_U / \text{Ker}(I_U \twoheadrightarrow I_{U,i})$. We shall refer to $\Pi_U^{(i\text{-sol})}$ as the maximal (geometrically) **cuspidally i -th step prosolvable quotient** of Π_U (with respect to the surjection $\Pi_U \twoheadrightarrow \Pi_X$).

2.1.2. Similarly, we have an exact sequence of absolute Galois groups $1 \rightarrow G_{\overline{k}.K_X} \rightarrow G_{K_X} \rightarrow G_k \rightarrow 1$, where $G_{\overline{k}.K_X} \stackrel{\text{def}}{=} \pi_1(\text{Spec}(\overline{k}.K_X), \eta)$, and $G_{K_X} \stackrel{\text{def}}{=} \pi_1(\text{Spec}(K_X), \eta)$. Let $G_{\overline{X}} \stackrel{\text{def}}{=} G_{\overline{k}.K_X}^\Sigma$ be the maximal pro- Σ quotient of $G_{\overline{k}.K_X}$, $G_X \stackrel{\text{def}}{=} G_{K_X} / \text{Ker}(G_{\overline{k}.K_X} \twoheadrightarrow G_{\overline{k}.K_X}^\Sigma)$, and $\mathcal{I}_X \stackrel{\text{def}}{=} \text{Ker}(G_X \twoheadrightarrow \Pi_X) = \text{Ker}(G_{\overline{X}} \twoheadrightarrow \Delta_X)$ the cuspidal subgroup of G_X (with respect to the surjection $G_X \twoheadrightarrow \Pi_X$). Let $\mathcal{I}_{X,i}$ be the maximal i -th step prosolvable quotient of \mathcal{I}_X . By pushing the exact sequence $1 \rightarrow \mathcal{I}_X \rightarrow G_X \rightarrow \Pi_X \rightarrow 1$ by the characteristic quotient $\mathcal{I}_X \twoheadrightarrow \mathcal{I}_{X,i}$ we obtain an exact sequence

$$(2.3) \quad 1 \rightarrow \mathcal{I}_{X,i} \rightarrow G_X^{(i\text{-sol})} \rightarrow \Pi_X \rightarrow 1.$$

We will refer to the quotient $G_X^{(i\text{-sol})}$ as the maximal (geometrically) **cuspidally i -th step prosolvable quotient** of G_X (with respect to the surjective homomorphism $G_X \twoheadrightarrow \Pi_X$). There exist natural isomorphisms

$$G_X^{(i\text{-sol})} \xrightarrow{\sim} \varprojlim_U \Pi_U^{(i\text{-sol})}, \quad \mathcal{I}_{X,i} \xrightarrow{\sim} \varprojlim_U I_{U,i},$$

where the limit is taken over all open subschemes $U \subseteq X$.

2.2. Let $i \geq 0$. We have a commutative diagram of exact sequences

$$(2.4) \quad \begin{array}{ccccccc} & & 1 & & 1 & & \\ & & \downarrow & & \downarrow & & \\ & & \mathcal{I}_X[i+1] & \xlongequal{\quad} & \mathcal{I}_X[i+1] & & \\ & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & G_{\overline{X},i+1} & \longrightarrow & G_X^{(i+1\text{-sol})} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & G_{\overline{X},i} & \longrightarrow & G_X^{(i\text{-sol})} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \\ & & 1 & & 1 & & \end{array}$$

and similarly, for $U \subseteq X$ nonempty open, we have the following commutative diagram

$$(2.5) \quad \begin{array}{ccccccc} & & 1 & & 1 & & \\ & & \downarrow & & \downarrow & & \\ & & I_U[i+1] & \xlongequal{\quad} & I_U[i+1] & & \\ & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & \Delta_{U,i+1} & \longrightarrow & \Pi_U^{(i+1-\text{sol})} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \Delta_{U,i} & \longrightarrow & \Pi_U^{(i-\text{sol})} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \\ & & 1 & & 1 & & \end{array}$$

(recall the definition of $\mathcal{I}_X[i+1]$ and $I_U[i+1]$ from §1).

Assume that the lower horizontal sequence in diagram (2.4) splits. Let $s : G_k \rightarrow G_X^{(i-\text{sol})}$ be a section of the projection $G_X^{(i-\text{sol})} \twoheadrightarrow G_k$, which induces a section $s_U : G_k \rightarrow \Pi_U^{(i-\text{sol})}$ of the projection $\Pi_U^{(i-\text{sol})} \twoheadrightarrow G_k$, $\forall U \subseteq X$ open.

The cuspidalisation problem for sections of cuspidally i -th step prosolvable arithmetic fundamental groups. *Let $i \geq 0$. Given a section $s_U : G_k \rightarrow \Pi_U^{(i-\text{sol})}$ as above, is it possible to construct a section $\tilde{s}_U : G_k \rightarrow \Pi_U^{(i+1-\text{sol})}$ of the projection $\Pi_U^{(i+1-\text{sol})} \twoheadrightarrow G_k$ which **lifts** the section s_U , i.e., which fits in a commutative diagram*

$$\begin{array}{ccc} G_k & \xrightarrow{\tilde{s}_U} & \Pi_U^{(i+1-\text{sol})} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s_U} & \Pi_U^{(i-\text{sol})} \end{array}$$

where the right vertical map is the natural surjection?

Similarly, is it possible to construct a section $\tilde{s} : G_k \rightarrow G_X^{(i+1-\text{sol})}$ of the projection $G_X^{(i+1-\text{sol})} \twoheadrightarrow G_k$ which **lifts** the section s , i.e., which fits in a commutative diagram

$$\begin{array}{ccc} G_k & \xrightarrow{\tilde{s}} & G_X^{(i+1-\text{sol})} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & G_X^{(i-\text{sol})} \end{array}$$

where the right vertical map is the natural surjection?

2.3. The above cuspidalisation problem in the case $i = 0$ has been investigated in [Saïdi]. Next, we will investigate this problem in the case $i \geq 1$.

We use the notations in 2.2. Recall the definition of the characteristic open subgroups $\{\Delta_{U,i}[j]\}_{j \geq 1}$ such that $\bigcap_{j \geq 1} \Delta_{U,i}[j] = \{1\}$ (cf. discussion before Proposition 1.1.1). Write $\widehat{\Pi}_U[i,j] \stackrel{\text{def}}{=} \widehat{\Pi}_U[i,j][s_U] \stackrel{\text{def}}{=} \Delta_{U,i}[j].s_U(G_k)$. Thus, $\widehat{\Pi}_U[i,j] \subseteq \Pi_U^{(i-\text{sol})}$

is an open subgroup which contains the image $s_U(G_k)$ of the section s_U . Write $\Pi_U[i, j] \stackrel{\text{def}}{=} \Pi_U[i, j][s_U]$ for the inverse image of $\widehat{\Pi}_U[i, j]$ in Π_U . Thus, $\Pi_U[i, j] \subseteq \Pi_U$ is an open subgroup which corresponds to an étale cover $V_{i,j} \rightarrow U$, where $V_{i,j}$ is a geometrically irreducible k -curve (since $\Pi_U[i, j]$ maps onto G_k via the natural projection $\Pi_U \twoheadrightarrow G_k$ by the very definition of $\Pi_U[i, j]$).

Write $X_{i,j}^U$ (resp. $\overline{X}_{i,j}^U$) for the smooth compactification of $V_{i,j}$ (resp. $\overline{V}_{i,j} \stackrel{\text{def}}{=} V_{i,j} \times_k \overline{k}$). We have an exact sequence $1 \rightarrow \pi_1(\overline{X}_{i,j}^U, \overline{\eta}_{i,j}) \rightarrow \pi_1(X_{i,j}^U, \eta_{i,j}) \rightarrow G_k \rightarrow 1$, where $\eta_{i,j}$ (resp. $\overline{\eta}_{i,j}$) is a geometric point naturally induced by η . Write $\pi_1(\overline{X}_{i,j}^U, \overline{\eta}_{i,j})^{\Sigma, \text{ab}}$ for the maximal pro- Σ abelian quotient of $\pi_1(\overline{X}_{i,j}^U, \overline{\eta}_{i,j})$, and $\pi_1(X_{i,j}^U, \eta_{i,j})^{(\Sigma, \text{ab})} \stackrel{\text{def}}{=} \pi_1(X_{i,j}^U, \eta_{i,j}) / \text{Ker}(\pi_1(\overline{X}_{i,j}^U, \overline{\eta}_{i,j}) \twoheadrightarrow \pi_1(\overline{X}_{i,j}^U, \overline{\eta}_{i,j})^{\Sigma, \text{ab}})$ for the geometrically pro- Σ abelian fundamental group of $X_{i,j}^U$. Consider the following pull-back diagram.

$$(2.6) \quad \begin{array}{ccccccc} 1 & \longrightarrow & I_U[i+1] & \longrightarrow & \mathcal{H}_{U,i} \stackrel{\text{def}}{=} \mathcal{H}_{U,i}[s_U] & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow s_U \\ 1 & \longrightarrow & I_U[i+1] & \longrightarrow & \Pi_U^{(i+1-\text{sol})} & \longrightarrow & \Pi_U^{(i-\text{sol})} \longrightarrow 1 \end{array}$$

Lemma 2.3.1. *The section $s_U : G_k \rightarrow \Pi_U^{(i-\text{sol})}$ lifts to a section $\tilde{s}_U : G_k \rightarrow \Pi_U^{(i+1-\text{sol})}$ of the projection $\Pi_U^{(i+1-\text{sol})} \twoheadrightarrow G_k$ if and only if the group extension $1 \rightarrow I_U[i+1] \rightarrow \mathcal{H}_{U,i} \rightarrow G_k \rightarrow 1$ splits.*

Proof. Follows immediately from the diagram (2.6). \square

Lemma 2.3.2. *Assume $i \geq 1$. Then we have natural identifications $I_U[i+1] \xrightarrow{\sim} \varprojlim_{j \geq 1} \pi_1(\overline{X}_{i,j}^U, \overline{\eta}_{i,j})^{\Sigma, \text{ab}}$, and $\mathcal{H}_{U,i} \xrightarrow{\sim} \varprojlim_{j \geq 1} \pi_1(X_{i,j}^U, \eta_{i,j})^{(\Sigma, \text{ab})}$.*

Proof. Follows from the various definitions and Proposition 1.1.1. \square

Definition 2.3.3. We say that the field k satisfies the condition (\mathbf{H}_Σ) if the following holds. The Galois cohomology groups $H^1(G_k, M)$ are *finite* for every finite G_k -module M whose cardinality is divisible only by primes in Σ .

Lemma 2.3.4. *Assume that $i \geq 1$, and k satisfies the condition (\mathbf{H}_Σ) . Then the group extension $1 \rightarrow I_U[i+1] \rightarrow \mathcal{H}_{U,i} \rightarrow G_k \rightarrow 1$ splits if and only if the group extensions $1 \rightarrow \pi_1(\overline{X}_{i,j}^U, \overline{\eta}_{i,j})^{\Sigma, \text{ab}} \rightarrow \pi_1(X_{i,j}^U, \eta_{i,j})^{(\Sigma, \text{ab})} \rightarrow G_k \rightarrow 1$ split, $\forall j \geq 1$.*

Proof. The only if part follows immediately from Lemma 2.3.2. Conversely, assume that the group extension $\pi_1(X_{i,j}^U, \eta_{i,j})^{(\Sigma, \text{ab})}$ splits, $\forall j \geq 1$. Write $\mathcal{H}_{U,i} = \varprojlim_G G$ as the projective limit of finite quotients G which insert into a commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & I_U[i+1] & \longrightarrow & \mathcal{H}_{U,i} \stackrel{\text{def}}{=} \mathcal{H}_{U,i}[s_U] & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \overline{G} & \longrightarrow & G & \longrightarrow & H \longrightarrow 1 \end{array}$$

where the vertical maps are surjective. Let \tilde{G} be the pull-back of the group extension $G \twoheadrightarrow H$ by the surjective homomorphism $G_k \twoheadrightarrow H$. Thus, we have an exact sequence

$1 \rightarrow \overline{G} \rightarrow \widetilde{G} \rightarrow G_k \rightarrow 1$, and $\mathcal{H}_{U,i} = \varprojlim_{\widetilde{G}} \widetilde{G}$. Given a (geometrically finite) quotient $\mathcal{H}_{U,i} \twoheadrightarrow \widetilde{G}$ as above, it factorizes as $\mathcal{H}_{U,i} \twoheadrightarrow \pi_1(X_{i,j}^U, \eta_{i,j})^{(\Sigma, \text{ab})} \twoheadrightarrow \widetilde{G}$ for some $j \geq 1$ (cf. Lemma 2.3.2). In particular, the group extension \widetilde{G} splits by our assumption that the group extensions $\pi_1(X_{i,j}^U, \eta_{i,j})^{(\Sigma, \text{ab})}$ split, $\forall j \geq 1$. The set $\text{Sect}(G_k, \mathcal{H}_{U,i})$ of continuous splittings of the group extension $\mathcal{H}_{U,i}$ is naturally identified with the inverse limit $\varprojlim_{\widetilde{G}} \text{Sect}(G_k, \widetilde{G})$ of the sets of continuous splittings of the group extensions \widetilde{G} as above. For a (geometrically finite) quotient \widetilde{G} of $\mathcal{H}_{U,i}$ as above the set $\text{Sect}(G_k, \widetilde{G})$ is non-empty (cf. above discussion) and is, up to conjugation by elements of \overline{G} , a torsor under the Galois cohomology group $H^1(G_k, \overline{G})$, which is finite by our assumption that k satisfies the condition (\mathbf{H}_Σ) . Thus, the set $\text{Sect}(G_k, \widetilde{G})$ is a non-empty finite set. Hence the set $\text{Sect}(G_k, \mathcal{H}_{U,i})$ is non-empty being the projective limit of non-empty finite sets. \square

For $j \geq 1$, let $J_{i,j}^U \stackrel{\text{def}}{=} \text{Pic}_k^0(X_{i,j}^U)$ be the jacobian of $X_{i,j}^U$, and $(J_{i,j}^1)^U \stackrel{\text{def}}{=} \text{Pic}_k^1(X_{i,j}^U)$. Thus, $(J_{i,j}^1)^U$ is a torsor under $J_{i,j}^U$.

Lemma 2.3.5. *The group extension $1 \rightarrow \pi_1(\overline{X}_{i,j}^U, \overline{\eta}_{i,j})^{\Sigma, \text{ab}} \rightarrow \pi_1(X_{i,j}^U, \eta_{i,j})^{(\Sigma, \text{ab})} \rightarrow G_k \rightarrow 1$ splits if and only if the class of $(J_{i,j}^1)^U$ in $H^1(G_k, J_{i,j}^U)$ lies in the maximal Σ -divisible subgroup of $H^1(G_k, J_{i,j}^U)$, i.e., the maximal subgroup of $H^1(G_k, J_{i,j}^U)$ which is divisible by integers whose prime factors are in Σ .*

Proof. This is a well known fact (cf. [Harari-Szamuely], Theorem 1.2, and Proposition 2.1). \square

Proposition 2.3.6. *We use the same notations as above. Assume that $i \geq 1$, and k satisfies the condition (\mathbf{H}_Σ) . Then the section $s_U : G_k \rightarrow \Pi_U^{(i-\text{sol})}$ lifts to a section $\tilde{s}_U : G_k \rightarrow \Pi_U^{(i+1-\text{sol})}$ of the projection $\Pi_U^{(i+1-\text{sol})} \twoheadrightarrow G_k$ if and only if the class of $(J_{i,j}^1)^U$ in $H^1(G_k, J_{i,j}^U)$ lies in the maximal Σ -divisible subgroup of $H^1(G_k, J_{i,j}^U)$, $\forall j \geq 1$.*

Proof. Follows from Lemmas 2.3.4 and 2.3.5. \square

Recall the section $s : G_k \rightarrow G_X^{(i-\text{sol})}$ of the projection $G_X^{(i-\text{sol})} \twoheadrightarrow G_k$ which induces the section $s_U : G_k \rightarrow \Pi_U^{(i-\text{sol})}$ of $\Pi_U^{(i-\text{sol})} \twoheadrightarrow G_k$, $\forall U \subseteq X$ open (cf. 2.2).

Lemma 2.3.7. *Assume that k satisfies the condition (\mathbf{H}_Σ) . Then the section $s : G_k \rightarrow G_X^{(i-\text{sol})}$ lifts to a section $\tilde{s} : G_k \rightarrow G_X^{(i+1-\text{sol})}$ of the projection $G_X^{(i+1-\text{sol})} \twoheadrightarrow G_k$ if and only if for each nonempty open subscheme $U \subseteq X$ the section $s_U : G_k \rightarrow \Pi_U^{(i-\text{sol})}$ lifts to a section $\tilde{s}_U : G_k \rightarrow \Pi_U^{(i+1-\text{sol})}$ of the projection $\Pi_U^{(i+1-\text{sol})} \twoheadrightarrow G_k$.*

Proof. Similar to the proof of Lemma 2.3.4, using the facts that k satisfies the condition (\mathbf{H}_Σ) , and $G_X^{(i-\text{sol})} \simeq \varprojlim_U \Pi_U^{(i-\text{sol})}$. \square

The following is our main result in this section.

Theorem 2.3.8. *We use the same notations as above. Assume that $i \geq 1$, and k satisfies the condition (\mathbf{H}_Σ) . Then the section $s : G_k \rightarrow G_X^{(i-\text{sol})}$ lifts to a section $\tilde{s} : G_k \rightarrow G_X^{(i+1-\text{sol})}$ of the projection $G_X^{(i+1-\text{sol})} \twoheadrightarrow G_k$ if and only if $\forall U \subseteq X$*

nonempty open subscheme the class of $(J_{i,j}^1)^U$ in $H^1(G_k, J_{i,j}^U)$ lies in the maximal Σ -divisible subgroup of $H^1(G_k, J_{i,j}^U)$, $\forall j \geq 1$.

Proof. Follows from Lemmas 2.3.4, 2.3.5, and 2.3.7. \square

§3. Lifting of sections to cuspidally $2/p$ -th step prosolvable arithmetic fundamental groups. In this section we investigate a certain mod- p variant of the cuspidalisation problem investigated in §2 (as well as in [Saïdi]). Throughout §3 we use the same notations as in §2. Let $p \in \Sigma$ be a prime integer.

3.1. Let $U \subseteq X$ be a nonempty open subscheme. Let $i \geq 0$, $t \geq 1$, be integers, and $I_{U,i/p^t}$ the maximal i/p^t -th step prosolvable quotient of I_U (cf. 1.2). By pushing the exact sequence (2.2) by the surjective homomorphism $I_U \twoheadrightarrow I_{U,i/p^t}$ we obtain an exact sequence $1 \rightarrow I_{U,i/p^t} \rightarrow \Pi_U^{(i/p^t-\text{sol})} \rightarrow \Pi_X \rightarrow 1$. We shall refer to $\Pi_U^{(i/p^t-\text{sol})}$ as the maximal (geometrically) **cuspidally i/p^t -th step prosolvable quotient** of Π_U (with respect to the surjection $\Pi_U \twoheadrightarrow \Pi_X$). We have a commutative diagram of exact sequence.

$$(3.1) \quad \begin{array}{ccccccc} & & 1 & & 1 & & \\ & & \downarrow & & \downarrow & & \\ & & I_{U,i/p^t} & \xlongequal{\quad} & I_{U,i/p^t} & & \\ & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & \Delta_{U,i/p^t} & \longrightarrow & \Pi_U^{(i/p^t-\text{sol})} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \\ & & 1 & & 1 & & \end{array}$$

Similarly, by pushing the exact sequence $1 \rightarrow \mathcal{I}_X \rightarrow G_X \rightarrow \Pi_X \rightarrow 1$ by the surjective homomorphism $\mathcal{I}_X \twoheadrightarrow \mathcal{I}_{X,i/p^t}$ we obtain an exact sequence $1 \rightarrow \mathcal{I}_{X,i/p^t} \rightarrow G_X^{(i/p^t-\text{sol})} \rightarrow \Pi_X \rightarrow 1$. We will refer to $G_X^{(i/p^t-\text{sol})}$ as the maximal (geometrically) **cuspidally i/p^t -th step prosolvable quotient** of G_X (with respect to the surjective homomorphism $G_X \twoheadrightarrow \Pi_X$). There exist natural isomorphisms

$$G_X^{(i/p^t-\text{sol})} \xrightarrow{\sim} \varprojlim_U \Pi_U^{(i/p^t-\text{sol})}, \quad \mathcal{I}_{X,i/p^t} \xrightarrow{\sim} \varprojlim_U I_{U,i/p^t},$$

where the limits are over all open subschemes $U \subseteq X$, and a commutative diagram.

$$\begin{array}{ccccccc}
& & 1 & & 1 & & \\
& & \downarrow & & \downarrow & & \\
& & \mathcal{I}_{X,i/p^t} & \xlongequal{\quad} & \mathcal{I}_{X,i/p^t} & & \\
& & \downarrow & & \downarrow & & \\
(3.2) & 1 & \longrightarrow & G_{\overline{X},i/p^t} & \longrightarrow & G_X^{(i/p^t-\text{sol})} & \longrightarrow & G_k & \longrightarrow & 1 \\
& & \downarrow & & \downarrow & & & \downarrow & & \\
& 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \longrightarrow & G_k & \longrightarrow & 1 \\
& & \downarrow & & \downarrow & & & & & \\
& & 1 & & 1 & & & & &
\end{array}$$

3.2. Assume that the lower horizontal exact sequence in diagram (3.2) splits. Let $s : G_k \rightarrow \Pi_X$ be a section of the projection $\Pi_X \rightarrow G_k$.

The lifting problem to sections of cuspidally $i + 1/p^t$ -th step prosolvable arithmetic fundamental groups. Let $i \geq 0$, $t \geq 1$, be integers. Given a section $s : G_k \rightarrow \Pi_X$ as above is it possible to construct a section $s_{U,i+1} : G_k \rightarrow \Pi_U^{(i+1/p^t-\text{sol})}$ of the projection $\Pi_U^{(i+1/p^t-\text{sol})} \rightarrow G_k$ which **lifts** the section s , i.e., which fits in a commutative diagram

$$\begin{array}{ccc}
G_k & \xrightarrow{s_{U,i+1}} & \Pi_U^{(i+1/p^t-\text{sol})} \\
\text{id} \downarrow & & \downarrow \\
G_k & \xrightarrow{s} & \Pi_X
\end{array}$$

where the right vertical map is the natural surjection?

Similarly, is it possible to construct a section $s_{i+1} : G_k \rightarrow G_X^{(i+1/p^t-\text{sol})}$ of the projection $G_X^{(i+1/p^t-\text{sol})} \rightarrow G_k$ which **lifts** the section s , i.e., which fits in a commutative diagram

$$\begin{array}{ccc}
G_k & \xrightarrow{s_{i+1}} & G_X^{(i+1/p^t-\text{sol})} \\
\text{id} \downarrow & & \downarrow \\
G_k & \xrightarrow{s} & \Pi_X
\end{array}$$

where the right vertical map is the natural surjection?

3.3. The quotients $G_X \rightarrow G_X^{(p,i+1)}$, $\Pi_U \rightarrow \Pi_U^{(p,i+1)}$, and lifting of sections. Next, recall the notations in 1.2 and the discussion therein, especially the definition of the quotient $\Delta_U \rightarrow \Delta_U^{p,i+1}$ (cf. the discussion after Lemma 1.2.3). The kernel of the surjective homomorphism $\Delta_U \rightarrow \Delta_U^{p,i+1}$ is a normal subgroup of Π_U (as one easily verifies). Write $\Pi_U^{(p,i+1)} \stackrel{\text{def}}{=} \Pi_U / \text{Ker}(\Delta_U \rightarrow \Delta_U^{p,i+1})$. Thus, we have an exact sequence

$$(3.3) \quad 1 \rightarrow \Delta_U^{p,i+1} \rightarrow \Pi_U^{(p,i+1)} \rightarrow G_k \rightarrow 1.$$

Recall the exact sequence $1 \rightarrow \varprojlim_{j \geq 1} ((\Delta'_{i+1,j})^U)_{i/p} \rightarrow \Delta_U^{p,i+1} \rightarrow \Delta_{U,1/p^{i+1}} \rightarrow 1$

(cf. loc. cit.). The quotient $\Pi_U \twoheadrightarrow \Pi_U^{(1/p^{i+1}\text{-sol})}$ (cf. 3.1) factorizes through $\Pi_U \twoheadrightarrow \Pi_U^{(p,i+1)}$ (cf. exact sequence (1.7)), and we have a commutative diagram of exact sequences.

$$(3.4) \quad \begin{array}{ccccccc} & & 1 & & 1 & & \\ & & \downarrow & & \downarrow & & \\ & & \varprojlim_{j \geq 1} ((\Delta'_{i+1,j})^U)_{i/p} & \xlongequal{\quad} & \varprojlim_{j \geq 1} ((\Delta'_{i+1,j})^U)_{i/p} & & \\ & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & \Delta_U^{p,i+1} & \longrightarrow & \Pi_U^{(p,i+1)} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \Delta_{U,1/p^{i+1}} & \longrightarrow & \Pi_U^{(1/p^{i+1}\text{-sol})} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \\ & & 1 & & 1 & & \end{array}$$

Similarly, $\text{Ker}(G_{\overline{X}} \twoheadrightarrow G_{\overline{X}}^{p,i+1})$ is a normal subgroup of G_X , and we have an exact sequence

$$(3.5) \quad 1 \rightarrow G_{\overline{X}}^{p,i+1} \rightarrow G_X^{(p,i+1)} \rightarrow G_k \rightarrow 1,$$

where $G_X^{(p,i+1)} \stackrel{\text{def}}{=} G_X / \text{Ker}(G_{\overline{X}} \twoheadrightarrow G_{\overline{X}}^{p,i+1})$. The exact sequence (1.8) induces a commutative diagram of exact sequences.

$$(3.6) \quad \begin{array}{ccccccc} & & 1 & & 1 & & \\ & & \downarrow & & \downarrow & & \\ & & \varprojlim_U (\varprojlim_{j \geq 1} ((\Delta'_{i+1,j})^U)_{i/p}) & \xlongequal{\quad} & \varprojlim_U (\varprojlim_{j \geq 1} ((\Delta'_{i+1,j})^U)_{i/p}) & & \\ & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & G_{\overline{X}}^{p,i+1} & \longrightarrow & G_X^{(p,i+1)} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & G_{\overline{X},1/p^{i+1}} & \longrightarrow & G_X^{(1/p^{i+1}\text{-sol})} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \\ & & 1 & & 1 & & \end{array}$$

Furthermore, $\Pi_U^{(i+1/p\text{-sol})}$ (resp. $G_X^{(i+1/p\text{-sol})}$) is a quotient of $\Pi_U^{(p,i+1)}$ (resp. of

$G_X^{(p,i+1)}$) (cf. Lemmas 1.2.4 and 1.2.5) and we have commutative diagrams

$$(3.7) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \Delta_U^{p,i+1} & \longrightarrow & \Pi_U^{(p,i+1)} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \Delta_{U,i+1/p} & \longrightarrow & \Pi_U^{(i+1/p-\text{sol})} & \longrightarrow & G_k \longrightarrow 1 \end{array}$$

resp.

$$(3.8) \quad \begin{array}{ccccccc} 1 & \longrightarrow & G_{\overline{X}}^{p,i+1} & \longrightarrow & G_X^{(p,i+1)} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & G_{\overline{X},i+1/p} & \longrightarrow & G_X^{(i+1/p-\text{sol})} & \longrightarrow & G_k \longrightarrow 1 \end{array}$$

where the left and middle vertical maps are surjective.

The lifting problem to sections of $\Pi_U^{(p,i+1)}$, and $G_X^{(p,i+1)}$. Given a section $s : G_k \rightarrow \Pi_X$ of the projection $\Pi_X \twoheadrightarrow G_k$ as in 3.2, is it possible to construct a section $\tilde{s}_{U,i+1} : G_k \rightarrow \Pi_U^{(p,i+1)}$ of the projection $\Pi_U^{(p,i+1)} \twoheadrightarrow G_k$ which **lifts** the section s , i.e., which fits in a commutative diagram

$$\begin{array}{ccc} G_k & \xrightarrow{\tilde{s}_{U,i+1}} & \Pi_U^{(p,i+1)} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \Pi_X \end{array}$$

where the right vertical map is the natural surjection?

Similarly, is it possible to construct a section $\tilde{s}_{i+1} : G_k \rightarrow G_X^{(p,i+1)}$ of the projection $G_X^{(p,i+1)} \twoheadrightarrow G_k$ which **lifts** the section s , i.e., which fits in a commutative diagram

$$\begin{array}{ccc} G_k & \xrightarrow{\tilde{s}_{i+1}} & G_X^{(p,i+1)} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \Pi_X \end{array}$$

where the right vertical map is the natural surjection?

Lemma 3.3.1. A positive answer to the lifting problem posed in 3.3 implies a positive answer to the lifting problem posed in 3.2 in the case $t = 1$.

Proof. Follows immediately from the above commutative diagrams (3.7) and (3.8). \square

3.4. In this section we investigate the lifting problem posed in 3.3 in the case $i = 1$, and draw consequences for the lifting problem posed in 3.2 in the case $t = i = 1$ (the only case we need for applications in §4). Let $s : G_k \rightarrow \Pi_X$ be a section of the projection $\Pi_X \twoheadrightarrow G_k$.

3.4.I. First, we investigate the problem of lifting the section s to a section $s'_{U,i+1} : G_k \rightarrow \Pi_U^{(1/p^{i+1}-\text{sol})}$ (resp. $s'_{i+1} : G_k \rightarrow G_X^{(1/p^{i+1}-\text{sol})}$) of the projection $\Pi_U^{(1/p^{i+1}-\text{sol})} \twoheadrightarrow$

G_k (resp. $G_X^{(1/p^{i+1}\text{-sol})} \twoheadrightarrow G_k$). Recall the notations in 2.3, and ($\forall U \subseteq X$ open) the sequence of characteristic open subgroups

$\dots \subseteq \Delta_X[j+1] \stackrel{\text{def}}{=} \Delta_{U,0}[j+1] \subseteq \Delta_X[j] \stackrel{\text{def}}{=} \Delta_{U,0}[j] \subseteq \dots \subseteq \Delta_X[1] \stackrel{\text{def}}{=} \Delta_{U,0}[1] = \Delta_X$
of $\Delta_{U,0} = \Delta_X$ with $\bigcap_{j \geq 1} \Delta_X[j] = \{1\}$. The sequence of (geometrically characteristic) open subgroups

$\dots \subseteq \Pi_X[j+1] \stackrel{\text{def}}{=} \Pi_{U,0}[j+1] \subseteq \Pi_X[j] \stackrel{\text{def}}{=} \Pi_{U,0}[j] \subseteq \dots \subseteq \Pi_X[1] \stackrel{\text{def}}{=} \Pi_{U,0}[1] = \Pi_X,$

where $\Pi_X[j] \stackrel{\text{def}}{=} \Delta_X[j].s(G_k)$, corresponds to a tower of finite (not necessarily Galois) étale covers

$$\dots \rightarrow X_{j+1} \stackrel{\text{def}}{=} X_{0,j+1}^U \rightarrow X_j \stackrel{\text{def}}{=} X_{0,j}^U \rightarrow \dots \rightarrow X \stackrel{\text{def}}{=} X_{0,1}^U.$$

Note that $\Pi_X[j]$ identifies naturally with $\Pi_{X_j} \stackrel{\text{def}}{=} \pi_1(X_j, \eta_j)^{(\Sigma)}$, where η_j is the base point induced by η . Moreover, the section s restricts to a section $s : G_k \rightarrow \Pi_{X_j}$ of the projection $\Pi_{X_j} \twoheadrightarrow G_k$, $\forall j \geq 1$.

Let $i \geq 0$, $j \geq 1$, be integers. Recall the Kummer sequence

$$1 \rightarrow \mu_{p^{i+1}} \rightarrow \mathbb{G}_m \xrightarrow{p^{i+1}} \mathbb{G}_m \rightarrow 1$$

in étale topology, which induces an exact sequence

$$0 \rightarrow \text{Pic}(X_j)/p^{i+1} \text{Pic}(X_j) \rightarrow H^2(X_j, \mu_{p^{i+1}}) \rightarrow_{p^{i+1}} \text{Br}(X_j) \rightarrow 0.$$

Here $\text{Pic} \stackrel{\text{def}}{=} H_{\text{et}}^1(\cdot, \mathbb{G}_m)$ is the Picard group, $\text{Br} \stackrel{\text{def}}{=} H_{\text{et}}^2(\cdot, \mathbb{G}_m)$ the Brauer-Grothendieck cohomological group, and ${}_{p^{i+1}}\text{Br} \subseteq \text{Br}$ the subgroup of Br which is annihilated by p^{i+1} . We identify $\text{Pic}(X_j)/p^{i+1} \text{Pic}(X_j)$ with its image in $H^2(X_j, \mu_{p^{i+1}})$ and refer to it as the Picard part of $H^2(X_j, \mu_{p^{i+1}})$. By pulling back cohomology classes via the section $s : G_k \rightarrow \Pi_{X_j}$, and bearing in mind the natural identification $H^2(\Pi_{X_j}, \mu_{p^{i+1}}) \xrightarrow{\sim} H^2(X_j, \mu_{p^{i+1}})$ (cf. [Mochizuki], Proposition 1.1), we obtain a restriction homomorphism $s_j^* : H^2(X_j, \mu_{p^{i+1}}) \rightarrow H^2(G_k, \mu_{p^{i+1}})$.

Observe that if k'/k is a finite extension, and $X_{k'} \stackrel{\text{def}}{=} X \times_k k'$, then we have a cartesian diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_{X_{k'}} & \longrightarrow & \Pi_{X_{k'}} & \longrightarrow & G_{k'} \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \longrightarrow & G_k \longrightarrow 1 \end{array}$$

and the section s induces a section $s_{k'} : G_{k'} \rightarrow \Pi_{X_{k'}}$ of the projection $\Pi_{X_{k'}} \twoheadrightarrow G_{k'}$.

Definition 3.4.1 (Sections with Cycle Classes Orthogonal to Pic mod- p^{i+1}). (Compare with [Saïdi], 1.4.)

(i) We say that the section s has a **cycle class orthogonal to Pic mod- p^{i+1}** if the homomorphism $s_j^* : H^2(X_j, \mu_{p^{i+1}}) \rightarrow H^2(G_k, \mu_{p^{i+1}})$ annihilates the Picard part $\text{Pic}(X_j)/p^{i+1} \text{Pic}(X_j)$ of $H^2(X_j, \mu_{p^{i+1}})$, $\forall j \geq 1$.

(ii) We say that the section s has a **cycle class uniformly orthogonal to Pic mod- p^{i+1}** (relative to the system of neighbourhoods $\{X_j\}_{j \geq 1}$ of s) if, for every finite extension k'/k , the induced section $s_{k'} : G_{k'} \rightarrow \Pi_{X_{k'}}$ has a cycle class orthogonal to Pic mod- p^{i+1} (relative to the system of neighbourhoods of $s_{k'}$ which is induced by the $\{X_j\}_{j \geq 1}$).

Definition 3.4.2. We say that the field k satisfies the condition $(\mathbf{H}_{\mathbf{p}^{i+1}})$ if the following holds. The Galois cohomology groups $H^1(G_k, M)$ are *finite* for every finite G_k -module M annihilated by p^{i+1} .

Theorem 3.4.3 (Lifting of Sections to Cuspidally mod- p^{i+1} abelian Arithmetic Fundamental Groups). *Assume that k satisfies the condition $(\mathbf{H}_{\mathbf{p}^{i+1}})$ (cf. Definition 3.4.2). Let $s : G_k \rightarrow \Pi_X$ be a section of the projection $\Pi_X \twoheadrightarrow G_k$. Assume that s has a **cycle class uniformly orthogonal to Pic mod- p^{i+1}** (cf. Definition 3.4.1(ii)). Let $U \subseteq X$ be a nonempty open subscheme. Then there exists a section $s'_{U,i+1} : G_k \rightarrow \Pi_U^{(1/p^{i+1}-\text{sol})}$ of the projection $\Pi_U^{(1/p^{i+1}-\text{sol})} \twoheadrightarrow G_k$ which **lifts** the section s , i.e., which inserts into the following commutative diagram.*

$$\begin{array}{ccc} G_k & \xrightarrow{s'_{U,i+1}} & \Pi_U^{(1/p^{i+1}-\text{sol})} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \Pi_X \end{array}$$

Proof. Similar to the proof of Theorem 2.3.3 in [Saïdi]. \square

Theorem 3.4.4 (Lifting of Sections to Cuspidally mod- p^{i+1} abelian Galois Groups). *Assume that the field k satisfies the condition $(\mathbf{H}_{\mathbf{p}^{i+1}})$ (cf. Definition 3.4.2). Let $s : G_k \rightarrow \Pi_X$ be a section of the projection $\Pi_X \twoheadrightarrow G_k$. Then s has a **cycle class uniformly orthogonal to Pic mod- p^{i+1}** (cf. Definition 3.4.1(ii)) **if and only if** there exists a section $s'_{i+1} : G_k \rightarrow G_X^{(1/p^{i+1}-\text{sol})}$ of the projection $G_X^{(1/p^{i+1}-\text{sol})} \twoheadrightarrow G_k$ which **lifts** the section s , i.e., which inserts in the following commutative diagram.*

$$\begin{array}{ccc} G_k & \xrightarrow{s'_{i+1}} & G_X^{(1/p^{i+1}-\text{sol})} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \Pi_X \end{array}$$

Proof. Similar to the proof of Theorem 2.3.5 in [Saïdi]. \square

3.4.II. Next, let $s' : G_k \rightarrow G_X^{(1/p^2-\text{sol})}$ be a section of the projection $G_X^{(1/p^2-\text{sol})} \twoheadrightarrow G_k$, which induces for every open subscheme $U \subseteq X$ a section $s'_U : G_k \rightarrow \Pi_U^{(1/p^2-\text{sol})}$ of the projection $\Pi_U^{(1/p^2-\text{sol})} \twoheadrightarrow G_k$. We investigate the problem of lifting the section s'_U (resp. s') to a section $\tilde{s}_U : G_k \rightarrow \Pi_U^{(p,2)}$ (resp. $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$) of the projection $\Pi_U^{(p,2)} \twoheadrightarrow G_k$ (resp. $G_X^{(p,2)} \twoheadrightarrow G_k$) (cf. diagrams (3.4) and (3.6)). Let $U \subseteq X$ be an open subscheme. Consider the following pull-back diagram.

$$(3.9) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \varprojlim_{j \geq 1} ((\Delta'_{2,j})^U)_{1/p} & \longrightarrow & \mathcal{H}_U^{(p,2)} & \longrightarrow & G_k & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & s'_U \downarrow & & \\ 1 & \longrightarrow & \varprojlim_{j \geq 1} ((\Delta'_{2,j})^U)_{1/p} & \longrightarrow & \Pi_U^{(p,2)} & \longrightarrow & \Pi_U^{(1/p^2-\text{sol})} & \longrightarrow & 1 \end{array}$$

Lemma 3.4.5. *The section s'_U lifts to a section $\tilde{s}_U : G_k \rightarrow \Pi_U^{(p,2)}$ of the projection $\Pi_U^{(p,2)} \rightarrow G_k$ if and only if the group extension $\mathcal{H}_U^{(p,2)}$ splits.*

Proof. Follows immediately from diagram (3.9). \square

Recall the discussion and notations after Lemma 1.2.2, especially the definition of the $\{\Delta_{U,1/p^{i+1}}[j]\}_{j \geq 1}$. For $j \geq 1$, write $\Pi_{U,1/p^2}[j] \stackrel{\text{def}}{=} \Delta_{U,1/p^2}[j] \cdot s'_U(G_k)$. Thus, $\Pi_{U,1/p^2}[j] \subseteq \Pi_U^{(1/p^2\text{-sol})}$ is an open subgroup corresponding to a (possibly tamely ramified) cover $\tilde{X}_j^U \rightarrow X$ between smooth, proper, and geometrically connected k -curves. The geometric point η determines a geometric point η_j of \tilde{X}_j^U . Write $\Pi_j^U = \Pi_j^U[s'_U] \stackrel{\text{def}}{=} \Pi_{\tilde{X}_j^U} \stackrel{\text{def}}{=} \pi_1(\tilde{X}_j^U, \eta_j)^{(\Sigma)}$, which inserts in the exact sequence $1 \rightarrow \Delta_j^U \rightarrow \Pi_j^U \rightarrow G_k \rightarrow 1$, where $\Delta_j^U \stackrel{\text{def}}{=} \Delta_{\tilde{X}_j^U \times_k \bar{k}}$. Further, consider the push-out diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_j^U & \longrightarrow & \Pi_j^U & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & (\Delta_j^U)_{1/p} & \longrightarrow & (\Pi_j^U)^{(1/p\text{-sol})} & \longrightarrow & G_k \longrightarrow 1 \end{array}$$

which defines the geometrically $1/p$ -th step solvable quotient $\Pi_j^U \twoheadrightarrow (\Pi_j^U)^{(1/p\text{-sol})}$ of Π_j^U .

Lemma 3.4.6. *There are natural isomorphisms $\varinjlim_{j \geq 1} ((\Delta'_{2,j})^U)_{1/p} \xrightarrow{\sim} \varinjlim_{j \geq 1} (\Delta_j^U)_{1/p}$, and $\mathcal{H}_U^{(p,2)} \xrightarrow{\sim} \varinjlim_{j \geq 1} (\Pi_j^U)^{(1/p\text{-sol})}$.*

Proof. Follows from the various definitions. \square

Note that $(\Delta'_{2,j})^U = \Delta_j^U$, we will in the sequel write Δ_j^U instead of $(\Delta'_{2,j})^U$.

Lemma 3.4.7. *Assume that k satisfies the condition (\mathbf{H}_p) (cf. Definition 3.4.2). Then the group extension $\mathcal{H}_U^{(p,2)}$ splits if and only if the group extension $(\Pi_j^U)^{(1/p\text{-sol})}$ splits, $\forall j \geq 1$.*

Proof. Similar to the proof of Lemma 2.3.4, using the fact that $H^1(G_k, (\Delta_j^U)_{1/p})$ is finite if k satisfies (\mathbf{H}_p) . \square

For $j \geq 1$, let $J_j[U] \stackrel{\text{def}}{=} \text{Pic}_k^0(\tilde{X}_j^U)$ be the jacobian of \tilde{X}_j^U , and $J_j^1[U] \stackrel{\text{def}}{=} \text{Pic}_k^1(\tilde{X}_j^U)$.

Lemma 3.4.8. *The group extension $(\Pi_j^U)^{(1/p\text{-sol})}$ splits if and only if the class of $J_j^1[U]$ in $H^1(G_k, J_j[U])$ is divisible by p .*

Proof. This fact is well-known, see [Harari-Szamuely] for instance. Strictly speaking loc. cit. treats the splittings of the group extension $(\Pi_j^U)^{(\text{ab})}$ = the geometrically abelian quotient of Π_j^U , but a similar argument leads to a mod- p variant as above for any prime $p \in \Sigma$. \square

Theorem 3.4.9. *With the above notations, assume that k satisfies the condition (\mathbf{H}_p) (cf. Definition 3.4.2). Then the section s'_U lifts to a section $\tilde{s}_U : G_k \rightarrow \Pi_U^{(p,2)}$ of the projection $\Pi_U^{(p,2)} \twoheadrightarrow G_k$ **if and only if** the class of $J_j^1[U]$ in $H^1(G_k, J_j[U])$ is divisible by p , $\forall j \geq 1$.*

Proof. Follows from Lemmas 3.4.5, 3.4.7, and 3.4.8. \square

Theorem 3.4.10. *With the above notations, assume that k satisfies the condition (\mathbf{H}_p) (cf. Definition 3.4.2). Then the section s' lifts to a section $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$ of the projection $G_X^{(p,2)} \twoheadrightarrow G_k$ **if and only if** the class of $J_j^1[U]$ in $H^1(G_k, J_j[U])$ is divisible by p , $\forall j \geq 1$, and $\forall U \subseteq X$ nonempty open subscheme as in the above discussion.*

Proof. Similar to the proof of Theorem 3.4.9. \square

The following is our main result in this section.

Theorem 3.4.11. *With the above notations, assume that the field k satisfies the condition (\mathbf{H}_{p^2}) (cf. Definition 3.4.2). Let $s : G_k \rightarrow \Pi_X$ be a section of the projection $\Pi_X \twoheadrightarrow G_k$. Then s lifts to a section $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$ (resp. $s' : G_k \rightarrow G_X^{(2/p)}$) of the projection $G_X^{(p,2)} \twoheadrightarrow G_k$ (resp. $G_X^{(2/p)} \twoheadrightarrow G_k$) **if and only if** (resp. **if**) the following two conditions occur.*

(i) *The section s has a cycle class uniformly orthogonal to $\text{Pic mod-}p^2$ (cf. Definition 3.4.1 (ii))*

(ii) *There exists a section $s' : G_k \rightarrow G_X^{(1/p^2-\text{sol})}$ of the projection $G_X^{(1/p^2-\text{sol})} \twoheadrightarrow G_k$ which **lifts** the section s (this holds if (i) holds by Theorem 3.4.4) such that the class of $J_j^1[U]$ in $H^1(G_k, J_j[U])$ is **divisible** by p , $\forall j \geq 1$, and $\forall U \subseteq X$ nonempty open subscheme.*

Proof. Follows from Theorems 3.4.4 and 3.4.10. The resp. assertion follows from diagram (3.8) (cf. Lemma 3.3.1). \square

§4. Geometric sections of arithmetic fundamental groups of p -adic curves.

In this section, applying the results in §3, we provide a characterisation of sections of (geometrically pro- Σ , $p \in \Sigma$) arithmetic fundamental groups of p -adic curves which arise from rational points. We use the notations in §2 and §3.

Let p be a prime integer. In this section k is a p -adic local field, i.e., k/\mathbb{Q}_p is a finite extension, and we assume $p \in \Sigma$. Let $s : G_k \rightarrow \Pi_X$ be a section of the projection $\Pi_X \twoheadrightarrow G_k$.

Definition 4.1. We say that the section s is *geometric* if the image $s(G_k)$ of s is contained (hence equal to) in the decomposition group $D_x \subset \Pi_X$ associated to a rational point $x \in X(k)$.

Recall the tower of finite étale covers $\dots \rightarrow X_{t+1} \rightarrow X_t \rightarrow \dots \rightarrow X_1 = X$ in 3.4.I, and the section $s : G_k \rightarrow \Pi_{X_t}$ of the projection $\Pi_{X_t} \twoheadrightarrow G_k$ induced by s , $\forall t \geq 1$. Assume that the section $s : G_k \rightarrow \Pi_X$ has a cycle class uniformly orthogonal to $\text{Pic mod-}p^2$ (cf. Definition 3.4.1). In particular, the induced section $s : G_k \rightarrow \Pi_{X_t}$ also has a cycle class uniformly orthogonal to $\text{Pic mod-}p^2$ (cf. loc. cit.). There exists, $\forall t \geq 1$, a section

$$s'_t : G_k \rightarrow G_{X_t}^{(1/p^2-\text{sol})}$$

of the projection $G_{X_t}^{(1/p^2-\text{sol})} \rightarrow G_k$ which lifts the section s (cf. Theorem 3.4.4). (Note that k satisfies the condition (\mathbf{H}_{p^2}) .) Given integers $t_1 \geq t_2 \geq 1$, we have a commutative diagram

$$\begin{array}{ccc} G_{X_{t_1}}^{(1/p^2-\text{sol})} & \longrightarrow & G_k \\ \downarrow & & \text{id} \downarrow \\ G_{X_{t_2}}^{(1/p^2-\text{sol})} & \longrightarrow & G_k \end{array}$$

where the left vertical map is induced by the scheme morphism $X_{t_1} \rightarrow X_{t_2}$. We say that the above sections $\{s'_t\}_{t \geq 1}$ are **compatible** if $\forall t_1 \geq t_2 \geq 1$ we have a commutative diagram.

$$\begin{array}{ccc} G_k & \xrightarrow{s'_{t_1}} & G_{X_{t_1}}^{(1/p^2-\text{sol})} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s'_{t_2}} & G_{X_{t_2}}^{(1/p^2-\text{sol})} \end{array}$$

Lemma 4.2. *With the above notations, let $s' = s'_1 : G_k \rightarrow G_X^{(1/p^2-\text{sol})}$ be a section of the projection $G_X^{(1/p^2-\text{sol})} \rightarrow G_k$ which lifts the section s . Then s' induces naturally compatible sections $s'_t : G_k \rightarrow G_{X_t}^{(1/p^2-\text{sol})}$ of the projections $G_{X_t}^{(1/p^2-\text{sol})} \rightarrow G_k$ which lift the section s , $\forall t \geq 1$.*

Proof. Follows from the fact that we have a commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathcal{I}_{X_t} & \longrightarrow & G_{X_t} & \longrightarrow & \Pi_{X_t} \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \mathcal{I}_X & \longrightarrow & G_X & \longrightarrow & \Pi_X \longrightarrow 1 \end{array}$$

where the right square is cartesian, and $\mathcal{I}_{X_t} = \mathcal{I}_X$. In particular, $\mathcal{I}_{X_t, 1/p^2} = \mathcal{I}_{X, 1/p^2}$ and $G_{X_t}^{(1/p^2-\text{sol})}$ is the pull back of the group extension $1 \rightarrow \mathcal{I}_{X, 1/p^2} \rightarrow G_X^{(1/p^2-\text{sol})} \rightarrow \Pi_X \rightarrow 1$ via the natural inclusion $\Pi_{X_t} \hookrightarrow \Pi_X$, $\forall t \geq 1$. \square

Next, recall the exact sequence (cf. diagram (3.6), the case $i = 1$)

$$1 \rightarrow \mathcal{I}_X[p, 2] \stackrel{\text{def}}{=} \varprojlim_U \left(\varinjlim_{j \geq 1} ((\Delta'_{2,j})^U)_{1/p} \right) \rightarrow G_X^{(p,2)} \rightarrow G_X^{(1/p^2-\text{sol})} \rightarrow 1.$$

We have a commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathcal{I}_{X_t}[p, 2] & \longrightarrow & G_{X_t}^{(p,2)} & \longrightarrow & G_{X_t}^{(1/p^2-\text{sol})} \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \mathcal{I}_X[p, 2] & \longrightarrow & G_X^{(p,2)} & \longrightarrow & G_X^{(1/p^2-\text{sol})} \longrightarrow 1 \end{array}$$

where the right square is cartesian (as one easily verifies). In particular, $\mathcal{I}_{X_t}[p, 2] = \mathcal{I}_X[p, 2]$. Let $s' : G_k \rightarrow G_X^{(1/p^2-\text{sol})}$ be a section of the projection $G_X^{(1/p^2-\text{sol})} \rightarrow G_k$ which lifts the section s , and $\{s'_t : G_k \rightarrow G_{X_t}^{(1/p^2-\text{sol})}\}_{t \geq 1}$ the induced compatible

sections as in Lemma 4.2 which lift the section s . Let $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$ be a section of the projection $G_X^{(p,2)} \twoheadrightarrow G_k$ which lifts the section s' . Then \tilde{s} induces sections $\tilde{s}_t : G_k \rightarrow G_{X_t}^{(p,2)}$ of the projections $G_{X_t}^{(p,2)} \twoheadrightarrow G_k$ which lift the section $s'_t, \forall t \geq 1$, and which are compatible in the sense that $\forall t_1 \geq t_2 \geq 1$ integers we have a commutative diagram (cf. above diagram whose right square is cartesian).

$$\begin{array}{ccc} G_k & \xrightarrow{s'_{t_1}} & G_{X_{t_1}}^{(p,2)} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s'_{t_2}} & G_{X_{t_2}}^{(p,2)} \end{array}$$

Also, recall the notations and definitions in 3.4.II, the case $i = 1$, relative to the sections $s'_U : G_k \rightarrow \Pi_U^{(1/p^2-\text{sol})}$ induced by s', \forall nonempty open subscheme $U \subseteq X$. Thus, the $\{\tilde{X}_j^U\}_{j \geq 1}$ are defined in this case $\forall U \subseteq X$ open (cf. loc. cit.); they form a system of neighbourhoods of the section s'_U . Suppose that the group extension $1 \rightarrow \pi_1(\tilde{X}_j^U \times_k \bar{k}, \bar{\eta}_j)^{1/p} \rightarrow \pi_1(\tilde{X}_j^U, \eta_j)^{(1/p)} \rightarrow G_k \rightarrow 1$ splits, or equivalently that the class of $J_j^1[U]$ in $H^1(G_k, J_j[U])$ is divisible by $p, \forall j \geq 1$, and $\forall U \subseteq X$ as above (cf. Lemma 3.4.8). Then the section $s' : G_k \rightarrow G_X^{(1/p^2-\text{sol})}$ lifts to a section $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$ of the projection $G_X^{(p,2)} \twoheadrightarrow G_k$ (cf. Theorem 3.4.10). Moreover, \tilde{s} induces compatible sections $\tilde{s}_t : G_k \rightarrow G_{X_t}^{(p,2)}$ of the projections $G_{X_t}^{(p,2)} \twoheadrightarrow G_k$ which lift the section $s'_t, \forall t \geq 1$ (cf. above discussion). In particular, the above sections \tilde{s}_t induce naturally sections

$$\rho_t : G_k \rightarrow G_{X_t}^{(2/p-\text{sol})}$$

of the projections $G_{X_t}^{(2/p-\text{sol})} \twoheadrightarrow G_k$ which lift the sections $s, \forall t \geq 1$ (cf. Lemma 3.3.1, as well as the diagrams (3.7) and (3.8)).

Definition 4.3. With the above notations, we say that the section s is **admissible** if the following two conditions hold.

A1) The section s has a **cycle class uniformly orthogonal to Pic mod- p^2** .

A2) There **exists** a section $s' : G_k \rightarrow G_X^{(1/p^2-\text{sol})}$ of the projection $G_X^{(1/p^2-\text{sol})} \twoheadrightarrow G_k$ which **lifts** the section s (this holds if condition A1 is satisfied by Theorem 3.4.4) such that the following holds. The class of $J_j^1[U]$ in $H^1(G_k, J_j[U])$ is **divisible** by $p, \forall U \subseteq X$ nonempty open subscheme, and $\forall j \geq 1$. Or, equivalently, the section s' **lifts** to a section $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$ of the projection $G_X^{(p,2)} \twoheadrightarrow G_k$ (cf. Theorem 3.4.10).

Lemma 4.4. *Let k'/k be a finite extension, $X_{k'} \stackrel{\text{def}}{=} X \times_k k'$, and $s_{k'} : G_{k'} \rightarrow \Pi_{X_{k'}}$ the section of the projection $\Pi_{X_{k'}} \twoheadrightarrow G_{k'}$ which is induced by s . Assume s is admissible then $s_{k'}$ is admissible.*

Proof. First, if s has a cycle class uniformly orthogonal to Pic mod- p^2 then so does $s_{k'}$ (cf. Definition 3.4.1(ii)). The second assertion follows from the various definitions. \square

The following is our main result in this section; it provides a characterisation of sections of (geometrically pro- $\Sigma, p \in \Sigma$) arithmetic fundamental groups of p -adic curves which are geometric.

Theorem 4.5. *We use the above notations. The section s is **admissible** (cf. Definition 4.3) **if and only if** s is **geometric** (cf. Definition 4.1).*

Proof. The if part follows easily from the various Definitions. We prove the only if part. Assume that s is admissible, and k contains a primitive p -th root ζ_p of 1. The section $s : G_k \rightarrow \Pi_{X_t}$ lifts to a section $\rho_t : G_k \rightarrow G_{X_t}^{(2/p-\text{sol})}$ of the projection $G_{X_t}^{(2/p-\text{sol})} \twoheadrightarrow G_k$ (cf. discussion before Definition 4.3), $\forall t \geq 1$. The section ρ_t induces a section $\tilde{\rho}_t : (G_k)_{2/p} \rightarrow (G_{X_t})_{2/p}$ of the projection $(G_{X_t})_{2/p} \twoheadrightarrow (G_k)_{2/p}$, where the $(\)_{2/p}$ of the various profinite groups are the second quotients of the $\mathbb{Z}/p\mathbb{Z}$ -derived series (cf. 1.2). The section $\tilde{\rho}_t$ is geometric and arises from a rational point $x_t \in X_t(k)$ by a result of Pop (cf. [Pop]). (Here one uses the fact that $\zeta_p \in k$.) In particular, $X_t(k) \neq \emptyset$, $\forall t \geq 1$. A well-known limit argument shows that s is geometric (cf. [Tamagawa], Proposition 2.1, (iv), see also the details of the proof of Theorem A in [Saïdi1]). In case $\zeta_p \notin k$, let $k' \stackrel{\text{def}}{=} k(\zeta_p)$. The section $s_{k'}$ is admissible (cf. Lemma 4.4), hence is geometric by the above discussion. One then verifies easily that s is geometric (cf. [Saïdi2], proof of Theorem B). \square

In the course of proving Theorem 4.5 we proved the following (cf. discussion before Definition 4.3, and the proof of Theorem 4.5).

Proposition 4.6. *Let $\tilde{s} : G_k \rightarrow G_X^{(p,2)}$ be a section of the projection $G_X^{(p,2)} \twoheadrightarrow G_k$, and $s : G_k \rightarrow \Pi_X$ the section of the projection $\Pi_X \twoheadrightarrow G_k$ which is induced by \tilde{s} . Then s is geometric.*

Remarks 4.7. 1) Theorem 4.5 above is stronger and more precise than Theorem A in [Saïdi1].

2) There are examples of sections $s : G_k \rightarrow \Pi_X$ as above, where $\Sigma = \{p\}$, which are *not* geometric (cf. [Hoshi]). These provide examples of sections s as above which are *not* admissible by Theorem 4.5 (where $\Sigma = \{p\}$). It would be interesting to know which of the conditions A1 and A2 in the definition of admissible sections fail to hold in Hoshi's example. In [Saïdi3] we observe that the section in Hoshi's example is orthogonal to Pic^0 in the sense that the map $s^* : H^2(\Pi_X, \mathbb{Z}_p) \rightarrow H^2(G_k, \mathbb{Z}_p)$ annihilates the image of $\text{Pic}^0(X)$.

The following is an application of our results to the absolute anabelian geometry of p -adic curves.

Theorem 4.8. *Let $p_X, p_Y \in \mathfrak{P}\text{rimes}$, and X (resp. Y) a proper smooth and geometrically connected hyperbolic curve over a p_X -adic local field k_X (respectively, p_Y -adic local field k_Y). Let $p_X \in \Sigma_X$ (resp. $p_Y \in \Sigma_Y$) be a non-empty set of prime integers of cardinality ≥ 2 , Π_X (resp. Π_Y) the geometrically pro- Σ_X (resp. pro- Σ_Y) arithmetic fundamental group of X (resp. Y), and $\varphi : \Pi_X \rightarrow \Pi_Y$ an isomorphism of profinite groups which fits in the following commutative diagram*

$$\begin{array}{ccc} G_X^{(p_X,2)} & \xrightarrow{\tilde{\varphi}} & G_Y^{(p_Y,2)} \\ \downarrow & & \downarrow \\ \Pi_X & \xrightarrow{\varphi} & \Pi_Y \end{array}$$

where $\tilde{\varphi}$ is an isomorphism of profinite groups, and the vertical maps are the natural projections. Then φ is geometric, i.e., arises from a uniquely determined isomorphism of schemes $X \xrightarrow{\sim} Y$.

Proof. The existence of the lifting $\tilde{\varphi}$ of φ implies, by Proposition 4.6, that φ preserves the decomposition groups at closed points. The statement follows then from [Mochizuki1], Corollary 2.9. \square

§5. Local sections of arithmetic fundamental groups of p -adic curves. We prove that a certain class of sections of arithmetic fundamental groups of p -adic curves are (uniformly) orthogonal to Pic^\wedge . We use the notations in §4.

5.1. Arithmetic fundamental groups of formal fibres of p -adic curves. Let \mathcal{O}_k be the valuation ring of k , and $\tilde{X} \rightarrow \text{Spec } \mathcal{O}_k$ a flat and proper model of X over \mathcal{O}_k with \tilde{X} normal. Let $x \in \tilde{X}^{\text{cl}}$ be a closed point, and $\hat{\mathcal{O}}_{\tilde{X},x}$ the completion of the local ring of \tilde{X} at x . We will refer to $\mathcal{X} \stackrel{\text{def}}{=} \text{Spec}(\hat{\mathcal{O}}_{\tilde{X},x} \otimes_{\mathcal{O}_k} k)$ as the *formal fibre* of \tilde{X} at x (or simply a formal fibre). Assume \mathcal{X} is geometrically connected, and write $\bar{\mathcal{X}} \stackrel{\text{def}}{=} \mathcal{X} \times_k \bar{k}$. Let $\bar{\beta}$ be a geometric point of $\bar{\mathcal{X}}$, which determines a geometric point β of \mathcal{X} . Write $\Delta_{\mathcal{X}} \stackrel{\text{def}}{=} \pi_1(\bar{\mathcal{X}}, \bar{\beta})^\Sigma$ for the maximal pro- Σ quotient of $\pi_1(\bar{\mathcal{X}}, \bar{\beta})$, and $\Pi_{\mathcal{X}} \stackrel{\text{def}}{=} \pi_1(\mathcal{X}, \beta) / \text{Ker}(\pi_1(\bar{\mathcal{X}}, \bar{\beta}) \twoheadrightarrow \pi_1(\bar{\mathcal{X}}, \bar{\beta})^\Sigma)$. We have a commutative diagram of exact sequences

$$(5.1) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \Delta_{\mathcal{X}} & \longrightarrow & \Pi_{\mathcal{X}} & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow \\ 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \longrightarrow & G_k \longrightarrow 1 \end{array}$$

where the middle vertical map (defined up to inner conjugation) is induced by the scheme morphism $\mathcal{X} \rightarrow X$.

For the rest of this section we assume that \mathcal{X} is a formal fibre as in 5.1, which is geometrically connected.

Definition 5.2. A section $\tilde{s} : G_k \rightarrow \Pi_{\mathcal{X}}$ of the projection $\Pi_{\mathcal{X}} \twoheadrightarrow G_k$ induces a section $s : G_k \rightarrow \Pi_X$ of the projection $\Pi_X \twoheadrightarrow G_k$ (cf. diagram (5.1)). We will refer to such a section s as a **local section** of the projection $\Pi_X \twoheadrightarrow G_k$.

Note that a geometric section (cf. Definition 4.1) is a local section in the above sense, as one easily verifies. Our main result is the following.

Theorem 5.3. *Let $s : G_k \rightarrow \Pi_X$ be a local section of the projection $\Pi_X \twoheadrightarrow G_k$. Then s has a cycle class uniformly orthogonal to Pic^\wedge in the sense of [Saïdi], Definition 1.4.1(i).*

Proof. Let $\mathcal{X} = \text{Spec}(\hat{\mathcal{O}}_{\tilde{X},x} \otimes_{\mathcal{O}_k} k)$ be as in 5.1, and $\tilde{s} : G_k \rightarrow \Pi_{\mathcal{X}}$ a section of the projection $\Pi_{\mathcal{X}} \twoheadrightarrow G_k$ which induces the section $s : G_k \rightarrow \Pi_X$. Let $\{X_t\}_{t \geq 1}$ be as in §4, $s_t : G_k \rightarrow \Pi_{X_t}$ the section of the projection $\Pi_{X_t} \twoheadrightarrow G_k$ which is induced by s , and $s_t^* : H^2(X_t, \hat{\mathbb{Z}}(1)^\Sigma) \rightarrow H^2(G_k, \hat{\mathbb{Z}}(1)^\Sigma)$ the retraction map induced by s_t , $\forall t \geq 1$. We show $s_t^*(\text{Pic}(X_t)^\wedge) = 0$ (cf. loc. cit. for the definition of $\text{Pic}(X_t)^\wedge$). Recall the continuous homomorphism $\phi : \Pi_{\mathcal{X}} \rightarrow \Pi_X$ (cf. the middle vertical map in diagram (5.1)). Then $\Pi_{\mathcal{X}_t} \stackrel{\text{def}}{=} \phi^{-1}(\Pi_{X_t})$ is an open subgroup containing $\tilde{s}(G_k)$, and corresponds to an étale cover $\mathcal{X}_t \rightarrow \mathcal{X}$ with \mathcal{X}_t geometrically connected (as $\Pi_{\mathcal{X}_t}$ projects onto G_k via the projection $\Pi_{\mathcal{X}} \twoheadrightarrow G_k$). Moreover, the section \tilde{s} induces a retraction $\tilde{s}_t^* : H^2(\Pi_{\mathcal{X}_t}, \hat{\mathbb{Z}}(1)^\Sigma) \rightarrow H^2(G_k, \hat{\mathbb{Z}}(1)^\Sigma)$ of the natural map $H^2(G_k, \hat{\mathbb{Z}}(1)^\Sigma) \rightarrow H^2(\Pi_{\mathcal{X}_t}, \hat{\mathbb{Z}}(1)^\Sigma)$ induced by the projection $\Pi_{\mathcal{X}_t} \twoheadrightarrow G_k$. Let

$A \stackrel{\text{def}}{=} \hat{\mathcal{O}}_{\tilde{X},x}$, and $A_k \stackrel{\text{def}}{=} \hat{\mathcal{O}}_{\tilde{X},x} \otimes_{\mathcal{O}_k} k$. The open subgroup $\Pi_{\mathcal{X}_t}$ corresponds to an étale cover $\mathcal{X}_t = \text{Spec } B_k \rightarrow \mathcal{X} = \text{Spec } A_k$, where B_k/A_k is an étale extension. Let B be the integral closure of A in B_k . Thus, B is a complete local ring of dimension 2, which dominates \mathcal{O}_k , and the residue field of B is finite. We have a scheme theoretic morphism $\mathcal{X}_t \rightarrow X_t$. Further, we have an injective homomorphism $H^2(\Pi_{\mathcal{X}_t}, \hat{\mathbb{Z}}(1)^\Sigma) \hookrightarrow H_{\text{ét}}^2(\mathcal{X}_t, \hat{\mathbb{Z}}(1)^\Sigma)$ arising from the Cartan-Leray spectral sequence (cf. [Serre], proof of Proposition 1), as well as an injective Kummer homomorphism $\text{Pic}(\mathcal{X}_t)^\wedge \hookrightarrow H_{\text{ét}}^2(\mathcal{X}_t, \hat{\mathbb{Z}}(1)^\Sigma)$, where $\text{Pic}^\wedge \stackrel{\text{def}}{=} \text{Pic} \otimes_{\mathbb{Z}} \hat{\mathbb{Z}}^\Sigma$. On the other hand we have a commutative diagram of homomorphisms

$$\begin{array}{ccc} H^2(\Pi_{\mathcal{X}_t}, \hat{\mathbb{Z}}(1)^\Sigma) & \xrightarrow{\tilde{s}_t^*} & H^2(G_k, \hat{\mathbb{Z}}(1)^\Sigma) \\ \psi_t \uparrow & & \text{id} \uparrow \\ H^2(\Pi_{X_t}, \hat{\mathbb{Z}}(1)^\Sigma) & \xrightarrow{s_t^*} & H^2(G_k, \hat{\mathbb{Z}}(1)^\Sigma) \end{array}$$

where ψ_t is induced by the map $\phi : \Pi_{\mathcal{X}_t} \rightarrow \Pi_{X_t}$.

We claim that $\psi_t(\text{Pic}(X_t)^\wedge)$ is *torsion*, from this it follows that $s_t^*(\text{Pic}(X_t)^\wedge) = \tilde{s}_t^*(\psi_t(\text{Pic}(X_t)^\wedge)) = 0$, since $H^2(G_k, \hat{\mathbb{Z}}(1)^\Sigma) \xrightarrow{\sim} \hat{\mathbb{Z}}^\Sigma$ is torsion free. Indeed, we have a pull-back morphism $\text{Pic}(X_t)^\wedge \rightarrow \text{Pic}(\mathcal{X}_t)^\wedge$, which fits in the commutative diagram

$$\begin{array}{ccc} \text{Pic}(\mathcal{X}_t)^\wedge & \longrightarrow & H_{\text{ét}}^2(\mathcal{X}_t, \hat{\mathbb{Z}}(1)^\Sigma) \\ \text{id} \uparrow & & \uparrow \\ \text{Pic}(\mathcal{X}_t)^\wedge & & H^2(\Pi_{\mathcal{X}_t}, \hat{\mathbb{Z}}(1)^\Sigma) \\ \uparrow & & \psi_t \uparrow \\ \text{Pic}(X_t)^\wedge & \longrightarrow & H^2(\Pi_{X_t}, \hat{\mathbb{Z}}(1)^\Sigma) \end{array}$$

where the horizontal maps are injective Kummer maps (recall the identification $H^2(\Pi_{X_t}, \hat{\mathbb{Z}}(1)^\Sigma) \xrightarrow{\sim} H_{\text{ét}}^2(X_t, \hat{\mathbb{Z}}(1)^\Sigma)$), and the upper right vertical map is the injective map discussed above. Our claim follows then from the following.

Proposition 5.4. *With the notations above $\text{Pic}(\mathcal{X}_t)$, and a fortiori $\text{Pic}(\mathcal{X}_t)^\wedge$, is finite.*

Proof of Proposition 5.4. This follows from the fact, proven by Shuji Saito, that $\text{Pic}(\text{Spec } B \setminus \{m_B\})$ is finite, where B is as in the above discussion and m_B is its maximal ideal (cf. [Saito], Theorem 0.11). \square

This finishes the proof of Theorem 5.3. \square

Finally, we provide the following characterisation of local sections which are geometric. We use the above notations.

Theorem 5.5. *Let $s : G_k \rightarrow \Pi_X$ be a **local** section of the projection $\Pi_X \rightarrow G_k$ (cf. Definition 5.2). Then s **lifts** to a section $\rho : G_k \rightarrow G_X^{(1-\text{sol})}$ (resp. $\rho_n : G_k \rightarrow G_X^{(1/p^n-\text{sol})}$) of the projection $G_X^{(1-\text{sol})} \twoheadrightarrow G_k$ (resp. $G_X^{(1/p^n-\text{sol})} \twoheadrightarrow G_k, \forall n \geq 1$). Moreover, the section s is **geometric if and only if** there exists a lifting of s to a section $\rho_2 : G_k \rightarrow G_X^{(1/p^2-\text{sol})}$ as above such that one of the following equivalent conditions hold.*

(i) With the notations in §4 (cf. the discussion before Definition 4.3), the class of $J_j^1[U]$ in $H^1(G_k, J_j[U])$ is **divisible** by p , $\forall U \subseteq X$ nonempty open subscheme, and $\forall j \geq 1$.

(ii) The section ρ_2 **lifts** to a section $\tilde{\rho}_2 : G_k \rightarrow G_X^{(p,2)}$ of the projection $G_X^{(p,2)} \twoheadrightarrow G_k$.

Proof. The first assertion follows from Theorem 5.3, Theorem 3.4.4, and Theorem 2.3.5 in [Saïdi]. The second assertion follows from Theorem 3.4.10, and Theorem 4.5. \square

References

- [Grothendieck] Grothendieck, A., Revêtements étales et groupe fondamental, Lecture Notes in Math. 224, Springer, Heidelberg, 1971.
- [Harari-Szamuely], Harari, D., Szamuely, T., Galois sections for abelianized fundamental groups, Math. Ann. (2009) 344, 779-800.
- [Hoshi] Hoshi, Y., Existence of nongeometric pro- p Galois sections of hyperbolic curves, Publ. Res. Inst. Math. Sci. 46 (2010), no. 4, 829-848.
- [Mochizuki] Mochizuki, S., Absolute anabelian cuspidalizations of proper hyperbolic curves, J. Math. Kyoto Univ. 47 (2007), no. 3, 451–539.
- [Mochizuki1] Mochizuki, S., Topics in absolute anabelian geometry II: decomposition groups and endomorphisms. J. Math. Sci. Univ. Tokyo, Vol. 20 (2013), No. 2, Page 171-269.
- [Pop] Pop, F., On the birational p -adic section Conjecture, Compos. Math. 146 (2010), no. 3, 621-637.
- [Saïdi] Saïdi, M., The cuspidalisation of sections of arithmetic fundamental groups, Advances in Mathematics, 230 (2012) 1931-1954.
- [Saïdi1] Saïdi, M., On the p -adic section conjecture, Journal of Pure and Applied Algebra, Volume 217, Issue 3, (2013), 583-584.
- [Saïdi2] Saïdi, M., On the section conjecture over function fields and finitely generated fields, Publ. RIMS Kyoto Univ. 52 (2016), 335-357.
- [Saïdi3] Saïdi, M., Arithmetic of p -adic curves and sections of geometrically abelian fundamental groups. Manuscript.
- [Saito] Saito, S., Arithmetic on two dimensional local rings, Invent. math. 85, 379-414 (1986).
- [Serre] Serre J.-P., Construction de revêtements de la droite affine en caractéristique $p > 0$, C. R. Acad. Sci. Paris 311 (1990), 341-346.
- [Tamagawa] Tamagawa, A., The Grothendieck conjecture for affine curves, Compositio Math. 109 (1997), no. 2, 135–194.

Mohamed Saïdi
 College of Engineering, Mathematics, and Physical Sciences
 University of Exeter
 Harrison Building
 North Park Road
 EXETER EX4 4QF
 United Kingdom
 M.Saïdi@exeter.ac.uk