

# Pure

## Scotland's Rural College

### The ENHANCED MAGNAPORTHE RESISTANCE 1 locus affects Ramularia leaf spot development in barley

McGrann, Graham R.D.; Miller, Steven; Havis, Neil D.

*Published in:*

European Journal of Plant Pathology

*DOI:*

[10.1007/s10658-019-01869-x](https://doi.org/10.1007/s10658-019-01869-x)

Print publication: 01/01/2020

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication](#)

*Citation for published version (APA):*

McGrann, G. R. D., Miller, S., & Havis, N. D. (2020). The *ENHANCED MAGNAPORTHE RESISTANCE 1* locus affects Ramularia leaf spot development in barley. *European Journal of Plant Pathology*, 156(1), 123-132. <https://doi.org/10.1007/s10658-019-01869-x>

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# The *ENHANCED MAGNAPORTHE RESISTANCE 1* locus affects *Ramularia* leaf spot development in barley

Graham R. D. McGrann · Steven Miller ·  
Neil D. Havis

Accepted: 10 October 2019  
© The Author(s) 2019

**Abstract** *Ramularia* leaf spot (RLS) is a newly-important disease of barley which is caused when the fungus *Ramularia collo-cygni* enters necrotrophic development during colonisation of the host. Mutant alleles at the barley *MILDEW LOCUS O*, *mlo*, locus confer broad spectrum durable resistance against the powdery mildew fungus, *Blumeria graminis* f. sp. *hordei*, but can enhance susceptibility to pathogens with necrotrophic development stages such as *R. collo-cygni*. Given the importance of *mlo* in spring barley breeding programmes, identifying loci that mitigate the effect of *mlo*-mediated susceptibility on necrotrophic disease development is an important target. Mutation of the *ENHANCED MAGNAPORTHE 1* (*emr1*) locus which can affect *mlo*-associated disease susceptibility, leads to a reduction in RLS symptoms on barley leaves but does not reduce *R. collo-cygni* accumulation. The effect of *emr1* on the transition of *R. collo-cygni* from endophyte

to necrotroph may relate to changes in reactive oxygen species in mutant plants which show reduced sensitivity to chloroplastic superoxide induced cell death and has lower relative chlorophyll content compared to *mlo* plants.

**Keywords** *Ramularia collo-cygni* · Disease resistance trade-off · Necrotroph · Endophyte · *emr1* · *Mlo*

## Introduction

*Ramularia* leaf spot (RLS) is an important disease of barley crops in many temperate countries (Havis et al. 2015; McGrann and Havis 2017). The disease is caused by the fungus *Ramularia collo-cygni*, which is transmitted through infected seed and by air-borne spores (Havis et al. 2014). *R. collo-cygni* has a long endophytic phase (Kaczmarek et al. 2017) with necrotrophic disease symptoms usually only visible once the crop has entered the reproductive phase (Schützendübel et al. 2008). Despite *R. collo-cygni* possessing the genetic capability to synthesise a range of secondary metabolites that could potentially affect disease severity (Dussart et al. 2018), expression of RLS appears to be linked with adverse environmental conditions (Makepeace et al. 2008; McGrann and Brown 2018; Peraldi et al. 2014). The vertical transmission by seeds and relationship between environmental conditions and symptom expression has led to the suggestion that *R. collo-cygni* may be an endophyte that only causes disease under specific circumstances (McGrann and Havis 2017). Resistance

---

S. Miller · N. D. Havis  
Crop Protection Team, Crop and Soil Systems, SRUC, West  
Mains Road, Edinburgh EH9 3JG, UK  
e-mail: grahammcgrann@googlemail.com

G. R. D. McGrann  
Potato Branch, SASA, Roddinglaw Road, Edinburgh EH12 9FJ,  
UK

*Present Address:*  
G. R. D. McGrann (✉)  
Crop Protection Team, Crop and Soil Systems, SRUC, West  
Mains Road, Edinburgh EH9 3JG, UK  
e-mail: grahammcgrann@googlemail.com

to RLS appears to be a quantitative trait although genetic studies in barley have indicated that mutant *MILDEW LOCUS O (mlo)* alleles increase susceptibility to RLS (McGrann et al. 2014), although this phenotype is not observed in all environments (Makepeace et al. 2007). Mutant *mlo* alleles are agriculturally important as they confer broad-spectrum resistance to all known field isolates of powdery mildew caused by *Blumeria graminis* f. sp. *hordei* in spring barley (Piffanelli et al. 2004). This resistance has been used to great effect for more than 40 years. The mutant allele is currently present in more than 70% of elite European varieties (Dreiseitl 2012; Jørgensen 1992) with most commercial spring barley with *mlo*-mediated mildew resistance carrying the *mlo11* allele although some older varieties contain *mlo9* (Jørgensen 1992).

The *MLO* gene encodes a seven-transmembrane domain protein (Büschges et al. 1997). Wild type *MLO* alleles are susceptibility factors for biotrophic powdery mildew fungi, which co-evolved before the divergence of monocot and dicot plants (Acevedo-Garcia et al. 2014; Appiano et al. 2015). However, introducing *mlo* in commercial spring barley varieties came with a number of undesirable pleiotropic effects. Plants containing *mlo* alleles developed spontaneous foliar necrosis spotting, tended to yield less (Kjaer et al. 1990; Wolter et al. 1993; Thomas et al. 1998) and were more susceptible to diseases with necrotrophic stages (Jarosch et al. 1999; Kumar et al. 2001; Jansen et al. 2005; McGrann et al. 2014). Plant breeding has managed to alleviate the necrotic spotting and yield penalties associated with *mlo* (Bjornstad and Aastveit 1990; Kjaer et al. 1990) but there are still concerns over increased disease susceptibility associated with this mutant allele. Mutant analyses have indicated that *mlo*-mediated susceptibility to the pathogens with necrotrophic growth stages is regulated by other genes. Two *REQUIRED FOR MLO RESISTANCE (ROR1 and ROR2)* genes are essential for the full expression of *mlo*-mediated resistance to powdery mildew (Freialdenhoven et al. 1996). Enhanced RLS symptom formation in *mlo* plants is reduced in *ror1* and *ror2* mutants although levels of *R. collo-cygni* DNA in colonised leaves are still elevated (McGrann et al. 2014). Mutant *ror1* plants show decreased sensitivity to toxins produced by *Bipolaris sorokiniana* compared to *mlo* plants with the wild type *ROR1* allele (Kumar et al. 2001). However, loss of *ROR1* function has no effect on barley susceptibility to *Magnaporthe oryzae* (Jarosch et al. 1999) suggesting *ROR* genes may

operate through mechanisms that differentially affect *mlo*-mediated susceptibility to specific fungal species. *NEC1* encodes a cyclic nucleotide gated channel 4 protein, which when defective, results in a spontaneous lesion mimic phenotype in barley (Rostoks et al. 2006). Mutant *nec1* alleles reduce lesion formation by *Fusarium culmorum* and *R. collo-cygni* in barley but these *nec1*-mediated phenotypes are differentially affected by *mlo* mutations. The presence of *nec1* mutation in an *mlo5* background was shown to have no effect on the development of RLS symptoms or the accumulation of *R. collo-cygni* DNA. However, the presence of *mlo5* in a *nec1* background compromises the reduction in *F. culmorum* lesion size usually seen in *nec1* mutants, although disease levels did not reach those seen in wild type *MLO* or *NEC1* plants or in *mlo5* mutants (McGrann et al. 2015a). A screen for restoring resistance to *M. oryzae* in *mlo*-barley identified two loss of function mutations, *ENHANCED MAGNAPORTHE RESISTANCE (EMR1 and EMR2)*, which showed partially restored resistance to *M. oryzae* with no compromise on mildew resistance (Jansen et al. 2007; Jansen and Schaffrath 2009). The gene responsible for the *emr1* or *emr2* phenotype have not yet been characterised. Genetic mapping of the *emr1* locus indicated that the *emr1* mutant contains at least one additional mutant gene, *3-KETOACYL-CoA SYNTHASE (HvKCS6)*, located on chromosome 4H that results in *emr1* plants having reduced lower levels of cuticular waxes (Weidenbach et al. 2014). These mutagenesis studies have shown that the negative effect of *mlo* on certain pathogens with necrotrophic colonisation stages can be mitigated. Here, the effect of the *emr1* mutation on RLS was examined to assess the potential of this locus to mitigate *mlo*-mediated susceptibility to this disease.

## Materials and methods

### Plant material

The *emr1mlo5* mutant was derived from NaN3 mutagenesis of IngridBC*mlo5* mutant (*mlo5*) and shown to restore resistance to the blast fungus, *Magnaporthe oryzae*, in the highly susceptible *mlo5* background (Jansen et al. 2007). Experiments included the *emr1mlo5* double mutant, the single mutant parental line *mlo5* and the wild type donor cultivar of both mutants cv. Ingrid. Seeds were sown in to Levington

M3 compost (ICL, Ipswich, UK) in rows of ten seeds per plastic tray (210 × 156 × 47 mm). Barley seedlings were cultivated in a MD1400 modular climate chamber (Snijders Labs, Tilberg, The Netherlands) at 18 °C with a 16:8 h light: dark cycle under 200  $\mu\text{mol m}^{-2} \text{s}^{-1}$  artificial light at 80% relative humidity.

#### *Ramularia collo-cygni* isolates and inoculation method

The *R. collo-cygni* isolate DK05 Rcc001 (McGrann et al. 2016) collected from Denmark in 2005 was used in this study. Fungal cultures were stored on potato dextrose agar (PDA) plates supplemented with streptomycin (100  $\mu\text{g mL}^{-1}$ ) at 15 °C. Liquid cultures were prepared by adding two 5 mm<sup>2</sup> plugs from 14 to 28 day old PDA storage plates to 200 mL potato dextrose broth (PDB) supplemented with streptomycin (100  $\mu\text{g mL}^{-1}$ ). PDB flasks with *R. collo-cygni* were incubated for 14 days at 15 °C in the dark on an orbital shaker set at 150 rpm. Barley seedlings were inoculated with *R. collo-cygni* as previously described (McGrann et al. 2014; Peraldi et al. 2014). Inoculum was prepared by fragmenting *R. collo-cygni* hyphal growth in the liquid cultures in a food processor (Kenwood Electronics, London, UK). A single drop of Tween20 (Sigma, Dorset, UK) was added to each batch of 50 mL inoculum that was used to inoculate approximately 240 barley plants at growth stage (GS) 12 (Zadoks et al. 1974). Plants were incubated in the dark for 48 h post inoculation and RLS lesions were scored as the proportion of the prophyll leaf covered with disease symptoms from 8 to 21 days post inoculation (dpi). Green leaf area (GLA) retention was assessed as the amount of prophyll leaf area not senescent at the same time point that RLS was scored in the *emr1* mutant seedling inoculation experiments. A minimum of three independent inoculation experiments were assessed for each seedling inoculation experiment.

#### Field trials

Barley cv. Ingrid, the near isogenic line *mlo5* and the *emr1mlo5* mutant were sown in tussock plots at a trial site in Lanark, Scotland in 2015 and 2016. The Lanark site was selected as it typically exhibits high levels of *R. collo-cygni* (Havis et al. 2014). To maintain green leaf area retention and control other foliar diseases during the growing season the fungicides Comet® 200 (0.4 L ha<sup>-1</sup> Pyraclostrobin [200 g L<sup>-1</sup>] BASF,

Cheshire, UK) plus Vegas® (0.25 L ha<sup>-1</sup> Cyflufenamid [50 g L<sup>-1</sup>] Certis, Cambridgeshire, UK) were applied as a mixture to each plot at both growth stage (GS) 25–20 and GS45. RLS symptoms were scored 2–3 times after the crop had flowered (GS60 onwards) as the proportion of leaf F-1 covered with disease lesions.

#### Quantification of in planta *Ramularia collo-cygni* DNA levels

Genomic DNA (gDNA) was extracted from five leaves of cv. Ingrid, *mlo5* and the *emr1mlo5* mutant 21 dpi with *R. collo-cygni* isolate DK05 using the DNeasy Plant Mini Kit (Qiagen, Hilden, Germany) following the manufacturer's protocol. *R. collo-cygni* DNA levels were quantified using a dilution series of *R. collo-cygni* DNA (50 ng to 0.1 pg) with qPCR following the method of Taylor et al. (2010). Data was collected from two independent inoculation experiments.

#### Dark-induced senescence assays

Differences in relative chlorophyll content and senescence between *emr1mlo5*, *mlo5* and cv. Ingrid were tested using a SPAD 502 Plus Chlorophyll meter (Konica Minolta, Warrington, UK) in a dark-induced senescence assay as previously described (McGrann et al. 2015b). Relative chlorophyll content was taken from three points across the blade of GS12 prophyll leaves after excision from the plant (day 0). Leaves were placed on damp tissue paper in clear plastic boxes covered with aluminium foil and stored at room temperature. Additional measurements were taken every two days up to and including day eight of the experiment. Three independent experiments were examined with 6–8 leaves of each line tested in each experiment.

#### Sensitivity to reactive oxygen species-induced cell death

The sensitivity of *emr1mlo5*, *mlo5* and cv. Ingrid to cell death induced by reactive oxygen species (ROS) was tested using detached prophyll leaves from barley plants at GS11–12 as described previously (McGrann et al. 2015b). Cell death was induced by the hydrogen peroxide donor alloxan (200 mM), the mitochondrial superoxide donor menadione (100 mM) and the chloroplastic superoxide donor methyl viologen (25  $\mu\text{M}$ ). Lesions were measured 96 h after treatment with each ROS

donor using ImageJ software (Abràmoff et al. 2004). Three independent experiments were performed with eight leaves of each line assessed for each ROS donor in each replicate experiment.

#### Data analysis

RLS scores over each seeding inoculation time course experiment were used to calculate the area disease under progress curve (AUDPC; Shaner and Finney 1977). AUDPC data was converted into the percentage of the maximum AUDPC (%maxAUDPC) across the entire seedling inoculation time course experiment. Prior to analysis %maxAUDPC data was LOGIT+ transformed (McGrann et al. 2014) and then analysed using a general linear model (GLM). Differences in seedling levels of RLS %maxAUDPC and *R. collo-cygni* gDNA in seedling prophyll leaves 21 dpi of the *emr1mlo5* mutant compared to *mlo5* and Ingrid were tested using GLM with line and experiment as factors. Differences in GLA retention in *emr1mlo5*, *mlo5* and cv. Ingrid was examined using linear mixed modelling of repeated measures. Differences in GLA retention were evaluated using the uniform correlation/split plot in time covariance matrix with dpi, line, experiment and the interactions between these factors set as fixed factors. The dpi by individual plant interaction term was set as the random factor in the model. Significant differences between *mlo5* and *emr1mlo5* or cv. Ingrid at specific dpi were assessed using a t-test. GLM was also used to assess the variation in RLS AUDPC in the field trials with line and year as factors following a Poisson transformation of the AUDPC data.

Variation in ROS-induced cell death between *emr1mlo5*, *mlo5* and cv. Ingrid was assessed using a separate GLM for each ROS donor. Each GLM assessed the contribution of experiment, line and the interactions between these terms on the observed phenotypes. Differences in dark-induced senescence were tested using linear mixed modelling with repeated measurements as described above for GLA.

## Results

### Development of Ramularia leaf spot on the spring barley *emr1mlo5* mutant

In seedling assays typical RLS symptoms were observed on leaves of barley cv. Ingrid, the *mlo5* single


mutant and the *emr1mlo5* double mutant (Fig. 1a) 21 dpi with lesions first visible 8–10 dpi. The *mlo5* mutant had increased RLS development compared to the wild type Ingrid ( $P < 0.001$ ) as previously reported (McGrann et al. 2014, 2015a), whilst the *emr1mlo5* mutant developed significantly less RLS than *mlo5* ( $P < 0.001$ ) but significantly more disease than cv. Ingrid ( $P = 0.007$ ; Fig. 1b). By 8 dpi both *mlo5* and *emr1mlo5* had significantly less GLA than cv. Ingrid ( $P < 0.01$ ) whereas *emr1mlo5* plants had significantly lower GLA than *mlo5* ( $P < 0.05$ ) at this time point (Fig. 1c). From 10 dpi onwards there were no significant differences in GLA between *mlo5* and *emr1mlo5* whereas both mutants had significantly lower levels of GLA compared to cv. Ingrid at all further time points ( $P < 0.001$ ; Fig. 1c). Despite the differences in RLS development between *emr1mlo5* and *mlo5* there was no significant difference in the accumulation of *R. collo-cygni* gDNA 21 dpi between the mutants ( $P = 0.98$ ) whereas cv. Ingrid leaves had significantly less fungal DNA than *mlo5* ( $P = 0.022$ ) and *emr1mlo5* ( $P = 0.039$ ) leaves (Fig. 1d).

In field grown plants no significant differences in RLS development were recorded between the two years of the field trials ( $P = 0.37$ ) nor were any differences observed between the lines ( $P = 0.74$ ). However, even though the differences in RLS development between field grown cv. Ingrid, *mlo5* and *emr1mlo5* were not statistically significant there was a trend for *mlo5* plants to exhibit more RLS symptoms than the other two lines (Fig. 1e).

### Effect of *emr1* mutation on relative chlorophyll content and dark-induced senescence

Processes linked with foliar senescence and chlorophyll breakdown have been linked with development of RLS (Schützendübel et al. 2008; McGrann et al. 2015b; McGrann and Brown 2018). Therefore, the effects of the *emr1* mutation on senescence were examined using a dark-induced senescence assay. Prophyll leaves of the *emr1mlo5* mutant at GS12 were a paler shade of green when compared to *mlo5* or cv. Ingrid (Fig. 2a). This may be related to the altered wax status of *emr1mlo5* plants (Weidenbach et al. 2014) or due to changes in chlorophyll content of the leaves associated with the *emr1* mutation. Relative chlorophyll content of the *emr1mlo5* mutant was significantly lower ( $P < 0.001$ ) at day 0 than either *mlo5* or cv. Ingrid. The relative chlorophyll

**Fig. 1** Effect of *emr1* mutation on *Ramularia* leaf spot (RLS) development. Disease phenotype on prophyll leaf of cv. Ingrid (*MLO*), *mlo5* and *emr1mlo5* plants 21 days post inoculation (dpi) with *R. collo-cygni* isolate DK05 Rcc001 (a). Development of RLS converted to the area under disease progress curve (AUDPC) and expressed as a proportion of the total AUDPC possible over the time course of the experiment (%maxAUDPC). Data was collected from five independent inoculation experiments with disease symptoms assessed on a minimum of eight plants of each line in each experiment (b). Green leaf area (GLA) retention over time following inoculation with *R. collo-cygni* isolate DK05 Rcc001. Data was collected from five independent inoculation experiments with GLA assessed on a minimum of eight plants of each line in each experiment (c). *R. collo-cygni* DNA levels in prophyll leaves 21 dpi measured by qPCR. Data was collected from two independent inoculation experiments with fungal DNA levels assessed from five replicates of each line in both experiments (d). Development of RLS in adult cv. Ingrid (*MLO*), *mlo5* and *emr1mlo5* plants. Field trials were conducted in two years with a single tussock plot scored for each line and the data from both years combined (e). Error bars indicate  $\pm 1$  SE. \*\*\* $P < 0.001$ ; \*  $0.01 < P < 0.05$  when *emr1mlo5* or cv. Ingrid (*MLO*) plants are compared to *mlo5*




content of *emr1mlo5* remained significantly lower than *mlo5* throughout the dark-induced senescence time course (day 2–6) but the SPAD readings for the two mutants were not significantly different by day 8 (Fig. 2b). Relative chlorophyll content of cv. Ingrid leaves was significantly higher than both *emr1mlo5* ( $P < 0.001$ ) and *mlo5* ( $P < 0.05$ ) across all time points.

### Sensitivity of *emr1mlo5* mutant to reactive oxygen species-induced cell death

Changes in ROS homeostasis associated with plant development and spontaneous cell death phenotypes have been linked with expression of RLS symptoms (McGrann and Brown 2018). No significant differences in the size of the alloxan-induced lesion ( $P = 0.49$ ) were

**Fig. 2** Effect of the *emr1* mutation on leaf phenotype and dark-induced senescence. Photograph of 14 day old prophyll leaf of cv. Ingrid (*MLO*), *mlo5* and *emr1mlo5* plants (a). Relative chlorophyll content for cv. Ingrid (*MLO*), *mlo5* and *emr1mlo5* plants during dark-induced senescence time course. Data was collected from three independent experiments with a minimum of six replicate plants tested in each experiment (b). Error bars indicate  $\pm 1$  SE




observed between *emr1mlo5*, *mlo5* and cv. Ingrid (Fig. 3a+d). No significant difference in the size of the lesions formed by the mitochondrial superoxide donor menadione were recorded between the lines (Fig. 3b+e;  $P = 0.07$ ) but the *emr1mlo5* mutant did produce a significantly smaller lesion compared to *mlo5* and cv. Ingrid ( $P < 0.001$ ) when treated with the chloroplastic superoxide donor methyl viologen (Fig. 3c+f).

## Discussion

Ramularia leaf spot has established itself as a serious threat to modern barley production across temperate climates (McGrann and Havis 2017; Havis et al. 2015;

Walters et al. 2008). Yield losses typically range from 5 to 10% but can be as high as 70% without considering potential losses to grain quality which can reduce the value of the crop further (Havis et al. 2015). Resistance against RLS is most probably a quantitatively controlled genetic trait and the strong genotype by environment effects observed for the expression of disease symptoms between field trials, has led to problems in breeding effective resistance against this disease (Havis et al. 2015). Durable powdery mildew resistance conferred by the *mlo* locus has resulted in mildew not being the threat it once was (Makepeace et al. 2007) and the importance of this resistance source is clear by the presence of the *mlo* allele in approximately 70% of European spring barley varieties (Dreiseitl 2012;



**Fig. 3** Sensitivity of *emr1mlo5* mutant to reactive oxygen species (ROS)-induced cell death. Photographs of ROS-induced lesions on cv. Ingrid (*MLO*), *mlo5* and *emr1mlo5* prophyll leaves 96 h after treatment with alloxan (a), menadione (b) or methyl viologen (c). Measurements of lesion size following treatment with alloxan

(d), menadione (e) or methyl viologen (f). Data was collected from three independent experiments with a minimum of five replicate plants tested in each experiment. Error bars indicate  $\pm 1$  SE. \*\*\*  $P < 0.001$  when *emr1mlo5* or cv. Ingrid (*MLO*) plants are compared to *mlo5*

Jørgensen 1992). Even though *mlo* alleles can increase the threat of emerging diseases such as RLS (McGrann et al. 2014) the overriding importance of this mutation to spring barley production means that it is essential to develop varieties that are able to mitigate the negative effect of *mlo* on RLS (Brown and Rant 2013).

Limiting the effect of *mlo* on the necrotrophic phase of fungal diseases is an important breeding target to protect the use of *mlo* in commercial spring barley. Recent studies by Aghnoum et al. (2019) suggest that the genetic background of barley varieties can mitigate *mlo*-mediated susceptibility to blast disease. Through mutation analysis of an experimental *mlo* barley line, Jansen et al. (2007) demonstrated the presence of genetic loci that can moderate the effects of *mlo* on necrotrophic diseases. The mutant alleles, *emr1* and *emr2*, both reduce *mlo*-mediated enhanced susceptibility to *M. oryzae* but have no effect on powdery mildew resistance (Jansen et al. 2007; Jansen and Schaffrath 2009). This suggests that there are genetic loci that have contrasting effects on the different pathways regulated by *mlo*, as previously suggested (McGrann et al. 2015a). Mutant *emr1mlo5* plants showed reduced RLS development, but this reduction in disease was not accompanied by a reduction in fungal biomass. Similar findings were

observed in barley *ror* mutants where both *ror1* and *ror2* reduce RLS expression in leaves but do not lower levels of *R. collo-cygni* biomass (McGrann et al. 2014). Not all genes that affect the expression of RLS interact with *mlo*. Mutant alleles of the cyclic nucleotide gated channel *NEC1* lower RLS symptoms in wild type *MLO* plants but do not affect disease levels in a *necl1 mlo* double mutant (McGrann et al. 2015a) indicating that multiple pathways affect the expression of RLS symptoms and that *mlo* may only affect some of these pathways.

The *emr1* mutant contains at least one other mutant locus, *iwa1*, which results in defective wax biosynthesis (Weidenbach et al. 2014). Whether or not altered surface wax composition affects RLS is currently unknown but *emr1mlo5* mutants had reduced relative chlorophyll content and were more sensitive to chloroplastic ROS. Defects in *ROR* genes and *NEC1* result in mis-regulated ROS and affect the transition of *R. collo-cygni* from asymptomatic growth to necrotrophic development. Changes in foliar chlorophyll content associated with the onset of senescence have been linked to RLS symptom expression (Schützendübel et al. 2008; McGrann and Brown 2018). Host responses associated with foliar senescence processes are regulated during *R. collo-*



*cygni* infection suggesting that this pathogen triggers premature senescence in the plant (Sjökvist et al. 2019). Transgenic barley plants that have delayed leaf senescence due to over-expression of a *Stress-induced NAC1* transcription factor show increased resistance against this disease (McGrann et al. 2015b). It is possible that defects in relative chlorophyll content, senescence and ROS balance in *emr1mlo5* plants affect the transition of *R. collo-cygni* entering its necrotrophic phase. Determining the function of the gene at the *EMRI* locus responsible for the effects on RLS symptom development may provide further insights into how plant genetics trigger this fungus to change growth habit and cause disease symptoms.

Whilst there is promise in modern commercial spring barley varieties for good RLS resistance in plants with *mlo* mildew resistance, a better understanding of the mechanisms and genetics controlling these phenotypes is required. Abiotic stress is known to affect expression of RLS (Peraldi et al. 2014; Makepeace et al. 2008; McGrann and Brown 2018) and Makepeace et al. (2007) suggested that the *mlo* effects on diseases such as RLS are influenced by environmental conditions. This finding is supported by the *emr1mlo5* field trial data, where statistically significant differences between *mlo*, wild type *MLO* (cv. Ingrid) and *emr1mlo5* were not recorded, despite the trend in RLS levels observed being similar to results obtained in the controlled environment tests. Assessing RLS development in varieties with good RLS resistance in *mlo* backgrounds in multiple environments is essential to confirm that the effect of genes moderating *mlo*-mediated susceptibility to RLS is not environmentally sensitive. This could be particularly important considering genetic loci such as *EMRI*, which are able to moderate the effect of *mlo* on RLS symptom expression, but do not appear to affect the accumulation of fungal biomass (McGrann et al. 2014, 2015a). As such *R. collo-cygni* DNA levels may remain high in spring barley varieties with *mlo* which have low levels of RLS. This could have negative consequences on grain quality and could result in epidemics when environmental conditions are favourable for the expression of RLS symptoms. The *emr1* mutation has potential to mitigate the effect of *mlo* on promoting RLS symptom production in spring barley but further experimental evidence is needed to evaluate how the environment influences the effects of genetic loci that interaction with the *mlo*-response against this disease.

**Acknowledgements** The authors thank Dr. Ulrich Schaffrath (RWTH Aachen, Germany) for providing the *emr1* mutant and associated parental lines. SRUC receives funding from the Scottish Government RESAS.

**Author contributions** All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by GMcG, SM and NDH. The first draft of the manuscript was written by GMcG and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

#### Compliance with ethical standards

**Disclosure of potential conflicts of interest** The authors declare that they have no conflict of interest.

**Research involving human participants and/or animals** This article does not contain any research with human or animal subjects performed by any of the authors.

**Informed consent** Not applicable.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Abràmoff, M. D., Magalhães, P. J., & Ram, S. J. (2004). Image processing with imageJ. *Biophotonics International*, 11, 36–41.
- Acevedo-Garcia, J., Kusch, S., & Panstruga, R. (2014). Magical mystery tour: MLO proteins in plant immunity and beyond. *New Phytologist*, 204, 273–281.
- Aghnoum, R., Bvindi, C., Menet, G., D'hoop, B., Maciel, J. L. N., & Niks, R. E. (2019). Host/nonhost status and genetics of resistance in barley against three pathotypes of *Magnaporthe* blast fungi. *Euphytica*, 215, 116.
- Appiano, M., Catalano, D., Martínez, M. S., Lotti, C., Zheng, Z., Visser, R. G. F., Luigi, R., Yuling, B., & Stefano, P. (2015). Monocot and dicot MLO powdery mildew susceptibility factors are functionally conserved in spite of the evolution of class-specific molecular features. *BMC Plant Biology*, 15, 257.
- Bjornstad, A., & Aastveit, K. (1990). Pleiotrophic effects on the *mlo-o* resistance gene in barley in different genetical backgrounds. *Euphytica*, 46, 217–226.
- Brown, J. K. M., & Rant, J. C. (2013). Fitness costs and trade-offs of disease resistance and their consequences for breeding arable crops. *Plant Pathology*, 62(S1), 83–95.

- Büschges, R., Hollricher, K., Panstruga, R., Simons, G., Wolter, M., Frijters, A., van Daelen, R., van der Lee, T., Diergaarde, P., Groenendijk, J., Töpsch, S., Vos, P., Salamini, F., & Schulze-Lefert, P. (1997). The barley *Mlo* gene: A novel control element of plant pathogen resistance. *Cell*, *88*, 695–705.
- Dreiseitl, A. (2012). Frequency of powdery mildew resistances in spring barley cultivars in Czech variety trials. *Plant Protection Science*, *48*, 17–20.
- Dussart, F., Douglas, R., Sjökvist, E., Hoebe, P. N., Spoel, S. H., & McGrann, G. R. D. (2018). Genome-based discovery of polyketide-derived secondary metabolism pathways in the barley pathogen *Ramularia collo-cygni*. *Molecular Plant-Microbe Interactions*, *31*, 962–975.
- Freialdenhoven, A., Peterhansel, C., Kurth, J., Kreuzaler, F., & Schulze-Lefert, P. (1996). Identification of genes required for the function of non-race-specific. *The Plant Cell*, *8*, 5–14.
- Havis, N. D., Nyman, M., & Oxley, S. J. P. (2014). Evidence for seed transmission and symptomless growth of *Ramularia collo-cygni* in barley (*Hordeum vulgare*). *Plant Pathology*, *63*, 929–936.
- Havis, N. D., Brown, J. K. M., Clemente, G., Frei, P., Jedryczka, M., Kaczmarek, J., Kaczmarek, M., Matusinsky, P., McGrann, G. R. D., Pereyra, S., Piotrowska, M., Sghyer, H., Tellier, A., & Hess, M. (2015). *Ramularia collo-cygni* - an emerging pathogen of barley crops. *Phytopathology*, *105*, 895–904.
- Jansen, M., & Schaffrath, U. (2009). The barley mutant *emr2* shows enhanced resistance against several fungal leaf pathogens. *Plant Breeding*, *128*, 124–129.
- Jansen, C., von Wettstein, D., Schäfer, W., Kogel, K.-H., Felk, A., & Maier, F. J. (2005). Infection patterns in barley and wheat spikes inoculated with wild-type and trichodiene synthase gene disrupted *Fusarium graminearum*. *Proceedings of the National Academy of Sciences of the United States of America*, *102*, 16892–16897.
- Jansen, M., Jarosch, B., & Schaffrath, U. (2007). The barley mutant *emr1* exhibits restored resistance against *Magnaporthe oryzae* in the hypersusceptible *mlo*-genetic background. *Planta*, *225*, 1381–1391.
- Jarosch, B., Kogel, K., & Schaffrath, U. (1999). The ambivalence of the barley *Mlo* Locus : mutations conferring resistance against powdery mildew (*Blumeria graminis* f. sp. *hordei*) enhance susceptibility to the rice blast fungus *Magnaporthe grisea*. *Molecular Plant-Microbe Interactions*, *12*, 508–514.
- Jørgensen, I. H. (1992). Discovery, characterization and exploitation of *Mlo* powdery mildew resistance in barley. *Euphytica*, *63*, 141–152.
- Kaczmarek, M., Piotrowska, M. J., Fountaine, J. M., Gorniak, K., McGrann, G. R. D., Armstrong, A., Wright, K. M., Newton, A. C., & Havis, N. D. (2017). Infection strategy of *Ramularia collo-cygni* and development of ramularia leaf spot on barley and alternative graminaceous hosts. *Plant Pathology*, *66*, 45–55.
- Kjaer, B., Jensen, H. P., Jensen, J., & Jørgensen, J. H. (1990). Associations between three *mlo* powdery mildew resistance genes and agronomic traits in barley. *Euphytica*, *46*, 185–193.
- Kumar, J., Hüchelhoven, R., Beckhove, U., Nagarajan, S., & Kogel, K. H. (2001). A compromised *Mlo* pathway affects the response of barley to the necrotrophic fungus *Bipolaris sorokiniana* (teleomorph: *Cochliobolus sativus*) and its toxins. *Phytopathology*, *91*, 127–133.
- Makepeace, J. C., Oxley, S. J. P., Havis, N. D., Hackett, R., Burke, J. I., & Brown, J. K. M. (2007). Associations between fungal and abiotic leaf spotting and the presence of *mlo* alleles in barley. *Plant Pathology*, *56*, 934–942.
- Makepeace, J. C., Havis, N. D., Burke, J. I., Oxley, S. J. P., & Brown, J. K. M. (2008). A method of inoculating barley seedlings with *Ramularia collo-cygni*. *Plant Pathology*, *57*, 991–999.
- McGrann, G. R. D., & Brown, J. K. M. (2018). The role of reactive oxygen in the development of *Ramularia* leaf spot disease in barley seedlings. *Annals of Botany*, *121*, 415–430.
- McGrann, G. R. D., & Havis, N. D. (2017). *Ramularia* leaf spot: A newly important threat to barley production. *Outlooks on Pest Management*, *28*, 65–70.
- McGrann, G. R. D., Stavrinides, A., Russell, J., Corbitt, M. M., Booth, A., Chartrain, L., Thomas, W. T. B., & Brown, J. K. M. (2014). A trade off between *mlo* resistance to powdery mildew and increased susceptibility of barley to a newly important disease, *Ramularia* leaf spot. *Journal of Experimental Botany*, *65*, 1025–1037.
- McGrann, G. R. D., Steed, A., Burt, C., Nicholson, P., & Brown, J. K. M. (2015a). Differential effects of lesion mimic mutants in barley on disease development by facultative pathogens. *Journal of Experimental Botany*, *66*, 3417–3428.
- McGrann, G. R. D., Steed, A., Burt, C., Goddard, R., Lachaux, C., Bansal, A., Corbitt, M., Gorniak, K., Nicholson, P., & Brown, J. K. M. (2015b). Contribution of the drought tolerance-related *Stress-responsive NAC1* transcription factor to resistance of barley to *Ramularia* leaf spot. *Molecular Plant Pathology*, *16*, 201–209.
- McGrann, G. R. D., Andongabo, A., Sjökvist, E., Trivedi, U., Dussart, F., Kaczmarek, M., Mackenzie, A., Fountaine, J. M., Taylor, J. M. G., Paterson, L. J., Gorniak, K., Burnett, F., Kanyuka, K., Hammond-Kosack, K. E., Rudd, J. J., Blaxter, M., & Havis, N. D. (2016). The genome of the emerging barley pathogen *Ramularia collo-cygni*. *BMC Genomics*, *17*, 584.
- Peraldi, A., Griffe, L. L., Burt, C., McGrann, G. R. D., & Nicholson, P. (2014). *Brachypodium distachyon* exhibits compatible interactions with *Oculimacula* spp. and *Ramularia collo-cygni*, providing the first pathosystem model to study eyespot and ramularia leaf spot diseases. *Plant Pathology*, *63*, 554–562.
- Piffanelli, P., Ramsay, L., Waugh, R., Benabdelmouna, A., D'Hont, A., Hollricher, K., Jørgensen, J. H., Schulze-Lefert, P., & Panstruga, R. (2004). A barley cultivation-associated polymorphism conveys resistance to powdery mildew. *Nature*, *430*, 887–891.
- Rostoks, N., Schmierer, D., Mudie, S., Drader, T., Brueggeman, R., Caldwell, D. G., Waugh, R., & Kleinhofs, A. (2006). Barley necrotic locus *necl* encodes the cyclic nucleotide-gated ion channel 4 homologous to the Arabidopsis HLM1. *Molecular Genetics and Genomics*, *275*, 159–168.
- Schützendübel, A., Stadler, M., Wallner, D., & Von Tiedemann, A. (2008). A hypothesis on physiological alterations during plant ontogenesis governing susceptibility of winter barley to ramularia leaf spot. *Plant Pathology*, *57*, 518–526.

- Shaner, G., & Finney, R. E. (1977). The effect of nitrogen fertilization on the expression of slow-mildewing resistance in Knox wheat. *Phytopathology*, *77*, 1051–1056.
- Sjökvist, E., Lemcke, R., Kamble, M., Turner, F., Blaxter, M., Havis, N. D., Lyngkjær, M. F., & Radutoiu, S. (2019). Dissection of *Ramularia* leaf spot disease by integrated analysis of barley and *Ramularia collo-cygni* transcriptome responses. *Molecular Plant-Microbe Interactions*, *32*, 176–193.
- Taylor, J. M. G., Paterson, L. J., & Havis, N. D. (2010). A quantitative real-time PCR assay for the detection of *Ramularia collo-cygni* from barley (*Hordeum vulgare*). *Letters in Applied Microbiology*, *50*, 493–499.
- Thomas, W. T. B., Baird, E., Fuller, J. D., Lawrence, P., Young, G. R., Russell, J., Ramsay, L., Waugh, R., & Powell, W. (1998). Identification of a QTL decreasing yield in barley linked to *Mlo* powdery mildew resistance. *Molecular Breeding*, *4*, 381–393.
- Walters, D. R., Havis, N. D., & Oxley, S. J. P. (2008). *Ramularia collo-cygni*: The biology of an emerging pathogen of barley. *FEMS Microbiology Letters*, *279*, 1–7.
- Weidenbach, D., Jansen, M., Franke, R. B., Hensel, G., Weissgerber, W., Ulferts, S., Jansen, I., Schreiber, L., Korzun, V., Pontzen, R., Kumlehn, J., Pillen, K., & Schaffrath, U. (2014). Evolutionary conserved function of barley and Arabidopsis 3-KETOACYL-CoA SYNTHASES in providing wax signals for germination of powdery mildew fungi. *Plant Physiology*, *166*, 1621–1633.
- Wolter, M., Hollricher, K., Salamini, F., & Schulze-Lefert, P. (1993). The *mlo* resistance alleles to powdery mildew infection in barley trigger a developmentally controlled defence mimic phenotype. *Molecular & General Genetics*, *239*, 122–128.
- Zadoks, J. C., Chang, T. T., & Konzak, C. F. (1974). A decimal code for the growth stages of cereals. *Weed Research*, *14*, 415–421.