

*Stomach Content of Three Tuna Species in The Eastern Indian Ocean (Setyadji, B. et al.)*

## STOMACH CONTENT OF THREE TUNA SPECIES IN THE EASTERN INDIAN OCEAN

**Bram Setyadji, Andi Bahtiar, and Dian Novianto**

*Research Station for Tuna Fisheries*

Received June 20-2011; Received in revised form December 6-2012; Accepted December 7-2012

E-Mail : [bram.setyadji@gmail.com](mailto:bram.setyadji@gmail.com)

### ABSTRACT

*Feeding habit of tuna in Indian Ocean has been described around Sri Lanka, Indian Waters, Andaman Sea, western Indian Ocean (Seychelles Islands), western equatorial Indian Ocean whereas the tunas feeding habit study in Eastern Indian Ocean is merely in existence. The purpose of this study is to investigate the stomach content of three tuna species (bigeye tuna, yellowfin tuna, and skipjack tuna), apex predator in the southern part of Eastern Indian Ocean. The study was conducted in March – April, 2010 on the basis of catches of commercial tuna longline vessel based in Port of Benoa. A total of 53 individual fishes were collected, consisting of bigeye tuna (*Thunnus obesus*), yellowfin tuna (*Thunnus albacores*), and skipjack tuna (*Katsuwonus pelamis*). Stomach specimens were collected and analyzed. Analysis was conducted on the basis of index of preponderance method. The diet of the three tuna species showed fishes as the main diet (56–82%), followed by cephalopods (squids) as the complementary diet (0–8%), and crustaceans (shrimps) as the additional diet (2–4%). Fish prey composed of 6 families i.e. Alepisauridae, Bramidae, Carangidae, Clupeidae, Engraulidae, and Scombridae.*

**Keywords:** *Stomach content, bigeye tuna, yellowfin tuna, skipjack tuna, Eastern Indian Ocean.*

### INTRODUCTION

Fish diets are frequently characterised by great diversity of prey species, which can be related to opportunistic predation (i.e. non-selective) (Menard *et al.*, 2006). Information related to food and feeding habit is important in understanding the life history of the species concerned, including growth, migration, and potentially useful for fisheries management (Effendie, 2002). The predator-prey interactions play an important part in the structure and the dynamics of multispecies communities (Notmoorn *et al.*, 2008). Considering the fast increase of tuna catches and related species in the Indian Ocean, especially in the southern Indian Ocean, it becomes necessary to assess the impact of such kind of fisheries in the pelagic ecosystems. Research activities of such kind leading to a better knowledge of tropic ecology of apex predators is important nowadays in the context of ecosystem in fisheries management.

Stomach content analyses are commonly used to study both fish feeding behavior and trophic conditions (Bertrand, 2002). However, the interpretation of such data depends on fish foraging behavior for a given environment and how representative the stomach contents are to the prey

distribution. The feeding habit of tuna in Indian Ocean has been described around Sri Lanka (Maldeniya, 1996; Dissanayake *et al.*, 2008), Indian Waters (John, 1998; Nootmorn *et al.*, 2008, Sivadas & Anasukoya, 1999; Rohit *et al.*, 2010), Andaman Sea (Panjarat, 2006), Western Indian Ocean (Seychelles Islands) (Potier *et al.*, 2004), Western Equatorial Indian Ocean (Potier *et al.*, 2007) whereas hardly any study has been conducted in Eastern Indian Ocean.

The purpose of this study is to analyse the stomach content of three tuna species (bigeye tuna, yellowfin tuna, and skipjack tuna) collected from fishing vessels operating in Eastern Indian Ocean.

### MATERIALS AND METHODS

The study was done during March – April, 2010 on board two commercial tuna longline vessels based in Port of Benoa. A total of 272 large pelagic fishes caught, 53 of them were sampled for observation, consisting of 32 bigeye tuna (*Thunnus obesus*) with size of 50 – 165 cm (FLT/Fork Length Tape), 10 yellowfin tuna (*Thunnus albacores*) of 43 – 165 cm (FLT), and 11 skipjack tuna (*Katsuwonus pelamis*) of 30 – 96 cm (FLT). Samples were taken during fishing operation in Eastern Indian Ocean as shown in Fig. 1.




Figure 1. Station distribution of samples collection (+ :bigeye tuna; “ : yellowfin tuna; ◇: skipjack tuna) in Eastern Indian Ocean

**Stomach Analysis Procedures**

Stomach analysis were conducted following this procedure: 1) the entire stomach was removed from the freshly caught fish when hauled on board; 2) size of the predator (fork length) and sex were recorded for each fish; 3) then a three-step activities were conducted: (a) total weight of the stomach contents was measured; (b) the stomach content was sorted by large categories (fish, mollusks, crustaceans); and (c) the weight of each category was recorded and; 4) classification was made to the lowest possible taxon using keys and descriptions of Nakabo (2002), FAO (1998) and Gloerfelt-Tarp and Kailola (1984).

**Data Analysis**

Index of Preponderance was used in order to analyze the main diet of an organism, developed by Natarjan & Jingran, (1962) after Mardlijah, (2008) with equation:

$$IP = \frac{(vi \times oi)}{\sum (vi \times oi)} \times 100\% \dots\dots\dots(1)$$

Where:

- IP : *Index of preponderance* for specific type of diet
- vi : Percentage of weight of one particular type of diet (%)
- oi : Frequency of Occurrence (FO) (%)

The value of *oi* or FO was obtained through the following equation (Mardlijah, 2008):

$$FO = \frac{A}{B} \times 100\% \dots\dots\dots(2)$$

Where:

- FO : Frequency of Occurrence (%)
- A : Degree of occurrence to one particular type of diet in organism
- B : Total number of organism with non-empty stomach

The calculation of IP was modified by replacing *vi* or percentage of coverage of particular type of diet with percentage of weight, so the result expected to be more accurate. Based on index of preponderance the result can be classified into three categories (Nikolsky, 1963):

- IP > 40% : Categorised as main diet.
- 4% ≥ IP ≥ 40% : Categorised as complementary diet.
- IP < 4% : Categorised as additional diet.

**RESULTS**

**Bigeye Tuna (*Thunnus obesus*)**

Of the total number of stomachs examined, 6 (11.5%) were empty. Out of 26 remaining stomachs (88.46%), 31.10% prey were digested and the rest (68.9%) as shown in Table 1. On a mass basis, undigested prey were recorded consisting of mackerel scad (family Carangidae) (29.0%), lancetfish (family Alepisauridae) (10.0%), sardines (family Clupeidae) (9.9%), sickle pomfret (family Bramidae) (3.3%), anchovy (family Engraulidae) (1.5%), unidentified fishes (2.6%), followed by squids (8.7%) and shrimps

(3.9%)(Fig. 2). Based on group of prey, fishes were likely favorable for main diet (48.31%), while cephalopods as the complementary diet (4.18%) and crustacean as the additional diet (2.57%).

Table 1. Number of bigeye, yellowfin, and skipjack stomachs sample with percentage of empty and non-empty stomachs.

Species	Number of sample	Length (FLT)	Empty stomach		Non-empty stomach	
			n	%	n	%
Bigeye tuna	52	50 - 165	6	11.5	46	88.5
Yellowfin tuna	16	43 - 165	-	-	16	100.0
Skipjack tuna	9	30 - 96	-	-	9	100.0




Figure 2. Diet composition of bigeye tuna (*Thunnus obesus*).

**Yellowfin tuna (*Thunnus albacares*)**

Disregarding the digested prey (15.5%), the diet proportion of yellowfin tunas showed a domination of mackerel scad (53.9%), followed by lancetfish (7.9%), sardines (7.5%), mackerel (3.8%), anchovy (1.0%), sickle pomfret (0.2%), and finally followed by shrimps (2.1%) and squids (8.1%). (Fig. 3). Fishes placed as main diet for yellowfin tuna (67.7%), while cephalopods (1.5%) and crustacean (0.3%) as additional diet.




Figure 3. Diet composition of yellowfin tuna (*Thunnus albacares*).

**Skipjack Tuna (*Katsuwonus pelamis*)**

Only 9 skipjack tunas were observed. Less varied composition which probably due to small number of samples. Unidentified fishes (39.74%) dominated the diet composition, followed by mackerel scad (27.95%), sardines (7.42%), lancetfish (6.99%), and shrimps (3.93%), while the rest was filled by group of digested prey (Fig. 4). Fishes were likely as main diet (71.17%) and crustacean as additional diet (0.97%).




Figure 4. Diet composition of skipjack tuna (*Katsuwonus pelamis*).

## DISCUSSION

Disregarding the proportion of digested food, the diet proportion of the three tuna species were dominated by group of fishes category as the main diet (56–82%), cephalopods (squids) as the complementary diet (0–8%), and crustaceans (shrimps) as the additional diet (2–4%). Similar result reported by Allain (2005a) in Western and Central Pacific for albacore, bluefin, yellowfin and skipjack tuna where the important prey groups found in the stomachs (measured by weight) were fish (64–88%), mollusks (6–25%) and crustaceans (0.2–9%). While in the Bay of Bengal (Notmoorn *et al.*, 2008) group of cephalopods were dominant in the diet composition for frigate tuna, skipjack tuna, bigeye, yellowfin and swordfish (60.7%), followed by group of fishes (38.8%). Observing stomach content on structure-associated aggregation conducted by Grubbs *et al.* (2001) found that bigeye tuna preyed on vertically migrating fishes rather than crustaceans on FAD-associated environment, while on seamount-associated environment, crustacean were likely became the main diet rather than teleost fishes.

Basically tunas are opportunistic predators (Menard *et al.*, 2006) feeding on a great variety of suitably sized forage fishes, crustaceans and squids (Collete & Nauen, 1983; Rohit *et al.*, 2010, Vaske & Castello, 1998; Blunt, 1960; Nakamura, 1965). In the western tropical Indian Ocean, crustaceans were almost the exclusive food source of surface-swimming bigeye tuna, in the meantime bigeye tuna fed predominantly on cephalopods and mesopelagic fish, for which this predator appeared to be the most active chaser (Potier, 2004).

Among three tuna species in this study, the yellowfin tuna showed to have the most diverse prey (Fig 2,3, & 4 for comparison). Percentage occurrence of major prey items from the previous research in Indian Ocean presented: a) **Arabian Sea (Southern)** – The prey items identified in 42 specimens in the order of preference are teleost fish (42.9%), squids (88.8%), crab (14.3%) and cuttle fish (4%); b) **Bay of Bengal** - The dominant prey items identified in 58 specimens are squids (39%), teleost fish (26.8%), crab (22%), shrimps (12.2%); c) **Andaman and Nicobar waters** – The important food items in 368 specimens identified are squids 45.1%), teleost fish (33.5%), crabs (17.8%), Octopus (2.1%), Cuttle fish (1.2%) and Stomatopods (0.3%) and; d) **Arabian Sea (Northern)** - The gut content studies of 850 specimens from Arabian Sea (Northern), method indicates preponderance of squids (52.8%) and fin-fishes (40.7%). The other components observed were Cuttle fish (3.1%), Crabs (2.4%) and

Octopus (1%)<sup>1</sup>. With few exceptions, previous stomach content studies concluded that yellowfin tuna are opportunistic predators (Collete & Nauen, 1983; Menard *et al.*, 2006) that feed on a tremendously diverse forage base, although the majority of the diet often comprises only a few families of epipelagic teleosts and crustaceans (Graham *et al.*, 2006). Crustaceans also became the main diet for yellowfin tuna, which is higher than fishes or cephalopods according to percentage index of relative importance for each food (IRI) as reported by Dissanayake *et al.* (2008) but by weight, fish prey was the most important prey with mackerel and lancetfish were likely became its main, ignoring sardines as bait. The diversity in food consumption in different sectors is indicative of the non-selective feeding nature of the species whereas the difference in the percentage composition of food items could be inferred as the availability of particular prey species rather than selection of preferred food items (Somvanshi, 2002).

Skipjack tuna landed in Bitung, North Sulawesi had Mackerel scad (57%) as the main prey item, followed by sardines and mackerel (Mardlijah, 2008). In Lakshadweep, Pakistan, the percentage composition by volume of the stomach contents excluding live baits show that fishes formed 70%, crustaceans 11 % and cephalopod 19% with *Decapterus* sp as one of the main prey (Sivadas & Anasukoya, 1999). While in Canary Islands Chub mackerel was the main prey, either as live bait or natural food (Ramos *et al.*, 1995) because of the abundance of this species in the area. Similar result appeared in this study, with lancetfish, anchovy, and sickle pomfret also found. The presence of lancetfish in all three tuna species' stomachs was interesting because it became bycatch in almost every longline fisheries in Indonesia (Nugraha & Wagiyono, 2006; Barata & Prisantoso, 2009; Prisantoso *et al.*, 2010; Nugraha & Triharyuni, 2009; Nugraha & Nurdin, 2006). This happened because lancetfish plays an important role on pelagic food chain i.e. as predator on micronekton organisms (Romanov *et al.*, 2008) alongside tunas (Bertrand *et al.*, 2001) and as prey for billfishes and tunas (Potier *et al.*, 2007). The multiplicity of prey found in this or previous studies indicate that perhaps skipjack or tunas in general are non-selective feeders and that stomach contents are probably determined by prey availability (Ramos *et al.*, 1995).

Diet studies provide information on basic biology and behaviour of the fish but they are also an important part of the parameterization of ecosystem models (Allain, 2005b); and information such as prey diversity,

size of the prey, composition of diet can be used as ecosystem indicators in conjunction with other indicators to detect changes in the ecosystem (Kirby et al., 2005).

## CONCLUSION

The diet proportion of three tuna species were dominated by group of fishes as the main diet (56–82%), followed by cephalopods (squids) as the complementary diet (0–8%), and crustaceans (shrimps) as the additional diet (2–4%). Fish prey composed of 6 families i.e. Alepisauridae, Bramidae, Carangidae, Clupeidae, Engraulidae, and Scombridae. Tunas in general are non-selective feeders and that stomach contents indicate by prey availability.

## ACKNOWLEDGEMENTS

This paper is part of authors' contribution in research on study of genetic diversity and reproductive biology of large pelagic (bigeye, *Thunnus obesus*, scombrids family) in the Indian Ocean conducted during 2010 under Research Institute for Marine Fisheries. Authors would also like to thank to all of scientific observers for their contribution in collecting onboard data throughout the years.

## REFERENCES

- Allain, V. 2005a. Diet of four tuna species of the Western and Central Pacific Ocean. *SPC Fisheries Newsletter* #114 – July/September 2005. 20 p.
- Allain V. 2005b. Ecopath model of the pelagic ecosystem of the WCPO. *WCPFC-SC1, EB-WP10*. Noumea, New Caledonia, 9–18 August 2005. p. 30-33.
- Barata, A & B.I. Prisantoso. 2009. Beberapa jenis ikan bawal (*Angel fish*, Bramidae) yang tertangkap dengan rawai tuna (*tuna long line*) di Samudera Hindia dan aspek penangkapannya. *BAWAL: 2* (5): 223-227.
- Bertrand, A., Josse, E., Bach, P & F. Gerlotto. 2001. Tuna catchability with a longline related to trophic and tuna habitat characteristics. *CM 2001/Q:03ICES Annual Science Conference Oslo*, 26-29 September, 2001. 11 p.
- Bertrand, A., Bard, F.-X. & E. Josse. 2002. Tuna food habits related to the micronekton distribution in French Polynesia. *Marine Biology* 140. p. 1023-1037.
- Blunt, C. E. 1960. Observations on the food habits of longline caught bigeye and yellowfin tuna from the tropical eastern Pacific. *California Fish and Game* 1960. 46. p. 69-80.
- Collette, B. B. & C. E. Nauen. FAO species 1983, catalogue. Vol. 2. Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. *FAO Fish. Synop.* 125 (2): 137 p.
- Dissanayake, D.C.T., Samaraweera, E.K.V. & C. Amarasiri. 2008. Fishery and feeding habits of yellowfin tuna (*Thunnus albacares*) targeted by coastal tuna longlining in the north western and north eastern coasts of Sri Lanka. *Sri Lanka J. Aquat. sci.* 13. p. 1-21.
- Effendie, M. I. 2002. *Biologi perikanan*. Yayasan Pustaka Nusatama. Yogyakarta. 157 p.
- Graham, B.S., Grubbs, D., Holland, K. & B.N. Popp. 2006. A rapid ontogenetic shift in the diet of juvenile yellowfin tuna from Hawaii. *Mar Biol* 150. p. 647-658.
- Grubbs, R.D., Holland, K.N. & D.G. Itano. 2001. Food habits and trophic dynamics of structure-associated aggregations of yellowfin and bigeye tuna (*Thunnus albacares* and *Thunnus obesus*) in the Hawaiian Islands: Project description, rationale and preliminary results. *Yellowfin Research Group – SCTB 14*. Noumea, New Caledonia, 9 – 16th August 2001. 5 p.
- John, M.E. 1998. A synoptic review of the biological studies on yellowfin tuna (*Thunnus albacares*) in the Indian Seas. *7<sup>th</sup> Expert Consultation on Indian Ocean Tunas, Victoria, Seychelles*, 9-14 November, 1998. p. 211-215.
- Kirby, D., Allain, V & B. Molony. 2005. Potential ecosystem indicators for the WCPO. *WCPFCSC1, EB-WP5*. Noumea, New Caledonia, 9–18 August 2005. 10 p.
- Mardijah, S. 2008. Analisis isi lambung ikan cakalang (*Katsuwonus pelamis*) dan ikan madidihang (*Thunnus albacares*) yang didaratkan di Bitung, Sulawesi Utara. *J. Lit. Perikan. Ind.* 14. p. 227 – 235.
- Maldeniya, R. 1996. Food consumption of yellowfin tuna, *Thunnus albacares*, in Sri Lankan waters. *Environmental Biology of Fishes* 47. p. 101 – 107.

- Ménard, F., Labrune, C., Shin, Y.J., Ah-Soy Asine, A.S. & F.X. Bard. 2006. Opportunistic predation in tuna: a size-based approach. *Mar. Ecol. Proc. Ser.* 323.p. 223-231.
- Nakamura, E. L. 1965. Food and feeding habits of skipjack tuna (*Katsuwonuspelamis*) from the Marquesas and Tuamotu Islands. *Transactions of the American Fisheries Society*. Vol. 94, Issue 3, p. 236 – 242.
- Nikolsky, G. V. 1963. *The ecology of fishes*. Academic Press. New York. xv+352 p.
- Nootmorn, P., Sumontha, M., Keereerut, P., Jayasinghe, R.P.P.K., Jagannath, N. & M. K. Sinha. 2008. *Stomach content of the three large pelagic fishes in Bay of Bengal*. IOTC-2008-WPEB-11. p. 1-13.
- Nugraha, B & E. Nurdin. 2006. Penangkapan tuna dengan menggunakan kapal riset M.V. SEAFDEC di perairan Samudera Hindia. *BAWAL: 1*(3). p.95 – 105.
- Nugraha, B & K. Wagiyono. 2006. Hasil tangkapan sampingan (*by-catch*) tuna *long line* di perairan Laut Banda. *BAWAL: 1* (2). p. 71 -75.
- Nugraha, B & S. Triharyuni. 2009. Pengaruh suhu dan kedalaman mata pancing rawai tuna (*tuna long line*) terhadap hasil tangkapan tuna di Samudera Hindia. *J. Lit. Perikan.Ind.* 15 (3).p. 239 – 247.
- Panjarat, S. 2006. Preliminary study on the stomach content of yellowfin tuna in the Andaman Sea. *In Preliminary results on the large pelagic fisheries resources survey in the Andaman Sea*. SEAFDEC TD/RES/99. p. 114-122.
- Potier, M., Marsac, F., Lucas, V., Sabatié, R., Hallier, J-P. & F. Ménard. 2004. Feeding Partitioning among Tuna Taken in Surface and Mid-water Layers: The Case of Yellowfin (*Thunnusalbacares*) and Bigeye (*T. obesus*) in the Western Tropical Indian Ocean. *Western Indian Ocean J. Mar. Sci.*3(1).p. 51–62.
- Potier, M., Marsac, F., Cherel, Y., Lucas, V., Sabatié, R., Maury, O & F. Ménard. 2007. Forage fauna in the diet of three large pelagic fishes (lancetfish, swordfish and yellowfin tuna) in the western equatorial Indian Ocean. *Fisheries Research* 83. p. 60–72.
- Prisantoso, B.I., Widodo, A.A., Mahiswara. & L. Sadiyah. 2010. Beberapa jenis hasil tangkap sampingan (*by-catch*) kapal rawai tuna di Samudera Hindia yang berbasis di Cilacap. *J. Lit. Perikan.Ind.* 16 (3).2010.p. 185 – 194.
- Ramos, A.G., Lorenzo, J.M. & J.G. Pajuelo. 1995. Food habits of bait-caught skipjack tuna *Katuwonuspelamis* off the Canary Islands. *SCI. MAR.*, 59(3-4).p. 365 – 369.
- Rohit, P., Rao, G. S. & K. Rammohan. 2010. Feeding strategies and diet composition of yellowfin tuna *Thunnusalbacares* (Bonnaterre, 1788) caught along Andhra Pradesh, east coast of India. *Indian J. Fish.*, 57(4).p. 13-19.
- Romanov, E.V., Ménard, F., Zamorov, V.V & M. Potier. 2008. Variability in conspecific predation among longnoselancetfish *Alepisaurusferox* in the western Indian Ocean. *Fisheries Science*; 74. p. 62–68.
- Sivadas, M. & A. Anasukoya. 1999. Observations on the food and feeding habits of the skipjack tuna, *Katsuwonuspelamis* (Linnaeus) from Minicoy, Lakshadweep. *The fourth Indian Fisheries Forum Proceedings*, 24-28 November, 1996. Konchi p. 5-7.
- Somvanshi, V.S. 2002. Review of biological aspects of yellowfin tuna (*Thunnusalbacares*) from the Indian Ocean. *IOTC Proceedings* no. 5. p. 420-426.
- Vaske Jr, T & J. P. Castello. 1998. Stomach contents of yellowfin tuna, *Thunnusalbacares*, during winter and spring in Southern Brazil. *Rev. Bras. Biol.* [online]. 58 (4), p. 639-647.